

THE HORMONES

PHYSIOLOGY, CHEMISTRY AND APPLICATIONS

VOLUME III

THE HORMONES

Physiology, Chemistry and Applications

Edited by

GREGORY PINCUS

*Worcester Foundation for Experimental
Biology, Shrewsbury, Massachusetts*

KENNETH V. THIMANN

*Harvard University
Cambridge, Massachusetts*

VOLUME III



1955

ACADEMIC PRESS INC. PUBLISHERS
NEW YORK

Copyright 1955 by
ACADEMIC PRESS INC
125 FIFTH 23RD STREET
NEW YORK 10 N Y

All Rights Reserved

NO PART OF THIS BOOK MAY BE REPRODUCED IN ANY FORM PHOTOSTAT
MICROFILM OR ANY OTHER MEANS WITHOUT WRITTEN PERMISSION
FROM THE PUBLISHERS

Library of Congress Catalog Card Number 48-0129

Contributors to Volume III

- T B ASTWOOD *New England Center Hospital, Boston Massachusetts*
 BEN BLOOM *National Institute of Arthritis and Metabolic Diseases National Institutes of Health Bethesda, Maryland*
 A T COWIE *National Institute for Research in Dairying Shinfield Reading England*
 RALPH I DOREMAN *Worcester Foundation for Experimental Biology, Shrewsbury Massachusetts*
 S J FOLLEY *National Institute for Research in Dairying Shinfield, Reading, England*
 ROY O GRIFF *Department of Anatomy Harvard School of Dental Medicine Boston, Massachusetts*
 EDWIN E HAYS *Research Department The Armour Laboratories Chicago Illinois*
 H HIRSCHMAN *Lakeside Hospital Cleveland, Ohio*
 ALEXANDER D KENY *Department of Pharmacology Harvard School of Dental Medicine Boston Massachusetts*
 B KETTERER *Department of Pharmacology Welsh National School of Medicine Cardiff, Wales*
 F W LANDGREBE *Department of Pharmacology Welsh National School of Medicine Cardiff Wales*
 A CARL LEOPOLD *Department of Horticulture, Agricultural Experiment Station Purdue University Lafayette Indiana*
 R L NOBLE *Department of Physiology University of Western Ontario, London Ontario Canada*
 H E PASCHKIS *Department of Physiology Jefferson Medical College of Philadelphia Philadelphia Pennsylvania*
 GREGORY PINCUS *Worcester Foundation for Experimental Biology Shrewsbury Massachusetts*
 A E RAKOFF *Department of Physiology Jefferson Medical College of Philadelphia Philadelphia Pennsylvania*
 JOSEPH E RALL *Memorial Center for Cancer and Allied Diseases, New York N Y*
 RULON W RAWSON *Memorial Center for Cancer and Allied Diseases New York N Y*
 BERTA SCHARER *Department of Anatomy Albert Einstein College of Medicine New York N Y*

- MARTIN SONENBERG, *Memorial Center for Cancer and Allied Diseases, New York, N. Y.*
- SANFORD I. STIFILMAN, *Research Department, The Armour Laboratories, Chicago, Illinois*
- DEWITT STETTIN, JR., *National Institute of Arthritis and Metabolic Diseases, National Institutes of Health, Bethesda, Maryland*
- KENNETH V. THIMANN, *The Biological Laboratories, Harvard University, Cambridge, Massachusetts*
- H. WARING, *Department of Zoology, University of Western Australia, Nedlands, Western Australia*
- JOHN H. WELSH, *The Biological Laboratories, Harvard University, Cambridge, Massachusetts*

Preface to Volume III

Seven years have passed since the appearance of Volume I of *The Hormones*. At that time it was suggested that the subject might have completed its first unfolding—virtually all the major hormones especially in the mammals and man having been discovered bioassayed and at least partially identified chemically. The subsequent period it was felt would be one first of consolidation, then of increasing emphasis upon physiological problems and the inner mechanism of the action of hormones. This prognostication, it now appears was true only in part. It was true for the hormones of the thyroid anterior pituitary and gonads which have always occupied so central a position in research and application. To a lesser extent it was true in other areas. But as regards the hormones of the invertebrates and insects and in the realms of the posterior pituitary and the nervous systems the appearance of new facts and ideas has opened new horizons. The concept of nerve as secreting and transmitting hormones as well as the long awaited isolation of the pure posterior pituitary hormones themselves are both notable. Even in the plants the discovery that auxin plays a role in flowering brings what was thought to be a growth hormone into relationship with the reproduction process. Not less to be regarded as new departures are the discoveries concerning the pituitary-adrenal axis and the widespread medical applications just touched on in Volume II which have followed.

The expected emphasis upon mechanism of action on the contrary has not yet developed to the degree expected and the major discoveries in this direction are still probably for the future although there have been interesting and suggestive developments. In the case of the animal hormones particularly intensive investigations of their metabolic fate have been considered necessary preludes to studies of their specific roles in biochemical reactions. In several instances it has been indicated that a metabolite of the secretory product rather than the secreted substance itself may be responsible for the presumed specific effects. Furthermore with the broadening of our knowledge of hormone interaction the need for distinction between primary and secondary effects of a given hormone has become more evident. For example the adrenocorticomimetic effects of estrogens are cited in the text the effects are presumably exerted through stimulating the secretion of ACTH from the anterior pituitary. Another complication

which has emerged with clarity is that mammals differ more widely in their physiology than was formerly thought. For this reason, any attempt to study mechanisms of action as problems in general physiology must recognize species differences in response to hormones, and the past septennium has unearthed additional remarkable hormonal responses occurring in one species and completely absent in another.

For all these reasons the present volume is not merely a supplement to Volumes I and II—that is, it is not merely a chronicle of recent experiments, extending in detail what has already been laid down in outline. Instead, some parts of the volume supplant their predecessors and certainly the majority of the chapters at least modify or recast the picture which had been presented. While part of its information is certainly supplementary, an important part must be regarded as revision or perhaps as reassessment.

The partition of subject matter between the authors and the planned content of the individual chapters has been somewhat revised from that of the previous volumes to allow of more unification and changed emphasis. Some of the authors are those who already reviewed their subjects in Volumes I and II, but the regrouping of the subject matter—as well as death, retirement, or preoccupation with other affairs—has necessitated a number of changes. In any event the chapters reflect individual viewpoints as much as ever. It is hoped that the elements both of variety and of uniformity will combine to make the book a useful tool in the difficult task of integrating modern biology.

G. Pincus

K. V. Thimann

June 1950

Contents

CONTRIBUTORS TO VOLUME III	v
PREFACE	vii
CONTENTS OF VOLUME I	xiii
CONTENTS OF VOLUME II	xiv
I Plant Growth Hormones By KENNETH V. THIMMANN AND A. CARL JOFOLD	1
I Introduction	1
II Assay Methods	2
III Natural Auxins	4
IV Synthetic Auxins	7
A The Transport of Auxin	16
VI The Role of Auxin in Physiological Inhibitions	20
VII The Formation and Destruction of Auxin	27
VIII Hormones and Reproduction	29
IX Auxins and Pathological Growth	35
A Mode of Action of Auxin	37
References	48
II Hormones in Invertebrates By BERTA SCHARRER	57
I Introduction	57
II Worms	58
III Crustaceans	60
IV Insects	67
A Additional Arthropods	77
VI Mollusks	78
VII Tunicates	79
VIII General Considerations	79
References	83
III Neurohormones By JOHN H. WEIGH	97
I Introduction	97
II The Diversity of Neurohormones and Their Distribution	99
III Production, Transport, Storage and Release of Neurohormones	105
IV Recent Advances in Understanding the Modes of Action of Neurohormones	118
V Summary	142
References	144
IV Physiology and Chemistry of the Parathyroids By ROY O. GREIF AND ALEXANDER D. KENNY	153
V The Hormones of the Islets of Langerhans By DEWITT STETTEN, JR. AND BEN BLOOM	175
I Insulin	175
II Glucagon	190
References	195
VI Chemistry of the Anterior Pituitary Hormones By EDWIN E. HAYE AND SANFORD L. STEELMAN	201
I Introduction	201
II Growth Hormone (GH) (STH) (PGH)	202

III	Thyrotropic Hormone (TSH)	206
IV	Lxophthalmos Producing Substance (LPS)	210
V	Prolactin (Luteotropin)	211
VI	Follicle Stimulating Hormone (FSH)	211
VII	Luteinizing Hormone (LSH)	219
VIII	The Chemistry of the Corticotropins (or the Adrenocorticotropins)	219
IX	Summary	230
	Addendum	230
	References	230
VII	Growth Hormone and Corticotropin By L. B. ASTWOOD	235
I	Introduction	235
II	Growth Hormone	236
III	Corticotropin	275
	References	296
VIII	Physiology of the Gonadotropins and the Lactogenic Hormone By A. T. COWIE AND S. J. GOLLEY	309
I	Introductory	309
II	The Gonadotropic Hormones	310
III	The Anterior Pituitary Lactogenic Hormone (Prolactin)	315
	References	379
IX	Hormones of the Posterior Pituitary By F. W. LANDGREBE, B. KETTERER AND H. WÄRING	383
I	Introduction	389
II	Neural Lobe	390
III	Intermediate Lobe	417
	References	425
X	The Chemistry and Physiology of the Thyroid By RILEY W. RAWSON, JOSEPH L. RALL AND MARTIN SONENBERG	433
I	Morphology of the Thyroid	434
II	Prethyroid Iodine Metabolism	439
III	Intrathyroidal Iodine Metabolism	441
IV	Postthyroid Iodine Metabolism	449
V	Agents Affecting Growth and Function of the Thyroid	455
VI	Actions of Thyroid Hormone	474
VII	Physiologic Aspects of Thyroid Tumors	498
	References	499
XI	Chemistry of Steroid Hormones By H. HIRSCHMANN	521
I	Nomenclature - Abbreviations	522
II	Stereochemistry	523
III	Total Syntheses	536
IV	Partial Syntheses Structures and Chemical Properties	539
V	Introduction of Isotopic Labels by Chemical Means	574
	References	578
XII	Steroid Hormone Metabolism By RALPH I. DORFMAN	589
I	Introduction	589
II	The Biosynthesis of the Steroid Nucleus	591
III	Estrogens	593
IV	Progesterone	601
V	Androgens	610
VI	Adrenocortical Hormones	622
VII	Mammalian Enzymes Influencing Steroid Reactions	644

VIII Generalizations	649
References	653
XIII The Physiology of Ovarian and Testis Hormones By GREGORY LINCOLN	66
I Introduction	663
II The Assay of Ovarian and Testis Hormones	666
III A Synopsis of the Physiological Effects of Ovarian Hormones	675
IV A Synopsis of the Physiological Effects of Testis Hormones	672
V Gonad Hormones in Embryogeny	674
VI Sexagens and Enzyme Systems	675
References	677
XIV Physiology of the Adrenal Cortex By R. I. NOBLE	685
I Introduction	687
II Morphology and Control of the Adrenal Cortex	690
III Adrenocortical Insufficiency	693
IV Adrenal Steroid Assay Procedures	701
V Adrenal Cortex and Kidney Function	704
VI Influence of the Adrenal Cortex on Sweat and Saliva	712
VII Influence of the Adrenal Cortex on the Cardiovascular System	713
VIII Influence of the Adrenal Cortex on the Nervous System	717
IX Adrenal Cortex and Muscle Function	720
X Adrenal Cortex and Carbohydrate Metabolism	721
XI Adrenal Cortex and Protein Metabolism	727
XII Adrenal Cortex and Fat Metabolism	732
XIII Adrenal Cortex and Vitamins	736
XIV Influence of the Adrenal Cortex on Lymphoid Tissue Reticuloendothelial and Mast Cells	742
XV Influence of the Adrenal Cortex on Blood and Bone Marrow	744
XVI Influence of the Adrenal Cortex on Malignancy	750
XVII Influence of the Adrenal Cortex on Bone	753
XVIII Influence of the Adrenal Cortex on Skin and Hair	754
XIX Influence of the Adrenal Cortex on Appetite and the Stomach	755
XX Adrenal Cortex and Resistance to Damage	756
XXI Adrenal Steroids and the Cellular Reaction to Inflammation and Injury	760
XXII Adrenal Steroids and Sensitivity Antigen Antibody Reactions	770
XXIII Adrenal Steroids and Susceptibility to Infection	776
XXIV Adrenal Thyroid Relationship	780
XXV Adrenal Gonad Relationship Lactation	783
References	784
XV Clinical Endocrinology By R. F. PASCHKIS AND A. F. RAKOFF	821
I Introduction	822
II Anterior Pituitary Gland	822
III Thyroid Gland	827
IV The Adrenal Gland	840
V Ovary	852
VI Placenta	861
VII Testes	864
VIII Puberty	868
References	870
AUTHOR INDEX	877
SUBJECT INDEX	960

Contents of Volume I

- I Historical Introduction By GREGORY PINCUS AND KENNETH V THIMANN
- II Plant Growth Hormones By KENNETH V THIMANN
- III Other Plant Hormones By KENNETH V THIMANN
- IV Hormones in Insects By BERTA SCHARFER
- V Hormones in Crustaceans By FRANK A BROWN JR
- VI Hormones of the Gastrointestinal Tract By HARRY GREENGARD
- VII The Physiology and Chemistry of the Parathyroid Hormone By ROY O GREEP
- VIII The Internal Secretion of the Pancreas By H JENSEN
- IX Assay of Ovarian Hormones By GREGORY PINCUS
 - X The Chemistry and Metabolism of the Estrogens By WILLIAM H PEARLMAN
 - XI Chemistry and Metabolism of Progesterone By WILLIAM H PEARLMAN
 - XII Biochemistry of Androgens By RALPH I DORFMAN
- XIII Chemistry and Metabolism of the Adrenal Cortical Hormones By R D H HEARD
- XIV Chemistry of Anterior Pituitary Hormones By CHOH HAO LI AND HERBERT M EVANS
- XV Hormonal Control of Mammary Growth By S J FOLLEY AND F H MALPRESS
- XVI Hormonal Control of Lactation By S J FOLLEY AND F H MALPRESS
- AUTHOR INDEX
- SUBJECT INDEX

Contents of Volume II

- I The Physiology of Ovarian Hormones By GREGORY LINCOLN
 - II Physiology of Androgens By RALPH I. DORFMAN
 - III Physiology of the Adrenal Cortex By R. J. NORLE
 - IV The Chemistry and Physiology of the Thyroid Hormone By WILLIAM T. SALTER
 - V The Control of Thyroid Activity By WILLIAM T. SALTER
 - VI Physiology of the Gonadotrophins By HERBERT M. IVANS AND MIRIAM F. SIMPSON
 - VII The Hypophysis and Diabetes Mellitus By IESUIF I. BENNETT AND HERBERT M. IVANS
 - VIII Hormones of the Posterior Pituitary By H. WARING AND F. W. LANDCRENE
 - IX Chemical Control of Nervous Activity
 - A Acetylcholine By DAVID NACHMANSOHN
 - B Adrenaline and Sympathin By H. BLASCHKO
 - C Neurohormones in Lower Vertebrates By GEORGE HOWARD PARKER
 - X Clinical Endocrinology By HARRY FREEMAN
- AUTHOR INDEX
SUBJECT INDEX

CHAPTER I

Plant Growth Hormones

By KENNETH A. THIMANN AND A. CARL HOFOLD

CONTENTS

	<i>Page</i>
I Introduction	1
II Assay Methods	2
III Natural Auxins	4
IV Synthetic Auxins	5
A Relation between Structure and Activity	8
B Phenomena of Antagonism and Synergism	14
C The Transport of Auxin	16
VI The Role of Auxin in Physiological Inhibitions	20
A Apical Dominance and the Inhibition of Buds	20
B Inhibition of Root Growth	23
C Agents Which Accentuate Auxin Action	24
D Auxin Inhibitors	25
VII The Formation and Destruction of Auxin	26
VIII Hormones and Reproduction	29
A The Humoral Stimulus Controlling Flowering	29
B Auxins and Flowering	30
C Auxins and Vernalization	32
D Hormones in Fruit Set and Fruit Development	33
IX Auxins and Pathological Growth	35
X Mode of Action of Auxin	37
A The Combination of Auxin with Its Substrate	3
B Metabolic Functions of Auxin	38
C Water Uptake	42
References	48

I Introduction

The progress of any field of science is highly irregular some parts may make a major break through whereas other parts—apparently no more complex—seem blocked by some difficulty of procedure or of concepts so that ideas and experiments proceed in spirals without advancing. The study of plant hormones has shown the expected irregularity of advance during the six years since the appearance of Volume I of *The Hormones*. Some aspects like that of the inhibition of the growth of buds have not undergone any really fundamental change whereas the role of auxin in flowering and the mechanism of auxin action in growth have developed with great rapidity. Recent evidence for an *in vitro* action of auxin and

But C is also equal to AD

Therefore each of the straight lines AE , C is equal to AD ,

so that AE is also equal to C

[CN 1]

Therefore given the two straight lines AB , C , from AB the greater AE has been cut off equal to C the less

(Being) what it was required to do

PROPOSITION 4

If two triangles have the two sides equal to two sides respectively and have the angles contained by the equal straight lines equal, they will also have the base equal to the base, the triangle will be equal to the triangle, and the remaining angles will be equal to the remaining angles respectively, namely those which the equal sides subtend

Let ABC , DEF be two triangles having the two sides AB , AC equal to the two sides DE , DF respectively, namely AB to DE and AC to DF , and the angle BAC equal to the angle EDF

I say that the base BC is also equal to the base EF the triangle ABC will be equal to the triangle DEF , and the remaining angles will be equal to the remaining angles respectively, namely those which the equal sides subtend that is, the angle ABC to the angle DEF and the angle ACB to the angle DFE

For, if the triangle ABC be applied to the triangle DEF

and if the point A be placed on the point D

and the straight line AB on DE

then the point B will also coincide with E because AB is equal to DE

Again, AB coinciding with DE

the straight line AC will also coincide with DF because the angle BAC is equal to the angle EDF

hence the point C will also coincide with the point F , because AC is again equal to DF

But B also coincided with E

hence the base BC will coincide with the base EF

[For if when B coincides with E and C with F the base BC does not coincide with the base EF two straight lines will enclose a space which is impossible

Therefore the base BC will coincide with EF] and will be equal to it [CN 4]
Thus the whole triangle ABC will coincide with the whole triangle DEF , and will be equal to it

And the remaining angles will also coincide with the remaining angles and will be equal to them

the angle ABC to the angle DEF
and the angle ACB to the angle DFE

Therefore etc

(Being) what it was required to prove


PROPOSITION 5

In isosceles triangles the angles at the base are equal to one another and if the equal straight lines be produced further the angles under the base will be equal to one another

Let ABC be an isosceles triangle having the side AB equal to the side AC ,

and let the straight lines BD , CE be produced further in a straight line with AB , AC [Post 2]

I say that the angle ABC is equal to the angle ACB and the angle CBD to the angle BCE



Let a point F be taken at random on BD , from AE the greater let AG be cut off equal to AF the less [I 3]

and let the straight lines FC , GB be joined [Post 1]

Then since AF is equal to AG and AB to AC , the two sides FA , AC are equal to the two sides GA , AB respectively,

and they contain a common angle, the angle FAG

Therefore the base FC is equal to the base GB

and the triangle AFC is equal to the triangle AGB

and the remaining angles will be equal to the remaining angles respectively namely those which the equal sides subtend

that is the angle ACF to the angle ABG

and the angle AFC to the angle AGB [I 4]

And, since the whole AF is equal to the whole AG

and in these AB is equal to AC

the remainder BF is equal to the remainder CG

But FC was also proved equal to GB

therefore the two sides BF , FC are equal to the two sides CG , GB respectively, and the angle BFC is equal to the angle CGB

while the base BC is common to them

therefore the triangle BFC is also equal to the triangle CGB and the remaining angles will be equal to the remaining angles respectively, namely those which the equal sides subtend

therefore the angle FBC is equal to the angle GCB ,

and the angle BCF to the angle CBG

Accordingly since the whole angle ABG was proved equal to the angle ACF

and in these the angle CBG is equal to the angle BCF

the remaining angle ABC is equal to the remaining angle ACB

and they are at the base of the triangle ABC

But the angle FBC was also proved equal to the angle GCB

and they are under the base

Therefore etc

Q E D

PROPOSITION 6

If in a triangle two angles be equal to one another the sides which subtend the equal angles will also be equal to one another

Let ABC be a triangle having the angle ABC equal to the angle ACB

I say that the side AB is also equal to the side AC

For if AB is unequal to AC one of them is greater

Let AB be greater and from AB the greater let DB be cut off equal to AC the less

let DC be joined

Then since DB is equal to AC

and BC is common,



the two sides DB BC are equal to the two sides AC , CB respectively,
 and the angle DBC is equal to the angle ACB ,
 therefore the base DC is equal to the base AB ,
 and the triangle DBC will be equal to the triangle ACB
 the less to the greater
 which is absurd

Therefore AB is not unequal to AC ,
 it is therefore equal to it


Therefore etc

Q E D

PROPOSITION 7

Given two straight lines constructed on a straight line (from its extremities) and meeting in a point, there cannot be constructed on the same straight line (from its extremities) and on the same side of it, two other straight lines meeting in another point and equal to the former two respectively, namely each to that which has the same extremity with it

For, if possible given two straight lines AC , CB constructed on the straight line AB and meeting at the point C , let two other straight lines AD DB be constructed on the same straight line AB , on the same side of it meeting in another point D and equal to the former two respectively namely each to that which has the same extremity with it so that CA is equal to DA which has the same extremity A with it and CB to DB which has the same extremity B with it, and let CD be joined



Then since AC is equal to AD ,
 the angle ACD is also equal to the angle ADC , [I 5]
 therefore the angle ADC is greater than the angle DCB
 therefore the angle CDB is much greater than the angle DCB

Again since CB is equal to DB
 the angle CDB is also equal to the angle DCB

But it was also proved much greater than it
 which is impossible

Therefore etc

Q E D

PROPOSITION 8

If two triangles have the two sides equal to two sides respectively and have also the base equal to the base they will also have the angles equal which are contained by the equal straight lines

Let ABC , DEF be two triangles having the two sides AB , AC equal to the two sides DE DF respectively namely AB to DE , and AC to DF and let them have the base BC equal to the base EF

I say that the angle BAC is also equal to the angle EDF



For if the triangle ABC be applied to the triangle DEF and if the point B be placed on the point E and the straight line BC on EF ,
 the point C will also coincide with F ,

because BC is equal to EF

Then BC coinciding with EF ,

BA AC will also coincide with ED DF ,

for if the base BC coincides with the base EF and the sides BA , AC do not coincide with ED DF but fall beside them as EG GF

then given two straight lines constructed on a straight line (from its extremities) and meeting in a point there will have been constructed on the same straight line (from its extremities) and on the same side of it two other straight lines meeting in another point and equal to the former two respectively namely each to that which has the same extremity with it

But they cannot be so constructed

[1 7]

Therefore it is not possible that if the base BC be applied to the base EF , the sides BA , AC should not coincide with ED DF

they will therefore coincide,

so that the angle BAC will also coincide with the angle EDF and will be equal to it

If therefore etc

Q E D

PROPOSITION 9

To bisect a given rectilineal angle



Let the angle BAC be the given rectilineal angle

Thus it is required to bisect it

Let a point D be taken at random on AB

let AE be cut off from AC equal to AD

[1 3]

let DE be joined and on DE let the equilateral triangle DEF be constructed

let AF be joined

I say that the angle BAC has been bisected by the straight line AF

For, since AD is equal to AE

and AF is common

the two sides DA AF are equal to the two sides EA AF respectively

And the base DF is equal to the base EF

therefore the angle DAF is equal to the angle EAF

[1 8]

Therefore the given rectilineal angle BAC has been bisected by the straight line AF

Q E F

PROPOSITION 10

To bisect a given finite straight line

Let AB be the given finite straight line

Thus it is required to bisect the finite straight line AB

Let the equilateral triangle ABC be constructed on it

[1 1]

and let the angle ACB be bisected by the straight line CD

[1 9]


I say that the straight line AB has been bisected at the point D

For since AC is equal to CB

and CD is common

the two sides AC CD are equal to the two sides BC , CD respectively

and the angle ACD is equal to the angle BCD



therefore the base AD is equal to the base BD [I 4]

Therefore the given finite straight line AB has been bisected at D Q E F

PROPOSITION 11

To draw a straight line at right angles to a given straight line from a given point on it

Let AB be the given straight line and C the given point on it

Thus it is required to draw from the point C a straight line at right angles to the straight line AB

Let a point D be taken at random on AC ,

let CE be made equal to CD , [I 3]

on DE let the equilateral triangle FDE be constructed, [I 1]

and let FC be joined

I say that the straight line FC has been drawn at right angles to the given straight line AB from C the given point on it

For, since DC is equal to CE ,

and CF is common,

the two sides DC CF are equal to the two sides EC CF respectively,

and the base DF is equal to the base FE ,


therefore the angle DCF is equal to the angle ECF [I 8]

and they are adjacent angles

But when a straight line set up on a straight line makes the adjacent angles equal to one another, each of the equal angles is right [Def 10]

therefore each of the angles DCF , FCE is right

Therefore the straight line CF has been drawn at right angles to the given straight line AB from the given point C on it Q E F



PROPOSITION 12

To a given infinite straight line, from a given point which is not on it to draw a perpendicular straight line

Let AB be the given infinite straight line and C the given point which is not on it

thus it is required to draw to the given infinite straight line AB , from the given point C which is not on it, a perpendicular straight line


For let a point D be taken at random on the other side of the straight line AB and with centre C and distance CD let the circle EFG be described [Post 3]
let the straight line EG be bisected at H [I 10]

and let the straight lines CG , CH CE be joined [Post 1]

I say that CH has been drawn perpendicular to the given infinite straight line AB from the given point C which is not on it

For since GH is equal to HE

and HC is common



the two sides GH , HC are equal to the two sides EH , HC respectively
and the base CG is equal to the base CE

therefore the angle CHG is equal to the angle EHC [I 8]

And they are adjacent angles

But when a straight line set up on a straight line makes the adjacent angles equal to one another each of the equal angles is right and the straight line standing on the other is called a perpendicular to that on which it stands

[Def 10]

Therefore CH has been drawn perpendicular to the given infinite straight line AB from the given point C which is not on it

Q E F

PROPOSITION 13

If a straight line set up on a straight line make angles it will make either two right angles or angles equal to two right angles

For let any straight line AB set up on the straight line CD make the angles CBA , ABD

I say that the angles CBA , ABD are either two right angles or equal to two right angles

Now if the angle CBA is equal to the angle ABD they are two right angles [Def 10]

But if not let BE be drawn from the point B at right angles to CD [I 11]

therefore the angles CBE , EBD are two right angles

Then since the angle CBE is equal to the two angles CBA , ABE ,
let the angle EBD be added to each
therefore the angles CBE , EBD are equal to the three angles CBA , ABE , EBD [C A 2]

Again since the angle DBA is equal to the two angles DBE , EBA
let the angle ABC be added to each
therefore the angles DBA , ABC are equal to the three angles DBE , EBA , ABC [C N 2]

But the angles CBE , EBD were also proved equal to the same three angles
and things which are equal to the same thing are also equal to one another [C V 1]

therefore the angles CBE , EBD are also equal to the angles DBA , ABC

But the angles CBE , EBD are two right angles

therefore the angles DBA , ABC are also equal to two right angles

Therefore etc

Q E D

PROPOSITION 14

If with any straight line and at a point on it two straight lines not lying on the same side make the adjacent angles equal to two right angles the two straight lines will be in a straight line with one another

For with any straight line AB and at the point B on it let the two straight lines BC , BD not lying on the same side make the adjacent angles ABC , ABD equal to two right angles

I say that BD is in a straight line with CB

For if BD is not in a straight line with BC , let BE be in a straight line with CB

Then since the straight line AB stands on the straight line CBE ,
the angles ABC , ABE are equal to two right angles [I 13]

But the angles ABC , ABD are also equal to two right angles

therefore the angles CBA , ABE are equal to the angles CBA , ABD [Post 4 and C N 1]

Let the angle CBA be subtracted from each,
therefore the remaining angle ABE is equal to the remaining angle ABD [C.N 3]

the less to the greater which is impossible

Therefore BE is not in a straight line with CB

Similarly we can prove that neither is any other straight line except BD

Therefore CB is in a straight line with BD

Therefore etc

Q E D

PROPOSITION 15

If two straight lines cut one another, they make the vertical angles equal to one another

For let the straight lines AB , CD cut one another at the point E ,

I say that the angle AEC is equal to the angle DEB ,
and the angle CEB to the angle AED

For since the straight line AE stands on the straight line CD , making the angles CEA , AED ,

the angles CEA , AED are equal to two right angles [I 13]

Again since the straight line DE stands on the straight line AB , making the angles AED , DEB

the angles AED , DEB are equal to two right angles [I 13]

But the angles CEA , AED were also proved equal to two right angles,
therefore the angles CEA , AED are equal to the angles AED , DEB

[Post 4 and C N 1]

Let the angle AED be subtracted from each

therefore the remaining angle CEA is equal to the remaining angle DEB [C N 3]

Similarly it can be proved that the angles CEB , DEA are also equal

Therefore etc

Q E D

[PORISM From this it is manifest that if two straight lines cut one another they will make the angles at the point of section equal to four right angles]

PROPOSITION 16

In any triangle if one of the sides be produced the exterior angle is greater than either of the interior and opposite angles



Let ABC be a triangle and let one side of it BC be produced to D ,

I say that the exterior angle ACD is greater than either of the interior and opposite angles CBA , BAC


Let AC be bisected at E [I 10] and let BE be joined and produced in a straight line to F

let EF be made equal to BE [I 3]

let FC be joined [Post 1] and let AC be drawn through to G [Post 2]



Then since AE is equal to EC and BE to EF ,
 the two sides AE EB are equal to the two sides CE EF respectively
 and the angle AEB is equal to the angle FEC
 for they are vertical angles [1 15]



Therefore the base AB is equal to the base FC
 and the triangle ABE is equal to the triangle CFE
 and the remaining angles are equal to the remain-
 ing angles respectively namely, those which the
 equal sides subtend [1 4]

therefore the angle BAE is equal to the angle ECF
 But the angle ECD is greater than the angle
 ECF [C 5]

therefore the angle ACD is greater than the angle BAE

Similarly also if BC be bisected the angle BCG that is the angle ACD
 [1 15] can be proved greater than the angle ABC as well

Therefore etc


Q E D

PROPOSITION 17

In any triangle two angles taken together in any manner are less than two right angles

Let ABC be a triangle

I say that two angles of the triangle ABC taken together in any manner are
 less than two right angles



For let BC be produced to D [Post 2]

Then since the angle ACD is an exterior
 angle of the triangle ABC

it is greater than the interior and opposite
 angle ABC [1 16]

Let the angle ACB be added to each

therefore the angles ACD ACB are greater than the angles ABC , BCA

But the angles ACD ACB are equal to two right angles [1 13]

Therefore the angles ABC BCA are less than two right angles

Similarly we can prove that the angles BAC ACB are also less than two
 right angles and so are the angles CAB ABC as well

Therefore etc


Q E D

PROPOSITION 18

In any triangle the greater side subtends the greater angle

For let ABC be a triangle having the side AC greater than AB

I say that the angle ABC is also greater than the angle BCA



For since AC is greater than AB let AD
 be made equal to AB [1 3] and let BD be
 joined

Then since the angle ADB is an exterior
 angle of the triangle BCD

it is greater than the interior and opposite
 angle DCB [1 16]

But the angle ADB is equal to the angle ABD

since the side AB is equal to AD

therefore the angle ABD is also greater than the angle ACB
 therefore the angle ABC is much greater than the angle ACB

Therefore etc

Q E D

PROPOSITION 19

In any triangle the greater angle is subtended by the greater side

Let ABC be a triangle having the angle ABC greater than the angle BCA ,
 I say that the side AC is also greater than the side AB

For, if not, AC is either equal to AB or less

Now AC is not equal to AB

for then the angle ABC would also have been equal to the angle ACB

[1 5]

but it is not,

therefore AC is not equal to AB

Neither is AC less than AB ,

for then the angle ABC would also have been less than the angle ACB

[1 18]

but it is not

therefore AC is not less than AB

And it was proved that it is not equal either

Therefore AC is greater than AB

Therefore etc

Q E D



PROPOSITION 20

In any triangle two sides taken together in any manner are greater than the remaining one

For let ABC be a triangle

I say that in the triangle ABC two sides taken together in any manner are greater than the remaining one namely

BA AC greater than BC

AB , BC greater than AC ,

BC CA greater than AB

For let BA be drawn through to the point D , let DA be made equal to CA ,
 and let DC be joined

Then since DA is equal to AC

the angle ADC is also equal to the angle ACD

[1 5]

therefore the angle BCD is greater than the angle ADC

[C A 5]

And since DCB is a triangle having the angle BCD greater
 than the angle BDC

and the greater angle is subtended by the greater side [1 19]

therefore DB is greater than BC


But DA is equal to AC

therefore BA AC are greater than BC

Similarly we can prove that AB BC are also greater than CA and BC CA
 than AB

Therefore etc

Q E D



PROPOSITION 21

If on one of the sides of a triangle from its extremities there be constructed two

straight lines meeting within the triangle, the straight lines so constructed will be less than the remaining two sides of the triangle, but will contain a greater angle

On BC , one of the sides of the triangle ABC from its extremities B C , let the two straight lines BD DC be constructed meeting within the triangle

I say that BD DC are less than the remaining two sides of the triangle BA , AC , but contain an angle BDC greater than the angle BAC

For let BD be drawn through to E

Then, since in any triangle two sides are greater than the remaining one [I 20]

therefore in the triangle ABE , the two sides AB , AE are greater than BE

Let EC be added to each

therefore BA AC are greater than BE EC

Again since in the triangle CED

the two sides CE ED are greater than CD

let DB be added to each

therefore CE , EB are greater than CD DB

But BA AC were proved greater than BE EC

therefore BA AC are much greater than BD DC

Again since in any triangle the exterior angle is greater than the interior and opposite angle [I 16]

therefore in the triangle CDE ,

the exterior angle BDC is greater than the angle CED

For the same reason moreover in the triangle ABE also

the exterior angle CEB is greater than the angle BAC

But the angle BDC was proved greater than the angle CEB

therefore the angle BDC is much greater than the angle BAC


Therefore etc

Q E D

PROPOSITION 22

Out of three straight lines which are equal to three given straight lines to construct a triangle thus it is necessary that two of the straight lines taken together in any manner should be greater than the remaining one [I 20]

Let the three given straight lines be A B C and of these let two taken together in any manner be greater than the remaining one



namely

A B greater than C

and

A C greater than B

B C greater than A

thus it is required to construct a triangle out of straight lines equal to A, B, C
 Let there be set out a straight line DE , terminated at D but of infinite length
 in the direction of E

and let DF be made equal to A FG equal to B , and GH equal to C [1 3]

With centre F and distance FD let the circle DKL be described

again with centre G and distance GH let the circle LKH be described

and let KF, KG be joined

I say that the triangle KFG has been constructed out of three straight lines
 equal to A, B, C

For since the point F is the centre of the circle DKL

FD is equal to FK

But FD is equal to A

therefore KF is also equal to A

Again since the point G is the centre of the circle LKH ,

GH is equal to GK

But GH is equal to C

therefore KG is also equal to C

And FG is also equal to B ,

therefore the three straight lines KF, FG, GK are equal to the three straight
 lines A, B, C

Therefore out of the three straight lines KF, FG, GK which are equal to the
 three given straight lines A, B, C , the triangle KFG has been constructed


Q E D

PROPOSITION 23

On a given straight line and at a point on it to construct a rectilineal angle equal to
 a given rectilineal angle

Let AB be the given straight line A the point on it and the angle DCE the
 given rectilineal angle

thus it is required to construct on the given straight line AB and at the
 point A on it, a rectilineal angle equal to the given rectilineal angle DCE



On the straight lines CD, CE respectively let the points D, E be taken at
 random,

let DE be joined

and out of three straight lines which are equal to the three straight lines CD ,
 DE, CE let the triangle AFG be constructed in such a way that CD is equal to
 AF, CE to AG and further DE to FG [1 22]

Then, since the two sides DC, CE are equal to the two sides FA, AG respec-
 tively,

and the base DE is equal to the base FG

the angle DCF is equal to the angle FAG

[1 8]

Therefore on the given straight line AB and at the point A on it the rectilinear angle FAG has been constructed equal to the given rectilinear angle DCE



Q E F

PROPOSITION 24

If two triangles have the two sides equal to two sides respectively but have the one of the angles contained by the equal straight lines greater than the other they will also have the base greater than the base

Let ABC , DEF be two triangles having the two sides AB , AC equal to the two sides DE , DF respectively namely AB to DE , and AC to DF , and let the angle at A be greater than the angle at D

I say that the base BC is also greater than the base EF



For since the angle BAC is greater than the angle EDF let there be constructed on the straight line DE , and at the point D on it, the angle EDG equal to the angle BAC , [I 23] let DG be made equal to either of the two straight lines AC , DF , and let EG FG be joined

Then since AB is equal to DE and AC to DG

the two sides BA AC are equal to the two sides ED DG , respectively

and the angle BAC is equal to the angle EDG

therefore the base BC is equal to the base EG [I 4]

Again since DF is equal to DG

the angle DGF is also equal to the angle DFG , [I 5]

therefore the angle DFG is greater than the angle EGF

Therefore the angle EFG is much greater than the angle EGF

And since EFG is a triangle having the angle EFG greater than the angle EGF

and the greater angle is subtended by the greater side [I 19]

the side LG is also greater than EF

But EG is equal to BC

Therefore BC is also greater than EF

Therefore etc

Q E D

PROPOSITION 25

If two triangles have the two sides equal to two sides respectively but have the base greater than the base they will also have the one of the angles contained by the equal straight lines greater than the other

Let ABC DEF be two triangles having the two sides AB AC equal to the two sides DE DF respectively namely AB to DE and AC to DF and let the base BC be greater than the base EF

I say that the angle BAC is also greater than the angle EDF


For if not it is either equal to it or less

Now the angle BAC is not equal to the angle EDF for then the base BC would also have been equal to the base EF [I 4]

but it is not,

therefore the angle BAC is not equal to the angle EDF

Neither again is the angle BAC less than the angle EDF for then the base BC would also have been less than the base EF , [1 24]



but it is not therefore the angle BAC is not less than the angle EDF
But it was proved that it is not equal either,

therefore the angle BAC is greater than the angle EDF

Therefore etc


Q E D

PROPOSITION 26

If two triangles have the two angles equal to two angles respectively and one side equal to one side namely either the side adjoining the equal angles or that subtending one of the equal angles they will also have the remaining sides equal to the remaining sides and the remaining angle to the remaining angle

Let ABC DEF be two triangles having the two angles ABC BCA equal to the two angles DEF EFD respectively namely the angle ABC to the angle DEF and the angle BCA to the angle EFD , and let them also have one side equal to one side first that adjoining the equal angles namely BC to EF

I say that they will also have the remaining sides equal to the remaining sides respectively namely AB to DE and AC to DF and the remaining angle to the remaining angle namely the angle BAC to the angle EDF



For if AB is unequal to DE one of them is greater

Let AB be greater and let BG be made equal to DE and let GC be joined
Then since BG is equal to DE and BC to EF

the two sides GB BC are equal to the two sides DE EF respectively,

and the angle GBC is equal to the angle DEF

therefore the base GC is equal to the base DF

and the triangle GBC is equal to the triangle DEF

and the remaining angles will be equal to the remaining angles namely those which the equal sides subtend [1 4]

therefore the angle GCB is equal to the angle DFE

But the angle DFE is by hypothesis equal to the angle BCA

therefore the angle BCG is equal to the angle BCA

the less to the greater which is impossible

Therefore AB is not unequal to DE

and is therefore equal to it

But BC is also equal to EF

therefore the two sides AB , BC are equal to the two sides DE , EF respectively,

and the angle ABC is equal to the angle DEF ,

therefore the base AC is equal to the base DF ,

and the remaining angle BAC is equal to the remaining angle EDF [I 4]

Again, let sides subtending equal angles be equal as AB to DE ,

I say again that the remaining sides will be equal to the remaining sides namely AC to DF and BC to EF , and further the remaining angle BAC is equal to the remaining angle EDF

For, if BC is unequal to EF , one of them is greater

Let BC be greater if possible and let BH be made equal to EF , let AH be joined

Then since BH is equal to EF , and AB to DE

the two sides AB , BH are equal to the two sides DE , EF respectively, and they contain equal angles

therefore the base AH is equal to the base DF

and the triangle ABH is equal to the triangle DEF

and the remaining angles will be equal to the remaining angles namely those which the equal sides subtend [I 4]

therefore the angle BHA is equal to the angle EFD

But the angle EFD is equal to the angle BCA

therefore, in the triangle AHC the exterior angle BHA is equal to the interior and opposite angle BCA

which is impossible

[I 16]

Therefore BC is not unequal to EF

and is therefore equal to it

But AB is also equal to DE

therefore the two sides AB , BC are equal to the two sides DE , EF respectively, and they contain equal angles,

therefore the base AC is equal to the base DF

the triangle ABC equal to the triangle DEF

and the remaining angle BAC equal to the remaining angle EDF [I 4]

Therefore etc

Q E D

PROPOSITION 27

If a straight line falling on two straight lines make the alternate angles equal to one another the straight lines will be parallel to one another

For let the straight line EF falling on the two straight lines AB , CD make the alternate angles AEF , EFD equal to one another

I say that AB is parallel to CD

For if not AB , CD when produced will meet either in the direction of B , D or towards A , C

Let them be produced and meet, in the direction of B , D at G

Then in the triangle GEF


the exterior angle AEF is equal to the interior and opposite angle EFG

which is impossible

[I 16]

Therefore AB , CD when produced will not meet in the direction of B , D

Similarly it can be proved that neither will they meet towards A , C



But straight lines which do not meet in either direction are parallel
[Def 23]

therefore AB is parallel to CD

Therefore etc

Q E D

PROPOSITION 28

If a straight line falling on two straight lines make the exterior angle equal to the interior and opposite angle on the same side or the interior angles on the same side equal to two right angles the straight lines will be parallel to one another

For let the straight line EF falling on the two straight lines AB CD make the exterior angle EGB equal to the interior and opposite angle CHD or the interior angles on the same side namely BGH , GHD , equal to two right angles, I say that AB is parallel to CD

For, since the angle EGB is equal to the angle GHD

while the angle EGB is equal to the angle AGH

[I 15]

the angle AGH is also equal to the angle GHD

and they are alternate

therefore AB is parallel to CD [I 27]

Again since the angles BGH GHD are equal to two right angles and the angles AGH , BGH are also equal to two right angles [I 13]

the angles AGH BGH are equal to the angles BGH GHD

Let the angle BGH be subtracted from each

therefore the remaining angle AGH is equal to the remaining angle GHD ,


and they are alternate

therefore AB is parallel to CD

[I 27]

Therefore etc

Q E D



PROPOSITION 29

A straight line falling on parallel straight lines makes the alternate angles equal to one another the exterior angle equal to the interior and opposite angle, and the interior angles on the same side equal to two right angles

For let the straight line EF fall on the parallel straight lines AB CD

I say that it makes the alternate angles AGH GHD equal, the exterior angle EGB equal to the interior and opposite angle GHD and the interior angles on the same side, namely BGH GHD equal to two right angles

For if the angle AGH is unequal to the angle GHD one of them is greater

Let the angle AGH be greater

Let the angle BGH be added to each


therefore the angles AGH BGH are greater than the angles BGH , GHD

But the angles AGH , BGH are equal to two right angles, [I 13]

therefore the angles BGH GHD are less than two right angles

But straight lines produced indefinitely from angles less than two right angles meet [Post 5]

therefore AB CD if produced indefinitely will meet,



but they do not meet, because they are by hypothesis parallel
Therefore the angle AGH is not unequal to the angle GHD ,
and is therefore equal to it

Again, the angle AGH is equal to the angle EGB , [I 15]

therefore the angle EGB is also equal to the angle GHD [C.N. 1]

Let the angle BGH be added to each,

therefore the angles EGB, BGH are equal to the angles BGH, GHD [C.N. 2]

But the angles EGB, BGH are equal to two right angles [I 13]

therefore the angles BGH, GHD are also equal to two right angles


Therefore etc

Q.E.D.

PROPOSITION 30

Straight lines parallel to the same straight line are also parallel to one another

Let each of the straight lines AB, CD be parallel to EF . I say that AB is also parallel to CD .

 For let the straight line GA fall upon them

Then, since the straight line GA has fallen on the parallel straight lines AB, EF ,

the angle AGK is equal to the angle GHE [I 27]

Again, since the straight line GA has fallen on the parallel straight lines EF, CD

the angle GHE is equal to the angle GKD [I 29]

But the angle AGK was also proved equal to the angle GHE

therefore the angle AGK is also equal to the angle GKD [C.N. 1]

and they are alternate


Therefore AB is parallel to CD

Q.E.D.

PROPOSITION 31

Through a given point to draw a straight line parallel to a given straight line

Let A be the given point and BC the given straight line
thus it is required to draw through the point A a straight line parallel to the straight line BC

 Let a point D be taken at random on BC
and let AD be joined, on the straight line DA ,
and at the point A on it let the angle DAE be
constructed equal to the angle ADC [I 23] and
let the straight line AF be produced in a straight
line with EA

Then, since the straight line AD falling on the two straight lines BC, EF has made the alternate angles EAD, ADC equal to one another

therefore DAF is parallel to BC [I 27]

Therefore through the given point A the straight line EAF has been drawn parallel to the given straight line BC

Q.E.D.

PROPOSITION 32

In any triangle, if one of the sides be produced the exterior angle is equal to the two interior and opposite angles and the three interior angles of the triangle are equal to two right angles

Let ABC be a triangle, and let one side of it BC be produced to D

I say that the exterior angle ACD is equal to the two interior and opposite angles $C \hat{B}A$, ABC , and the three interior angles of the triangle ABC , BCA , CAB are equal to two right angles

For let CE be drawn through the point C parallel to the straight line AB [I 31]

Then since AB is parallel to CE ,
and AC has fallen upon them
the alternate angles BAC , ACE are equal to one another [I 29]

Again, since AB is parallel to CE
and the straight line BD has fallen upon them
the exterior angle ECD is equal to the interior and opposite angle ABC [I 29]

But the angle ACE was also proved equal to the angle $B \hat{C}A$,
therefore the whole angle ACD is equal to the two interior and opposite angles BAC ABC

Let the angle ACB be added to each
therefore the angles ACD ACB are equal to the three angles ABC , BCA , CAB

But the angles ACD ACB are equal to two right angles [I 13]
therefore the angles ABC , BCA , CAB are also equal to two right angles

Therefore etc

Q E D

PROPOSITION 33

The straight lines joining equal and parallel straight lines (at the extremities which are) in the same directions (respectively) are themselves also equal and parallel

Let AB CD be equal and parallel and let the straight lines AC , BD join them (at the extremities which are) in the same directions (respectively),

I say that AC BD are also equal and parallel

Let BC be joined

Then since AB is parallel to CD and BC has fallen upon them
the alternate angles ABC , BCD are equal to one another [I 29]

And since AB is equal to CD

and BC is common

the two sides AB BC are equal to the two sides DC CB

and the angle ABC is equal to the angle BCD

therefore the base AC is equal to the base BD

and the triangle ABC is equal to the triangle DCB ,

and the remaining angles will be equal to the remaining angles respectively
namely those which the equal sides subtend [I 4]

therefore the angle ACB is equal to the angle CBD

And since the straight line BC falling on the two straight lines AC , BD has made the alternate angles equal to one another



AC is parallel to BD

[I 27]

And it was also proved equal to it

Therefore etc

Q E D




PROPOSITION 34

In parallelogrammic areas the opposite sides and angles are equal to one another, and the diameter bisects the areas

Let $ACDB$ be a parallelogrammic area, and BC its diameter

I say that the opposite sides and angles of the parallelogram $ACDB$ are equal to one another and the diameter BC bisects it



For since AB is parallel to CD , and the straight line BC has fallen upon them the alternate angles ABC BCD are equal to one another [I 29]

Again, since AC is parallel to BD and BC has fallen upon them,

the alternate angles ACB CBD are equal to one another [I 29]

Therefore ABC DCB are two triangles having the two angles ABC , BCA equal to the two angles DCB , CBD respectively, and one side equal to one side namely that adjoining the equal angles and common to both of them, BC , therefore they will also have the remaining sides equal to the remaining sides respectively, and the remaining angle to the remaining angle [I 26]

therefore the side AB is equal to CD ,

and AC to BD ,

and further the angle BAC is equal to the angle CDB

And, since the angle ABC is equal to the angle BCD

and the angle CBD to the angle ACB

the whole angle ABD is equal to the whole angle ACD [C.N 2]

And the angle BAC was also proved equal to the angle CDB

Therefore in parallelogrammic areas the opposite sides and angles are equal to one another

I say, next, that the diameter also bisects the areas

For, since AB is equal to CD ,

and BC is common,

the two sides AB BC are equal to the two sides DC , CB respectively,

and the angle ABC is equal to the angle BCD ,

therefore the base AC is also equal to DB

and the triangle ABC is equal to the triangle DCB [I 4]


Therefore the diameter BC bisects the parallelogram $ACDB$ Q.E.D.

PROPOSITION 35

Parallelograms which are on the same base and in the same parallels are equal to one another

Let $ABCD$ $EBCF$ be parallelograms on the same base BC and in the same parallels AF BC ,

I say that $ABCD$ is equal to the parallelogram $EBCF$



For since $ABCD$ is a parallelogram

AD is equal to BC [I 34]

For the same reason also

EF is equal to BC

so that AD is also equal to EF [C.N 1]

and DE is common,

therefore the whole AF is equal to the whole DF [C. 1. 2]

But AB is also equal to DC , [1. 34]

therefore the two sides EA, AB are equal to the two sides FD, DC respectively,

and the angle FDC is equal to the angle EAB ,
the exterior to the interior, [1. 29]

therefore the base FB is equal to the base FC ,

and the triangle FAB will be equal to the triangle FDC [1. 4]

Let DGE be subtracted from each

therefore the trapezium $ABGD$ which remains is equal to the trapezium $EGCF$
which remains [C. 1. 3]

Let the triangle GBC be added to each

therefore the whole parallelogram $ABCD$ is equal to the whole parallelogram
 $FBCF$ [C. 1. 2]

Therefore etc

Q. E. D.

PROPOSITION 36

Parallelograms which are on equal bases and in the same parallels are equal to one another

Let $ABCD, EFGH$ be parallelograms which are on equal bases BC, FG and in the same parallels AH, BG

I say that the parallelogram $ABCD$ is equal to $EFGH$

For let BE, CH be joined

Then since BC is equal to FG ,
while

FG is equal to EH

BC is also equal to FH [C. 1. 1]

But they are also parallel

And EB, HC join them

but straight lines joining equal and parallel straight lines (at the extremities which are) in the same directions (respectively) are equal and parallel [1. 33]

Therefore $EBCH$ is a parallelogram

[1. 34]

And it is equal to $ABCD$

for it has the same base BC with it and is in the same parallels BC, AH with it [1. 35]

For the same reason also $EFGH$ is equal to the same $EBCH$ [1. 35]

so that the parallelogram $ABCD$ is also equal to $EFGH$ [C. 1. 1]

Therefore etc

Q. E. D.

PROPOSITION 37

Triangles which are on the same base and in the same parallels are equal to one another


Let ABC, DBC be triangles on the same base BC and in the same parallels AD, BC

I say that the triangle ABC is equal to the triangle DBC

Let AD be produced in both directions to E, F

through B let BE be drawn parallel to CA

[1. 31]



and through C let CF be drawn parallel to BD [I 31]

Then each of the figures $EBCA$, $DBCF$ is a parallelogram, and they are equal

for they are on the same base BC and in the same parallels BC EF [I 35]

Moreover the triangle ABC is half of the parallelogram $EBCA$, for the diameter AB bisects it [I 34]

And the triangle DBC is half of the parallelogram $DBCF$ for the diameter DC bisects it [I 34]

[But the halves of equal things are equal to one another]

Therefore the triangle ABC is equal to the triangle DBC

Therefore etc

Q E D

PROPOSITION 38

Triangles which are on equal bases and in the same parallels are equal to one another

Let ABC DEF be triangles on equal bases BC EF and in the same parallels BC AD ,

I say that the triangle ABC is equal to the triangle DEF

For let AD be produced in both directions to G H

through B let BG be drawn parallel to CA , [I 31]

and through F let FH be drawn parallel to DE

Then each of the figures $GBCA$, $DEFH$ is a parallelogram

and $GBCA$ is equal to $DEFH$,

for they are on equal bases BC , EF and in the same parallels BF GH [I 36]

Moreover the triangle ABC is half of the parallelogram $GBCA$ for the diameter AB bisects it [I 34]

And the triangle FED is half of the parallelogram $DEFH$ for the diameter DF bisects it [I 34]

[But the halves of equal things are equal to one another]

Therefore the triangle ABC is equal to the triangle DEF

Therefore etc

Q E D

PROPOSITION 39

Equal triangles which are on the same base and on the same side are also in the same parallels

Let ABC DBC be equal triangles which are on the same base BC and on the same side of it

[I say that they are also in the same parallels]

And [For] let AD be joined I say that AD is parallel to BC



For if not let AE be drawn through the point A parallel to the straight line BC [I 31]

and let EC be joined

Therefore the triangle ABC is equal to the triangle EBC ,

for it is on the same base BC with it and in the same parallels [I 37]

But ABC is equal to DBC



therefore DBC is also equal to FBC [C.A. 1]

the greater to the less which is impossible

Therefore AF is not parallel to BC

Similarly we can prove that neither is any other straight line except AD ,
therefore AD is parallel to BC

Therefore etc

Q E D

PROPOSITION 10

Equal triangles which are on equal bases and on the same side are also in the same parallels

Let ABC, CDE be equal triangles on equal bases BC, CE and on the same side

I say that they are also in the same parallels

I or let AD be joined

I say that AD is parallel to BF

I or if not let AF be drawn through A parallel to BF

[I 31] and let FE be joined

Therefore the triangle ABC is equal to the triangle FCE ,
[I 31]

for they are on equal bases BC, CE and in the same parallels BE, AF [I 35]

But the triangle ABC is equal to the triangle DCE ,

therefore the triangle DCF is also equal to the triangle FCE ,

[C.A. 1]

the greater to the less which is impossible

Therefore AF is not parallel to BE

Similarly we can prove that neither is any other straight line except AD ,
therefore AD is parallel to BE

Therefore etc

Q E D

PROPOSITION 41

If a parallelogram have the same base with a triangle and be in the same parallels the parallelogram is double of the triangle

For let the parallelogram $ABCD$ have the same base BC with the triangle EBC and let it be in the same parallels BC, AE

I say that the parallelogram $ABCD$ is double of the triangle EBC

For let AC be joined

Then the triangle ABC is equal to the triangle EBC ,
for it is on the same base BC with it and in the same
parallels BC, AE [I 37]

But the parallelogram $ABCD$ is double of the triangle ABC

for the diameter AC bisects it

[I 34]

so that the parallelogram $ABCD$ is also double of the triangle EBC



Therefore etc


Q E D

PROPOSITION 42

To construct in a given rectilineal angle a parallelogram equal to a given triangle

Let ABC be the given triangle and D the given rectilineal angle
thus it is required to construct in the rectilineal angle D a parallelogram equal





to the triangle ABC

Let BC be bisected at E , and let AE be joined,
on the straight line EC , and at the point
 E on it let the angle CEF be constructed
equal to the angle D , [I 23]

through A let AG be drawn parallel to EC and [I 31]
through C let CG be drawn parallel to EF

Then $FECG$ is a parallelogram

And since BE is equal to EC ,

the triangle ABE is also equal to the triangle AEC

for they are on equal bases BE, EC and in the same parallels BC, AG , [I 38]

therefore the triangle ABC is double of the triangle AEC

But the parallelogram $FECG$ is also double of the triangle AEC , for it has
the same base with it and is in the same parallels with it, [I 41]

therefore the parallelogram $FECG$ is equal to the triangle ABC

And it has the angle CEF equal to the given angle D

Therefore the parallelogram $FECG$ has been constructed equal to the given
triangle ABC in the angle CEF which is equal to D Q E F

PROPOSITION 43

In any parallelogram the complements of the parallelograms about the diameter are equal to one another

Let $ABCD$ be a parallelogram and AC its diameter

and about AC let EH, FG be parallelograms and BK, KD the so-called complements,

I say that the complement BK is equal to the complement KD

For since $ABCD$ is a parallelogram, and AC its diameter,

the triangle ABC is equal to the triangle ACD [I 34]

Again since EH is a parallelogram, and AK is
its diameter,

the triangle AEK is equal to the triangle AHK

For the same reason

the triangle KFC is also equal to KGC

Now since the triangle AEK is equal to the
triangle AHK ,

and KFC to KGC

the triangle AEK together with KGC is equal to the triangle AHK together
with KFC [C.N 2]

And the whole triangle ABC is also equal to the whole ADC ,
therefore the complement BK which remains is equal to the complement KD
which remains [C.N 3]

Therefore etc

Q E D

PROPOSITION 44

To a given straight line to apply in a given rectilineal angle a parallelogram equal to a given triangle

Let AB be the given straight line C the given triangle and D the given recti-
lineal angle,

thus it is required to apply to the given straight line AB in an angle equal to the angle D a parallelogram equal to the given triangle C

Let the parallelogram $BLFG$ be constructed equal to the triangle C in the angle FLG which is equal to D [I 42] let it be placed so that BE is in a straight line with AB let FC be drawn through to H , and let AH be drawn through A parallel to either BG or FF [I 31]

Let HB be joined

Then since the straight line HF falls upon the parallels AH IF , the angles AHF HFF are equal to two right angles [I 29]

Therefore the angles BHG GFF are less than two right angles, and straight lines produced indefinitely from angles less than two right angles meet, [Post 5]

therefore HB , FE , when produced will meet

Let them be produced and meet at K through the point K let KL be drawn parallel to either FA or FH , [I 31]

and let HA GB be produced to the points L , M

Then $HLKF$ is a parallelogram
 HA is its diameter and AG ME are parallelograms, and LB BF the so-called complements about HA

therefore LB is equal to BF [I 43]

But BF is equal to the triangle C

therefore LB is also equal to C [C.N. 1]

And, since the angle GBE is equal to the angle ABM [I 15]

while the angle GBE is equal to D

the angle ABM is also equal to the angle D



Therefore the parallelogram LB equal to the given triangle C has been applied to the given straight line AB in the angle ABM which is equal to D

Q E F

PROPOSITION 45

To construct in a given rectilinear angle a parallelogram equal to a given rectilinear figure

Let $ABCD$ be the given rectilinear figure and E the given rectilinear angle thus it is required to construct in the given angle E , a parallelogram equal to the rectilinear figure $ABCD$



Let DB be joined and let the parallelogram FH be constructed equal to the triangle ABD in the angle HKF which is equal to E , [I 42]
 let the parallelogram GM equal to the triangle DBC be applied to the straight

line GH in the angle GHM which is equal to E [I 44]

Then since the angle E is equal to each of the angles HAF GHM
the angle HAF is also equal to the angle GHM [CN 1]

Let the angle AHG be added to each,

therefore the angles FAH AHG are equal to the angles KHG , GHM

But the angles FAH , KHG are equal to two right angles, [I 29]

therefore the angles AHG GHM are also equal to two right angles

Thus with a straight line GH , and at the point H on it two straight lines KH HM not lying on the same side make the adjacent angles equal to two right angles,

therefore AH is in a straight line with HM [I 14]

And, since the straight line HG falls upon the parallels AM , FG the alternate angles MHG HGF are equal to one another [I 29]

Let the angle HGL be added to each

therefore the angles MHG HGL are equal to the angles HGF , HGL [CN 2]

But the angles MHG HGL are equal to two right angles, [I 29]

therefore the angles HGF , HGL are also equal to two right angles [CN 1]

Therefore FG is in a straight line with GL [I 14]

And since FA is equal and parallel to HG , [I 34]

and HG to ML also

AF is also equal and parallel to ML , [CN 1 I 30]

and the straight lines AM FL join them (at their extremities), therefore AM FL are also equal and parallel [I 33]

Therefore $AFLM$ is a parallelogram

And since the triangle ABD is equal to the parallelogram FH
and DBC to GM


the whole rectilineal figure $ABCD$ is equal to the whole parallelogram $AFLM$

Therefore the parallelogram $AFLM$ has been constructed equal to the given rectilineal figure $ABCD$ in the angle FAM which is equal to the given angle E

Q E F

PROPOSITION 46

On a given straight line to describe a square



Let AB be the given straight line thus it is required to describe a square on the straight line AB

Let AC be drawn at right angles to the straight line AB from the point A on it [I 11] and let AD be made equal to AB

through the point D let DE be drawn parallel to AB
and through the point B let BE be drawn parallel to AD [I 31]

Therefore $ADFB$ is a parallelogram

therefore AB is equal to DE and AD to BE [I 34]

But AB is equal to AD

therefore the four straight lines BA AD DL EB are equal to one another,
therefore the parallelogram $ADEB$ is equilateral

I say next that it is also right angled

For since the straight line AD falls upon the parallels AB DE
the angles BAD ADE are equal to two right angles

But the angle BAD is right,

therefore the angle ADE is also right

And in parallelogrammic areas the opposite sides and angles are equal to one another, [1 34]

therefore each of the opposite angles ABE , BED is also right

Therefore $ADEB$ is right-angled

And it was also proved equilateral

Therefore it is a square, and it is described on the straight line AB $Q E R$

PROPOSITION 47

In right-angled triangles the square on the side subtending the right angle is equal to the squares on the sides containing the right angle

Let ABC be a right angled triangle having the angle BAC right,

I say that the square on BC is equal to the squares on BA , AC

For let there be described on BC the square $BDEC$, and on BA , AC the squares GB , HC [1 46]

through A let AL be drawn parallel to either BD or CE and let AD , FC be joined

Then since each of the angles BAC , BAG is right it follows that with a straight line BA and at the point A on it the two straight lines AC , AG not lying on the same side make the adjacent angles equal to two right angles,

therefore CA is in a straight line with AG [1 14]

For the same reason

BA is also in a straight line with AH

And since the angle DBC is equal to the angle FBA for each is right

let the angle ABC be added to each

therefore the whole angle DBA is equal to the whole angle FBC [CN 2]

And since DB is equal to BC and FB to BA

the two sides AB , BD are equal to the two sides FB , BC respectively,

and the angle ABD is equal to the angle FBC

therefore the base AD is equal to the base FC ,

and the triangle ABD is equal to the triangle FBC [1 4]

Now the parallelogram BL is double of the triangle ABD for they have the same base BD and are in the same parallels BD , AL [1 41]

And the square GB is double of the triangle FBC

for they again have the same base FB and are in the same parallels FB , GC [1 41]

[But the doubles of equals are equal to one another]


Therefore the parallelogram BL is also equal to the square GB

Similarly if AE , BA be joined

the parallelogram CL can also be proved equal to the square HC ,

therefore the whole square $BDEC$ is equal to the two squares GB , HC [CN 2]

And the square $BDEC$ is described on BC ,



and the squares GB, HC on BA, AC

Therefore the square on the side BC is equal to the squares on the sides BA, AC

Therefore etc

Q E D

PROPOSITION 48

If in a triangle the square on one of the sides be equal to the squares on the remaining two sides of the triangle, the angle contained by the remaining two sides of the triangle is right

For in the triangle ABC let the square on one side BC be equal to the squares on the sides BA, AC ,

I say that the angle BAC is right

For let AD be drawn from the point A at right angles to the straight line BC , let AD be made equal to BA , and let DC be joined

Since DA is equal to AB ,

the square on DA is also equal to the square on AB

Let the square on AC be added to each

therefore the squares on DA, AC are equal to the squares on BA, AC

But the square on DC is equal to the squares on DA, AC , for the angle DAC is right, [1 47]

and the square on BC is equal to the squares on BA, AC , for this is the hypothesis

therefore the square on DC is equal to the square on BC

so that the side DC is also equal to BC

And since DA is equal to AB ,

and AC is common,

the two sides DA, AC are equal to the two sides BA, AC

and the base DC is equal to the base BC

therefore the angle DAC is equal to the angle BAC [1 8]

But the angle DAC is right,

therefore the angle BAC is also right

Therefore etc

Q E D



BOOK TWO

DEFINITIONS

1 Any rectangular parallelogram is said to be *contained* by the two straight lines containing the right angle

2 And in any parallelogrammic area let any one whatever of the parallelograms about its diameter with the two complements be called a *gnomon*

BOOK II PROPOSITIONS

PROPOSITION 1

If there be two straight lines and one of them be cut into any number of segments whatever the rectangle contained by the two straight lines is equal to the rectangles contained by the uncut straight line and each of the segments

Let A BC be two straight lines, and let BC be cut at random at the points D E

I say that the rectangle contained by A BC is equal to the rectangle contained by A BD that contained by A DE and that contained by A EC

For let BF be drawn from B at right angles to BC ,

[1 11]


let BG be made equal to A

[1 3]

through G let GH be drawn parallel to BC

[1 31]

and through D E C let DK EL , CH be drawn parallel to BG



Then BH is equal to BK DL EH

Now BH is the rectangle A BC for it is contained by GB BC and BG is equal to A

BK is the rectangle A BD for it is contained by GB BD and BG is equal to A

and DL is the rectangle A DE for DK that is BG is equal to A [1 34]

Similarly also LH is the rectangle A EC

Therefore the rectangle A BC is equal to the rectangle A , BD the rectangle A , DE and the rectangle A EC

Therefore etc

Q E D


PROPOSITION 2

If a straight line be cut at random the rectangle contained by the whole and both of the segments is equal to the square on the whole

For let the straight line AB be cut at random at the point C

I say that the rectangle contained by AB BC together with the rectangle contained by BA AC is equal to the square on AB

For let the square $ADEB$ be described on AB [I 46] and let CF be drawn through C parallel to either AD or BE [I 31]



Then AE is equal to AF , CE

Now AE is the square on AB ,

AF is the rectangle contained by BA , AC for it is contained by DA , AC , and AD is equal to AB ,

and CE is the rectangle AB , BC for BE is equal to AB

Therefore the rectangle BA , AC together with the rectangle AB , BC is equal to the square on AB

Therefore etc


Q E D

PROPOSITION 3

If a straight line be cut at random the rectangle contained by the whole and one of the segments is equal to the rectangle contained by the segments and the square on the aforesaid segment

For let the straight line AB be cut at random at C ,

I say that the rectangle contained by AB , BC is equal to the rectangle contained by AC , CB together with the square on BC



For let the square $CDEB$ be described on CB , [I 46]

let ED be drawn through to F ,

and through A let AF be drawn parallel to either CD or BE [I 31]

Then AE is equal to AD , CE

Now AE is the rectangle contained by AB , BC for it is contained by AB , BE and BE is equal to BC

AD is the rectangle AC , CB for DC is equal to CB , and DB is the square on CB

Therefore the rectangle contained by AB , BC is equal to the rectangle contained by AC , CB together with the square on BC

Therefore etc

Q E D

PROPOSITION 4

If a straight line be cut at random the square on the whole is equal to the squares on the segments and twice the rectangle contained by the segments

For let the straight line AB be cut at random at C ,

I say that the square on AB is equal to the squares on AC , CB and twice the rectangle contained by AC , CB

For let the square $ADEB$ be described on AB

[I 46]

let BD be joined

through C let CF be drawn parallel to either AD or EB ,

and through G let HK be drawn parallel to either AB or DE [I 31]


Then since CF is parallel to AD and BD has fallen on them

the exterior angle CGB is equal to the interior and opposite angle ADB [I 29]

But the angle ADB is equal to the angle ABD

since the side BA is also equal to AD ,

[I 5]



therefore the angle CGB is also equal to the angle GBC ,

so that the side BC is also equal to the side CG

[1 6]

But CB is equal to GA , and CG to AB ,

[1 34]

therefore GA is also equal to AB ,

therefore $CGAB$ is equilateral

I say next that it is also right-angled

For since CG is parallel to BA

the angles ABC GCB are equal to two right angles

[1 29]

But the angle ABC is right

therefore the angle BCG is also right

so that the opposite angles CGA , GAB are also right

[1 34]

Therefore $CGAB$ is right angled,

and it was also proved equilateral,

therefore it is a square

and it is described on CB

For the same reason

HF is also a square,

and it is described on HG that is AC

[1 31]

Therefore the squares HF AC are the squares on AC , CB

Now, since AG is equal to GE

and AG is the rectangle AC , CB , for GC is equal to CB ,

therefore GE is also equal to the rectangle AC , CB

Therefore AG , GE are equal to twice the rectangle AC , CB

But the squares HF , AC are also the squares on AC , CB therefore the four areas HF AC , AG GE are equal to the squares on AC , CB and twice the rectangle contained by AC , CB

But HF , AC , AG , GE are the whole $ADFB$

which is the square on AB

Therefore the square on AB is equal to the squares on AC , CB and twice the rectangle contained by AC , CB

Therefore etc

Q E D

PROPOSITION 5

If a straight line be cut into equal and unequal segments the rectangle contained by the unequal segments of the whole together with the square on the straight line between the points of section is equal to the square on the half

For let a straight line AB be cut into equal segments at C and into unequal segments at D ,

I say that the rectangle contained by AD DB together with the square on CD is equal to the square on CB

For let the square $CEFB$ be described on CB ,

[1 46]

and let BE be joined

through D let DG be drawn parallel to either CE or BF


through H again let AM be drawn parallel to either AB or EF

and again through A let AK be drawn parallel to either CL or BM

[1 31]

Then, since the complement CH is equal to the complement HT ,

[1 43]



let DM be added to each,
therefore the whole CM is equal to the whole DF
But CM is equal to AL ,

since AC is also equal to CB

therefore AL is also equal to DF

[I 36]

Let CH be added to each,

therefore the whole AH is equal to the gnomon NOP

But AH is the rectangle AD DB , for DH is equal to DB ,

therefore the gnomon NOP is also equal to the rectangle AD DB

Let LG , which is equal to the square on CD , be added to each
therefore the gnomon NOP and LG are equal to the rectangle contained by AD , DB and the square on CD

But the gnomon NOP and LG are the whole square $CEFB$ which is described on CB ,

therefore the rectangle contained by AD DB together with the square on CD is equal to the square on CB

Therefore etc

Q E D

PROPOSITION 6

If a straight line be bisected and a straight line be added to it in a straight line the rectangle contained by the whole with the added straight line and the added straight line together with the square on the half is equal to the square on the straight line made up of the half and the added straight line

For let a straight line AB be bisected at the point C , and let a straight line BD be added to it in a straight line,

I say that the rectangle contained by AD , DB together with the square on CB is equal to the square on CD

For let the square $CEFD$ be described on CD

[I 46]


and let DE be joined,

through the point B let BG be drawn parallel to either EC or DF

through the point H let AM be drawn parallel to either AB or EF

and further through A let AK be drawn parallel to either CL or DM

[I 31]



Then since AC is equal to CB ,

AL is also equal to CH

[I 36]

But CH is equal to HF

[I 43]

Therefore AI is also equal to HF

Let CM be added to each,

therefore the whole AM is equal to the gnomon NOP

But AM is the rectangle AD DB ,

for DM is equal to DB ,

therefore the gnomon NOP is also equal to the rectangle AD DB

Let LG which is equal to the square on BC be added to each,

therefore the rectangle contained by AD DB together with the square on CB is equal to the gnomon NOP and LG

But the gnomon NOP and LG are the whole square $CEFD$ which is described on CD ,

therefore the rectangle contained by AD , DB together with the square on CB is equal to the square on CD

Therefore etc

Q E D

PROPOSITION 7

If a straight line be cut at random the square on the whole and that on one of the segments both together are equal to twice the rectangle contained by the whole and the said segment and the square on the remaining segment

For let a straight line AB be cut at random at the point C ,

I say that the squares on AB , BC are equal to twice the rectangle contained by AB , BC and the square on CA

For let the square $ADEB$ be described on AB ,

[I 46]

and let the figure be drawn

Then since AG is equal to GE [I 43] let CF be added to each,

therefore the whole AF is equal to the whole CE

Therefore AF , CE are double of AG

But AF , CE are the gnomon KLM and the square CF therefore the gnomon KLM and the square CF are double of AG

But twice the rectangle AB , BC is also double of AG for BF is equal to BC ,

therefore the gnomon KLM and the square CF are equal to twice the rectangle AB , BC

Let DG which is the square on AC , be added to each, therefore the gnomon KLM and the squares BG , GD are equal to twice the rectangle contained by AB , BC and the square on AC


But the gnomon KLM and the squares BG , GD are the whole $ADEB$ and CF ,

which are squares described on AB , BC ,

therefore the squares on AB , BC are equal to twice the rectangle contained by AB , BC together with the square on AC

Therefore etc

Q E D



PROPOSITION 8

If a straight line be cut at random four times the rectangle contained by the whole and one of the segments together with the square on the remaining segment is equal to the square described on the whole and the aforesaid segment as on one straight line

For let a straight line AB be cut at random at the point C

I say that four times the rectangle contained by AB , BC together with the square on AC is equal to the square described on AB , BC as on one straight line

For let [the straight line] BD be produced in a straight line [with AB] and let BD be made equal to CB ,

let the square $AEFD$ be described on AD and let the figure be drawn

Then since CB is equal to BD while CB is equal to GK , and BD to KN , therefore GK is also equal to KN

For the same reason

QR is also equal to RP

And, since BC is equal to BD and GK to KN

therefore CH is also equal to AD and GR to RN [I 36]

But CA is equal to RN for they are complements of the parallelogram CP [I 43]

therefore AD is also equal to GR

therefore the four areas DK CK GR RN are equal to one another
Therefore the four are quadruple of CA

Again since CB is equal to BD

while BD is equal to BA that is CG

and CB is equal to GK that is GQ

therefore CG is also equal to GQ

And since CG is equal to GQ and QR to RP ,

AG is also equal to MQ and QL to RF [I 36]

But MQ is equal to QL for they are complements of the parallelogram ML [I 43]

therefore AG is also equal to RF ,

therefore the four areas AG MQ QL RF are equal to one another

Therefore the four are quadruple of AG

But the four areas CA AD GR RN were proved to be quadruple of CK
therefore the eight areas which contain the gnomon STU , are quadruple of AK

Now, since AK is the rectangle AB BD for BK is equal to BD

therefore four times the rectangle AB BD is quadruple of AK

But the gnomon STU was also proved to be quadruple of AK

therefore four times the rectangle AB , BD is equal to the gnomon STU

Let OH which is equal to the square on AC be added to each

therefore four times the rectangle AB BD together with the square on AC is equal to the gnomon STU and OH

But the gnomon STU and OH are the whole square $AEFD$

which is described on AD

therefore four times the rectangle AB BD together with the square on AC is equal to the square on AD

But BD is equal to BC

therefore four times the rectangle contained by AB BC together with the square on AC is equal to the square on AD that is to the square described on AB and BC as on one straight line

Therefore etc

Q E D

PROPOSITION 9

If a straight line be cut into equal and unequal segments the squares on the unequal segments of the whole are double of the square on the half and of the square on the straight line between the points of section

For let a straight line AB be cut into equal segments at C and into unequal segments at D

I say that the squares on AD DB are double of the squares on AC CD

For let CE be drawn from C at right angles to AB and let it be made equal to either AC or CB

let EA EB be joined

let DF be drawn through D parallel to EC

and FG through F parallel to AB

and let AF be joined

Then, since AC is equal to CF

the angle EAC is also equal to the angle AEC

And since the angle at C is right

the remaining angles EAC AEC are equal to one right angle [I 32]

And they are equal,

therefore each of the angles CEA , CAE is half a right angle

For the same reason

each of the angles CFB , EBC is also half a right angle

therefore the whole angle AEB is right

And since the angle GEF is half a right angle

and the angle EGF is right for it is equal to the interior and opposite angle ECB

the remaining angle EFG is half a right angle

therefore the angle GEF is equal to the angle EFG ,

so that the side FG is also equal to GF

[I 29]

[I 32]

[I 6]

Again since the angle at B is half a right angle

and the angle FDB is right, for it is again equal to the interior and opposite angle ECB ,

the remaining angle BFD is half a right angle

therefore the angle at B is equal to the angle DFB ,

so that the side FD is also equal to the side DB

[I 29]

[I 32]

[I 6]

Now since AC is equal to CE

the square on AC is also equal to the square on CE

therefore the squares on AC CE are double of the square on AC

But the square on EA is equal to the squares on AC CE for the angle ACE is right

[I 47]

therefore the square on EA is double of the square on AC

Again since EG is equal to GF

the square on EG is also equal to the square on GF ,

therefore the squares on EG GF are double of the square on GF

But the square on EF is equal to the squares on EG GF

therefore the square on EF is double of the square on GF

But GF is equal to CD

[I 34]

therefore the square on EF is double of the square on CD

But the square on EA is also double of the square on AC

therefore the squares on AE , EF are double of the squares on AC , CD

And the square on AF is equal to the squares on AE EF for the angle AEF is right

[I 47]

therefore the square on AF is double of the squares on AC CD

But the squares on AD , DF are equal to the square on AF for the angle at D is right,

[I 47]


therefore the squares on AD DF are double of the squares on AC CD

And DF is equal to DB

therefore the squares on AD , DB are double of the squares on AC CD

Therefore etc

Q E D




PROPOSITION 10

If a straight line be bisected and a straight line be added to it in a straight line the square on the whole with the added straight line and the square on the added straight line both together are double of the square on the half and of the square described on the straight line made up of the half and the added straight line as on one straight line

For let a straight line AB be bisected at C , and let a straight line BD be added to it in a straight line,

I say that the squares on AD , DB are double of the squares on AC , CD



For let CE be drawn from the point C at right angles to AB [I 11] and let it be made equal to either AC or CB [I 3],

let EA , EB be joined

through E let EF be drawn parallel to AD and through D let FD be drawn parallel to CE [I 31]

Then since a straight line EF falls on the parallel straight lines EC , FD

the angles CEF , EFD are equal to two right angles [I 29]

therefore the angles FEB , EFD are less than two right angles

But straight lines produced from angles less than two right angles meet [I Post 5]

therefore EB , FD if produced in the direction B , D will meet

Let them be produced and meet at G

and let AG be joined

Then since AC is equal to CE ,

the angle EAC is also equal to the angle AEC , [I 5]

and the angle at C is right

therefore each of the angles EAC , AEC is half a right angle [I 32]

For the same reason

each of the angles CEB , EBC is also half a right angle

therefore the angle AEB is right

And since the angle EBC is half a right angle

the angle DBG is also half a right angle [I 15]

But the angle BDG is also right

for it is equal to the angle DCE they being alternate, [I 29]

therefore the remaining angle DGB is half a right angle, [I 32]

therefore the angle DGB is equal to the angle DBG

so that the side BD is also equal to the side GD [I 6]

Again, since the angle EGF is half a right angle

and the angle at F is right for it is equal to the opposite angle the angle at C [I 34]

the remaining angle FEG is half a right angle [I 32]

therefore the angle EGF is equal to the angle FEG

so that the side GF is also equal to the side EF [I 6]

Now since the square on EC is equal to the square on CA

the squares on EC , CA are double of the square on CA

But the square on EA is equal to the squares on EC , CA , [I 47]

therefore the square on EA is double of the square on AC [C.N. 1]
 Again since FG is equal to EF ,

the square on FG is also equal to the square on FE ,

therefore the squares on GF, FE are double of the square on EF

But the square on EG is equal to the squares on GF, FE , [1 47]

therefore the square on EG is double of the square on EF

And EF is equal to CD [1 34]

therefore the square on EG is double of the square on CD

But the square on EA was also proved double of the square on AC ,

therefore the squares on AF, EG are double of the squares on AC, CD

And the square on AG is equal to the squares on AE, EG [1 47]

therefore the square on AG is double of the squares on AC, CD

But the squares on AD, DG are equal to the square on AG , [1 47]

therefore the squares on AD, DG are double of the squares on AC, CD

And DG is equal to DB

therefore the squares on AD, DB are double of the squares on AC, CD

Therefore etc

Q E D

PROPOSITION 11

To cut a given straight line so that the rectangle contained by the whole and one of the segments is equal to the square on the remaining segment

Let AB be the given straight line,
 thus it is required to cut AB so that the rectangle contained by the whole and one of the segments is equal to the square on the remaining segment

For let the square $ABDC$ be described on AB , [1 46]

let AC be bisected at the point E and let BF be joined

let CA be drawn through to F and let EF be made equal to BE

let the square FHI be described on AF and let GHI be drawn through to K

I say that AB has been cut at H so as to make the rectangle contained by AB, BH equal to the square on AH

For since the straight line AC has been bisected at E , and FA is added to it

the rectangle contained by CF, FA together with the square on AE is equal to the square on EF [11 6]

But EF is equal to EB

therefore the rectangle CF, FA together with the square on AE is equal to the square on EB

But the squares on BA, AE are equal to the square on EB for the angle at A is right [1 47]

therefore the rectangle CF, FA together with the square on AE is equal to the squares on BA, AE

Let the square on AE be subtracted from each


therefore the rectangle CF, FA which remains is equal to the square on AB

Now the rectangle CF, FA is FK for AF is equal to FG ,

and the square on AB is AD

therefore FK is equal to AD

Let AK be subtracted from each,



therefore FH which remains is equal to HD

And HD is the rectangle $AB \cdot BH$, for AB is equal to BD ,

and FH is the square on AH

therefore the rectangle contained by $AB \cdot BH$ is equal to the square on HA
therefore the given straight line AB has been cut at H so as to make the rectangle contained by AB, BH equal to the square on HA Q E F

PROPOSITION 12

In obtuse-angled triangles the square on the side subtending the obtuse angle is greater than the squares on the sides containing the obtuse angle by twice the rectangle contained by one of the sides about the obtuse angle namely that on which the perpendicular falls, and the straight line cut off outside by the perpendicular towards the obtuse angle

Let ABC be an obtuse-angled triangle having the angle BAC obtuse, and let BD be drawn from the point B perpendicular to CA produced,

I say that the square on BC is greater than the squares on BA, AC by twice the rectangle contained by $CA \cdot AD$

For, since the straight line CD has been cut at random at the point A ,
the square on DC is equal to the squares on $CA \cdot AD$
and twice the rectangle contained by CA, AD [I 4]

Let the square on DB be added to each
therefore the squares on CD, DB are equal to the squares on $CA \cdot AD \cdot DB$ and twice the rectangle CA, AD

But the square on CB is equal to the squares on CD, DB , for the angle at D is right, [I 47]

and the square on AB is equal to the squares on $AD \cdot DB$ [I 47]
therefore the square on CB is equal to the squares on $CA \cdot AB$ and twice the rectangle contained by $CA \cdot AD$,

so that the square on CB is greater than the squares on $CA \cdot AB$ by twice the rectangle contained by $CA \cdot AD$

Therefore etc

Q E D


PROPOSITION 13

In acute angled triangles the square on the side subtending the acute angle is less than the squares on the sides containing the acute angle by twice the rectangle contained by one of the sides about the acute angle namely that on which the perpendicular falls, and the straight line cut off within by the perpendicular towards the acute angle

Let ABC be an acute-angled triangle having the angle at B acute, and let AD be drawn from the point A perpendicular to BC ,

I say that the square on AC is less than the squares on $CB \cdot BA$ by twice the rectangle contained by $CB \cdot BD$

For since the straight line CB has been cut at random at D
the squares on $CB \cdot BD$ are equal to twice the rectangle contained by $CB \cdot BD$ and the square on DC



Let the square on DA be added to each,
therefore the squares on CB BD DA are equal to twice the rectangle contained by CB , BD and the squares on AD DC

But the square on AB is equal to the squares on BD , DA , for the angle at D is right, [1 47]

and the square on AC is equal to the squares on AD , DC ,
therefore the squares on CB , BD are equal to the square on AC and twice the rectangle CB , BD ,
so that the square on AC alone is less than the squares on CB , BA by twice the rectangle contained by CB BD

Therefore etc

Q E D

PROPOSITION 14

To construct a square equal to a given rectilinear figure

I let A be the given rectilinear figure,
thus it is required to construct a square equal to the rectilinear figure A

For let there be constructed
the rectangular parallelogram BD
equal to the rectilinear figure A

[1 45]

Then, if BE is equal to ED ,
that which was enjoined will
have been done for a square BD

has been constructed equal to the rectilinear figure A

But if not one of the straight lines BE ED is greater

Let BE be greater, and let it be produced to F ,

let EF be made equal to ED and let BF be bisected at G

With centre G and distance one of the straight lines GB GF let the semi-circle BHF be described let DE be produced to H and let GH be joined

Then since the straight line BF has been cut into equal segments at G and into unequal segments at E

the rectangle contained by BE EF together with the square on EG is equal to the square on GF [11 5]

But GF is equal to GH ,
therefore the rectangle BE , EF together with the square on GE is equal to the square on GH

But the squares on HE EG are equal to the square on GH , [1 47]
therefore the rectangle BE , EF together with the square on GE is equal to the squares on HE , EG

Let the square on GE be subtracted from each
therefore the rectangle contained by BE , EF which remains is equal to the square on EH

But the rectangle BE EF is BD for EF is equal to ED


therefore the parallelogram BD is equal to the square on HE

And BD is equal to the rectilinear figure A

Therefore the rectilinear figure A is also equal to the square which can be described on EH

Therefore a square namely that which can be described on EH , has been constructed equal to the given rectilinear figure A

Q E F



BOOK THREE

DEFINITIONS

1 *Equal circles* are those the diameters of which are equal or the radii of which are equal

2 A straight line is said to *touch a circle* which, meeting the circle and being produced, does not cut the circle

3 *Circles* are said to *touch one another* which meeting one another do not cut one another

4 In a circle straight lines are said to be *equally distant from the centre* when the perpendiculars drawn to them from the centre are equal

5 And that straight line is said to be *at a greater distance* on which the greater perpendicular falls

6 A *segment of a circle* is the figure contained by a straight line and a circumference of a circle

7 An *angle of a segment* is that contained by a straight line and a circumference of a circle

8 An *angle in a segment* is the angle which when a point is taken on the circumference of the segment and straight lines are joined from it to the extremities of the straight line which is the *base of the segment*, is contained by the straight lines so joined

9 And when the straight lines containing the angle cut off a circumference, the angle is said to *stand upon* that circumference

10 A *sector of a circle* is the figure which when an angle is constructed at the centre of the circle, is contained by the straight lines containing the angle and the circumference cut off by them

11 *Similar segments of circles* are those which admit equal angles or in which the angles are equal to one another

BOOK III PROPOSITIONS

PROPOSITION 1

To find the centre of a given circle

Let ABC be the given circle

thus it is required to find the centre of the circle ABC

Let a straight line AB be drawn through it at random, and let it be bisected at the point D

from D let DC be drawn at right angles to AB and let it be drawn through to E let CE be bisected at F

I say that F is the centre of the circle ABC

For suppose it is not but if possible let G be the centre

and let G 1, GD GB be joined

Then since AD is equal to DB and DG is common
the two sides AD DG are equal to the two sides BD DG respectively,
and the base GA is equal to the base GB , for they are
radii,

therefore the angle ADG is equal to the angle GDB

[1 8]

But when a straight line set up on a straight line
makes the adjacent angles equal to one another each of
the equal angles is right,

[1 Def 10]

therefore the angle GDB is right

But the angle FDB is also right

therefore the angle FDB is equal to the angle GDB ,
the greater to the less which is impossible


Therefore G is not the centre of the circle ABC

Similarly we can prove that neither is any other point except F

Therefore the point F is the centre of the circle ABC

PROPOSITION From this it is manifest that if in a circle a straight line cut a
straight line into two equal parts and at right angles, the centre of the circle is
on the cutting straight line

Q E F



PROPOSITION 2

If on the circumference of a circle two points be taken at random, the straight line
joining the points will fall within the circle

Let ABC be a circle and let two points A B be taken at random on its cir-
cumference

I say that the straight line joined from A to B will fall within the circle

For suppose it does not but if possible let it fall outside as AEB ,
let the centre of the circle ABC be taken [III 1], and let it be D , let DA DB be
joined and let DFE be drawn through

Then since DA is equal to DB

the angle DAE is also equal to the angle DBE [1 5]

And since one side AEB of the triangle DAE is pro-
duced

the angle DEB is greater than the angle DAE [1 16]

But the angle DAF is equal to the angle DBF

therefore the angle DEB is greater than the angle
 DBE

And the greater angle is subtended by the greater
side,

[1 19]

therefore DB is greater than DE

But DB is equal to DF

therefore DF is greater than DE ,


the less than the greater which is impossible

Therefore the straight line joined from A to B will not fall outside the circle

Similarly we can prove that neither will it fall on the circumference itself
therefore it will fall within

Therefore etc

Q E D




PROPOSITION 3

If in a circle a straight line through the centre bisect a straight line not through the centre, it also cuts it at right angles and if it cut it at right angles it also bisects it

Let ABC be a circle and in it let a straight line CD through the centre bisect a straight line AB not through the centre at the point F ,

I say that it also cuts it at right angles

For let the centre of the circle ABC be taken and let it be E let EA , EB be joined



Then, since AF is equal to FB and FE is common,
two sides are equal to two sides,

and the base EA is equal to the base EB ,

therefore the angle AFE

is equal to the angle BFE [I 8]

But when a straight line set up on a straight line
makes the adjacent angles equal to one another each
of the equal angles is right, [I Def 10]

therefore each of the angles AFE , BFE is right

Therefore CD which is through the centre and
bisects AB which is not through the centre also cuts it at right angles

Again let CD cut AB at right angles

I say that it also bisects it that is that AF is equal to FB

For with the same construction

since EA is equal to EB ,

the angle EAF is also equal to the angle EBF [I 5]

But the right angle AFE is equal to the right angle BFE therefore EAF
 EBF are two triangles having two angles equal to two angles and one side
equal to one side namely EF which is common to them and subtends one of
the equal angles,

therefore they will also have the remaining sides equal to the remaining sides
[I 26]

therefore AF is equal to FB

Therefore etc

Q E D

PROPOSITION 4

If in a circle two straight lines cut one another which are not through the centre they do not bisect one another

Let $ABCD$ be a circle and in it let the two straight lines AC BD which are
not through the centre cut one another at E

I say that they do not bisect one another

For if possible let them bisect one another so that


AE is equal to EC and BE to ED ,

let the centre of the circle $ABCD$ be taken [III 1]
and let it be F let FE be joined

Then since a straight line FE through the centre
bisects a straight line AC not through the centre

it also cuts it at right angles [III 3]

therefore the angle FEA is right



Again, since a straight line FE bisects a straight line BD ,
 it also cuts it at right angles,
 therefore the angle FEB is right

[III 3]

But the angle FEA was also proved right,
 therefore the angle FFA is equal to the angle FEB the less to the greater
 which is impossible

Therefore AC , BD do not bisect one another

Therefore etc

Q E D

PROPOSITION 5

If two circles cut one another they will not have the same centre

For let the circles ABC , CDG cut one another at the points B , C ,

I say that they will not have the same centre

For if possible let it be E , let EC be joined, and let EFG be drawn through at random

Then, since the point E is the centre of the circle ABC ,

EC is equal to EF [I Def 15]


Again, since the point E is the centre of the circle CDG ,

EC is equal to EG

But EC was proved equal to EF also
 therefore EF is also equal to EG , the less to the
 greater which is impossible

Therefore the point E is not the centre of the circles ABC , CDG

Therefore etc



Q E D

PROPOSITION 6

If two circles touch one another they will not have the same centre

For let the two circles ABC , CDE touch one another at the point C

I say that they will not have the same centre

For if possible let it be F let FC be joined and let FLB be drawn through at random

Then since the point F is the centre of the circle ABC

FC is equal to FB


Again since the point F is the centre of the circle CDE ,

FC is equal to FE

But FC was proved equal to FB
 therefore FE is also equal to FB the less to the
 greater which is impossible

Therefore F is not the centre of the circles ABC , CDE

Therefore etc



Q E D

PROPOSITION 7

If on the diameter of a circle a point be taken which is not the centre of the circle and from the point straight lines fall upon the circle that will be greatest on which the centre is the remainder of the same diameter will be least and of the rest the

nearer to the straight line through the centre is always greater than the more remote and only two equal straight lines will fall from the point on the circle one on each side of the least straight line

Let $ABCD$ be a circle and let AD be a diameter of it, on AD let a point F be taken which is not the centre of the circle let E be the centre of the circle

and from F let straight lines FB , FC , FG fall upon the circle $ABCD$,
I say that FA is greatest FD is least, and of the rest FB is greater than FC , and FC than FG

For let BE CE GE be joined

Then, since in any triangle two sides are greater than the remaining one

[I 20]

EB EF are greater than BF

But AE is equal to BE

therefore AF is greater than BF

Again since BE is equal to CE and FE is common

the two sides BE EF are equal to the two sides CE EF

But the angle BEF is also greater than the angle CEF

therefore the base BF is greater than the base CF

[I 24]

For the same reason

CF is also greater than FG

Again since GF , FE are greater than EG

and EG is equal to ED

GF , FE are greater than ED

Let EF be subtracted from each,

therefore the remainder GF is greater than the remainder FD

Therefore FA is greatest FD is least, and FB is greater than FC and FC than FG

I say also that from the point F only two equal straight lines will fall on the circle $ABCD$, one on each side of the least FD

For on the straight line EF and at the point E on it let the angle FEH be constructed equal to the angle GEF [I 23] and let FH be joined

Then since GE is equal to EH

and EF is common

the two sides GE EF are equal to the two sides HE EF

and the angle GEF is equal to the angle HEF

therefore the base GF is equal to the base FH

[I 4]

I say again that another straight line equal to FG will not fall on the circle from the point F


For if possible let FK so fall

Then since FA is equal to FG and FH to FG

FK is also equal to FH

the nearer to the straight line through the centre being thus equal to the more remote which is impossible

Therefore another straight line equal to GF will not fall from the point F upon the circle



therefore only one straight line will so fall

Therefore etc

Q. E. D.

PROPOSITION 8

If a point be taken outside a circle and from the point straight lines be drawn through to the circle one of which is through the centre and the others are drawn at random, then of the straight lines which fall on the concave circumference that through the centre is greatest while of the rest the nearer to that through the centre is always greater than the more remote but of the straight lines falling on the convex circumference that between the point and the diameter is least while of the rest the nearer to the least is always less than the more remote and only two equal straight lines will fall on the circle from the point one on each side of the least

Let ABC be a circle and let a point D be taken outside ABC , let there be drawn through from it straight lines DA, DE, DF, DC , and let DA be through the centre,

I say that, of the straight lines falling on the concave circumference $AEFC$ the straight line DA through the centre is greatest

while DE is greater than DF and DF than DC ,
but of the straight lines falling on the convex circumference $HLKG$ the straight line DG between the point and the diameter AG is least and the nearer to the least DG is always less than the more remote namely DA than DL and DI than DH

For let the centre of the circle ABC be taken [III 1] and let it be M , let ME, MF, MC, MA, ML, MH be joined

Then since AM is equal to EM , let MD be added to each

therefore AD is equal to $FM + MD$

But EM, MD are greater than ED

therefore AD is also greater than ED

Again since ME is equal to MF

and MD is common

therefore $EM + MD$ are equal to $FM + MD$

and the angle EMD is greater than the angle FMD

therefore the base ED is greater than the base FD

[I 24]

Similarly we can prove that FD is greater than CD therefore DA is greatest while DL is greater than DF and DF than DC

Next since MA, AD are greater than MD

[I 20]

and MG is equal to MA

therefore the remainder AD is greater than the remainder GD

so that GD is less than AD

And since on MD one of the sides of the triangle MLD two straight lines MA, AD were constructed meeting within the triangle

therefore MA, AD are less than $ML + LD$

[I 21]


and MA is equal to ML ,

therefore the remainder DA is less than the remainder DL

Similarly we can prove that DL is also less than DH

therefore DG is least while DA is less than DL and DL than DH

I say also that only two equal straight lines will fall from the point D on the



circle, one on each side of the least DG

On the straight line MD , and at the point M on it, let the angle DMB be constructed equal to the angle KMD , and let DB be joined

Then, since MA is equal to MB ,

and MD is common

the two sides KM , MD are equal to the two sides BM , MD respectively,

and the angle KMD is equal to the angle BMD ,

therefore the base DA is equal to the base DB [I 4]

I say that no other straight line equal to the straight line DA will fall on the circle from the point D

For, if possible let a straight line so fall and let it be DN

Then, since DK is equal to DN ,

while DK is equal to DB

DB is also equal to DN ,

that is, the nearer to the least DG equal to the more remote which was proved impossible

Therefore no more than two equal straight lines will fall on the circle ABC from the point D , one on each side of DG the least

Therefore etc

Q E D

PROPOSITION 9

If a point be taken within a circle, and more than two equal straight lines fall from the point on the circle, the point taken is the centre of the circle

Let ABC be a circle and D a point within it, and from D let more than two equal straight lines, namely DA , DB , DC , fall on the circle ABC ,

I say that the point D is the centre of the circle ABC

For let AB , BC be joined and bisected at the points E , F , and let ED , FD be joined and drawn through to the points G , K , H , L

Then since AE is equal to EB and ED is common

the two sides AE , ED are equal to the two sides BE , ED

and the base DA is equal to the base DB

therefore the angle AED is equal to the angle BED [I 8]

Therefore each of the angles AED , BED is right [I Def 10]

therefore GK cuts AB into two equal parts and at right angles

And since if in a circle a straight line cut a straight line into two equal parts and at right angles the centre of the circle is on the cutting straight line [III 1 Por]

the centre of the circle is on GK

For the same reason


the centre of the circle ABC is also on HL

And the straight lines GK , HL have no other point common but the point D

therefore the point D is the centre of the circle ABC

Therefore etc

Q E D



PROPOSITION 10

A circle does not cut a circle at more points than two

For if possible let the circle ABC cut the circle DEF at more points than two namely B, C, F, H ,

let BH, BG be joined and bisected at the points K, L ,
and from K, L let AC, LM be drawn at right angles to BH, BG and carried through to the points A, E

Then since in the circle ABC a straight line AC cuts a straight line BH into two equal parts and at right angles

the centre of the circle ABC is on AC

[III 1, Por]

Again since in the same circle ABC a straight line MO cuts a straight line BG into two equal parts and at right angles


the centre of the circle ABC is on MO

But it was also proved to be on AC , and
the straight lines AC, MO meet at no point
except at P ,
therefore the point P is the centre of the
circle ABC

Similarly we can prove that P is also the centre of the circle DEF ,
therefore the two circles ABC, DEF which cut one another have the same centre P which is impossible

Therefore etc

[III 5]
Q E D



PROPOSITION 11

If two circles touch one another internally and their centres be taken the straight line joining their centres, if it be also produced will fall on the point of contact of the circles

For let the two circles ABC, ADE touch one another internally at the point A and let the centre F of the circle ABC and the centre G of ADE , be taken,

I say that the straight line joined from G to F
and produced will fall on A

For suppose it does not but if possible let it fall as FGH and let AF, AG be joined

Then since AG, GF are greater than FA that
is than FH

let FG be subtracted from each
therefore the remainder AG is greater than the
remainder GH


But AG is equal to GD ,

therefore GD is also greater than GH
the less than the greater which is impossible

Therefore the straight line joined from F to
 G will not fall outside,

therefore it will fall at A on the point of contact

Therefore etc



Q E D

PROPOSITION 12

If two circles touch one another externally, the straight line joining their centres will pass through the point of contact

For let the two circles ABC , ADE touch one another externally at the point A , and let the centre F of ABC and the centre G of ADE , be taken

I say that the straight line joined from F to G will pass through the point of contact at A

For suppose it does not but, if possible, let it pass as $FCDG$, and let AF , AG be joined

Then since the point F is the centre of the circle ABC ,

FA is equal to FC

Again, since the point G is the centre of the circle ADE

GA is equal to GD

But FA was also proved equal to FC , therefore FA , AG are equal to FC , GD so that the whole FG is greater than FA , AG ,

but it is also less [I 20] which is impossible

Therefore the straight line joined from F to G will not fail to pass through the point of contact at A ,

therefore it will pass through it

Therefore etc

Q E D

PROPOSITION 13

A circle does not touch a circle at more points than one whether it touch it internally or externally

For, if possible let the circle $ABDC$ touch the circle $EBFD$ first internally, at more points than one namely D , B

Let the centre G of the circle $ABDC$ and the centre H of $EBFD$ be taken

Therefore the straight line joined from G to H will fall on B , D [III 11]

Let it so fall as $BGHD$

Then since the point G is the centre of the circle $ABCD$

BG is equal to GD ,

therefore BG is greater than HD

therefore BH is much greater than HD

Again since the point H is the centre of the circle $EBFD$



BH is equal to HD ,

but it was also proved much greater than it which is impossible

Therefore a circle does not touch a circle internally at more points than one

I say further that neither does it so touch it externally

For if possible let the circle ACH touch the circle $ABDC$ at more points than one namely A , C ,



and let AC be joined

Then, since on the circumference of each of the circles $ABDC$, ACK two points A , C have been taken at random, the straight line joining the points will fall within each circle, [III 2]

but it fell within the circle $ABDC$ and outside ACK [III Def 3] which is absurd

Therefore a circle does not touch a circle externally at more points than one
And it was proved that neither does it so touch it internally

Therefore etc

Q E D

PROPOSITION 14

In a circle equal straight lines are equally distant from the centre and those which are equally distant from the centre are equal to one another

Let $ABDC$ be a circle and let AB , CD be equal straight lines in it,
I say that AB , CD are equally distant from the centre

For let the centre of the circle $ABDC$ be taken [III 1] and let it be E , from E let EF , EG be drawn perpendicular to AB , CD and let $1F$ EC be joined

Then since a straight line EF through the centre cuts a straight line AB not through the centre at right angles it also bisects it [III 3]

Therefore AF is equal to FB ,
therefore AB is double of AF

For the same reason

CD is also double of CG ,

and AB is equal to CD

therefore AF is also equal to CG

And since AE is equal to EC ,

the square on AE is also equal to the square on EC

But the squares on AF EF are equal to the square on AE for the angle at F is right,

and the squares on EG GC are equal to the square on EC , for the angle at G is right, [I 47]

therefore the squares on AF FE are equal to the squares on CG GF ,
of which the square on AF is equal to the square on CG for AF is equal to CG ,

therefore the square on FE which remains is equal to the square on EG

therefore EF is equal to EG

But in a circle straight lines are said to be equally distant from the centre when the perpendiculars drawn to them from the centre are equal [III Def 4],

therefore AB CD are equally distant from the centre

Next let the straight lines AB CD be equally distant from the centre that is let EF be equal to EG

I say that AB is also equal to CD


For with the same construction we can prove similarly, that AB is double of AF and CD of CG

And since AE is equal to CE

the square on AE is equal to the square on CE

But the squares on EF FA are equal to the square on AE and the squares on EG GC equal to the square on CE [I 47]

Therefore the squares on EF FA are equal to the squares on EG GC ,
of which the square on EF is equal to the square on EG for EF is equal to EG ,



therefore the square on AF which remains is equal to the square on CG ,
 therefore AF is equal to CG
 And AB is double of AF , and CD double of CG
 therefore AB is equal to CD

Therefore etc

Q E D

PROPOSITION 15

Of straight lines in a circle the diameter is greatest and of the rest the nearer to the centre is always greater than the more remote

Let $ABCD$ be a circle, let AD be its diameter and E the centre, and let BC be nearer to the diameter AD , and FG more remote

I say that AD is greatest and BC greater than FG

For from the centre E let EH , EA be drawn perpendicular to BC FG

Then since BC is nearer to the centre and FG more remote EK is greater than EH [III Def 5]

Let EL be made equal to EH through L let IM be drawn at right angles to EA and carried through to N , and let ME EN , FE , EG be joined

Then since EH is equal to EL

BC is also equal to MN

[III 14]

Again since AE is equal to EM and ED to EN ,

AD is equal to ME , EN

But ME , EN are greater than MN ,

[I 20]

and MN is equal to BC ,

therefore AD is greater than BC

And since the two sides ME EN are equal to the two sides FE EG

and the angle MEN greater than the angle FEG

therefore the base MN is greater than the base FG

[I 24]

But MN was proved equal to BC

Therefore the diameter AD is greatest and BC greater than FG

Therefore etc

Q E D

PROPOSITION 16

The straight line drawn at right angles to the diameter of a circle from its extremity will fall outside the circle and into the space between the straight line and the circumference another straight line cannot be interposed further the angle of the semi circle is greater and the remaining angle less, than any acute rectilineal angle

Let ABC be a circle about D as centre and AB as diameter,

I say that the straight line drawn from A at right angles to AB from its extremity will fall outside the circle

For suppose it does not but if possible let it fall within as CA and let DC be joined

Since DA is equal to DC

the angle DAC is also equal to the angle ACD

[I 5]

But the angle DAC is right

therefore the angle ACD is also right

thus, in the triangle ACD the two angles DAC ACD are equal to two right angles which is impossible

[I 17]

Therefore the straight line drawn from the point *A* at right angles to *BA* will not fall within the circle

Similarly we can prove that neither will it fall on the circumference,

therefore it will fall outside

Let it fall as *AE*,

I say next that into the space between the straight line *AE* and the circumference *CHA* another straight line cannot be interposed

For if possible let another straight line be so interposed, as *FA* and let *DG* be drawn from the point *D* perpendicular to *FA*

Then, since the angle *AGD* is right,

and the angle *DAG* is less than a right angle,

AD is greater than *DG*

{ 19 }

But *DA* is equal to *DH*,

therefore *DH* is greater than *DG*, the less than the greater which is impossible

Therefore another straight line cannot be interposed into the space between the straight line and the circumference

I say further that the angle of the semicircle contained by the straight line *BA* and the circumference *CHA* is greater than any acute rectilineal angle and the remaining angle contained by the circumference *CHA* and the straight line *AE* is less than any acute rectilineal angle

For if there is any rectilineal angle greater than the angle contained by the straight line *BA* and the circumference *CHA* and any rectilineal angle less than the angle contained by the circumference *CHA* and the straight line *AE*, then into the space between the circumference and the straight line *AE* a straight line will be interposed such as will make an angle contained by straight lines which is greater than the angle contained by the straight line *BA* and the circumference *CHA*, and another angle contained by straight lines which is less than the angle contained by the circumference *CHA* and the straight line *AE*

But such a straight line cannot be interposed

therefore there will not be any acute angle contained by straight lines which is greater than the angle contained by the straight line *BA* and the circumference *CHA* nor yet any acute angle contained by straight lines which is less than the angle contained by the circumference *CHA* and the straight line *AE* —

PROB. From this it is manifest that the straight line drawn at right angles to the diameter of a circle from its extremity touches the circle Q E D


PROPOSITION 17

From a given point to draw a straight line touching a given circle

Let *A* be the given point and *BCD* the given circle thus it is required to draw from the point *A* a straight line touching the circle *BCD*

For let the centre *F* of the circle be taken, [III 1]
let *AE* be joined and with centre *F* and distance *FA* let the circle *AFG* be described,

from *D* let *DF* be drawn at right angles to *EA*,
and let *EF* *AB* be joined



I say that AB has been drawn from the point A touching the circle BCD
 For, since E is the centre of the circles BCD AFG

EA is equal to EF , and ED to EB ,

therefore the two sides AE , EB are equal to the two sides FE , ED ,
 and they contain a common angle, the angle at E ,
 therefore the base DF is equal to the base AB ,
 and the triangle DEF is equal to the triangle BEA ,
 and the remaining angles to the remaining angles

[I 4]

therefore the angle EDF is equal to the angle EBA

But the angle EDF is right,

therefore the angle EBA is also right

Now EB is a radius

and the straight line drawn at right angles to the di-
 ameter of a circle from its extremity, touches the
 circle,

[III 16 Por]

therefore AB touches the circle BCD

Therefore from the given point A the straight line AB has been drawn
 touching the circle BCD

Q E F

PROPOSITION 18

If a straight line touch a circle, and a straight line be joined from the centre to the
 point of contact the straight line so joined will be perpendicular to the tangent

For let a straight line DE touch the circle ABC at the point C , let the centre
 F of the circle ABC be taken, and let FC be joined from F to C ,

I say that FC is perpendicular to DE

For, if not let FG be drawn from F perpen-
 dicular to DE

Then since the angle FGC is right

the angle FCG is acute [I 17]

and the greater angle is subtended by the great-
 er side, [I 19]

therefore FC is greater than FG

But FC is equal to FB ,

therefore FB is also greater than FG

the less than the greater which is impossible

Therefore FG is not perpendicular to DE

Similarly we can prove that neither is any other straight line except FC ,
 therefore FC is perpendicular to DE

Therefore etc

Q E D

PROPOSITION 19

If a straight line touch a circle and from the point of contact a straight line be
 drawn at right angles to the tangent the centre of the circle will be on the straight
 line so drawn

For let a straight line DE touch the circle ABC at the point C and from C
 let CA be drawn at right angles to DE

I say that the centre of the circle is on AC

For suppose it is not but if possible let F be the centre

and let CF be joined

Since a straight line DE touches the circle ABC ,

and FC has been joined from the centre to the point of contact,

FC is perpendicular to DE , [III 18]

therefore the angle FCE is right


But the angle ACE is also right,
therefore the angle FCE is equal to the angle ACE ,

the less to the greater which is impossible

Therefore F is not the centre of the circle ABC

Similarly we can prove that neither is any other point except a point on AC

Therefore etc



PROPOSITION 20

In a circle the angle at the centre is double of the angle at the circumference when the angles have the same circumference as base

Let ABC be a circle let the angle BEC be an angle at its centre, and the angle BAC an angle at the circumference and let them have the same circumference BC as base

I say that the angle BEC is double of the angle BAC

For let AE be joined and drawn through to F

Then since EA is equal to EB ,

the angle EAB is also equal to the angle EBA , [I 5]
therefore the angles EAB , EBA are double of the angle EAB

But the angle BEF is equal to the angles EAB , EBA [I 32]
therefore the angle BEF is also double of the angle EAB

For the same reason

the angle FEC is also double of the angle EAC


Therefore the whole angle BEC is double of the whole angle BAC

Again let another straight line be inflected and let there be another angle BDC , let DE be joined and produced to G

Similarly then we can prove that the angle GEC is double of the angle EDC ,
of which the angle GEB is double of the angle EDB

therefore the angle BEC which remains is double of the angle BDC

Therefore etc




PROPOSITION 21

In a circle the angles in the same segment are equal to one another

Let $ABCD$ be a circle and let the angles BAD , BED be angles in the same segment $BAED$

I say that the angles BAD , BED are equal to one another

For let the centre of the circle $ABCD$ be taken and let it be F , let BF , FD be joined



Now, since the angle BFD is at the centre,
and the angle BAD at the circumference
and they have the same circumference BCD as base
therefore the angle BFD is double of the angle
 BAD [III 20]

For the same reason
the angle BFD is also double of the angle BFD
therefore the angle BAD is equal to the angle BED
Therefore etc Q E D

PROPOSITION 22

The opposite angles of quadrilaterals in circles are equal to two right angles

Let $ABCD$ be a circle and let $ABCD$ be a quadrilateral in it,
I say that the opposite angles are equal to two right angles

Let AC BD be joined

Then since in any triangle the three angles are equal
to two right angles [I 32]

the three angles CAB ABC , BCA of the triangle ABC
are equal to two right angles

But the angle CAB is equal to the angle BDC for they
are in the same segment $BADC$ [III 21]

and the angle ACB is equal to the angle ADB for they
are in the same segment $ADCB$

therefore the whole angle ADC is equal to the angles BAC ACB

Let the angle ABC be added to each

therefore the angles ABC , BAC ACB are equal to the angles ABC ADC

But the angles ABC BAC , ACB are equal to two right angles,

therefore the angles ABC ADC are also equal to two right angles


Similarly we can prove that the angles BAD , DCB are also equal to two
right angles

Therefore etc Q E D

PROPOSITION 23

*On the same straight line there cannot be constructed two similar and unequal
segments of circles on the same side*

For if possible on the same straight line AB let two similar and unequal
segments of circles ACB ADB be constructed on the
same side



let ACD be drawn through and let CB DB be joined

Then since the segment ACB is similar to the seg-
ment ADB ,

and similar segments of circles are those which admit equal angles [III Def 11]
the angle ACB is equal to the angle ADB , the exterior to the interior which is
impossible [I 16]

Therefore etc Q E D

PROPOSITION 21

Similar segments of circles on equal straight lines are equal to one another

For let AEB , CFD be similar segments of circles on equal straight lines AB , CD ,

I say that the segment AEB is equal to the segment CFD

I or, if the segment AEB be applied to CFD and if the point A be placed on C and the straight line AB on CD ,

the point B will also coincide with the point D , because AB is equal to CD ,



and AB coinciding with CD ,

the segment AEB will also coincide with CFD

For if the straight line AB coincide with CD but the segment AEB do not coincide with CFD

it will either fall within it or outside it

or it will fall away as CGD and a circle cuts a circle at more points than two which is impossible [III 10]

Therefore if the straight line AB be applied to CD the segment AEB will not fail to coincide with CFD also,

therefore it will coincide with it and will be equal to it

Therefore etc

Q E D

PROPOSITION 22

Given a segment of a circle to describe the complete circle of which it is a segment

Let ABC be the given segment of a circle, thus it is required to describe the complete circle belonging to the segment ABC that is of which it is a segment

For let AC be bisected at D , let DB be drawn from the point D at right angles to AC and let AB be joined

the angle ABD is then greater than equal to or less than the angle BAD

First let it be greater

and on the straight line BA and at the point A on it, let the angle BAE be constructed equal to the angle ABD , let DB be drawn through to E and let EC be joined

Then since the angle ABE is equal to the angle BAE the straight line EB is also equal to EA [I 6]


And since AD is equal to DC and DE is common the two sides AD , DE are equal to the two sides CD , DE respectively

and the angle ADE is equal to the angle CDE for each is right,

therefore the base AE is equal to the base CE

But AE was proved equal to BE

therefore BE is also equal to CE ,




therefore the three straight lines AE , EB , EC are equal to one another

Therefore the circle drawn with centre E and distance one of the straight lines AE , EB , EC will also pass through the remaining points and will have been completed

[III 9]

Therefore, given a segment of a circle, the complete circle has been described



And it is manifest that the segment ABC is less than a semicircle because the centre E happens to be outside it

Similarly, even if the angle ABD be equal to the angle BAD ,

AD being equal to each of the two BD , DC the three straight lines DA , DB , DC will be equal to one another,

D will be the centre of the completed circle

and ABC will clearly be a semicircle

But, if the angle ABD be less than the angle BAD , and if we construct on the straight line BA and at the point A on it an angle equal to the angle ABD the centre will fall on DB within the segment ABC , and the segment ABC will clearly be greater than a semicircle



Therefore, given a segment of a circle, the complete circle has been described

Q E F



PROPOSITION 26

In equal circles equal angles stand on equal circumferences, whether they stand at the centres or at the circumferences

Let ABC , DEF be equal circles, and in them let there be equal angles, namely at the centres the angles BGC , EHF and at the circumferences the angles BAC , EDF

I say that the circumference BAC is equal to the circumference ELF

For let BC , EF be joined



Now, since the circles ABC , DEF are equal

the radii are equal

Thus the two straight lines BG , GC are equal to the two straight lines EH , HF , and the angle at G is equal to the angle at H

therefore the base BC is equal to the base EF [I 4]

And since the angle at A is equal to the angle at D

the segment BAC is similar to the segment EDF [III Def 11]

and they are upon equal straight lines

But similar segments of circles on equal straight lines are equal to one another, [III 24]

therefore the segment BAC is equal to EDF

But the whole circle ABC is also equal to the whole circle DEF , therefore the circumference BAC which remains is equal to the circumference ELF

Therefore etc

Q E D

PROPOSITION 27

In equal circles angles standing on equal circumferences are equal to one another, whether they stand at the centres or at the circumferences

For in equal circles ABC , DEF , on equal circumferences BC EF , let the angles BGC , EHF stand at the centres G H , and the angles BAC , EDF at the circumferences,

I say that the angle BGC is equal to the angle EHF ,
and the angle BAC is equal to the angle EDF

For, if the angle BGC is unequal to the angle EHF ,

one of them is greater

Let the angle BGC be greater and on the straight line BG , and at the point G on it let the angle BGA be constructed equal to the angle EHF [I 23]

Now equal angles stand on equal circumferences, when they are at the centres

therefore the circumference BA is equal to the circumference EF [III 26]

But EF is equal to BC ,
therefore BA is also equal to BC the less to the greater which is impossible

Therefore the angle BGC is not unequal to the angle EHF ,

therefore it is equal to it

And the angle at A is half of the angle BGC

and the angle at D half of the angle EHF , [III 20]

therefore the angle at A is also equal to the angle at D

Therefore etc

Q E D

PROPOSITION 28

In equal circles equal straight lines cut off equal circumferences the greater equal to the greater and the less to the less

Let ABC , DEF be equal circles and in the circles let AB , DE be equal straight lines cutting off ACB , DFE as greater circumferences and AGB , DHE as lesser

I say that the greater circumference ACB is equal to the greater circumference DFE and the less circumference AGB to DHE

For let the centres K L of the circles be taken and let AK KB DL LE be joined

Now since the circles are equal
the radii are also equal
therefore the two sides AK KB are equal
to the two sides DL LE ,





and the base AB is equal to the base DE

therefore the angle AKB is equal to the angle DLE [I 8]

But equal angles stand on equal circumferences when they are at the centres [III 26]

therefore the circumference AGB is equal to DHE

And the whole circle ABC is also equal to the whole circle DEF ,



therefore the circumference ACB which remains is also equal to the circumference DFE which remains

Therefore etc

Q E D

PROPOSITION 29


In equal circles equal circumferences are subtended by equal straight lines

Let ABC DEF be equal circles and in them let equal circumferences BGC EHF be cut off and let the straight lines BC , EF be joined

I say that BC is equal to EF

For let the centres of the circles be taken and let them be K L let BK KC EL LF be joined

Now, since the circumference BGC is equal to the circumference EHF , the angle BKC is also equal to the angle ELF [III 27]



And since the circles ABC DEF are equal

the radii are also equal

therefore the two sides BK , KC are equal to the two sides EL LF and they contain equal angles

therefore the base BC is equal to the base EF [I 4]

Therefore etc


Q E D

PROPOSITION 30

To bisect a given circumference

Let ADB be the given circumference,

thus it is required to bisect the circumference ADB



Let AB be joined and bisected at C from the point C let CD be drawn at right angles to the straight line AB and let AD DB be joined

Then since AC is equal to CB and CD is common

the two sides AC CD are equal to the two sides BC CD

and the angle ACD is equal to the angle BCD for each is right

therefore the base AD is equal to the base DB [I 4]

But equal straight lines cut off equal circumferences the greater equal to the greater, and the less to the less [III 25]

and each of the circumferences AD DB is less than a semicircle

therefore the circumference AD is equal to the circumference DB

Therefore the given circumference has been bisected at the point D

Q E F

PROPOSITION 31

In a circle the angle in the semicircle is right that in a greater segment less than a right angle and that in a less segment greater than a right angle and further the angle of the greater segment is greater than a right angle and the angle of the less segment less than a right angle


Let $ABCD$ be a circle let BC be its diameter and E its centre and AC , AD DC be joined

I say that the angle BAC in the semicircle BAC is right
the angle ABC in the segment ABC greater than the semicircle is less than a
right angle.

and the angle ADC in the segment ADC less than the semicircle is greater than a right angle

Let AE be joined, and let BA be carried through to F

Then since BE is equal to EA
the angle ABE is also equal to the angle BAE [1 5]



Again since CE is equal to EA ,
the angle ACE is also equal to the angle CAE

Therefore the whole angle BAC is equal to the two angles ABC, ACB
But the angle FAC exterior to the triangle ABC is also equal to the two angles ABC, ACB .

therefore the angle BAC is also equal to the angle FAC ,
therefore each is right. [I Def 10]

therefore the angle BAC in the semicircle BAC is right

Next since in the triangle ABC the two angles ABC, BAC are less than two right angles

and the angle BAC is a right angle,

the angle ABC is less than a right angle.

and it is the angle in the segment ABC greater than the semicircle

Next, since $ABCD$ is a quadrilateral in a circle and the opposite angles of quadrilaterals in circles are equal to two right angles

while the angle ABC is less than a right angle

therefore the angle ADC which remains is greater than a right angle

and it is the angle in the segment ADC less than the semicircle

I say further that the angle of the greater segment namely that contained by the circumference ABC and the straight line AC , is greater than a right angle

and the angle of the less segment namely that contained by the circumference ADC and the straight line AC is less than a right angle

This is at once manifest

For since the angle contained by the straight lines BA, AC is right the angle contained by the circumference ABC and the straight line AC is greater than a right angle

Again since the angle contained by the straight lines AC AF is right the angle contained by the straight line CA and the circumference ADC is less than a right angle

Therefore etc

Q E D

PROPOSITION 32

If a straight line touch a circle and from the point of contact there be drawn across, in the circle, a straight line cutting the circle, the angles which it makes with the tangent will be equal to the angles in the alternate segments of the circle

For let a straight line EF touch the circle $ABCD$ at the point B , and from

the point B let there be drawn across, in the circle $ABCD$ a straight line BD cutting it,

I say that the angles which BD makes with the tangent EF will be equal to the angles in the alternate segments of the circle that is that the angle FBD is equal to the angle constructed in the segment BAD and the angle EBD is equal to the angle constructed in the segment DCB

For let BA be drawn from B at right angles to EF ,
let a point C be taken at random on the circumference BD ,

and let AD DC , CB be joined

Then, since a straight line EF touches the circle $ABCD$ at B ,

and BA has been drawn from the point of contact at right angles to the tangent the centre of the circle $ABCD$ is on BA [III 19]

Therefore BA is a diameter of the circle $ABCD$

therefore the angle ADB , being an angle in a semicircle is right [III 31]

Therefore the remaining angles BAD ABD are equal to one right angle [I 32]

But the angle ABF is also right,

therefore the angle ABF is equal to the angles BAD ABD

Let the angle ABD be subtracted from each

therefore the angle DBF which remains is equal to the angle BAD in the alternate segment of the circle

Next since $ABCD$ is a quadrilateral in a circle,

its opposite angles are equal to two right angles [III 22]

But the angles DBF DBE are also equal to two right angles,

therefore the angles DBF DBE are equal to the angles BAD BCD ,

of which the angle BAD was proved equal to the angle DBF ,

therefore the angle DBE which remains is equal to the angle DCB in the alternate segment DCB of the circle

Therefore etc

Q E D

PROPOSITION 33



On a given straight line to describe a segment of a circle admitting an angle equal to a given rectilineal angle

Let AB be the given straight line and the angle at C the given rectilineal angle,

thus it is required to describe on the given straight line AB a segment of a circle admitting an angle equal to the angle at C

The angle at C is then acute or right or obtuse

First let it be acute and as in the first figure on the straight line AB and at the point A let the angle BAD be constructed equal to the angle at C



therefore the angle BAD is also acute

Let AE be drawn at right angles to DA , let AB be bisected at F , let FG be drawn from the point F at right angles to AB , and let GB be joined

Then, since AF is equal to FB

and FG is common

the two sides AF FG are equal to the two sides BF FG ,

and the angle AFG is equal to the angle BFG ,

therefore the base AG is equal to the base BG [I 4]

Therefore the circle described with centre G and distance GA will pass through B also

Let it be drawn, and let it be ABE ,

let EB be joined

Now since AD is drawn from A the extremity of the diameter AE , at right angles to AE

therefore AD touches the circle ABE [III 16 Por]

Since then a straight line AD touches the circle ABE and from the point of contact at A a straight line AB is drawn across in the circle ABE

the angle DAB is equal to the angle AEB in the alternate segment of the circle [III 32]

But the angle DAB is equal to the angle at C ,

therefore the angle at C is also equal to the angle AEB

Therefore on the given straight line AB the segment AEB of a circle has been described admitting the angle AEB equal to the given angle the angle at C

Next let the angle at C be right and let it be again required to describe on AB a segment of a circle admitting an angle equal to the right angle at C

Let the angle BAD be constructed equal to the right angle at C as is the case in the second figure

let AB be bisected at F and with centre F and distance either FA or FB let the circle AEB be described

Therefore the straight line AD touches the circle ABE because the angle at A is right [III 16 Por]

And the angle BAD is equal to the angle in the segment AEB for the latter too is itself a right angle being an angle in a semicircle [III 31]

But the angle BAD is also equal to the angle at C


Therefore the angle AEB is also equal to the angle at C

Therefore again the segment AEB of a circle has been described on AB admitting an angle equal to the angle at C


Next let the angle at C be obtuse

and on the straight line AB and at the point A let the angle BAD be constructed equal to it as is the case in the third figure

let AB be drawn at right angles to AD let AB be again bisected at F let FG be drawn at right angles to AB and let GB be joined



Then, since AF is again equal to FB ,
and FG is common.



the two sides AF, FG are equal to
the two sides BF, FG ,
and the angle AFG is equal to the
angle BFG ,
therefore the base AG is equal to
the base BG [I. 4]

Therefore the circle described with centre G and distance GA will pass through B also, let it so pass, as AEB

Now, since AD is drawn at right angles to the diameter AE from its extremity,

AD touches the circle AEB [III 16, Por.]

And AB has been drawn across from the point of contact at A ,
therefore the angle BAD is equal to the angle constructed in the alternate seg-
ment AHB of the circle

But the angle BAD is equal to the angle at C

Therefore the angle in the segment AHB is also equal to the angle at C


Therefore on the given straight line AB the segment AHB of a circle has been described admitting an angle equal to the angle at C Q E F

PROPOSITION 34

From a given circle to cut off a segment admitting an angle equal to a given rectilineal angle

Let ABC be the given circle and the angle at D the given rectilineal angle, thus it is required to cut off from the circle ABC a segment admitting an angle equal to the given rectilineal angle, the angle at D .

Let EF be drawn touching ABC at the point B and on the straight line FB and at the point B on it let the angle FBC be constructed equal to the angle



Then since a straight line EF touches the circle ABC ,

and BC has been drawn across from the point of contact at B

the angle FBC is equal to the angle constructed in the alternate segment BAC (III 32)

But the angle FBC is equal to the angle at D .

therefore the angle in the segment BAC is equal to the angle at D

Therefore from the given circle ABC the segment BAC has been cut off admitting an angle equal to the given rectilineal angle the angle at D $Q E F$

PROPOSITION 35

If in a circle two straight lines cut one another the rectangle contained by the segments of the one is equal to the rectangle contained by the segments of the other

For in the circle $ABCD$ let the two straight lines AC BD cut one another at the point E

I say that the rectangle contained by AE , EC is equal to the rectangle contained by DE , EB

If now AC , BD are through the centre so that E is the centre of the circle $ABCD$

it is manifest that, AE EC DE EB being equal the rectangle contained by AE , EC is also equal to the rectangle contained by DE , EB

Next let AC DB not be through the centre, let the centre of $ABCD$ be taken, and let it be F from F let FG , FH be drawn perpendicular to the straight lines AC , DB , and let FB FC , FE be joined

Then since a straight line GF through the centre cuts a straight line AC not through the centre at right angles

it also bisects it, [III 3]

therefore AG is equal to GC

Since then, the straight line AC has been cut into equal parts at G and into unequal parts at E the rectangle contained by AE EC together with the square on EG is equal to the square on GC , [II 5]

Let the square on GF be added therefore the rectangle AE , EC together with the squares on GF , GF is equal to the squares on CG GF

But the square on FE is equal to the squares on EG GF , and the square on FC is equal to the squares on CG GF , [I 47] therefore the rectangle AE , EC together with the square on FE is equal to the square on FC

And FC is equal to FB therefore the rectangle AE EC together with the square on EF is equal to the square on FB



For the same reason also the rectangle DE EB together with the square on FE is equal to the square on FB

But the rectangle AE EC together with the square on FE was also proved equal to the square on FB , therefore the rectangle AE EC together with the square on FE is equal to the rectangle DE EB together with the square on FE

Let the square on FE be subtracted from each, therefore the rectangle contained by AE EC which remains is equal to the rectangle contained by DE EB

Therefore etc

Q E D




PROPOSITION 36

If a point be taken outside a circle and from it there fall on the circle two straight lines and if one of them cut the circle and the other touch it the rectangle contained by the whole of the straight line which cuts the circle and the straight line intercepted on it outside between the point and the convex circumference will be equal to the square on the tangent

For let a point D be taken outside the circle ABC and from D let the two

straight lines DCA , DB fall on the circle ABC , let DCA cut the circle ABC and let DB touch it

I say that the rectangle contained by AD DC is equal to the square on DB



Then DCA is either through the centre or not through the centre

First let it be through the centre and let F be the centre of the circle ABC let FB be joined,

therefore the angle FBD is right [III 18]

And, since AC has been bisected at F and CD is added to it,

the rectangle AD , DC together with the square on FC is equal to the square on FD [II 6]

But FC is equal to FB ,
therefore the rectangle AD , DC together with the square on FB is equal to the square on FD

And the squares on FB , BD are equal to the square on FD [I 47]
therefore the rectangle AD , DC together with the square on FB is equal to the squares on FB , BD

Let the square on FB be subtracted from each
therefore the rectangle AD DC which remains is equal to the square on the tangent DB

Again let DCA not be through the centre of the circle ABC ,

let the centre E be taken, and from E let EF be drawn perpendicular to AC ,

let EB , EC , ED be joined

Then the angle EBD is right [III 18]

And since a straight line EF through the centre cuts a straight line AC not through the centre at right angles,

it also bisects it [III 3]

therefore AF is equal to FC

Now, since the straight line AC has been bisected at the point F and CD is added to it

the rectangle contained by AD DC together with the square on FC is equal to the square on FD [II 6]

Let the square on FE be added to each
therefore the rectangle AD , DC together with the squares on CF FE is equal to the squares on FD FE

But the square on EC is equal to the squares on CF FE for the angle EFC is right [I 47]

and the square on ED is equal to the squares on DF FE
therefore the rectangle AD , DC together with the square on EC is equal to the square on ED

And EC is equal to EB
therefore the rectangle AD DC together with the square on EB is equal to the square on ED

But the squares on EB BD are equal to the square on ED for the angle EBD is right [I 47]

therefore the rectangle AD, DC together with the square on EB is equal to the squares on EB, BD

Let the square on EB be subtracted from each,

therefore the rectangle AD, DC which remains is equal to the square on DB


Therefore etc

Q E D

PROPOSITION 37

If a point be taken outside a circle and from the point there fall on the circle two straight lines, if one of them cut the circle, and the other fall on it and if further the rectangle contained by the whole of the straight line which cuts the circle and the straight line intercepted on it outside between the point and the convex circumference be equal to the square on the straight line which falls on the circle the straight line which falls on it will touch the circle

For let a point D be taken outside the circle ABC , from D let the two straight lines DCA, DB fall on the circle ABC , let DCA cut the circle and DB fall on it and let the rectangle AD, DC be equal to the square on DB



I say that DB touches the circle ABC

For let DF be drawn touching ABC , let the centre of the circle ABC be taken, and let it be F let FE, FB, FD be joined

Thus the angle FED is right

[III 18]

Now since DE touches the circle ABC and DCA cuts it the rectangle AD, DC is equal to the square on DE

[III 36]

But the rectangle AD, DC was also equal to the square on DB

therefore the square on DE is equal to the square on DB

therefore DE is equal to DB

And FE is equal to FB ,

therefore the two sides DE, EF are equal to the two sides DB, BF ,

and FD is the common base of the triangles,

therefore the angle DEF is equal to the angle DBF

[I 8]

But the angle DEF is right

therefore the angle DBF is also right

And FB produced is a diameter

and the straight line drawn at right angles to the diameter of a circle from its extremity touches the circle

[III 16, Por]

therefore DB touches the circle

Similarly this can be proved to be the case even if the centre be on AC

Therefore etc

Q E D

BOOK FOUR

DEFINITIONS

1 A rectilineal figure is said to be *inscribed in a rectilineal figure* when the respective angles of the inscribed figure lie on the respective sides of that in which it is inscribed

2 Similarly a figure is said to be *circumscribed about a figure* when the respective sides of the circumscribed figure pass through the respective angles of that about which it is circumscribed

3 A rectilineal figure is said to be *inscribed in a circle* when each angle of the inscribed figure lies on the circumference of the circle

4 A rectilineal figure is said to be *circumscribed about a circle*, when each side of the circumscribed figure touches the circumference of the circle

5 Similarly a circle is said to be *inscribed in a figure* when the circumference of the circle touches each side of the figure in which it is inscribed

6 A circle is said to be *circumscribed about a figure* when the circumference of the circle passes through each angle of the figure about which it is circumscribed

7 A straight line is said to be *fitted into a circle* when its extremities are on the circumference of the circle

BOOK IV PROPOSITIONS

PROPOSITION I

Into a given circle to fit a straight line equal to a given straight line which is not greater than the diameter of the circle

Let ABC be the given circle, and D the given straight line not greater than the diameter of the circle,


thus it is required to fit into the circle ABC a straight line equal to the straight line D

Let a diameter BC of the circle ABC be drawn

Then if BC is equal to D that which was enjoined will have been done for BC has been fitted into the circle ABC equal to the straight line D

But if BC is greater than D
let CE be made equal to D , and with centre C and distance CE let the circle EAF be described
let CA be joined

Then since the point C is the centre of the circle EAF ,



CA is equal to CE

But CE is equal to D ,

therefore D is also equal to CA

Therefore into the given circle ABC there has been fitted CA equal to the given straight line D $Q\ E\ F$

PROPOSITION 2

In a given circle to inscribe a triangle equiangular with a given triangle

Let ABC be the given circle, and DEF the given triangle, thus it is required to inscribe in the circle ABC a triangle equiangular with the triangle DEF

Let GH be drawn touching the circle ABC at A [III 16 Por], on the straight line AH , and at the point A on it let the angle HAC be constructed equal to the angle DEF ,

and on the straight line AG and at the point A on it, let the angle GAB be constructed equal to the angle DFE , [I 23]

let BC be joined

Then since a straight line AH touches the circle ABC , and from the point of contact at A the straight line AC is drawn across in the circle,

therefore the angle HAC is equal to the angle ABC in the alternate segment of the circle [III 32]

But the angle HAC is equal to the angle DEF ,

therefore the angle ABC is also equal to the angle DEF

For the same reason

the angle ACB is also equal to the angle DFE ,

therefore the remaining angle BAC is also equal to the remaining angle EDF [I 32]

Therefore in the given circle there has been inscribed a triangle equiangular with the given triangle $Q\ E\ F$

PROPOSITION 3



About a given circle to circumscribe a triangle equiangular with a given triangle

Let ABC be the given circle and DEF the given triangle, thus it is required to circumscribe about the circle ABC a triangle equiangular with the triangle DEF

Let EF be produced in both directions to the points $G\ H$

let the centre K of the circle ABC be taken [III 1] and let the straight line KB be drawn across at random,

on the straight line KB and at the point K on it, let the angle BKA be constructed equal to the angle DEG , and the angle BKC equal to the angle DHF , [I 23]



and through the points A, B, C let LAM, MBN, NCL be drawn touching the circle ABC [III 16, Por.]

Now since LM, MN, NL touch the circle ABC at the points A, B, C , and LA, KB, AC have been joined from the centre K to the points A, B, C , therefore the angles at the points A, B, C are right [III 18]

And since the four angles of the quadrilateral $AMBA$ are equal to four right angles,asmuch as $AMBA$ is in fact divisible into two triangles, and the angles KAM, KBM are right

therefore the remaining angles KAB, KMB are equal to two right angles

But the angles DEG, DEF are also equal to two right angles [I 13]

therefore the angles KAB, AMB are equal to the angles DEG, DEF ,

of which the angle KAB is equal to the angle DEG ,

therefore the angle AMB which remains is equal to the angle DEF which remains

Similarly it can be proved that the angle LNB is also equal to the angle DFE ,

therefore the remaining angle MLN is equal to the angle EDF [I 32]

Therefore the triangle LMN is equiangular with the triangle DEF , and it has been circumscribed about the circle ABC

Therefore about a given circle there has been circumscribed a triangle equiangular with the given triangle Q E F

PROPOSITION 4

In a given triangle to inscribe a circle

Let ABC be the given triangle

thus it is required to inscribe a circle in the triangle ABC

Let the angles ABC, ACB be bisected by the straight lines BD, CD [I 9] and let these meet one another at the point D , from D let DE, DF, DG be drawn perpendicular to the straight lines AB, BC, CA

Now since the angle ABD is equal to the angle CBD

and the right angle BED is also equal to the right angle BFD ,

EBD, FBD are two triangles having two angles equal to two angles and one side equal to one side namely that subtending one of the equal angles which is BD common to the triangles

therefore they will also have the remaining sides equal to the remaining sides [I 26]


therefore DE is equal to DF

For the same reason

DG is also equal to DF

Therefore the three straight lines DE, DF, DG are equal to one another, therefore the circle described with centre D and distance one of the straight lines DE, DF, DG will pass also through the remaining points and will touch the straight lines AB, BC, CA because the angles at the points E, F, G are right

For, if it cuts them the straight line drawn at right angles to the d



the circle from its extremity will be found to fall within the circle which was proved absurd, [iii 16]

therefore the circle described with centre D and distance one of the straight lines DE DF , DG will not cut the straight lines AB , BC CA , therefore it will touch them and will be the circle inscribed in the triangle ABC [iv Def 5]

Let it be inscribed as IGI

Therefore in the given triangle ABC the circle EFG has been inscribed




Q E F

PROPOSITION 5

About a given triangle to circumscribe a circle

Let ABC be the given triangle

thus it is required to circumscribe a circle about the given triangle ABC



Let the straight lines AB AC be bisected at the points D , E [i 10], and from the points D E let DF EF be drawn at right angles to AB , AC , they will then meet within the triangle ABC , or on the straight line BC , or outside BC

First let them meet within at F , and let FB , FC FA be joined

Then, since AD is equal to DB ,

and DF is common and at right angles,

therefore the base AF is equal to the base FB

[i 4]

Similarly we can prove that

CF is also equal to AF ,

so that FB is also equal to FC

therefore the three straight lines FA FB FC are equal to one another

Therefore the circle described with centre F and distance one of the straight lines FA FB FC will pass also through the remaining points and the circle will have been circumscribed about the triangle ABC

Let it be circumscribed as ABC

Next let DF EF meet on the straight line BC at F , as is the case in the second figure, and let AF be joined

Then similarly we shall prove that the point F is the centre of the circle circumscribed about the triangle ABC

Again let DF , EF meet outside the triangle ABC at F , as is the case in the third figure and let AF BF CF be joined

Then again since AD is equal to DB ,

and DF is common and at right angles,

therefore the base AF is equal to the base BF

[i 4]

Similarly we can prove that

CF is also equal to AF

so that BF is also equal to FC ,

therefore the circle described with centre F and distance one of the straight lines FA, FB, FC will pass also through the remaining points and will have been circumscribed about the triangle ABC

Therefore about the given triangle a circle has been circumscribed

Q E F


And it is manifest that, when the centre of the circle falls within the triangle the angle BAC , being in a segment greater than the semicircle is less than a right angle,
when the centre falls on the straight line BC , the angle BAC , being in a semicircle is right,
and when the centre of the circle falls outside the triangle, the angle BAC , being in a segment less than the semicircle, is greater than a right angle [III 31]

PROPOSITION 6

In a given circle to inscribe a square

Let $ABCD$ be the given circle,

thus it is required to inscribe a square in the circle $ABCD$



Let two diameters AC, BD of the circle $ABCD$ be drawn at right angles to one another, and let AB, BC, CD, DA be joined

Then since BE is equal to ED for E is the centre and EA is common and at right angles
therefore the base AB is equal to the base AD [I 4]

For the same reason
each of the straight lines BC, CD is also equal to each of the straight lines AB, AD ,

therefore the quadrilateral $ABCD$ is equilateral

I say next that it is also right-angled

For, since the straight line BD is a diameter of the circle $ABCD$

therefore BAD is a semicircle

therefore the angle BAD is right

[III 31]

For the same reason

each of the angles ABC, BCD, CDA is also right,

therefore the quadrilateral $ABCD$ is right angled

But it was also proved equilateral

therefore it is a square

[I Def 22]

and it has been inscribed in the circle $ABCD$

Therefore in the given circle the square $ABCD$ has been inscribed Q E F

PROPOSITION 7

About a given circle to circumscribe a square

Let $ABCD$ be the given circle

thus it is required to circumscribe a square about the circle $ABCD$

Let two diameters AC, BD of the circle $ABCD$ be drawn at right angles to one another and through the points A, B, C, D let FG, GH, HA, KF be drawn touching the circle $ABCD$ [III 16 Por]

Then since FG touches the circle $ABCD$

and EA has been joined from the centre E to the point of contact at A

therefore the angles at A are right

[III 18]

For the same reason

the angles at the points B C D are also right

Now, since the angle AEB is right

and the angle EBG is also right

therefore GH is parallel to AC [1 28]

For the same reason

AC is also parallel to FK ,

so that GH is also parallel to FK [1 30]

Similarly we can prove that

each of the straight lines GF HA is parallel to BED

Therefore GA , GC AK , FB , BK are parallelograms,

therefore GF is equal to HA , and GH to FK [1 34]

And since AC is equal to BD

and AC is also equal to each of the straight lines GH FK ,

while BD is equal to each of the straight lines GF , HA , [1 34]

therefore the quadrilateral $FGHA$ is equilateral

I say next that it is also right-angled

For, since $GBEA$ is a parallelogram

and the angle AEB is right,

therefore the angle AGB is also right [1 34]

Similarly we can prove that

the angles at H , A , F are also right


Therefore $FGHA$ is right angled

But it was also proved equilateral,

therefore it is a square

and it has been circumscribed about the circle $ABCD$

Therefore about the given circle a square has been circumscribed $Q.E.D.$



PROPOSITION 8

In a given square to inscribe a circle

Let $ABCD$ be the given square

thus it is required to inscribe a circle in the given square $ABCD$

Let the straight lines AD , AB be bisected at the points E , F respectively [1 10]

through E let EH be drawn parallel to either AB or CD , and through F let FK be drawn parallel to either AD or BC , [1 31]

therefore each of the figures AK KB AH HD AG GC BG GD is a parallelogram and their opposite sides are evidently equal [1 34]

Now since AD is equal to AB

and AE is half of AD , and AF half of AB ,


therefore AE is equal to AF ,

so that the opposite sides are also equal

therefore FG is equal to GE

Similarly we can prove that each of the straight lines GH GA is equal to each of the straight lines FG GE

therefore the four straight lines GE GF GH GA are equal to one another



Therefore the circle described with centre G and distance one of the straight lines GE, GF, GH, GK will pass also through the remaining points

And it will touch the straight lines AB, BC, CD, DA , because the angles at E, F, H, K are right

For, if the circle cuts AB, BC, CD, DA , the straight line drawn at right angles to the diameter of the circle from its extremity will fall within the circle which was proved absurd, [III 16]

therefore the circle described with centre G and distance one of the straight lines GE, GF, GH, GK will not cut the straight lines AB, BC, CD, DA

Therefore it will touch them and will have been inscribed in the square $ABCD$

Therefore in the given square a circle has been inscribed

Q E F

PROPOSITION 9

About a given square to circumscribe a circle

Let $ABCD$ be the given square,

thus it is required to circumscribe a circle about the square $ABCD$

For let AC, BD be joined and let them cut one another at E

Then, since DA is equal to AB and AC is common

therefore the two sides DA, AC are equal to the two sides BA, AC ,


and the base DC is equal to the base BC ,

therefore the angle DAC is equal to the angle BAC

[I 8]

Therefore the angle DAB is bisected by AC

Similarly we can prove that each of the angles ABC, BCD, CDA is bisected by the straight lines AC, DB



Now, since the angle DAB is equal to the angle ABC

and the angle EAB is half the angle DAB ,

and the angle EBA half the angle ABC ,

therefore the angle EAB is also equal to the angle EBA ,

so that the side EA is also equal to EB

[I 6]

Similarly we can prove that each of the straight lines EA, EB, EC, ED is equal to each of the straight lines EC, ED

Therefore the four straight lines EA, EB, EC, ED are equal to one another

Therefore the circle described with centre E and distance one of the straight lines EA, EB, EC, ED will pass also through the remaining points,

and it will have been circumscribed about the square $ABCD$

Let it be circumscribed as $ABCD$

Therefore about the given square a circle has been circumscribed Q E F

PROPOSITION 10

To construct an isosceles triangle having each of the angles at the base double of the remaining one

Let any straight line AB be set out and let it be cut at the point C so that the rectangle contained by AB, BC is equal to the square on CA [II 11]

with centre A and distance AB let the circle BDE be described and let there be fitted in the circle BDE the straight line BD equal to the straight line AC which is not greater than the diameter of the circle BDE

Let AD , DC be joined and let the circle ACD be circumscribed about the triangle ACD [IV 5]

Then, since the rectangle AB , BC is equal to the square on AC ,

and AC is equal to BD

therefore the rectangle AB , BC is equal to the square on BD

And since a point B has been taken outside the circle ACD

and from B the two straight lines BA , BD have fallen on the circle ACD , and one of them cuts it while the other falls on it

and the rectangle AB , BC is equal to the square on BD ,

therefore BD touches the circle ACD [III 37]

Since, then BD touches it and DC is drawn across from the point of contact at D

therefore the angle BDC is equal to the angle DAC in the alternate segment of the circle [III 32]

Since, then the angle BDC is equal to the angle DAC ,

let the angle CDA be added to each,

therefore the whole angle BDA is equal to the two angles CDA , DAC

But the exterior angle BCD is equal to the angles CDA , DAC [I 32]

therefore the angle BDA is also equal to the angle BCD

But the angle BDA is equal to the angle CBD since the side AD is also equal to AB [I 5]

so that the angle DBA is also equal to the angle BCD

Therefore the three angles BDA , DBA , BCD are equal to one another

And since the angle DBC is equal to the angle BCD

the side BD is also equal to the side DC [I 6]

But BD is by hypothesis equal to CA

therefore CA is also equal to CD

so that the angle CDA is also equal to the angle DAC , [I 5]

therefore the angles CDA , DAC are double of the angle DAC

But the angle BCD is equal to the angles CDA , DAC

therefore the angle BCD is also double of the angle CAD

But the angle BCD is equal to each of the angles BDA , DBA ,

therefore each of the angles BDA , DBA is also double of the angle DAB

Therefore the isosceles triangle ABD has been constructed having each of the angles at the base DB double of the remaining one Q.E.F.

PROPOSITION 11


In a given circle to inscribe an equilateral and equiangular pentagon

Let $ABCDE$ be the given circle

thus it is required to inscribe in the circle $ABCDE$ an equilateral and equiangular pentagon

Let the isosceles triangle FGH be set out having each of the angles at G , H double of the angle at F [IV 10]

let there be inscribed in the circle $ABCDE$ the triangle ACD equiangular with



the triangle FGH , so that the angle CAD is equal to the angle at F and the angles at G, H respectively equal to the angles ACD, CDA , [iv 2]
therefore each of the angles ACD, CDA is also double of the angle CAD



Now let the angles ACD, CDA be bisected respectively by the straight lines CE, DB [i 9] and let AB, BC, DE, EA be joined

Then since each of the angles ACD, CDA is double of the angle CAD and they have been bisected by the straight lines CE, DB ,

therefore the five angles DAC, ACE, ECD, CDB, BDA are equal to one another

But equal angles stand on equal circumferences [iii 26]
therefore the five circumferences AB, BC, CD, DE, EA are equal to one another

But equal circumferences are subtended by equal straight lines [iii 29]
therefore the five straight lines AB, BC, CD, DE, EA are equal to one another
therefore the pentagon $ABCDE$ is equilateral

I say next that it is also equiangular

For since the circumference AB is equal to the circumference DE , let BCD be added to each
therefore the whole circumference $ABCD$ is equal to the whole circumference $EDCB$

And the angle AED stands on the circumference $ABCD$ and the angle BAE on the circumference $EDCB$,

therefore the angle BAE is also equal to the angle AED [iii 27]

For the same reason
each of the angles ABC, BCD, CDE is also equal to each of the angles BAE, AED

therefore the pentagon $ABCDE$ is equiangular

But it was also proved equilateral
therefore in the given circle an equilateral and equiangular pentagon has been inscribed Q E F

PROPOSITION 12

About a given circle to circumscribe an equilateral and equiangular pentagon

Let $ABCDE$ be the given circle
thus it is required to circumscribe an equilateral and equiangular pentagon about the circle $ABCDE$

Let A, B, C, D, E be conceived to be the angular points of the inscribed pentagon so that the circumferences AB, BC, CD, DE, EA are equal [iv 11]
through A, B, C, D, E let GH, HK, KL, LM, MG be drawn touching the circle, [iii 16 Por.]

let the centre F of the circle $ABCDE$ be taken [iii 1] and let FB, FK, FC, FL, FD be joined

Then since the straight line KL touches the circle $ABCDE$ at C
and FC has been joined from the centre F to the point of contact at C
therefore FC is perpendicular to KL , [iii 18]

therefore each of the angles at C is right

For the same reason

the angles at the points B, D are also right

And, since the angle FCA is right

therefore the square on FA is equal to the squares on FC, CA

For the same reason

the square on FA is also equal to the squares on FB, BA ,

so that the squares on FC, CA are equal to the squares on FB, BA ,

of which the square on FC is equal to the square on FB

therefore the square on CA which remains is equal to the square on BA

Therefore BA is equal to CA

And, since FB is equal to FC

and FA common

the two sides BF, FA are equal to the two sides CF, FA , and the base BA equal to the base CA ,

therefore the angle BFA is equal to the angle AFC ,

and the angle BAF to the angle FAC

Therefore the angle BFC is double of the angle AFC ,

and the angle BAC of the angle FAC

For the same reason

the angle CFD is also double of the angle CFL ,

and the angle DLC of the angle FLC

Now since the circumference BC is equal to CD

the angle BFC is also equal to the angle CFD

[III 27]

And the angle BFC is double of the angle AFC , and the angle DFC of the angle LFC

therefore the angle AFC is also equal to the angle LFC

But the angle FCA is also equal to the angle FCL ,
therefore FAC, FLC are two triangles having two angles equal to two angles and one side equal to one side, namely FC which is common to them,
therefore they will also have the remaining sides equal to the remaining sides and the remaining angle to the remaining angle,

[I 26]

therefore the straight line AC is equal to CL ,

and the angle FAC to the angle FLC ,

And, since AC is equal to CL

therefore AL is double of AC

For the same reason it can be proved that

HA is also double of BA

And BA is equal to AC ,


therefore HA is also equal to AL

Similarly each of the straight lines HG, GM, ML can also be proved equal to each of the straight lines HA, AL

therefore the pentagon $GHALM$ is equilateral

I say next that it is also equiangular

For, since the angle FAC is equal to the angle FLC ,



and the angle HKL was proved double of the angle FAC ,
 and the angle KLM double of the angle FLC
 therefore the angle HKL is also equal to the angle KLM

Similarly each of the angles KHG , HGM , GML can also be proved equal to each of the angles HKL , KLM ,
 therefore the five angles GHL , HKL , KLM , LMG , MGH are equal to one another

Therefore the pentagon $GHLKM$ is equiangular

And it was also proved equilateral, and it has been circumscribed about the circle $ABCDE$ Q E F

PROPOSITION 13

In a given pentagon, which is equilateral and equiangular, to inscribe a circle

Let $ABCDE$ be the given equilateral and equiangular pentagon,
 thus it is required to inscribe a circle in the pentagon $ABCDE$

For let the angles BCD , CDE be bisected by the straight lines CF , DF respectively, and from the point F at which the straight lines CF , DF meet one another, let the straight lines FB , FA , FE be joined

Then since BC is equal to CD and CF common,

the two sides BC , CF are equal to the two sides DC , CF ,

and the angle BCF is equal to the angle DCF ,

therefore the base BF is equal to the base DF

and the triangle BCF is equal to the triangle DCF ,
 and the remaining angles will be equal to the remaining angles namely those which the equal sides subtend [14]

Therefore the angle CBF is equal to the angle CDF

And since the angle CDE is double of the angle CDI ,

and the angle CDE is equal to the angle ABC ,

while the angle CDF is equal to the angle CBF

therefore the angle CBA is also double of the angle CBF ,

therefore the angle ABI is equal to the angle FBC

therefore the angle ABC has been bisected by the straight line BF

Similarly it can be proved that

the angles BAE , AED have also been bisected by the straight lines FA , FE respectively

Now let FG , FH , FK , FL , FM be drawn from the point F perpendicular to the straight lines AB , BC , CD , DE , EA

Then, since the angle HCF is equal to the angle KCF

and the right angle FHC is also equal to the angle FKC

FHC , FKC are two triangles having two angles equal to two angles and one side equal to one side, namely FC which is common to them and subtends one of the equal angles

therefore they will also have the remaining sides equal to the remaining sides [15]

therefore the perpendicular FH is equal to the perpendicular FK

Similarly it can be proved that

each of the straight lines FL FM FG is also equal to each of the straight lines FH FK ,

therefore the five straight lines FG FH , FK , FL FM are equal to one another

Therefore the circle described with centre F and distance one of the straight lines FG FH FK FL FM will pass also through the remaining points and it will touch the straight lines AB BC , CD , DE , EA , because the angles at the points G , H , K , L , M are right

For if it does not touch them, but cuts them, it will result that the straight line drawn at right angles to the diameter of the circle from its extremity falls within the circle which was proved absurd

[III 16]

Therefore the circle described with centre F and distance one of the straight lines FG FH , FK , FL , FM will not cut the straight lines AB , BC , CD , DE , EA , therefore it will touch them

Let it be described as $GHIKIM$

Therefore in the given pentagon which is equilateral and equiangular a circle has been inscribed

Q E F


PROPOSITION 14

About a given pentagon which is equilateral and equiangular, to circumscribe a circle

Let $ABCDE$ be the given pentagon which is equilateral and equiangular, thus it is required to circumscribe a circle about the pentagon $ABCDE$

Let the angles BCD CDE be bisected by the straight lines CF , DF respectively, and from the point F at which the straight lines meet let the straight lines FB FA FE be joined to the points B , A , E

Then in manner similar to the preceding it can be proved that the angles CBA BAE AED have also been bisected by the straight lines FB , FA FE respectively



Now, since the angle BCD is equal to the angle CDE ,
and the angle FCD is half of the angle BCD
and the angle CDF half of the angle CDE ,
therefore the angle FCD is also equal to the angle CDF ,
so that the side FC is also equal to the side FD

[I 6]

Similarly it can be proved that each of the straight lines FB FA , FE is also equal to each of the straight lines FC , FD

therefore the five straight lines FA FB FC FD FE are equal to one another

Therefore the circle described with centre F and distance one of the straight lines FA FB FC FD FE will pass also through the remaining points and will have been circumscribed

Let it be circumscribed and let it be $ABCDE$

Therefore about the given pentagon which is equilateral and equiangular, a circle has been circumscribed

Q E F

PROPOSITION 15

In a given circle to inscribe an equilateral and equiangular hexagon

Let $ABCDEF$ be the given circle,
thus it is required to inscribe an equilateral and equiangular hexagon in the circle $ABCDEF$

Let the diameter AD of the circle $ABCDEF$ be drawn,

let the centre G of the circle be taken and with centre D and distance DG let the circle $EGCH$ be described
let EG CG be joined and carried through to the points B , F ,

and let AB BC , CD DE , EF , FA be joined

I say that the hexagon $ABCDEF$ is equilateral and equiangular

For since the point G is the centre of the circle $ABCDEF$,

GE is equal to GD

Again, since the point D is the centre of the circle GCH ,

DF is equal to DG

But GE was proved equal to GD ,

therefore GH is also equal to ED ,

therefore the triangle EGD is equilateral

and therefore its three angles EGD , GDE DEG are equal to one another inasmuch as in isosceles triangles the angles at the base are equal to one another [I 5]

And the three angles of the triangle are equal to two right angles, [I 32]

therefore the angle EGD is one third of two right angles

Similarly the angle DGC can also be proved to be one-third of two right angles

And since the straight line CG standing on EB makes the adjacent angles EGC , CGB equal to two right angles

therefore the remaining angle CGB is also one-third of two right angles

Therefore the angles EGD DGC CGB are equal to one another
so that the angles vertical to them the angles BGA AGF FGE are equal [I 15]

Therefore the six angles EGD DGC CGB , BGA AGF , FGE are equal to one another

But equal angles stand on equal circumferences [III 26]
therefore the six circumferences AB BC CD DE EF FA are equal to one another

And equal circumferences are subtended by equal straight lines [III 29]

therefore the six straight lines are equal to one another

therefore the hexagon $ABCDEF$ is equilateral


I say next that it is also equiangular

For, since the circumference FA is equal to the circumference ED ,

let the circumference $ABCD$ be added to each

therefore the whole $FABCD$ is equal to the whole $EDCBA$

and the angle FED stands on the circumference $FABCD$



and the angle AFE on the circumference $EDCBA$,
therefore the angle AFE is equal to the angle DEF [III 27]

Similarly it can be proved that the remaining angles of the hexagon $ABCDEF$
are also severally equal to each of the angles AFE , FED ,
therefore the hexagon $ABCDEF$ is equiangular

But it was also proved equilateral
and it has been inscribed in the circle $ABCDEF$

Therefore in the given circle an equilateral and equiangular hexagon has
been inscribed Q E F

PROPOSITION 16 From this it is manifest that the side of the hexagon is equal to the
radius of the circle

And in like manner as in the case of the pentagon if through the points of
division on the circle we draw tangents to the circle, there will be circumscribed
about the circle an equilateral and equiangular hexagon in conformity with
what was explained in the case of the pentagon

And further by means similar to those explained in the case of the pentagon
we can both inscribe a circle in a given hexagon and circumscribe one about it

Q E F

PROPOSITION 16

In a given circle to inscribe a fifteen-angled figure which shall be both equilateral
and equiangular

Let $ABCD$ be the given circle,
thus it is required to inscribe in the circle $ABCD$ a fifteen angled figure which
shall be both equilateral and equiangular

In the circle $ABCD$ let there be in-
scribed a side AC of the equilateral tri-
angle inscribed in it and a side AB of an
equilateral pentagon
therefore of the equal segments of which
there are fifteen in the circle $ABCD$ there
will be five in the circumference ABC which
is one-third of the circle and there will be
three in the circumference AB which is
one-fifth of the circle


therefore in the remainder BC there will
be two of the equal segments

Let BC be bisected at E , [III 30]
therefore each of the circumferences BE EC is a fifteenth of the circle $ABCD$

If therefore we join BE EC and fit into the circle $ABCD$ straight lines equal
to them and in contiguity a fifteen angled figure which is both equilateral and
equiangular will have been inscribed in it Q E F

And, in like manner as in the case of the pentagon if through the points of
division on the circle we draw tangents to the circle there will be circumscribed
about the circle a fifteen angled figure which is equilateral and equi-
angular

And further by proofs similar to those in the case of the pentagon we can
both inscribe a circle in the given fifteen angled figure and circumscribe one
about it Q E F



BOOK FIVE

DEFINITIONS

1 A magnitude is a *part* of a magnitude, the less of the greater, when it measures the greater

2 The greater is a *multiple* of the less when it is measured by the less

3 A *ratio* is a sort of relation in respect of size between two magnitudes of the same kind

4 Magnitudes are said to *have a ratio* to one another which are capable, when multiplied, of exceeding one another

5 Magnitudes are said to be in the *same ratio*, the first to the second and the third to the fourth when if any equimultiples whatever be taken of the first and third, and any equimultiples whatever of the second and fourth the former equimultiples alike exceed are alike equal to, or alike fall short of, the latter equimultiples respectively taken in corresponding order

6 Let magnitudes which have the same ratio be called *proportional*

7 When of the equimultiples the multiple of the first magnitude exceeds the multiple of the second, but the multiple of the third does not exceed the multiple of the fourth, then the first is said to *have a greater ratio* to the second than the third has to the fourth

8 A proportion in three terms is the least possible

9 When three magnitudes are proportional the first is said to have to the third the *duplicate ratio* of that which it has to the second

10 When four magnitudes are <continuously> proportional, the first is said to have to the fourth the *triplicate ratio* of that which it has to the second, and so on continually whatever be the proportion

11 The term *corresponding magnitudes* is used of antecedents in relation to antecedents and of consequents in relation to consequents

12 *Alternate ratio* means taking the antecedent in relation to the antecedent and the consequent in relation to the consequent

13 *Inverse ratio* means taking the consequent as antecedent in relation to the antecedent as consequent

14 *Composition of a ratio* means taking the antecedent together with the consequent as one in relation to the consequent by itself

15 *Separation of a ratio* means taking the excess by which the antecedent exceeds the consequent in relation to the consequent by itself

16 *Conversion of a ratio* means taking the antecedent in relation to the excess by which the antecedent exceeds the consequent

17 A ratio *ex aequali* arises when there being several magnitudes and another set equal to them in multitude which taken two and two are in the same proportion as the first is to the last among the first magnitudes, so is the first to the last among the second magnitudes,

Or in other words, it means taking the extreme terms by virtue of the removal of the intermediate terms

18 A *perturbed proportion* arises when, there being three magnitudes and another set equal to them in multitude, as antecedent is to consequent among the first magnitudes so is antecedent to consequent among the second magnitudes while as the consequent is to a third among the first magnitudes so is a third to the antecedent among the second magnitudes


BOOK V PROPOSITIONS

PROPOSITION 1

If there be any number of magnitudes whatever which are respectively equimultiples of any magnitudes equal in multitude, then, whatever multiple one of the magnitudes is of one that multiple also will all be of all

Let any number of magnitudes whatever AB CD be respectively equimultiples of any magnitudes E F equal in multitude,

I say that whatever multiple AB is of E that multiple will AB , CD also be of E F



For since AB is the same multiple of E that CD is of F , as many magnitudes as there are in AB equal to E so many also are there in CD equal to F

Let AB be divided into the magnitudes AG , GB equal to E ,
and CD into CH HD equal to F

then the multitude of the magnitudes AG GB will be equal to the multitude of the magnitudes CH HD

Now since AG is equal to E , and CH to F

therefore AG is equal to E , and AG , CH to E F

For the same reason

GB is equal to F and GB HD to E F

therefore as many magnitudes as there are in AB equal to E so many also are there in AB CD equal to E F

therefore whatever multiple AB is of E , that multiple will AB CD also be of E F


Therefore etc

Q E D

PROPOSITION 2

If a first magnitude be the same multiple of a second that a third is of a fourth and a fifth also be the same multiple of the second that a sixth is of the fourth the sum of the first and fifth will also be the same multiple of the second that the sum of the third and sixth is of the fourth

Let a first magnitude AB be the same multiple of a second C , that a third DE is of a fourth F , and let a fifth BG also be the same multiple of the second C that a sixth EH is of the fourth F



I say that the sum of the first and fifth AG will be the same multiple of the

second, C , that the sum of the third and sixth, DH is of the fourth F

For since AB is the same multiple of C that DE is of F therefore, as many magnitudes as there are in AB equal to C , so many also are there in DE equal to F

For the same reason also

as many as there are in BG equal to C so many are there also in EH equal to F , therefore as many as there are in the whole AG equal to C , so many also are there in the whole DH equal to F

Therefore whatever multiple AG is of C , that multiple also is DH of F

Therefore the sum of the first and fifth AG is the same multiple of the second C that the sum of the third and sixth, DH is of the fourth, F

Therefore etc

Q E D

PROPOSITION 3

If a first magnitude be the same multiple of a second that a third is of a fourth, and if equimultiples be taken of the first and third then also ex aequali the magnitudes taken will be equimultiples respectively the one of the second and the other of the fourth

Let a first magnitude A be the same multiple of a second B that a third C is of a fourth D , and let equimultiples EF , GH be taken of A , C ,

I say that EF is the same multiple of B that GH is of D

For, since EF is the same multiple of A that GH is of C , therefore as many magnitudes as there are in EF equal to A so many also are there in GH equal to C

Let EF be divided into the magnitudes EK , KF equal to A and GH into the magnitudes GL , LH equal to C

then the multitude of the magnitudes EK , KF will be equal to the multitude of the magnitudes GL , LH

A —————

B ———

E ————— K ————— F

C ———

D ———

G ————— L ————— H

And since A is the same multiple of B that C is of D ,

while EK is equal to A and GL to C , therefore EK is the same multiple of B that GL is of D

For the same reason

KF is the same multiple of B that LH is of D

Since then a first magnitude EK is the same multiple of a second B that a third GL is of a fourth D

and a fifth KF is also the same multiple of the second B that a sixth LH is of the fourth D

therefore the sum of the first and fifth EF is also the same multiple of the second B that the sum of the third and sixth GH is of the fourth D [v 2]

Therefore etc

Q E D

PROPOSITION 4

If a first magnitude have to a second the same ratio as a third to a fourth any equimultiples whatever of the first and third will also have the same ratio to any equimultiples whatever of the second and fourth respectively taken in corresponding order

For let a first magnitude A have to a second B the same ratio as a third C to a fourth D , and let equimultiples E F be taken of A , C , and G , H other, chance, equimultiples of B , D ,

I say that as E is to G , so is F to H


For let equimultiples K , L be taken of E , F , and other, chance equimultiples M N of G , H

Since E is the same multiple of A that F is of C , and equimultiples K , L of E , F have been taken, therefore K is the same multiple of A that L is of C

[v 3]

For the same reason M is the same multiple of B that N is of D

And since as A is to B , so is C to D



and of A C equimultiples K L have been taken,
and of B D other chance equimultiples M N
therefore if K is in excess of M , L also is in excess of N ,
if it is equal equal, and if less less [v Def 5]
And K L are equimultiples of E , F ,
and M N other chance equimultiples of G , H ,
therefore as E is to G so is F to H [v Def 5]

Therefore etc


Q E D

PROPOSITION 5

If a magnitude be the same multiple of a magnitude that a part subtracted is of a part subtracted the remainder will also be the same multiple of the remainder that the whole is of the whole

For let the magnitude AB be the same multiple of the magnitude CD that the part AE subtracted is of the part CF subtracted,

I say that the remainder EB is also the same multiple of the remainder FD that the whole AB is of the whole CD



For whatever multiple AE is of CF let EB be made that multiple of CG

Then, since AE is the same multiple of CF that EB is of GC

therefore AE is the same multiple of CF that AB is of GF [v 1]

But by the assumption AE is the same multiple of CF that AB is of CD

Therefore AB is the same multiple of each of the magnitudes GF CD ,

therefore GF is equal to CD

Let CF be subtracted from each,

therefore the remainder GC is equal to the remainder FD

And, since AE is the same multiple of CF that EB is of GC ,

and GC is equal to DF

therefore AE is the same multiple of CF that EB is of FD

But, by hypothesis

AE is the same multiple of CF that AB is of CD ,

therefore EB is the same multiple of FD that AB is of CD

That is the remainder EB will be the same multiple of the remainder FD that the whole AB is of the whole CD


Therefore etc

Q E D

PROPOSITION 6

If two magnitudes be equimultiples of two magnitudes, and any magnitudes subtracted from them be equimultiples of the same, the remainders also are either equal to the same or equimultiples of them

For let two magnitudes AB , CD be equimultiples of two magnitudes E , F , and let AG , CH subtracted from them be equimultiples of the same two E , F ,



I say that the remainders also GB , HD , are either equal to E , F or equimultiples of them

For, first let GB be equal to E ,

I say that HD is also equal to F

For let KA be made equal to F

Since AG is the same multiple of E that CH is of F ,

while GB is equal to E and KC to F ,

therefore AB is the same multiple of E that KH is of F [v 2]

But by hypothesis AB is the same multiple of E that CD is of F ,

therefore KH is the same multiple of F that CD is of F

Since then each of the magnitudes KH , CD is the same multiple of F ,

therefore KH is equal to CD

Let CH be subtracted from each,

therefore the remainder KC is equal to the remainder HD

But F is equal to KC ,

therefore HD is also equal to F

Hence if GB is equal to E HD is also equal to F

Similarly we can prove that, even if GB be a multiple of E , HD is also the same multiple of F

Therefore etc

Q E D

PROPOSITION 7

Equal magnitudes have to the same the same ratio as also has the same to equal magnitudes

Let A , B be equal magnitudes and C any other chance, magnitude,

I say that each of the magnitudes A , B has the same ratio to C , and C has the same ratio to each of the magnitudes A , B

For let equimultiples D , E of A , B be taken, and of C another, chance multiple F


Then since D is the same multiple of A that E is of B , while A is equal to B , therefore D is equal to E

But F is another, chance, magnitude

If therefore D is in excess of F , E is also in excess of F , if equal to it, equal, and if less less

And D , E are equimultiples of A , B , while F is another, chance multiple of C , therefore as A is to C , so is B to C

[v Def 5]



I say next that C also has the same ratio to each of the magnitudes A , B . For, with the same construction, we can prove similarly that D is equal to E , and F is some other magnitude

If therefore F is in excess of D it is also in excess of E if equal equal, and if less, less

And F is a multiple of C , while D , E are other chance equimultiples of A , B , therefore as C is to A , so is C to B [v Def 5]

Therefore etc

PORISM From this it is manifest that if any magnitudes are proportional they will also be proportional inversely

Q E D

PROPOSITION 8

Of unequal magnitudes the greater has to the same a greater ratio than the less has and the same has to the less a greater ratio than it has to the greater

Let AB , C be unequal magnitudes, and let AB be greater let D be another chance magnitude

I say that AB has to D a greater ratio than C has to D and D has to C a greater ratio than it has to AB

For since AB is greater than C let BE be made equal to C

then the less of the magnitudes AE , EB if multiplied will sometime be greater than D [v Def 4]

First let AE be less than EB let AE be multiplied and let

FG be a multiple of it which is greater than D ,

then whatever multiple FG is of AE let GH be made the same multiple of EB and K of C

and let L be taken double of D , M triple of it and successive multiples increasing by one until what is taken is a multiple of D and the first that is greater than K . Let it be taken and let it be N which is quadruple of D and the first multiple of it that is greater than K

Then since K is less than N first

therefore K is not less than M

And since FG is the same multiple of AE that GH is of EB


therefore FG is the same multiple of AE that FH is of AB [v 1]

But FG is the same multiple of AE that K is of C

therefore FH is the same multiple of AB that K is of C ,

therefore FH , K are equimultiples of AB , C

Again since GH is the same multiple of EB that K is of C and EB is equal to C



therefore GH is equal to K

But K is not less than M

therefore neither is GH less than M

And FG is greater than D ,

therefore the whole FH is greater than D M together

But D M together are equal to N inasmuch as M is triple of D , and M , D together are quadruple of D while N is also quadruple of D , whence M D together are equal to N

But FH is greater than M D ,

therefore FH is in excess of N

while K is not in excess of N

And FH , K are equimultiples of AB , C , while N is another chance multiple of D

therefore AB has to D a greater ratio than C has to D [v Def 7]

I say next that D also has to C a greater ratio than D has to AB

For with the same construction we can prove similarly that N is in excess of K while N is not in excess of FH

And N is a multiple of D ,

while FH K are other, chance equimultiples of AB C ,

therefore D has to C a greater ratio than D has to AB [v Def 7]

Again let AE be greater than EB

Then the less EB , if multiplied will sometime be greater than D [v Def 4]

Let it be multiplied and let GH be a multiple of EB and greater than D

and whatever multiple GH is of EB let FG be made the same multiple of AE and K of C

Then we can prove similarly that FH K are equimultiples of AB C

and similarly let N be taken a multiple of D but the first that is greater than FG , so that FG is again not less than M

But GH is greater than D

therefore the whole FH is in excess of D M that is of N

Now K is not in excess of N inasmuch as FG also which is greater than GH that is than K is not in excess of N

And in the same manner by following the above argument we complete the demonstration

Therefore etc

Q E D

PROPOSITION 9


Magnitudes which have the same ratio to the same are equal to one another and magnitudes to which the same has the same ratio are equal

For let each of the magnitudes A B have the same ratio to C

I say that A is equal to B

For otherwise each of the magnitudes A B would not have had the same ratio to C but it has [v 8]

therefore A is equal to B



Again, let C have the same ratio to each of the magnitudes A, B ,

I say that A is equal to B

For otherwise, C would not have had the same ratio to each of the magnitudes A, B , [v 5]

but it has,

therefore A is equal to B

Therefore etc


Q E D

PROPOSITION 10

Of magnitudes which have a ratio to the same, that which has a greater ratio is greater and that to which the same has a greater ratio is less

For let A have to C a greater ratio than B has to C ,

I say that A is greater than B



For if not A is either equal to B or less

Now A is not equal to B ,

for in that case each of the magnitudes A, B would have had the same ratio to C , [v 7]

but they have not,

therefore A is not equal to B

Nor again is A less than B

for in that case A would have had to C a less ratio than B has to C , [v 8]

but it has not

therefore A is not less than B

But it was proved not to be equal either

therefore A is greater than B

Again, let C have to B a greater ratio than C has to A ,

I say that B is less than A

For, if not it is either equal or greater

Now B is not equal to A

for in that case C would have had the same ratio to each of the magnitudes A, B , [v 7]

but it has not

therefore A is not equal to B

Nor again is B greater than A

for in that case C would have had to B a less ratio than it has to A , [v 8]

but it has not

therefore B is not greater than A

But it was proved that it is not equal either

therefore B is less than A

Therefore etc

Q E D

PROPOSITION 11

Ratios which are the same with the same ratio are also the same with one another

For, as A is to B so let C be to D

and as C is to D so let E be to F ,

I say that as A is to B , so is E to F

For of A, C, E let equmultiples G, H, K be taken and of B, D, F other,

A _____	C _____	E _____
B _____	D _____	F _____
G _____	H _____	K _____
L _____	M _____	N _____

chance, equmultiples L, M, N

Then since, as A is to B , so is C to D ,

and of A, C equmultiples G, H have been taken,

and of B, D other, chance equmultiples L, M

therefore, if G is in excess of L , H is also in excess of M ,

if equal equal,

and if less less

Again, since as C is to D , so is E to F ,

and of C, E equmultiples H, K have been taken

and of D, F other, chance equmultiples M, N

therefore if H is in excess of M , K is also in excess of N ,

if equal equal

and if less less

But we saw that if H was in excess of M , G was also in excess of L , if equal equal, and if less less

so that, in addition, if G is in excess of L , K is also in excess of N ,

if equal equal

and if less less

And G, K are equmultiples of A, E ,

while L, N are other, chance equmultiples of B, F ,

therefore as A is to B , so is E to F

Therefore etc

Q E D

PROPOSITION 12

If any number of magnitudes be proportional, as one of the antecedents is to one of the consequents, so will all the antecedents be to all the consequents

Let any number of magnitudes A, B, C, D, E, F be proportional, so that, as A is to B so is C to D and E to F ,

I say that as A is to B so are A, C, E to B, D, F

A _____	B _____	C _____
D _____	E _____	F _____
G _____	L _____	
H _____	M _____	
K _____	N _____	

For of A, C, E let equmultiples G, H, K be taken

and of B, D, F other, chance equmultiples L, M, N

Then since as A is to B , so is C to D , and E to F ,

and of A, C, E equmultiples G, H, K have been taken

and of B, D, F other, chance equmultiples L, M, N ,

therefore if G is in excess of L , H is also in excess of M and K of N ,

if equal equal,

and if less less,
 so that in addition
 if G is in excess of L then G, H, K are in excess of L, M, N ,
 if equal, equal
 and if less less

Now G and G, H, K are equmultiples of A and A, C, E since, if any number of magnitudes whatever are respectively equmultiples of any magnitudes equal in multitude whatever multiple one of the magnitudes is of one that multiple also will all be of all [v 1]

For the same reason

L and L, M, N are also equmultiples of B and B, D, F ,
 therefore as A is to B , so are A, C, E to B, D, F [v Def 5]

Therefore etc


Q E D

PROPOSITION 13

If a first magnitude have to a second the same ratio as a third to a fourth and the third have to the fourth a greater ratio than a fifth has to a sixth the first will also have to the second a greater ratio than the fifth to the sixth

For let a first magnitude A have to a second B the same ratio as a third C has to a fourth D and let the third C have to the fourth D a greater ratio than a fifth E has to a sixth F ,

I say that the first A will also have to the second B a greater ratio than the fifth E to the sixth F



For since there are some equmultiples of C, E ,
 and of D, F other chance, equmultiples such that the multiple of C is in excess of the multiple of D

while the multiple of E is not in excess of the multiple of F , [v Def 7]
 let them be taken

and let G, H be equmultiples of C, E ,

and K, L other chance equmultiples of D, F

so that G is in excess of K but H is not in excess of L

and whatever multiple G is of C let M be also that multiple of A ,

and, whatever multiple K is of D let N be also that multiple of B

Now, since as A is to B so is C to D ,

and of A, C equmultiples M, G have been taken

and of B, D other chance equmultiples N, K

therefore if M is in excess of N, G is also in excess of K ,

if equal equal

and if less less

[v Def 5]

But G is in excess of A therefore M is also in excess of N But H is not in excess of L and M, H are equimultiples of A, E ,and N, L other chance equimultiples of B, F ,therefore A has to B a greater ratio than E has to F [v Def 7]

Therefore etc

Q E D

PROPOSITION 14

If a first magnitude have to a second the same ratio as a third has to a fourth, and the first be greater than the third the second will also be greater than the fourth if equal equal and if less less

For let a first magnitude A have the same ratio to a second B as a third C has to a fourth D , and let A be greater than C ,

I say that B is also greater than D

A —————

C —————

B —————

D —————

For, since A is greater than C and B is another, chance magnitudetherefore A has to B a greater ratio than C has to B [v 8]But as A is to B so is C to D ,therefore C has also to D a greater ratio than C has to B [v 13]

But that to which the same has a greater ratio is less, [v 10]

therefore D is less than B so that B is greater than D Similarly we can prove that if A be equal to C B will also be equal to D and if A be less than C , B will also be less than D

Therefore etc

Q E D

PROPOSITION 15

Parts have the same ratio as the same multiples of them taken in corresponding order

For let AB be the same multiple of C that DE is of F I say that as C is to F so is AB to DE

A — G — H — B

C —————

C — K — L — E

F —————

For since AB is the same multiple of C that DE is of F as many magnitudes as there are in AB equal to C so many are there also in DE equal to F

Let AB be divided into the magnitudes AG, GH, HB equal to C and DE into the magnitudes DK, KL, LE equal to F then the multitude of the magnitudes AG, GH, HB will be equal to the multitude of the magnitudes DK, KL, LE And since AG, GH, HB are equal to one anotherand DK, KL, LE are also equal to one anothertherefore as AG is to DK so is GH to KL and HB to LE [v 7]

Therefore as one of the antecedents is to one of the consequents so will all the antecedents be to all the consequents [v 12]

therefore as AG is to DK so is AB to DE

But AG is equal to C and DA to F ,
therefore, as C is to F , so is AB to DE

Therefore etc

Q E D


PROPOSITION 16

If four magnitudes be proportional, they will also be proportional alternately

Let A, B, C, D be four proportional magnitudes

so that as A is to B , so is C to D ,

I say that they will also be so alternately, that is, as A is to C , so is B to D



For of A, B let equimultiples E, F be taken

and of C, D other, chance, equimultiples G, H

Then, since E is the same multiple of A that F is of B ,

and parts have the same ratio as the same multiples of them [v 15]

therefore as A is to B , so is E to F

But as A is to B so is C to D ,

therefore also as C is to D so is E to F [v 11]

Again, since G, H are equimultiples of C, D ,

therefore, as C is to D , so is G to H [v 15]

But as C is to D so is E to F ,

therefore also as F is to E so is G to H [v 11]

But if four magnitudes be proportional and the first be greater than the third,

the second will also be greater than the fourth,

if equal equal

and if less less

[v 14]

Therefore if F is in excess of G , F is also in excess of H ,

if equal equal,

and if less less

Now E, F are equimultiples of A, B

and G, H other, chance equimultiples of C, D ,

therefore as A is to C so is B to D

[v Def 5]

Therefore etc

Q E D

PROPOSITION 17

If magnitudes be proportional componendo they will also be proportional separando


Let AB, BE, CD, DF be magnitudes proportional componendo so that as AB is to BE so is CD to DF ,

I say that they will also be proportional separando, that is as AE is to EB , so is CF to DF

For of AE, EB, CF, FD let equimultiples GH, HK, LM, MN be taken,

and of EB, FD other chance equimultiples KO, NP

And, since GH is the same multiple of AE that HA is of EB ,
 therefore GH is the same multiple of AE that GA is of AB [v 1]
 But GH is the same multiple of AE that LM is of CF
 therefore GA is the same multiple of AB that LM is of CF



Again, since LM is the same multiple of CF that MN is of FD
 therefore LM is the same multiple of CF that LN is of CD [v 1]
 But LM was the same multiple of CF that GK is of AB
 therefore GK is the same multiple of AB that LN is of CD
 Therefore GK LN are equimultiples of AB , CD
 Again, since HK is the same multiple of EB that MN is of FD ,
 and KO is also the same multiple of EB that NP is of FD ,
 therefore the sum HO is also the same multiple of EB that MP is of FD [v 2]

And, since as AB is to BE so is CD to DF
 and of AB , CD equimultiples GK , LN have been taken
 and of EB , FD equimultiples HO MP
 therefore, if GA is in excess of HO , LN is also in excess of MP ,
 if equal, equal
 and if less, less

Let GH be in excess of KO ,
 then if HA be added to each,
 GA is also in excess of HO

But we saw that if GK was in excess of HO LN was also in excess of MP ,
 therefore LN is also in excess of MP
 and if MN be subtracted from each,
 LM is also in excess of NP

so that if GH is in excess of KO LM is also in excess of NP

Similarly we can prove that

if GH be equal to KO LM will also be equal to NP
 and if less less

And GH LM are equimultiples of AE CF

while KO NP are other chance equimultiples of EB , FD ,

therefore, as AL is to EB so is CF to FD

Therefore etc

Q E D

PROPOSITION 18


If magnitudes be proportional *separando* they will also be proportional *componendo*

Let AE EB CF FD be magnitudes proportional *separando*, so that as AE is to EB so is CF to FD

I say that they will also be proportional *componendo*, that is as AB is to BE so is CD to FD

For, if CD be not to DF as AB to BE ,
then, as AB is to BE , so will CD be either to some magnitude less than DF or
to a greater

First, let it be in that ratio to a less mag
nitude DG

Then, since as AB is to BE , so is CD to C  D
 DG ,

they are magnitudes proportional *componendo*,
so that they will also be proportional *separando* [v 17]

Therefore as AE is to EB so is CG to GD

But also by hypothesis

as AE is to EB , so is CF to FD

Therefore also as CG is to GD , so is CF to FD [v 11]

But the first CG is greater than the third CF ,

therefore the second GD is also greater than the fourth FD [v 14]

But it is also less which is impossible

Therefore as AB is to BE so is not CD to a less magnitude than FD

Similarly we can prove that neither is it in that ratio to a greater,
it is therefore in that ratio to FD itself

Therefore etc

Q E D

PROPOSITION 19

If as a whole is to a whole so is a part subtracted to a part subtracted the remainder
will also be to the remainder as whole to whole

For as the whole AB is to the whole CD , so let the part AE subtracted be to
the part CF subtracted,

I say that the remainder EB will also be to the remainder FD as the whole AB to the whole CD

For since as AB is to CD , so is AE to CF ,

alternately also as BA is to AE , so is DC to CF [v 16]

And since the magnitudes are proportional *componendo* they will also be
proportional *separando*, [v 17]

that is as BE is to EA so is DF to CF

and alternately

as BE is to DF so is EA to FC [v 16]

But as AE is to CF so by hypothesis is the whole AB to the whole CD

Therefore also the remainder EB will be to the remainder FD as the whole
 AB is to the whole CD [v 11]

Therefore etc

[PORISM From this it is manifest that if magnitudes be proportional *componendo* they will also be proportional *convertendo*]

Q E D

PROPOSITION 20

If there be three magnitudes and others equal to them in multitude which taken
two and two are in the same ratio and if ex aequali the first be greater than the
third the fourth will also be greater than the sixth if equal equal and if less less

Let there be three magnitudes A B C and others D E , F equal to them in
multitude which taken two and two are in the same ratio so that,

as A is to B so is D to E

as B is to C so is E to F

and let A be greater than C *ex aequali*,

I say that D will also be greater than F , if A is equal to C , equal, and if less, less

A —————

D —————

B ———

E ———

C —————

F —————

For since A is greater than C ,
and B is some other magnitude
and the greater has to the same a greater
ratio than the less has [v 8]
therefore A has to B a greater ratio
than C has to B

But as A is to B , so is D to E ,

and as C is to B inversely so is F to E ,

therefore D has also to E a greater ratio than F has to E [v 13]

But of magnitudes which have a ratio to the same, that which has a greater ratio is greater, [v 10]

therefore D is greater than F

Similarly we can prove that, if A be equal to C , D will also be equal to F , and if less less

Therefore etc

Q E D

PROPOSITION 21

If there be three magnitudes and others equal to them in multitude which taken two and two together are in the same ratio and the proportion of them be perturbed then if *ex aequali* the first magnitude is greater than the third the fourth will also be greater than the sixth if equal equal and if less less

Let there be three magnitudes A, B, C and others D, E, F equal to them in multitude which taken two and two are in the same ratio and let the proportion of them be perturbed so that,

as A is to B so is E to F

and

as B is to C so is D to E ,

and let A be greater than C *ex aequali*,

I say that D will also be greater than F if A is equal to C equal, and if less less

A —————

D —————

B ———

E —————

C —————

F —————

For since A is greater than C
and B is some other magnitude,
therefore A has to B a greater
ratio than C has to B [v 8]

But as A is to B so is E to F

and as C is to B inversely so is E to D

Therefore also E has to F a greater ratio than E has to D [v 13]

But that to which the same has a greater ratio is less [v 10]

therefore F is less than D

therefore D is greater than F

Similarly we can prove that if A be equal to C D will also be equal to F and if less less

Therefore etc

Q E D

PROPOSITION 22

If there be any number of magnitudes whatever, and others equal to them in multitude which taken two and two together are in the same ratio they will also be in the same ratio *ex aequali*

Let there be any number of magnitudes A, B, C , and others D, E, F equal to them in multitude which taken two and two together are in the same ratio, so that,

as A is to B so is D to E

and, as B is to C so is E to F ,


I say that they will all be in the same ratio *ex aequali*,

<that is as A is to C so is D to F >

For of A, D let equimultiples G, H be taken

and of B, E other chance, equimultiples K, L ,

and, further of C, F other chance, equimultiples M, N



Then, since as A is to B so is D to E ,

and of A, D equimultiples G, H have been taken,

and of B, E other chance equimultiples K, L ,

therefore as G is to K , so is H to L

[v. 4]

For the same reason also

as K is to M so is L to N

Since then there are three magnitudes G, K, M and others H, L, N equal to them in multitude, which taken two and two together are in the same ratio,

therefore *ex aequali* if G is in excess of M , H is also in excess of N

if equal, equal and if less less

[v. 20]

And G, H are equimultiples of A, D

and M, N other chance equimultiples of C, F

Therefore as A is to C so is D to F

[v. Def 5]

Therefore etc

Q E D

PROPOSITION 23

If there be three magnitudes and others equal to them in multitude which taken two and two together are in the same ratio and the proportion of them be perturbed they will also be in the same ratio *ex aequali*

Let there be three magnitudes A, B, C , and others equal to them in multitude, which taken two and two together are in the same proportion, namely D, E, F and let the proportion of them be perturbed so that,

as A is to B so is F to E

and, as B is to C so is D to E

I say that as A is to C so is D to F


Of A, B, D let equimultiples G, H, K be taken

and of C, E, F other chance equimultiples L, M, N

Then since G, H are equimultiples of A, B

and parts have the same ratio as the same multiples of them, [v 15]
therefore, as A is to B , so is G to H

For the same reason also



as E is to F , so is M to N

And as A is to B , so is E to F ,
therefore also, as G is to H , so is M to N [v 11]

Next, since as B is to C , so is D to E ,
alternately, also, as B is to D , so is C to E [v 16]

And since H K are equimultiples of B , D
and parts have the same ratio as their equimultiples
therefore as B is to D , so is H to K [v 15]

But as B is to D so is C to E ,
therefore also as H is to K , so is C to E [v 11]

Again, since L , M are equimultiples of C , E ,
therefore as C is to E , so is L to M [v 15]

But, as C is to E so is H to K ,
therefore also as H is to K , so is L to M [v 11]

and alternately as H is to L so is K to M [v 16]

But it was also proved that,

as G is to H , so is M to N

Since then there are three magnitudes G H L and others equal to them in multitude K , M , N which taken two and two together are in the same ratio, and the proportion of them is perturbed,

therefore, *ex aequali*, if G is in excess of L , K is also in excess of N ,
if equal equal, and if less, less [v 21]

And G , K are equimultiples of A D
and L N of C , F

Therefore as A is to C , so is D to F

Therefore etc

Q E D

PROPOSITION 24

If a first magnitude have to a second the same ratio as a third has to a fourth and also a fifth have to the second the same ratio as a sixth to the fourth the first and fifth added together will have to the second the same ratio as the third and sixth have to the fourth

Let a first magnitude AB have to a second C the same ratio as a third DE has to a fourth F

A ————— B ————— G

C —————

D ————— E ————— H

F —————

and let also a fifth BG have to the second C the same ratio as a sixth EH has to the fourth F ,

I say that the first and fifth added together, AG will have to the second C the same ratio

as the third and sixth DH has to the fourth F

For since as BG is to C , so is EH to F

inversely as C is to BG , so is F to EH

Since then, as AB is to C , so is DE to F ,

and as C is to BG so is F to EH ,

therefore *ex aequali* as AB is to BG , so is DE to EH [v 22]

And, since the magnitudes are proportional *separando*, they will also be proportional *componendo*, [v 18]

therefore, as AG is to GB , so is DH to HE

But also, as BG is to C , so is EH to F ,

therefore, *ex aequali* as AG is to C , so is DH to F [v 22]

Therefore etc

Q E D


PROPOSITION 25

If four magnitudes be proportional the greatest and the least are greater than the remaining two

Let the four magnitudes AB , CD , E , F be proportional so that as AB is to CD , so is E to F and let AB be the greatest of them and F the least

I say that AB , F are greater than CD , E

For let AG be made equal to E , and CH equal to F



Since as AB is to CD , so is E to F ,

and E is equal to AG and F to CH ,

therefore as AB is to CD so is AG to CH

And since as the whole AB is to the whole CD so is the part AG subtracted to the part CH subtracted

the remainder GB will also be to the remainder HD as the whole AB is to the whole CD [v 19]

But AB is greater than CD

therefore GB is also greater than HD

And since AG is equal to E and CH to F ,

therefore AG , F are equal to CH , E

And if GB , HD being unequal and GB greater AG , F be added to GB and CH , E be added to HD ,

it follows that AB , F are greater than CD , E

Therefore etc

Q E D

BOOK SIX

DEFINITIONS

1 *Similar rectilineal figures* are such as have their angles severally equal and the sides about the equal angles proportional

2 A straight line is said to have been *cut in extreme and mean ratio* when, as the whole line is to the greater segment so is the greater to the less

3 The *height* of any figure is the perpendicular drawn from the vertex to the base

BOOK VI PROPOSITIONS

PROPOSITION 1

Triangles and parallelograms which are under the same height are to one another as their bases

Let ABC ACD be triangles and EC CF parallelograms under the same height,

I say that as the base BC is to the base CD so is the triangle ABC to the triangle ACD and the parallelogram EC to the parallelogram CF

For let BD be produced in both directions to the points H , L and let [any number of straight lines] BG , GH be made equal to the base BC , and any number of straight lines DK , KL equal to the base CD ,

let AG , AH , AK , AL be joined

Then since CB , BG , GH are equal to one another

the triangles ABC AGB AHG are also equal to one another [I 38]

Therefore whatever multiple the base HC is of the base BC that multiple also is the triangle AHC of the triangle ABC

For the same reason


whatever multiple the base LC is of the base CD that multiple also is the triangle AIC of the triangle ACD

and if the base HC is equal to the base CL the triangle AHC is also equal to the triangle ACL

if the base HC is in excess of the base CL the triangle AHC is also in excess of the triangle ACL

and if less less

Thus there being four magnitudes two bases BC CD and two triangles ABC ACD



equimultiples have been taken of the base BC and the triangle ABC , namely the base HC and the triangle AHC and of the base CD and the triangle ADC other chance equimultiples namely the base LC and the triangle ALC ,

and it has been proved that

if the base HC is in excess of the base CL , the triangle AHC is also in excess of the triangle ALC ,

if equal equal, and, if less less

Therefore as the base BC is to the base CD , so is the triangle ABC to the triangle ACD [v Def 5]

Next, since the parallelogram EC is double of the triangle ABC , [i 41] and the parallelogram FC is double of the triangle ACD

while parts have the same ratio as the same multiples of them [v 15] therefore as the triangle ABC is to the triangle ACD so is the parallelogram EC to the parallelogram FC

Since then it was proved that as the base BC is to CD so is the triangle ABC to the triangle ACD

and as the triangle ABC is to the triangle ACD so is the parallelogram EC to the parallelogram FC

therefore also as the base BC is to the base CD so is the parallelogram EC to the parallelogram FC [v 11]

Therefore etc

Q E D

PROPOSITION 2

If a straight line be drawn parallel to one of the sides of a triangle it will cut the sides of the triangle proportionally and if the sides of the triangle be cut proportionally the line joining the points of section will be parallel to the remaining side of the triangle

For let DE be drawn parallel to BC one of the sides of the triangle ABC ,

I say that as BD is to DA so is CE to EA

For let BE CD be joined

Therefore the triangle BDE is equal to the triangle CDE

for they are on the same base DE and in the same parallels DE BC [i 38]

And the triangle ADE is another area

But equals have the same ratio to the same [v 7]

therefore as the triangle BDE is to the triangle ADE so is the triangle CDE to the triangle ADE

But as the triangle BDE is to ADE so is BD to DA

for being under the same height the perpendicular drawn from E to AB they are to one another as their bases [vi 1]


For the same reason also

as the triangle CDE is to ADE so is CE to EA

Therefore also as BD is to DA so is CE to EA [v 11]

Again let the sides AB AC of the triangle ABC be cut proportionally so that as BD is to DA so is CE to EA and let DE be joined

I say that DE is parallel to BC



For, with the same construction

since as BD is to DA so is CE to EA

but, as BD is to DA , so is the triangle BDE to the triangle ADE

and as CE is to EA , so is the triangle CDE to the triangle ADE , [vi 1]

therefore also,

as the triangle BDE is to the triangle ADE , so is the triangle CDE to the triangle ADE [v 11]

Therefore each of the triangles BDE CDE has the same ratio to ADE

Therefore the triangle BDE is equal to the triangle CDE , [v 9]

and they are on the same base DF

But equal triangles which are on the same base are also in the same parallels [i 39]

Therefore DE is parallel to BC


Therefore etc

Q E D

PROPOSITION 3

If an angle of a triangle be bisected and the straight line cutting the angle cut the base also the segments of the base will have the same ratio as the remaining sides of the triangle and, if the segments of the base have the same ratio as the remaining sides of the triangle the straight line joined from the vertex to the point of section will bisect the angle of the triangle

Let ABC be a triangle and let the angle BAC be bisected by the straight line AD ,



I say that as BD is to CD so is BA to AC

For let CE be drawn through C parallel to DA , and let BA be carried through and meet it at E

Then since the straight line AC falls upon the parallels AD EC

the angle ACE is equal to the angle CAD

[i 29]

But the angle CAD is by hypothesis equal to the angle BAD ,

therefore the angle BAD is also equal to the angle ACE

Again since the straight line BAE falls upon the parallels AD EC ,

the exterior angle BAD is equal to the interior angle AEC [i 29]

But the angle ACE was also proved equal to the angle BAD

therefore the angle ACE is also equal to the angle AEC ,

so that the side AE is also equal to the side AC

[i 6]

And since AD has been drawn parallel to EC one of the sides of the triangle BCF

therefore proportionally as BD is to DC , so is BA to AE

But AE is equal to AC ,

[vi 2]

therefore as BD is to DC so is BA to AC

Again let BA be to AC as BD to DC , and let AD be joined,

I say that the angle BAC has been bisected by the straight line AD

For with the same construction,

since as BD is to DC , so is BA to AC

and also as BD is to DC so is BA to AE for AD has been drawn parallel to

EC, one of the sides of the triangle *BCE*

therefore also, as *BA* is to *AC*, so is *BA* to *AE*

{vi 2}

{v 11}

{v 9}

{i 5}

{i 29}

{id}

Therefore *AC* is equal to *AE*

so that the angle *AEC* is also equal to the angle *ACE*

But the angle *AEC* is equal to the exterior angle *BAD*

and the angle *ACE* is equal to the alternate angle *CAD*,

therefore the angle *BAD* is also equal to the angle *CAD*

Therefore the angle *BAC* has been bisected by the straight line *AD*

Therefore etc

Q E D

PROPOSITION 4

In equiangular triangles the sides about the equal angles are proportional, and those are corresponding sides which subtend the equal angles

Let *ABC*, *DCE* be equiangular triangles having the angle *ABC* equal to the angle *DCE* the angle *BAC* to the angle *CDE* and further the angle *ACB* to the angle *CED*,

I say that in the triangles *ABC*, *DCE* the sides about the equal angles are proportional, and those are corresponding sides which subtend the equal angles

For let *BC* be placed in a straight line with *CE*

Then since the angles *ABC* *ACB* are less than two right angles [i 17]

and the angle *ACB* is equal to the angle *DEC*,

therefore the angles *ABC* *DEC* are less than two right angles

therefore *BA*, *ED* when produced, will meet [i Post 5]

Let them be produced and meet at *F*

Now since the angle *DCE* is equal to the angle *ABC*,

BF is parallel to *CD*

{i 28}

Again since the angle *ACB* is equal to the angle *DEC*

AC is parallel to *FE*

{i 28}

Therefore *FACD* is a parallelogram,

therefore *FA* is equal to *DC*, and *AC* to *FD*

{i 34}

And since *AC* has been drawn parallel to *FE*, one side of the triangle *FBE*,

therefore as *BA* is to *AF* so is *BC* to *CE*

{vi 2}

But *AF* is equal to *CD*

therefore as *BA* is to *CD*, so is *BC* to *CE*,

and alternately as *AB* is to *BC*, so is *DC* to *CE*

{v 16}

Again since *CD* is parallel to *BF*,

therefore as *BC* is to *CE* so is *FD* to *DE*

{vi 2}

But *FD* is equal to *AC*,

therefore as *BC* is to *CE* so is *AC* to *DE*,

and alternately as *BC* is to *CA*, so is *CE* to *ED*

{v 16}

Since then it was proved that

as *AB* is to *BC* so is *DC* to *CE*


and, as *BC* is to *CA* so is *CE* to *ED*,

therefore *ex aequali* as *BA* is to *AC*, so is *CD* to *DE*

{v 22}

Therefore etc

Q E D



PROPOSITION 5

If two triangles have their sides proportional the triangles will be equiangular and will have those angles equal which the corresponding sides subtend

Let ABC , DEF be two triangles having their sides proportional, so that

as AB is to BC , so is DE to EF ,

as BC is to CA , so is EF to FD ,

and further, as BA is to AC , so is ED to DF ,

I say that the triangle ABC is equiangular with the triangle DEF , and they will have those angles equal which the corresponding sides subtend namely the angle ABC to the angle DEF the angle BCA to the angle EFD and further the angle BAC to the angle EDF

For on the straight line EF , and at the points E , F on it, let there be constructed the angle FEG equal to the angle ABC , and the angle EFG equal to the angle ACB ,

[I 23]

therefore the remaining angle at A is equal to the remaining angle at G

[I 32]

Therefore the triangle ABC is equiangular with the triangle GEF

Therefore in the triangles ABC GEF the sides about the equal angles are proportional and those are corresponding sides which subtend the equal angles

[VI 4]

therefore as AB is to BC , so is GE to EF

But as AB is to BC , so by hypothesis is DE to EF ,

therefore as DE is to EF , so is GE to EF

[V 11]

Therefore each of the straight lines DE GE has the same ratio to EF

therefore DE is equal to GE

[V 9]

For the same reason

DF is also equal to GF

Since then DE is equal to EG

and EF is common,

the two sides DE , EF are equal to the two sides GE EF ,

and the base DF is equal to the base FG

therefore the angle DEF is equal to the angle GEF ,

[I 8]

and the triangle DEF is equal to the triangle GEF ,

and the remaining angles are equal to the remaining angles namely those which the equal sides subtend

[I 4]

Therefore the angle DFE is also equal to the angle GFE ,

and the angle EDF to the angle EGF

And, since the angle FED is equal to the angle GEF

while the angle GEF is equal to the angle ABC

therefore the angle ABC is also equal to the angle DEF

For the same reason

the angle ACB is also equal to the angle DFE

and further the angle at A to the angle at D

therefore the triangle ABC is equiangular with the triangle DEF

Therefore etc

Q E D



PROPOSITION 6

If two triangles have one angle equal to one angle and the sides about the equal angles proportional, the triangles will be equiangular and will have those angles equal which the corresponding sides subtend

Let ABC DEF be two triangles having one angle BAC equal to one angle EDF and the sides about the equal angles proportional so that,
as BA is to AC , so is ED to DF ,

I say that the triangle ABC is equiangular with the triangle DEF , and will have the angle ABC equal to the angle DEF , and the angle ACB to the angle DFE

For on the straight line DF and at the points D, F on it, let there be constructed the angle FDG equal to either of the angles BAC, EDF , and the angle DFG equal to the angle ACB , [I 23]

therefore the remaining angle at B is equal to the remaining angle at G [I 32]


Therefore the triangle ABC is equiangular with the triangle DGF

Therefore proportionally as BA is to AC , so is GD to DF [VI 4]

But by hypothesis as BA is to AC so also is ED to DF ,
therefore also as ED is to DF so is GD to DF [v 11]

Therefore ED is equal to DG , [v 9]
and DF is common

therefore the two sides ED, DF are equal to the two sides GD, DF , and the angle EDF is equal to the angle GDF ,



therefore the base EF is equal to the base GF ,

and the triangle DEF is equal to the triangle DGF

and the remaining angles will be equal to the remaining angles namely those which the equal sides subtend [I 4]

Therefore the angle DFG is equal to the angle DFE

and the angle DGF to the angle DEF

But the angle DFG is equal to the angle ACB ,

therefore the angle ACB is also equal to the angle DFE

And by hypothesis the angle BAC is also equal to the angle EDF ,
therefore the remaining angle at B is also equal to the remaining angle at E , [I 32]

therefore the triangle ABC is equiangular with the triangle DEF

Therefore etc


Q E D

PROPOSITION 7

If two triangles have one angle equal to one angle the sides about other angles proportional and the remaining angles either both less or both not less than a right angle, the triangles will be equiangular and will have those angles equal, the sides about which are proportional

Let ABC, DEF be two triangles having one angle equal to one angle the angle BAC to the angle EDF , the sides about other angles ABC, DEF proportional so that as AB is to BC so is DE to EF , and first each of the remaining angles at C, F less than a right angle,

I say that the triangle ABC is equiangular with the triangle DEF , the angle ABC will be equal to the angle DEF , and the remaining angle namely the angle at C , equal to the remaining angle the angle at F



For if the angle ABC is unequal to the angle DEF , one of them is greater

Let the angle ABC be greater,
and on the straight line AB , and at the point B on it, let the angle ABG be constructed equal to the angle DEF [I 23]

Then, since the angle A is equal to D ,
and the angle ABG to the angle DEF ,

therefore the remaining angle AGB is equal to the remaining angle DFE [I 32]

Therefore the triangle ABG is equiangular with the triangle DEF

Therefore, as AB is to BG , so is DE to EF [VI 4]

But as DE is to EF so by hypothesis is AB to BC ,

therefore AB has the same ratio to each of the straight lines BC, BG , [V 11]

therefore BC is equal to BG [V 9]

so that the angle at C is also equal to the angle BGC [I 5]

But, by hypothesis the angle at C is less than a right angle

therefore the angle BGC is also less than a right angle,

so that the angle AGB adjacent to it is greater than a right angle [I 13]

And it was proved equal to the angle at F ,

therefore the angle at F is also greater than a right angle

But it is by hypothesis less than a right angle which is absurd

Therefore the angle ABC is not unequal to the angle DEF ,

therefore it is equal to it

But the angle at A is also equal to the angle at D

therefore the remaining angle at C is equal to the remaining angle at F [I 32]

Therefore the triangle ABC is equiangular with the triangle DEF

But again, let each of the angles at C, F be supposed not less than a right angle,

I say again that in this case too the triangle ABC is equiangular with the triangle DEF

For, with the same construction, we can prove similarly that

BC is equal to BG

so that the angle at C is also equal to the angle BGC [I 5]

But the angle at C is not less than a right angle,

therefore neither is the angle BGC less than a right angle

Thus in the triangle BGC the two angles are not less than two right angles which is impossible [I 17]

Therefore once more the angle ABC is not unequal to the angle DEF ,

therefore it is equal to it


But the angle at A is also equal to the angle at D

therefore the remaining angle at C is equal to the remaining angle at F [I 32]

Therefore the triangle ABC is equiangular with the triangle DEF

Therefore etc

Q E D



PROPOSITION 8

If in a right angled triangle a perpendicular be drawn from the right angle to the base, the triangles adjoining the perpendicular are similar both to the whole and to one another

Let ABC be a right-angled triangle having the angle BAC right, and let AD be drawn from A perpendicular to BC ,

I say that each of the triangles ABD , ADC is similar to the whole ABC and further, they are similar to one another

For, since the angle BAC is equal to the angle ADB , for each is right and the angle at B is common to the two triangles ABC and ABD

therefore the remaining angle ACB is equal to the remaining angle BAD , [I 32]

therefore the triangle ABC is equiangular with the triangle ABD

Therefore as BC which subtends the right angle in the triangle ABC is to BA which subtends the right angle in the triangle ABD , so is AB itself which subtends the angle at C in the triangle ABC to BD which subtends the equal angle BAD in the triangle ABD and so also is AC to AD which subtends the angle at B common to the two triangles [VI 4]

Therefore the triangle ABC is both equiangular to the triangle ABD and has the sides about the equal angles proportional

Therefore the triangle ABC is similar to the triangle ABD [VI Def 1]

Similarly we can prove that

the triangle ABC is also similar to the triangle ADC

therefore each of the triangles ABD , ADC is similar to the whole ABC

I say next that the triangles ABD , ADC are also similar to one another

For since the right angle BDA is equal to the right angle ADC ,

and moreover the angle BAD was also proved equal to the angle at C

therefore the remaining angle at B is also equal to the remaining angle DAC , [I 32]

therefore the triangle ABD is equiangular with the triangle ADC

Therefore as BD which subtends the angle BAD in the triangle ABD is to DA which subtends the angle at C in the triangle ADC equal to the angle BAD , so is AD itself which subtends the angle at B in the triangle ABD to DC which subtends the angle DAC in the triangle ADC equal to the angle at B , and so also is BA to AC these sides subtending the right angles, [VI 4]

therefore the triangle ABD is similar to the triangle ADC [VI Def 1]

Therefore etc


PORISM From this it is clear that if in a right angled triangle a perpendicular be drawn from the right angle to the base the straight line so drawn is a mean proportional between the segments of the base Q E D

PROPOSITION 9

From a given straight line to cut off a prescribed part


Let AB be the given straight line,

thus it is required to cut off from AB a prescribed part



Let the third part be that prescribed

Let a straight line AC be drawn through from A containing with AB any angle,



let a point D be taken at random on AC , and let DE, EC be made equal to AD [I 3]

Let BC be joined, and through D let DF be drawn parallel to it [I 31]

Then since FD has been drawn parallel to BC one of the sides of the triangle ABC , therefore, proportionally, as CD is to DA , so is BF to FA [VI 2]

But CD is double of DA

therefore BF is also double of FA ,

therefore BA is triple of AF

Therefore from the given straight line AB the prescribed third part AF has been cut off Q E F

PROPOSITION 10

To cut a given uncut straight line similarly to a given cut straight line

Let AB be the given uncut straight line and AC the straight line cut at the points D, E , and let them be so placed as to contain any angle,

let CB be joined and through D, E let DF, EG be drawn parallel to BC , and through D let DH, AK be drawn parallel to AB [I 31]

Therefore each of the figures FH, HB is a parallelogram,

therefore DH is equal to FG and HK to GB

[I 34]

Now since the straight line HE has been drawn parallel to KC , one of the sides of the triangle DKC

therefore, proportionally, as CE is to ED , so is KH to HD [VI 2]

But KH is equal to BG and HD to GF ,

therefore, as CE is to ED so is BG to GF

Again since FD has been drawn parallel to GE , one of the sides of the triangle AGE

therefore proportionally, as ED is to DA so is GF to FA [VI 2]

But it was also proved that

as CE is to ED , so is BG to GF ,

therefore as CE is to ED so is BG to GF

and as ED is to DA so is GF to FA

Therefore the given uncut straight line AB has been cut similarly to the given cut straight line AC Q E F

PROPOSITION 11

To two given straight lines to find a third proportional

Let BA, AC be the two given straight lines and let them be placed so as to contain any angle

thus it is required to find a third proportional to BA, AC

For let them be produced to the points D, E , and let BD be made equal to AC ,


let BC be joined, and through D let DE be drawn parallel to it [I 3]

Since then, BC has been drawn parallel to DE , one of the sides of the triangle ADE , proportionally as AB is to BD , so is AC to CE [I 31]

But BD is equal to AC ,

therefore as AB is to AC , so is AC to CE [VI 2]

Therefore to two given straight lines AB, AC a third proportional to them CE , has been found Q E F





PROPOSITION 12

To three given straight lines to find a fourth proportional

Let A, B, C be the three given straight lines,

thus it is required to find a fourth proportional to A, B, C



Let two straight lines DE, DF be set out containing any angle EDF , let DG be made equal to A , GE equal to B , and further DH equal to C ,

let GH be joined, and let EF be drawn through E parallel to it [I 31]

Since then, GH has been drawn parallel to EF , one of the sides of the triangle DEF ,

therefore as DG is to GE so is DH to HF [VI 2]

But DG is equal to A , GE to B and DH to C ,

therefore as A is to B so is C to HF

Therefore to the three given straight lines A, B, C a fourth proportional HF has been found Q E F


PROPOSITION 13

To two given straight lines to find a mean proportional

Let AB, BC be the two given straight lines, thus it is required to find a mean proportional to AB, BC

Let them be placed in a straight line, and let the semicircle ADC be described on AC

let BD be drawn from the point B at right angles to the straight line AC and let AD, DC be joined



Since the angle ADC is an angle in a semicircle it is right [III 31]

And, since in the right-angled triangle ADC DB has been drawn from the right angle perpendicular to the base,

therefore DB is a mean proportional between the segments of the base, AB
 BC [vi 8 For]

Therefore to the two given straight lines AB, BC a mean proportional DB
 has been found Q E F

PROPOSITION 14

In equal and equiangular parallelograms the sides about the equal angles are reciprocally proportional and equiangular parallelograms in which the sides about the equal angles are reciprocally proportional are equal

Let AB, BC be equal and equiangular parallelograms having the angles at B equal and let DB, BE be placed in a straight line

therefore FB, BG are also in a straight line

[I 14]

I say that in AB, BC the sides about the equal angles are reciprocally proportional that is to say, that, as DB is to BE , so is GB to BF

For let the parallelogram FE be completed

Since then, the parallelogram AB is equal to the parallelogram BC

and FE is another area

therefore as AB is to FE , so is BC to FE

[v 7]

But, as AB is to FE so is DB to BE

[vi 1]

and as BC is to FE so is GB to BF

[id]

therefore also as DB is to BE so is GB to BF

[v 11]

Therefore in the parallelograms AB, BC the sides about the equal angles are reciprocally proportional

Next let GB be to BF as DB to BE

I say that the parallelogram AB is equal to the parallelogram BC

For since as DB is to BE so is GB to BF

while as DB is to BE so is the parallelogram AB to the parallelogram FE

[vi 1]

and as GB is to BF so is the parallelogram BC to the parallelogram FE , [vi 1]

therefore also as AB is to FE so is BC to FE

[v 11]

therefore the parallelogram AB is equal to the parallelogram BC [v 9]

Therefore etc

Q E D

PROPOSITION 15

In equal triangles which have one angle equal to one angle the sides about the equal angles are reciprocally proportional and those triangles which have one angle equal to one angle and in which the sides about the equal angles are reciprocally proportional are equal

Let ABC, ADE be equal triangles having one angle equal to one angle namely the angle BAC to the angle DAE

I say that in the triangles ABC, ADE the sides about the equal angles are reciprocally proportional that is to say that

as CA is to AD so is EA to AB

For let them be placed so that CA is in a straight line with AD

therefore EA is also in a straight line with AB

Q E D

Let BD be joined

Since, then the triangle ABC is equal to the triangle ADE , and BAD is an other area

therefore as the triangle CAB is to the triangle BAD so is the triangle EAD to the triangle BAD [v 7]

But, as CAB is to BAD , so is CA to AD [vi 1]

and, as EAD is to BAD so is EA to AB [id]

Therefore also as CA is to AD , so is EA to AB [v 11]

Therefore in the triangles ABC , ADE the sides about the equal angles are reciprocally proportional

Next, let the sides of the triangles ABC ADE be reciprocally proportional that is to say, let EA be to AB as CA to AD ,

I say that the triangle ABC is equal to the triangle ADE

For if BD be again joined

since as CA is to AD , so is EA to AB ,

while, as CA is to AD , so is the triangle ABC to the triangle BAD ,

and as EA is to AB , so is the triangle EAD to the triangle BAD , [vi 1]


therefore as the triangle ABC is to the triangle BAD , so is the triangle EAD to the triangle BAD [v 11]

Therefore each of the triangles ABC , EAD has the same ratio to BAD

Therefore the triangle ABC is equal to the triangle EAD [v 9]

Therefore etc

Q E D




PROPOSITION 16

If four straight lines be proportional, the rectangle contained by the extremes is equal to the rectangle contained by the means and if the rectangle contained by the extremes be equal to the rectangle contained by the means the four straight lines will be proportional

Let the four straight lines AB CD , E F be proportional so that, as AB is to CD so is E to F

I say that the rectangle contained by AB F is equal to the rectangle contained by CD , E



E ———



F ———

Let AG CH be drawn from the points A C at right angles to the straight lines AB , CD and let AG be made equal to F and CH equal to E

Let the parallelograms BG DH be completed

Then since, as AB is to CD so is E to F

while E is equal to CH and F to AG

therefore as AB is to CD , so is CH to AG

Therefore in the parallelograms BG , DH the sides about the equal angles are reciprocally proportional

But those equiangular parallelograms in which the sides about the equal angles are reciprocally proportional are equal [vi 14]

therefore the parallelogram BG is equal to the parallelogram DH
 And BG is the rectangle AB, F for AG is equal to F
 and DH is the rectangle CD, E , for E is equal to CH ,
 therefore the rectangle contained by AB, F is equal to the rectangle contained
 by CD, E

Next let the rectangle contained by AB, F be equal to the rectangle contained by CD, E ,

I say that the four straight lines will be proportional so that, as AB is to CD so is E to F

For, with the same construction,

since the rectangle AB, F is equal to the rectangle CD, E ,

and the rectangle AB, F is BG , for AG is equal to F

and the rectangle CD, E is DH , for CH is equal to E

therefore BG is equal to DH

And they are equiangular

But in equal and equiangular parallelograms the sides about the equal angles are reciprocally proportional [VI 14]

Therefore, as AB is to CD so is CH to AG

But CH is equal to E , and AG to F ,

therefore, as AB is to CD , so is E to F

Therefore etc

Q E D

PROPOSITION 17

If three straight lines be proportional the rectangle contained by the extremes is equal to the square on the mean and, if the rectangle contained by the extremes be equal to the square on the mean the three straight lines will be proportional

Let the three straight lines A, B, C be proportional, so that as A is to B , so is B to C ,

I say that the rectangle contained by A, C is equal to the square on B

A —————

Let D be made equal to B

B —————

D —————

Then since as A is to B , so is B

C —————

and B is equal to D

therefore as A is to B so is D to C

But if four straight lines be proportional the rectangle contained by the extremes is equal to the rectangle contained by the means [VI 16]

Therefore the rectangle A, C is equal to the rectangle B, D

But the rectangle B, D is the square on B for B is equal to D ,

therefore the rectangle contained by A, C is equal to the square on B

Next let the rectangle A, C be equal to the square on B I say that as A is to B , so is B to C

For with the same construction

since the rectangle A, C is equal to the square on B

while the square on B is the rectangle B, D for B is equal to D

therefore the rectangle A, C is equal to the rectangle B, D

But if the rectangle contained by the extremes be equal to that contained by the means the four straight lines are proportional [VI 16]

Therefore as A is to B so is D to C

But B is equal to D ,
therefore as A is to B , so is B to C

Therefore etc

Q E D

PROPOSITION 18

On a given straight line to describe a rectilineal figure similar and similarly situated to a given rectilineal figure

Let AB be the given straight line and CE the given rectilineal figure thus it is required to describe on the straight line AB a rectilineal figure similar and similarly situated to the rectilineal figure CE

Let DF be joined and on the straight line AB , and at the points A, B on it, let the angle GAB be constructed equal to the angle at C , and the angle ABG equal to the angle CDF [I 23]

Therefore the remaining angle CFD is equal to the angle AGB ,

therefore the triangle FCD is equiangular with the triangle GAB

Therefore proportionally, as FD is to GB , so is FC to GA and CD to AB

Again, on the straight line BG , and at the points B, G on it, let the angle BGH be constructed equal to the angle DFE and the angle GBH equal to the angle FDE [I 23]

Therefore the remaining angle at E is equal to the remaining angle at H , [I 3rd]

therefore the triangle FDE is equiangular with the triangle GBH , therefore, proportionally, as FD is to GB , so is FE to GH , and ED to HB [VI 4]

But it was also proved that as FD is to GB so is FC to GA , and CD to AB , therefore also, as FC is to GA so is CD to AB , and FE to GH , and further ED to HB

And, since the angle CFD is equal to the angle AGB ,

and the angle DFE to the angle BGH ,

therefore the whole angle CFE is equal to the whole angle AGH

For the same reason

the angle CDE is also equal to the angle ABH

And the angle at C is also equal to the angle at A

and the angle at E to the angle at H

Therefore AH is equiangular with CE

and they have the sides about their equal angles proportional

therefore the rectilineal figure AH is similar to the rectilineal figure CE [VI Def 1]

Therefore on the given straight line AB the rectilineal figure AH has been described similar and similarly situated to the given rectilineal figure CE

Q E F


PROPOSITION 19

Similar triangles are to one another in the duplicate ratio of the corresponding sides

Let ABC, DEF be similar triangles having the angle at B equal to the angle

at E , and such that as AB is to BC , so is DE to EF so that BC corresponds to EF [v Def 11]

I say that the triangle ABC has to the triangle DEF a ratio duplicate of that which BC has to EF



For let a third proportional BG be taken to BC , EF , so that as BC is to EF , so is EF to BG , [vi 11]

and let AG be joined

Since then as AB is to BC , so is DE to EF

therefore alternately, as AB is to DE so is BC to EF [v 16]

But, as BC is to EF , so is EF to BG ,

therefore also as AB is to DE , so is EF to BG [v 11]

Therefore in the triangles ABG , DEF the sides about the equal angles are reciprocally proportional

But those triangles which have one angle equal to one angle, and in which the sides about the equal angles are reciprocally proportional are equal, [vi 15]

therefore the triangle ABG is equal to the triangle DEF

Now since, as BC is to EF so is EF to BG , and if three straight lines be proportional the first has to the third a ratio duplicate of that which it has to the second [v Def 9]

therefore BC has to BG a ratio duplicate of that which CB has to EF

But, as CB is to BG so is the triangle ABC to the triangle ABG , [vi 1] therefore the triangle ABC also has to the triangle ABG a ratio duplicate of that which BC has to EF

But the triangle ABG is equal to the triangle DEF therefore the triangle ABC also has to the triangle DEF a ratio duplicate of that which BC has to EF

Therefore etc

ΠΟΡΙΣΜ From this it is manifest that if three straight lines be proportional then, as the first is to the third so is the figure described on the first to that which is similar and similarly described on the second Q E D

PROPOSITION 20

Similar polygons are divided into similar triangles and into triangles equal in multitude and in the same ratio as the wholes and the polygon has to the polygon a ratio duplicate of that which the corresponding side has to the corresponding side

Let $ABCDE$ $FGHKL$ be similar polygons and let AB correspond to FG

I say that the polygons $ABCDE$ $FGHKL$ are divided into similar triangles and into triangles equal in multitude and in the same ratio as the wholes and the polygon $ABCDE$ has to the polygon $FGHKL$ a ratio duplicate of that which AB has to FG

Let BE EC , GL LH be joined

Now, since the polygon $ABCDE$ is similar to the polygon $FGHKL$,

the angle BAE is equal to the angle GFL

and as BA is to AE so is GF to FL [vi Def 1]


Since then ABE FGL are two triangles having one angle equal to one angle

and the sides about the equal angles proportional,

therefore the triangle ABE is equiangular with the triangle FGL , [VI 6]
so that it is also similar, [VI 4 and Def 1]

therefore the angle ABE is equal to the angle FGL

But the whole angle ABC is also equal to the whole angle FGH because of the similarity of the polygons,
therefore the remaining angle EBC is equal to the angle LGH



And, since because of the similarity of the triangles ABE , FGI

as EB is to BA , so is LG to GF ,

and moreover also because of the similarity of the polygons

as AB is to BC so is FG to GH ,

therefore, *ex aequali* as EB is to BC , so is LG to GH , [v 2^o]

that is the sides about the equal angles EBC , LGH are proportional,

therefore the triangle EBC is equiangular with the triangle LGH , [VI 6]

so that the triangle EBC is also similar to the triangle LGH [VI 4 and Def 1]

For the same reason

the triangle ECD is also similar to the triangle LHK

Therefore the similar polygons $ABCDE$ $FGHKL$ have been divided into similar triangles and into triangles equal in multitude

I say that they are also in the same ratio as the wholes that is in such manner that the triangles are proportional and ABE , EBC , ECD are antecedents while FGL LGH LHK are their consequents and that the polygon $ABCDE$ has to the polygon $FGHAL$ a ratio duplicate of that which the corresponding side has to the corresponding side that is AB to FG

For let AC FH be joined

Then since because of the similarity of the polygons

the angle ABC is equal to the angle FGH ,

and as AB is to BC so is FG to GH

the triangle ABC is equiangular with the triangle FGH [VI 6]

therefore the angle BAC is equal to the angle GFH

and the angle BCA to the angle GHF

And since the angle BAM is equal to the angle GFN ,

and the angle ABM is also equal to the angle FGN ,

therefore the remaining angle AMB is also equal to the remaining angle FNG , [I 32]

therefore the triangle ABM is equiangular with the triangle FGN

Similarly we can prove that

the triangle BMC is also equiangular with the triangle GNH

Therefore proportionally as AM is to MB so is FN to NG

and as BM is to MC so is GN to NH ,

so that in addition *ex aequali*

as AM is to MC , so is FN to NH

But, as AM is to MC so is the triangle ABM to MBC , and AME to EMC , for they are to one another as their bases [VI 1]

Therefore also as one of the antecedents is to one of the consequents, so are all the antecedents to all the consequents, [v 12]

therefore, as the triangle AMB is to BMC , so is ABE to CBE

But as AMB is to BMC , so is AM to MC ,

therefore also as AM is to MC , so is the triangle ABE to the triangle EBC

For the same reason also

as FN is to NH , so is the triangle FGL to the triangle GLH

And as AM is to MC , so is FN to NH ,

therefore also, as the triangle ABE is to the triangle BEC , so is the triangle FGL to the triangle GLH ,

and, alternately, as the triangle ABE is to the triangle FGL so is the triangle BEC to the triangle GLH

Similarly we can prove if BD , GA be joined, that as the triangle BEC is to the triangle LGH , so also is the triangle ECD to the triangle LHA

And since as the triangle ABE is to the triangle FGL , so is EBC to LGH , and further ECD to LHA ,

therefore also as one of the antecedents is to one of the consequents, so are all the antecedents to all the consequents, [v 12]

therefore as the triangle ABE is to the triangle FGL

so is the polygon $ABCDE$ to the polygon $FGHAL$

But the triangle ABE has to the triangle FGL a ratio duplicate of that which the corresponding side AB has to the corresponding side FG , for similar triangles are in the duplicate ratio of the corresponding sides [vi 19]

Therefore the polygon $ABCDE$ also has to the polygon $FGHAL$ a ratio duplicate of that which the corresponding side AB has to the corresponding side FG

Therefore etc

PORISM Similarly also it can be proved in the case of quadrilaterals that they are in the duplicate ratio of the corresponding sides And it was also proved in the case of triangles therefore also generally similar rectilineal figures are to one another in the duplicate ratio of the corresponding sides


Q E D

PROPOSITION 21

Figures which are similar to the same rectilineal figure are also similar to one another

For let each of the rectilineal figures A , B be similar to C , I say that A is also similar to B

For since A is similar to C ,



it is equiangular with it and has the sides about the equal angles proportional [vi Def 1]

Again since B is similar to C


it is equiangular with it and has the sides about the equal angles proportional
 Therefore each of the figures A, B is equiangular with C and with C has the sides about the equal angles proportional,
 therefore A is similar to B Q E D

PROPOSITION 22

If four straight lines be proportional the rectilineal figures similar and similarly described upon them will also be proportional and if the rectilineal figures similar and similarly described upon them be proportional the straight lines will themselves also be proportional

Let the four straight lines AB, CD, EF, GH be proportional,
 so that as AB is to CD , so is EF to GH ,
 and let there be described on AB, CD the similar and similarly situated rectilineal figures KAB, LCD
 and on EF, GH the similar and similarly situated rectilineal figures MF, NH ,
 I say that as KAB is to LCD , so is MF to NH
 For let there be taken a third proportional O to AB, CD , and a third proportional P to EF, GH [vi 11]

Then since as AB is to CD , so is EF to GH
 and, as CD is to O so is GH to P ,
 therefore *ex aequali* as AB is to O so is EF to P [v 22]
 But as AB is to O so is KAB to LCD , [vi 19 Por]
 and as EF is to P , so is MF to NH ,
 therefore also as KAB is to LCD so is MF to NH [v 11]



Next let MF be to NH as KAB is to LCD ,
 I say also that as AB is to CD so is EF to GH
 For if EF is not to GH as AB to CD
 let EF be to QR as AB to CD [vi 12]
 and on QR let the rectilineal figure SR be described similar and similarly situated to either of the two MF, NH [vi 18]
 Since then as AB is to CD so is EF to QR
 and there have been described on AB, CD the similar and similarly situated figures KAB, LCD
 and on EF, QR the similar and similarly situated figures MF, SR ,
 therefore as KAB is to LCD so is MF to SR
 But also by hypothesis
 as KAB is to LCD so is MF to NH
 therefore also as MF is to SR so is MF to NH [v 11]

Therefore MF has the same ratio to each of the figures NH , SR ,

therefore NH is equal to SR

[v 9]

But it is also similar and similarly situated to it

therefore GH is equal to QR

And, since, as AB is to CD so is EF to QR ,

while QR is equal to GH

therefore as AB is to CD , so is EF to GH

Therefore etc


Q E D

PROPOSITION 23

Equangular parallelograms have to one another the ratio compounded of the ratios of their sides

Let AC CF be equangular parallelograms having the angle BCD equal to the angle ECG ,

I say that the parallelogram AC has to the parallelogram CF the ratio compounded of the ratios of the sides



For let them be placed so that BC is in a straight line with CG ,

therefore DC is also in a straight line with CE

Let the parallelogram DG be completed,

let a straight line A be set out and let it be contrived that

as BC is to CG , so is A to L

and, as DC is to CE so is L to M

[vi 12]

Then the ratios of K to L and of L to M are the same as the ratios of the sides namely of BC to CG and of DC to CE

But the ratio of A to M is compounded of the ratio of A to L and of that of L to M ,

so that A has also to M the ratio compounded of the ratios of the sides

Now since, as BC is to CG , so is the parallelogram AC to the parallelogram CH ,

[vi 1]

while as BC is to CG so is K to L ,

therefore also as A is to L so is AC to CH

[v 11]

Again since as DC is to CE , so is the parallelogram CH to CF

[vi 1]

while as DC is to CE so is L to M ,

therefore also, as L is to M so is the parallelogram CH to the parallelogram CF

[v 11]

Since then it was proved that, as A is to L so is the parallelogram AC to the parallelogram CH

and as L is to M so is the parallelogram CH to the parallelogram CF

therefore, *ex aequali* as A is to M so is AC to the parallelogram CF

But A has to M the ratio compounded of the ratios of the sides

therefore AC also has to CF the ratio compounded of the ratios of the sides

Therefore etc

Q E D

PROPOSITION 24

In any parallelogram the parallelograms about the diameter are similar both to the whole and to one another

Let $ABCD$ be a parallelogram and AC its diameter, and let EG, HK be parallelograms about AC ,

I say that each of the parallelograms EG, HK is similar both to the whole $ABCD$ and to the other

For, since EF has been drawn parallel to BC one of the sides of the triangle ABC ,

proportionally, as BE is to EA , so is CF to FA

[VI 2]

Again since FG has been drawn parallel to CD , one of the sides of the triangle ACD ,

proportionally as CF is to FA , so is DG to GA [VI 2]

But it was proved that

as CF is to FA , so also is BE to EA ,

therefore also as BE is to EA so is DG to GA ,

and therefore *componendo*

as BA is to AE , so is DA to AG , [I 18]

and alternately,

as BA is to AD , so is EA to AG [I 16]

Therefore in the parallelograms $ABCD, EG$, the sides about the common angle BAD are proportional

And, since GF is parallel to DC ,

the angle AFG is equal to the angle DCA ,

and the angle DAC is common to the two triangles ADC, AGF ,

therefore the triangle ADC is equiangular with the triangle AGF

For the same reason

the triangle ACB is also equiangular with the triangle AFE

and the whole parallelogram $ABCD$ is equiangular with the parallelogram EG

Therefore proportionally

as AD is to DC , so is AG to GF

as DC is to CA so is GF to FA ,

as AC is to CB , so is AF to FE

and further as CB is to BA , so is FE to EA

And since it was proved that

as DC is to CA so is GF to FA ,

and,

as AC is to CB , so is AF to FE ,

therefore *ex aequali* as DC is to CB so is GF to FE [V 22]

Therefore in the parallelograms $ABCD, EG$ the sides about the equal angles are proportional,

therefore the parallelogram $ABCD$ is similar to the parallelogram EG

[VI Def 1]

For the same reason

the parallelogram $ABCD$ is also similar to the parallelogram HK

therefore each of the parallelograms EG, HK is similar to $ABCD$


But figures similar to the same rectilineal figure are also similar to one another [VI 21]

therefore the parallelogram EG is also similar to the parallelogram HK
Therefore etc Q E D

PROPOSITION 25

To construct one and the same figure similar to a given rectilineal figure and equal to another given rectilineal figure

Let ABC be the given rectilineal figure to which the figure to be constructed must be similar and D that to which it must be equal
thus it is required to construct one and the same figure similar to ABC and equal to D



Let there be applied to BC the parallelogram BE equal to the triangle ABC [I 44] and to CF the parallelogram CM equal to D in the angle FCE which is equal to the angle CBL [I 45]

Therefore BC is in a straight line with CF and LE with EM

Now let GH be taken a mean proportional to BC , CF [VI 13] and on GH let AKH be described similar and similarly situated to ABC [VI 18]

Then since, as BC is to GH , so is GH to CF ,
and if three straight lines be proportional as the first is to the third so is the figure on the first to the similar and similarly situated figure described on the second [VI 19 Per]

therefore as BC is to CF so is the triangle ABC to the triangle AKH

But, as BC is to CF so also is the parallelogram BE to the parallelogram EF [VI 1]

Therefore also as the triangle ABC is to the triangle AKH so is the parallelogram BE to the parallelogram EF ,

therefore alternately as the triangle ABC is to the parallelogram BE so is the triangle AKH to the parallelogram EF [V 16]

But the triangle ABC is equal to the parallelogram BE

therefore the triangle AKH is also equal to the parallelogram EF

But the parallelogram EF is equal to D

therefore AKH is also equal to D

And AKH is also similar to ABC

Therefore one and the same figure AKH has been constructed similar to the given rectilineal figure ABC and equal to the other given figure D Q E D

PROPOSITION 26

If from a parallelogram there be taken away a parallelogram similar and similarly situated to the whole and having a common angle with it it is about the same diameter with the whole

For from the parallelogram $ABCD$ let there be taken away the parallelo-

gram AF similar and similarly situated to $ABCD$, and having the angle DAB common with it,

I say that $ABCD$ is about the same diameter with AF

For suppose it is not but, if possible let AHC be the diameter $<$ of $ABCD >$, let GF be produced and carried through to H , and let HK be drawn through H parallel to either of the straight lines AD, BC

[I 31]

Since, then $ABCD$ is about the same diameter with AG , therefore as DA is to AB , so is GA to AK [VI 24]

But also because of the similarity of $ABCD$ EG ,
as DA is to AB , so is GA to AE ,

therefore also as GA is to AK so is GA to AE [V 11]

Therefore GA has the same ratio to each of the straight lines AK, AE

Therefore AE is equal to AK [V 9] the less to the greater which is impossible

Therefore $ABCD$ cannot but be about the same diameter with AF , therefore the parallelogram $ABCD$ is about the same diameter with the parallelogram AF

Therefore etc

Q E D


PROPOSITION 27

Of all the parallelograms applied to the same straight line and deficient by parallelogrammic figures similar and similarly situated to that described on the half of the straight line that parallelogram is greatest which is applied to the half of the straight line and is similar to the defect

Let AB be a straight line and let it be bisected at C , let there be applied to the straight line AB the parallelogram AD deficient by the parallelogrammic figure DB described on the half of AB , that is CB ,

I say that of the parallelograms applied to AB and deficient by parallelogrammic figures similar and similarly situated to DB AD is greatest

For let there be applied to the straight line AB the parallelogram AF deficient by the parallelogrammic figure FB similar and similarly situated to DB ,



I say that AD is greater than AF

For since the parallelogram DB is similar to the parallelogram FB ,
they are about the same diameter [VI 26]

Let their diameter DB be drawn and let the figure be described

Then, since CF is equal to FE [I 43]

and FB is common

therefore the whole CH is equal to the whole AE

But CH is equal to CG since AC is also equal to CB [I 36]

Therefore GC is also equal to EK

Let CF be added to each

therefore the whole AF is equal to the gnomon LMN ,

so that the parallelogram DB , that is, AD , is greater than the parallelogram AF

Therefore etc

Q E D

PROPOSITION 28


To a given straight line to apply a parallelogram equal to a given rectilineal figure and deficient by a parallelogrammic figure similar to a given one thus the given rectilineal figure must not be greater than the parallelogram described on the half of the straight line and similar to the defect

Let AB be the given straight line C the given rectilineal figure to which the figure to be applied to AB is required to be equal not being greater than the parallelogram described on the half of AB and similar to the defect and D the parallelogram to which the defect is required to be similar, thus it is required to apply to the given straight line AB a parallelogram equal to the given rectilineal figure C and deficient by a parallelogrammic figure which is similar to D

Let AB be bisected at the point E , and on EB let $EBFG$ be described similar and similarly situated to D , [VI 18]

let the parallelogram AG be completed

If then AG is equal to C that which was enjoined will have been done for there has been applied to the given straight line AB the parallelogram AG equal to the given rectilineal figure C and deficient by a parallelogrammic figure GB which is similar to D



But if not let HE be greater than C

Now HE is equal to GB ,

therefore GB is also greater than C

Let $KLMN$ be constructed at once equal to the excess by which GB is greater than C and similar and similarly situated to D [VI 25]

But D is similar to GB ,

therefore KM is also similar to GB

[VI 21]

Let then, KL correspond to GE and LM to GF

Now since GB is equal to C KM ,

therefore GB is greater than KM

therefore also GE is greater than KL and GF than LM

Let GO be made equal to KL and GP equal to LM and let the parallelogram $OGPQ$ be completed

therefore it is equal and similar to KM

Therefore GQ is also similar to GB

[VI 21]

therefore GQ is about the same diameter with GB [VI 26]
 Let GQB be their diameter, and let the figure be described
 Then, since BG is equal to C , AM ,
 and in them GQ is equal to AM
 therefore the remainder the gnomon UVV , is equal to the remainder C
 And since PR is equal to OS

let QB be added to each
 therefore the whole PB is equal to the whole OB
 But OB is equal to TE since the side AE is also equal to the side EB , [I 36]
 therefore TE is also equal to PB


Let OS be added to each,
 therefore the whole TS is equal to the whole the gnomon UVU
 But the gnomon UVU was proved equal to C ,
 therefore TS is also equal to C

Therefore to the given straight line AB there has been applied the parallelogram ST equal to the given rectilinear figure C and deficient by a parallelogrammic figure QB which is similar to D Q E F

PROPOSITION 29

To a given straight line to apply a parallelogram equal to a given rectilinear figure and exceeding by a parallelogrammic figure similar to a given one

Let AB be the given straight line, C the given rectilinear figure to which the figure to be applied to AB is required to be equal, and D that to which the excess is required to be similar
 thus it is required to apply to the straight line AB a parallelogram equal to the rectilinear figure C and exceeding by a parallelogrammic figure similar to D



Let AB be bisected at E
 let there be described on EB the parallelogram BF similar and similarly situated to D
 and let GH be constructed at once equal to the sum of BF C and similar and similarly situated to D [VI 25]

Let AH correspond to FL and AG to FE

Now since GH is greater than FB

therefore AH is also greater than FL and AG than FE

Let FL , FE be produced

let FLM be equal to AH and FEN to AG

and let MN be completed

therefore MN is both equal and similar to GH

But GH is similar to EL ,

therefore MN is also similar to EL

[VI 21]

therefore EL is about the same diameter with MN

[VI 26]

Let their diameter FO be drawn, and let the figure be described

Since GH is equal to EL , C ,

while GH is equal to MN

therefore MN is also equal to EL , C

Let EL be subtracted from each

therefore the remainder, the gnomon $VH\Lambda$ is equal to C

Now, since AE is equal to EB ,

AN is also equal to NB [I 36] that is, to LP

[I 43]

Let EO be added to each,

therefore the whole AO is equal to the gnomon $VH\Lambda$

But the gnomon $VH\Lambda$ is equal to C

therefore AO is also equal to C

Therefore to the given straight line AB there has been applied the parallelogram AO equal to the given rectilineal figure C and exceeding by a parallelogram figure QP which is similar to D since PQ is also similar to EL [VI 24]


Q E F

PROPOSITION 30

To cut a given finite straight line in extreme and mean ratio

Let AB be the given finite straight line

thus it is required to cut AB in extreme and mean ratio



On AB let the square BC be described and let there be applied to AC the parallelogram CD equal to BC and exceeding by the figure AD similar to BC

[VI 29]

Now BC is a square

therefore AD is also a square

And since BC is equal to CD

let CE be subtracted from each,

therefore the remainder BF is equal to the remainder AD

But it is also equiangular with it

therefore in BF , AD the sides about the equal angles are reciprocally proportional,

[VI 14]

therefore as FE is to ED , so is AE to EB

But FE is equal to AB and ED to AE

Therefore, as BA is to AE so is AE to EB

And AB is greater than AE ,

therefore AE is also greater than EB

Therefore the straight line AB has been cut in extreme and mean ratio at E and the greater segment of it is AE

Q E F

PROPOSITION 31

In right-angled triangles the figure on the side subtending the right angle is equal to the similar and similarly described figures on the sides containing the right angle

Let ABC be a right angled triangle having the angle BAC right

I say that the figure on BC is equal to the similar and similarly described figures on BA AC

Let AD be drawn perpendicular

Then since in the right-angled triangle ABC , AD has been drawn from the right angle at A perpendicular to the base BC , the triangles ABD , ADC adjoining the perpendicular are similar both to the whole ABC and to one another [vi 8]

And since ABC is similar to ABD , therefore as CB is to BA so is AB to BD [vi Def 1]

And since three straight lines are proportional, as the first is to the third so is the figure on the first to the similar and similarly described figure on the second [vi 19 Por]

Therefore as CB is to BD , so is the figure on CB to the similar and similarly described figure on BA

For the same reason also,

as BC is to CD so is the figure on BC to that on CA ,
so that, in addition

as BC is to BD , DC so is the figure on BC to the similar and similarly described figures on BA , AC

But BC is equal to BD , DC , therefore the figure on BC is also equal to the similar and similarly described figures on BA , AC

Therefore etc

Q E D

PROPOSITION 32

If two triangles having two sides proportional to two sides be placed together at one angle so that their corresponding sides are also parallel, the remaining sides of the triangles will be in a straight line

Let ABC , DCE be two triangles having the two sides BA , AC proportional to the two sides DC , DE , so that as AB is to AC so is DC to DE , and AB parallel to DC , and AC to DE

I say that BC is in a straight line with CE

For since AB is parallel to DC and the straight line AC has fallen upon them

the alternate angles BAC , ACD are equal to one another [i 20]

For the same reason

the angle CDE is also equal to the angle ACD ,



so that the angle BAC is equal to the angle CDE

And, since ABC , DCE are two triangles having one angle the angle at A , equal to one angle the angle at D

and the sides about the equal angles proportional,

so that as BA is to AC so is CD to DE

therefore the triangle ABC is equiangular with the triangle DCE , [vi 6]



therefore the angle ABC is equal to the angle DCE

But the angle ACD was also proved equal to the angle BAC ,

therefore the whole angle ACE is equal to the two angles ABC, BAC

Let the angle ACB be added to each,

therefore the angles ACE, ACB are equal to the angles BAC, ACB, CBA

But the angles BAC, ABC, ACB are equal to two right angles, [I 32]

therefore the angles ACE, ACB are also equal to two right angles

Therefore with a straight line AC , and at the point C on it the two straight lines BC, CE not lying on the same side make the adjacent angles ACE, ACB equal to two right angles,

therefore BC is in a straight line with CE

[I 14]

Therefore etc



Q E D

PROPOSITION 33

In equal circles angles have the same ratio as the circumferences on which they stand whether they stand at the centres or at the circumferences

Let ABC, DEF be equal circles and let the angles BGC, EHF be angles at their centres G, H , and the angles BAC, EDF angles at the circumferences,

I say that, as the circumference BC is to the circumference EF so is the angle BGC to the angle EHF , and the angle BAC to the angle EDF



For let any number of consecutive circumferences CK, KL be made equal to the circumference BC

and any number of consecutive circumferences FM, MN equal to the circumference EF

and let GK, GL, HM, HN be joined

Then since the circumferences BC, CK, KL are equal to one another

the angles BGC, CGK, KGL are also equal to one another [III 27]

therefore whatever multiple the circumference BL is of BC , that multiple also is the angle BGL of the angle BGC

For the same reason also

whatever multiple the circumference NE is of EF that multiple also is the angle NHE of the angle EHF

If then the circumference BL is equal to the circumference EN the angle BGL is also equal to the angle EHN [III 27]

if the circumference BL is greater than the circumference EN the angle BGL is also greater than the angle EHN

and if less less

There being then four magnitudes two circumferences BC, EF and two angles BGC, EHF

there have been taken of the circumference BC and the angle BGC equimultiples namely the circumference BL and the angle BGL

and of the circumference EF and the angle EHF equimultiples namely the circumference EN and the angle EHN

And it has been proved that

if the circumference BL is in excess of the circumference EN

the angle BGL is also in excess of the angle EHN ,
 if equal, equal,
 and if less less

I therefore as the circumference BC is to EF , so is the angle BGC to the angle EHF [v Def 5]

But as the angle BGC is to the angle EHF , so is the angle BAC to the angle EDF , for they are doubles respectively

Therefore also, as the circumference BC is to the circumference EF , so is the angle BGC to the angle EHF , and the angle BAC to the angle EDF

Therefore etc

Q E D

BOOK SEVEN

DEFINITIONS

- 1 An *unit* is that by virtue of which each of the things that exist is called one
- 2 A *number* is a multitude composed of units
- 3 A number is a *part* of a number the less of the greater, when it measures the greater
- 4 but *parts* when it does not measure it
- 5 The greater number is a *multiple* of the less when it is measured by the less
- 6 An *even number* is that which is divisible into two equal parts
- 7 An *odd number* is that which is not divisible into two equal parts or that which differs by an unit from an even number
- 8 An *even times even number* is that which is measured by an even number according to an even number
- 9 An *even times odd number* is that which is measured by an even number according to an odd number
- 10 An *odd times odd number* is that which is measured by an odd number according to an odd number
- 11 A *prime number* is that which is measured by an unit alone
- 12 Numbers *prime to one another* are those which are measured by an unit alone as a common measure
- 13 A *composite number* is that which is measured by some number
- 14 Numbers *composite to one another* are those which are measured by some number as a common measure
- 15 A number is said to *multiply* a number when that which is multiplied is added to itself as many times as there are units in the other and thus some number is produced
- 16 And when two numbers having multiplied one another make some number the number so produced is called *plane* and its *sides* are the numbers which have multiplied one another
- 17 And when three numbers having multiplied one another make some number the number so produced is *solid* and its *sides* are the numbers which have multiplied one another
- 18 A *square number* is equal multiplied by equal, or a number which is contained by two equal numbers
- 19 And a *cube* is equal multiplied by equal and again by equal or a number which is contained by three equal numbers
- 20 Numbers are *proportional* when the first is the same multiple or the same part or the same parts of the second that the third is of the fourth

21 *Similar plane and solid numbers* are those which have their sides proportional

22 A *perfect number* is that which is equal to its own parts

BOOK VII PROPOSITIONS

PROPOSITION 1

Two unequal numbers being set out and the less being continually subtracted in turn from the greater if the number which is left never measures the one before it until an unit is left the original numbers will be prime to one another

For the less of two unequal numbers AB , CD being continually subtracted from the greater let the number which is left never measure the one before it until an unit is left,

I say that AB , CD are prime to one another that is that an unit alone measures AB , CD

For, if AB , CD are not prime to one another, some number will measure them

Let a number measure them, and let it be E , let CD , measuring BF , leave F less than itself

let AF measuring DG leave GC less than itself,

and let GC measuring FH leave an unit HA

Since then E measures CD and CD measures BF ,

therefore E also measures BF

But it also measures the whole BA ,

therefore it will also measure the remainder AF

But AF measures DG ,

therefore E also measures DG

But it also measures the whole DC

therefore it will also measure the remainder CG

But CG measures FH


therefore E also measures FH

But it also measures the whole FA

therefore it will also measure the remainder the unit HA , though it is a number which is impossible

Therefore no number will measure the numbers AB , CD therefore AB , CD are prime to one another

[VII Def 12]
Q E D



PROPOSITION 2

Given two numbers not prime to one another to find their greatest common measure

Let AB , CD be the two given numbers not prime to one another

Thus it is required to find the greatest common measure of AB , CD

If now CD measures AB —and it also measures itself— CD is a common measure of CD , AB

And it is manifest that it is also the greatest, for no greater number than CD will measure CD

But if CD does not measure AB then the less of the numbers AB , CD being continually subtracted from the greater, some number will be left which will measure the one before it

For an unit will not be left, otherwise AB, CD will be prime to one another [VII 1] which is contrary to the hypothesis

Therefore some number will be left which will measure the one before it

Now let CD , measuring BE , leave EA less than itself,

let EA , measuring DF , leave FC less than itself,

and let CF measure AE

Since then CF measures AE , and AE measures DF ,

therefore CF will also measure DF

But it also measures itself,

therefore it will also measure the whole CD

But CD measures BE ,

therefore CF also measures BE

But it also measures EA ,

therefore it will also measure the whole BA

But it also measures CD

therefore CF measures AB, CD

Therefore CF is a common measure of AB, CD

I say next that it is also the greatest

For if CF is not the greatest common measure of AB, CD , some number which is greater than CF will measure the numbers AB, CD

Let such a number measure them and let it be G

Now, since G measures CD while CD measures BE , G also measures BE

But it also measures the whole BA ,

therefore it will also measure the remainder AE

But AE measures DF ,

therefore G will also measure DF

But it also measures the whole DC ,

therefore it will also measure the remainder CF that is, the greater will measure the less which is impossible

Therefore no number which is greater than CF will measure the numbers AB, CD

therefore CF is the greatest common measure of AB, CD

PORISM From this it is manifest that if a number measure two numbers it will also measure their greatest common measure Q E D

PROPOSITION 3

Given three numbers not prime to one another to find their greatest common measure

Let A, B, C be the three given numbers not prime to one another

thus it is required to find the greatest common measure of A, B, C

For let the greatest common measure D of the two numbers A, B be taken [VII 2]

then D either measures or does not measure C

First let it measure it

But it measures A, B also,

therefore D measures A, B, C

therefore D is a common measure of A, B, C

I say that it is also the greatest

For if D is not the greatest common measure of A, B, C some number which

21 *Similar plane and solid numbers* are those which have their sides proportional

22 A *perfect number* is that which is equal to its own parts

BOOK VII PROPOSITIONS

PROPOSITION 1

Two unequal numbers being set out and the less being continually subtracted in turn from the greater if the number which is left never measures the one before it until an unit is left the original numbers will be prime to one another

For the less of two unequal numbers AB , CD being continually subtracted from the greater let the number which is left never measure the one before it until an unit is left,

I say that AB , CD are prime to one another, that is that an unit alone measures AB , CD

For if AB , CD are not prime to one another, some number will measure them

Let a number measure them, and let it be E , let CD , measuring BF , leave FA less than itself

let AF , measuring DG , leave GC less than itself

and let GC measuring FH , leave an unit HA

Since then E measures CD and CD measures BF ,

therefore E also measures BF

But it also measures the whole BA

therefore it will also measure the remainder AF

But AF measures DG ,

therefore E also measures DG

But it also measures the whole DC

therefore it will also measure the remainder CG

But CG measures FH

therefore E also measures FH


But it also measures the whole FA ,

therefore it will also measure the remainder the unit HA though it is a number which is impossible

Therefore no number will measure the numbers AB , CD , therefore AB , CD are prime to one another

[VII Def 12]

Q E D



PROPOSITION 2

Given two numbers not prime to one another to find their greatest common measure

Let AB CD be the two given numbers not prime to one another

Thus it is required to find the greatest common measure of AB CD

If now CD measures AB —and it also measures itself— CD is a common measure of CD , AB

And it is manifest that it is also the greatest, for no greater number than CD will measure CD

But if CD does not measure AB then the less of the numbers AB CD being continually subtracted from the greater some number will be left which will measure the one before it

For an unit will not be left otherwise AB, CD will be prime to one another [vii 1] which is contrary to the hypothesis

Therefore some number will be left which will measure the one before it

Now let CD measuring BE , leave EA less than itself,

let EA measuring DF , leave FC less than itself,

and let CF measure AE

Since then CF measures AE , and AE measures DF ,

therefore CF will also measure DF

But it also measures itself

therefore it will also measure the whole CD

But CD measures BE ,

therefore CF also measures BE

But it also measures EA

therefore it will also measure the whole BA

But it also measures CD ,

therefore CF measures AB, CD

Therefore CF is a common measure of AB, CD

I say next that it is also the greatest

For if CF is not the greatest common measure of AB, CD some number which is greater than CF will measure the numbers AB, CD

Let such a number measure them and let it be G

Now since G measures CD while CD measures BE G also measures BE

But it also measures the whole BA

therefore it will also measure the remainder AE

But AE measures DF ,

therefore G will also measure DF

But it also measures the whole DC

therefore it will also measure the remainder CF that is the greater will measure the less which is impossible

Therefore no number which is greater than CF will measure the numbers AB, CD

therefore CF is the greatest common measure of AB, CD

PROBLEM From this it is manifest that if a number measure two numbers it will also measure their greatest common measure Q E D

PROPOSITION 3

Given three numbers not prime to one another to find their greatest common measure

Let A, B, C be the three given numbers not prime to one another

thus it is required to find the greatest common measure of A, B, C

For let the greatest common measure D of the two numbers A, B be taken [vii 2]

then D either measures or does not measure C

First let it measure it

But it measures A, B also

therefore D measures A, B, C

therefore D is a common measure of A, B, C

I say that it is also the greatest

For if D is not the greatest common measure of A, B, C some number which

is greater than D will measure the numbers A, B, C

Let such a number measure them and let it be E

Since then E measures A, B, C

it will also measure $A, B,$

therefore it will also measure the greatest common measure of A, B

[VII 2, Por]

But the greatest common measure of A, B is $D,$

therefore E measures D , the greater the less which is impossible¹

Therefore no number which is greater than D will measure the numbers $A, B, C,$

therefore D is the greatest common measure of A, B, C

Next let D not measure $C,$

I say first that C, D are not prime to one another

For since A, B, C are not prime to one another some number will measure them

Now that which measures A, B, C will also measure $A, B,$ and will measure D the greatest common measure of A, B

[VII 2 Por]

But it measures C also,

therefore some number will measure the numbers $D, C,$

therefore D, C are not prime to one another

Let then their greatest common measure E be taken

[VII 2]

Then since E measures D

and D measures $A, B,$

therefore E also measures A, B

But it measures C also

therefore E measures A, B, C

therefore E is a common measure of A, B, C

I say next that it is also the greatest

For if E is not the greatest common measure of $A, B, C,$ some number which is greater than E will measure the numbers A, B, C

Let such a number measure them and let it be F

Now since F measures A, B, C

it also measures A, B

therefore it will also measure the greatest common measure of A, B

[VII 2 Por]

But the greatest common measure of A, B is D

therefore F measures D

And it measures C also

therefore F measures D, C

therefore it will also measure the greatest common measure of D, C

[VII 2, Por]

But the greatest common measure of D, C is $E,$

therefore F measures E the greater the less which is impossible

Therefore no number which is greater than E will measure the numbers A, B, C

therefore E is the greatest common measure of A, B, C

Q E D

PROPOSITION 4

Any number is either a part or parts of any number, the less of the greater

Let A, BC be two numbers and let BC be the less,

I say that BC is either a part or parts of A

For A, BC are either prime to one another or not

First, let A, BC be prime to one another

Then if BC be divided into the units in it, each unit of those in BC will be some part of A , so that BC is parts of A

Next let A, BC not be prime to one another, then BC either measures or does not measure A

If now BC measures A , BC is a part of A

But, if not let the greatest common measure D of A, BC be taken [VII 2] and let BC be divided into the numbers equal to D , namely BL, EF, FC

Now since D measures A , D is a part of A

But D is equal to each of the numbers BE, EF, FC ,

therefore each of the numbers BE, EF, FC is also a part of A , so that BC is parts of A

Therefore etc

Q E D

PROPOSITION 5

If a number be a part of a number and another be the same part of another, the sum will also be the same part of the sum that the one is of the one

For let the number A be a part of BC

and another D the same part of another EF that A is of BC

I say that the sum of A, D is also the same part of the sum of BC, EF that A is of BC

For since, whatever part A is of BC , D is also the same part of EF ,

therefore, as many numbers as there are in BC equal to A so many numbers are there also in EF equal to D

Let BC be divided into the numbers equal to A namely BG, GC ,

and EF into the numbers equal to D namely EH, HF ,

then the multitude of BG, GC will be equal to the multitude of EH, HF

And since BG is equal to A , and EH to D ,

therefore BG, EH are also equal to A, D

For the same reason

GC, HF are also equal to A, D

Therefore as many numbers as there are in BC equal to A so many are there also in BC, EF equal to A, D

Therefore whatever multiple BC is of A , the same multiple also is the sum of BC, EF of the sum of A, D

Therefore whatever part A is to BC , the same part also is the sum of A, D of the sum of BC, EF

Q E D

PROPOSITION 6

If a number be parts of a number and another be the same parts of another, the sum will also be the same parts of the sum that the one is of the one

For let the number AB be parts of the number C , and another, DE , the same parts of another, F , that AB is of C ,

I say that the sum of AB , DE is also the same parts of the sum of C , F that AB is of C

For since whatever parts AB is of C , DE is also the same parts of F ,

therefore, as many parts of C as there are in AB , so many parts of F are there also in DE

Let AB be divided into the parts of C , namely AG , GB , and DE into the parts of F , namely DH , HE ,

thus the multitude of AG , GB will be equal to the multitude of DH , HE

And since, whatever part AG is of C , the same part is DH of F also, therefore, whatever part AG is of C , the same part also is the sum of AG , DH of the sum of C , F [VII 5]

For the same reason

whatever part GB is of C , the same part also is the sum of GB , HE of the sum of C , F

Therefore, whatever parts AB is of C , the same parts also is the sum of AB , DE of the sum of C , F Q E D

PROPOSITION 7

If a number be that part of a number, which a number subtracted is of a number subtracted the remainder will also be the same part of the remainder that the whole is of the whole

For let the number AB be that part of the number CD which AE subtracted is of CF subtracted

I say that the remainder EB is also the same part of the remainder FD that the whole AB is of the whole CD



For whatever part AE is of CF , the same part also let EB be of CG
 Now since whatever part AE is of CF the same part also is EB of CG
 therefore whatever part AE is of CF the same part also is AB of GF [VII 5]
 But, whatever part AE is of CF , the same part also, by hypothesis, is AB of CD

therefore whatever part AB is of GF , the same part is it of CD also
 therefore GF is equal to CD

Let CF be subtracted from each

therefore the remainder GC is equal to the remainder FD

Now since whatever part AE is of CF , the same part also is EB of GC , while GC is equal to FD

therefore whatever part AE is of CF the same part also is EB of FD

But whatever part AE is of CF the same part also is AB of CD , therefore also the remainder EB is the same part of the remainder FD that the whole AB is of the whole CD Q E D

PROPOSITION 8

If a number be the same parts of a number that a number subtracted is of a number subtracted, the remainder will also be the same parts of the remainder that the whole is of the whole


For let the number AB be the same parts of the number CD that AE subtracted is of CF subtracted,

I say that the remainder EB is also the same parts of the remainder FD that the whole AB is of the whole CD

For let GH be made equal to AB

Therefore whatever parts GH is of CD , the same parts also is AE of CF

Let GH be divided into the parts of CD namely GK , KH , and AE into the parts of CF , namely AL , LE



thus the multitude of GK , KH will be equal to the multitude of AL , LE

Now since whatever part GK is of CD the same part also is AL of CF , while CD is greater than CF

therefore GK is also greater than AL

Let GM be made equal to AL

Therefore, whatever part GK is of CD , the same part also is GM of CF therefore also the remainder MA is the same part of the remainder FD that the whole GK is of the whole CD [II 7]

Again since whatever part KH is of CD the same part also is EL of CF , while CD is greater than CF ,

therefore KH is also greater than EL

Let KN be made equal to FL

Therefore whatever part KH is of CD , the same part also is KN of CF , therefore also the remainder NH is the same part of the remainder FD that the whole KH is of the whole CD [II 7]

But the remainder MK was also proved to be the same part of the remainder FD that the whole GK is of the whole CD , therefore also the sum of MK , NH is the same parts of DF that the whole HG is of the whole CD

But the sum of MA , NH is equal to EB ,

and HG is equal to BA ,

therefore the remainder EB is the same parts of the remainder FD that the whole AB is of the whole CD Q E D

PROPOSITION 9

If a number be a part of a number and another be the same part of another alternately also whatever part or parts the first is of the third the same part or the same parts will the second also be of the fourth

For let the number A be a part of the number BC , and another D the same part of another, EF that A is of BC ,

I say that alternately also whatever part or parts A is of D the same part or parts is BC of EF also

For since whatever part A is of BC the same part also is D of EF



therefore as many numbers as there are in BC equal to A , so many also are there in EF equal to D

Let BC be divided into the numbers equal to A , namely BG GC ,
and EF into those equal to D namely EH , HF ,

thus the multitude of BG GC will be equal to the multitude of EH , HF

Now, since the numbers BG , GC are equal to one another, and the numbers EH HF are also equal to one another

while the multitude of BG , GC is equal to the multitude of EH HF ,¹
therefore whatever part or parts BG is of EH , the same part or the same parts is GC of HF also,

so that in addition whatever part or parts BG is of EH , the same part also or the same parts is the sum BC of the sum EF [VII 5 6]

But BG is equal to A , and EH to D ,
therefore whatever part or parts A is of D the same part or the same parts is BC of EF also Q E D

PROPOSITION 10

If a number be parts of a number and another be the same parts of another alternately also whatever parts or part the first is of the third the same parts or the same part will the second also be of the fourth

For let the number AB be parts of the number C and another, DE , the same parts of another, F ,

I say that alternately also whatever parts or part AB is of DE the same parts or the same part is C of F also

For since whatever parts AB is of C the same parts also is DE of F

therefore as many parts of C as there are in AB so many parts also of F are there in DE

Let AB be divided into the parts of C namely AG , GB
and DE into the parts of F , namely DH , HE

thus the multitude of AG GB will be equal to the multitude of DH , HE

Now since whatever part AG is of C the same part also is DH of F ,

alternately also whatever part or parts AG is of DH ,


the same part or the same parts is C of F also [VII 9]

For the same reason also,

whatever part or parts GB is of HE , the same part or the same parts is C of F also

so that in addition whatever parts or part AB is of DE , the same parts also or the same part is C of F [VII 5 6]

Q E D



PROPOSITION 11

If as whole is to whole so is a number subtracted to a number subtracted the remainder will also be to the remainder as whole to whole

As the whole AB is to the whole CD so let AE subtracted be to CF subtracted,

I say that the remainder EB is also to the remainder FD as the whole AB to the whole CD

Since, as AB is to CD so is AE to CF ,

$\left. \begin{array}{l} A \\ E \\ B \end{array} \right\} \left. \begin{array}{l} C \\ F \\ D \end{array} \right\}$ whatever part or parts AB is of CD , the same part or the same parts is $4E$ of CF also, [VII Def 20]
 Therefore also the remainder EB is the same part or parts of FD that $1B$ is of CD [VII 7, 8]
 Therefore as EB is to FD , so is AB to CD [VII Def 20]
 Q E D

PROPOSITION 12

If there be as many numbers as we please in proportion, then as one of the antecedents is to one of the consequents, so are all the antecedents to all the consequents

Let A, B, C, D be as many numbers as we please in proportion so that, as A is to B , so is C to D

$\left. \begin{array}{l} A \\ B \\ C \\ D \end{array} \right\}$ I say that as A is to B so are $4, C$ to B, D
 For since, as A is to B so is C to D
 whatever part or parts A is of B , the same part or parts is C of D also [VII Def 20]
 Therefore also the sum of A, C is the same part or the same parts of the sum of B, D that A is of B [VII 5, 6]
 Therefore as A is to B , so are A, C to B, D [VII Def 20]
 Q E D

PROPOSITION 13

If four numbers be proportional they will also be proportional alternately

Let the four numbers A, B, C, D be proportional so that as A is to B so is C to D

I say that they will also be proportional alternately so that as A is to C so will B be to D

$\left. \begin{array}{l} A \\ B \\ C \\ D \end{array} \right\}$ For since as A is to B so is C to D
 therefore whatever part or parts A is of B the same part or the same parts is C of D also [VII Def 20]
 Therefore, alternately whatever part or parts A is of C the same part or the same parts is B of D also [VII 10]
 Therefore as A is to C so is B to D [VII Def 20]
 Q E D

PROPOSITION 14

If there be as many numbers as we please and others equal to them in multitude, which taken two and two are in the same ratio they will also be in the same ratio *ex aequali*

Let there be as many numbers as we please A, B, C and others equal to them in multitude D, E, F which taken two and two are in the same ratio so that

$\frac{A}{B} = \frac{D}{E}$ as 4 is to B so is D to E
 $\frac{B}{C} = \frac{E}{F}$ and as B is to C , so is E to F
 I say that *ex aequali*
 as A is to C so also is D to F

For since as A is to B so is D to E

therefore alternately

as A is to D so is B to E

[VII 13]

Again since as B is to C so is E to F

therefore alternately,

as B is to E , so is C to F

[vii 13]

But as B is to F so is A to D ,

therefore also, as A is to D , so is C to F

Therefore alternately


as A is to C , so is D to F

[id]

PROPOSITION 15

If an unit measure any number, and another number measure any other number the same number of times alternately also the unit will measure the third number the same number of times that the second measures the fourth

For let the unit A measure any number BC ,
and let another number D measure any
other number EF the same number of
times,



I say that alternately also the unit A
measures the number D the same number
of times that BC measures EF

For, since the unit A measures the number BC the same number of times
that D measures EF

therefore as many units as there are in BC , so many numbers equal to D are
there in EF also

Let BC be divided into the units in it BG GH , HC ,

and EF into the numbers EK , KL LF equal to D

Thus the multitude of BG , GH HC will be equal to the multitude of EK ,
 KL , LF

And since the units BG , GH HC are equal to one another

and the numbers EK , KL , LF are also equal to one another,

while the multitude of the units BG , GH , HC is equal to the multitude of the
numbers EK KL , LF ,

therefore as the unit BG is to the number EK , so will the unit GH be to the
number KL and the unit HC to the number LF

Therefore also as one of the antecedents is to one of the consequents so will
all the antecedents be to all the consequents [vii 1^o]

therefore as the unit BG is to the number EK so is BC to EF

But the unit BG is equal to the unit A ,

and the number EK to the number D

Therefore as the unit A is to the number D so is BC to EF

Therefore the unit A measures the number D the same number of times that
 BC measures EF

Q E D

PROPOSITION 16

*If two numbers by multiplying one another make certain numbers the numbers so
produced will be equal to one another*

Let A B be two numbers and let A by multiplying B make C , and B by
multiplying A make D ,

I say that C is equal to D

For since A by multiplying B has made C

therefore B measures C according to the units in A

But the unit *E* also measures the number *A* according to the units in it
therefore the unit *E* measures *A* the same number of times that *B* measures *C*

_____ *A*
 _____ *B*
C _____
D _____
 _____ *E*

Therefore alternately, the unit *E*
measures the number *B* the same num-
ber of times that *A* measures *C* [VII 15]

Again, since *B* by multiplying *A* has
made *D*,
therefore *A* measures *D* according to the
units in *B*

But the unit *E* also measures *B* according to the units in it
therefore the unit *E* measures the number *B* the same number of times that *A*
measures *D*

But the unit *E* measured the number *B* the same number of times that *A*
measures *C*,

therefore *A* measures each of the numbers *C*, *D* the same number of times

Therefore *C* is equal to *D*

Q E D

PROPOSITION 17

If a number by multiplying two numbers make certain numbers the numbers so produced will have the same ratio as the numbers multiplied

For let the number *A* be multiplying the two numbers *B* *C* make *D* *E*

I say that as *B* is to *C*, so is *D* to *E*

For since *A* by multiplying *B* has made *D*

therefore *B* measures *D* according to the units in *A*

But the unit *F* also meas-
ures the number *A* according
to the units in it

therefore the unit *F* measures
the number *A* the same num-
ber of times that *B* measures *D*

_____ *A*
B _____ *C* _____
 _____ *D* _____ *E* _____
 _____ *F*

Therefore as the unit *F* is to the number *A* so is *B* to *D* [VII Def 20]

For the same reason

as the unit *F* is to the number *A* so also is *C* to *E*

therefore also as *B* is to *D* so is *C* to *E*

Therefore alternately as *B* is to *C* so is *D* to *E*

[VII 13]

Q E D

PROPOSITION 18

If two numbers by multiplying any number make certain numbers the numbers so produced will have the same ratio as the multipliers

For let two numbers *A* *B* by multiplying any number *C* make *D* *E*

I say that as *A* is to *B* so is *D* to *E*

C _____
 _____ *A* _____
 _____ *B* _____
 _____ *D* _____
 _____ *E* _____

For since *A* by multiplying *C* has made *D*
therefore also *C* by multiplying *A* has made *D*

[VII 16]

For the same reason also

C by multiplying *B* has made *E*

Therefore the number *C* by multiplying the two numbers *A* *B* has made *D* *E*

Therefore as *A* is to *B* so is *D* to *E*

[VII 17]

Q E D

PROPOSITION 19

If four numbers be proportional the number produced from the first and fourth will be equal to the number produced from the second and third and, if the number produced from the first and fourth be equal to that produced from the second and third the four numbers will be proportional — —

Let A, B, C, D be four numbers in proportion so that

as A is to B , so is C to D .

and let A by multiplying D make E , and let B by multiplying C make F ,
I say that E is equal to F

For let A by multiplying C make G

Since then A by multiplying C has made G and by multiplying D has made E

the number A by multiplying the two numbers C D
has made G E

Therefore, as C is to D so is G to E [vii 17]

But as C is to D , so is A to B ,

therefore also as A is to B , so is G to E

Again, since A by multiplying C has made G but further, B has also by multiplying C made F , the two numbers A, B by multiplying a certain number C have made G, F .

Therefore as A is to B , so is G to F [VII 18]

But further as *A* is to *B*, so is *G* to *E* also,

therefore also, as G is to E , so is G to F

Therefore G has to each of the numbers E, F the same ratio,
therefore E is equal to F

[cf v 9]

Again let E be equal to F

I say that as 4 is to B so is C to D

For, with the same construction

since E is equal to F

therefore as G is to L , so is G to F

[cf. V. 4]

But as G is to E so is C to D .

[VI 17]

and as G is to F so is A to B

[vii 18]

Therefore also, as A is to B so is C to D

Q E D

PROPOSITION 20

The least numbers of those which have the same ratio with them measure those which have the same ratio the same number of times the greater the greater and the less the less

For let CD EF be the least numbers of those which have the same ratio with A , B .

I say that CD measures A the same number of times that EF measures B

Now CD is not parts of A

For, if possible let it be so

therefore EF is also the same parts of B that CD is of A

[vii 13 and Def 20]

Therefore as many parts of A as there are in CD so many parts of B are there also in EF

Let CD be divided into the parts of A namely CG GD and EF into the parts of B namely EH , HF

thus the multitude of CG , GD will be equal to the multitude of EH HF

Now since the numbers CG , GD are equal to one another and the numbers EH , HF are also equal to one another while the multitude of CG , GD is equal to the multitude of EH , HF ,

therefore as CG is to EH so is GD to HF
Therefore also as one of the antecedents is to one of the consequents so will all the antecedents be to all the consequents [VII 12]

Therefore as CG is to EH , so is CD to EF

Therefore CG EH are in the same ratio with CD , EF , being less than they which is impossible, for by hypothesis CD EF are the least numbers of those which have the same ratio with them

Therefore CD is not parts of A ,

therefore it is a part of it [VII 4]

And EF is the same part of B that CD is of A [VII 13 and Def 20]

therefore CD measures A the same number of times that EF measures B

Q E D

PROPOSITION 21

Numbers prime to one another are the least of those which have the same ratio with them

Let A B be numbers prime to one another
I say that A , B are the least of those which have the same ratio with them
For if not there will be some numbers less than A B which are in the same ratio with A , B
Let them be C , D

Since then the least numbers of those which have the same ratio measure those which have the same ratio the same number of times the greater the greater and the less the less that is, the antecedent the antecedent and the consequent the consequent, [VII 20]

therefore C measures A the same number of times that D measures B

Now, as many times as C measures A so many units let there be in E

Therefore D also measures B according to the units in E

And since C measures A according to the units in E

therefore E also measures A according to the units in C [VII 16]

For the same reason

E also measures B according to the units in D [VII 16]

Therefore E measures A , B which are prime to one another which is impossible [VII Def 12]

Therefore there will be no numbers less than A , B which are in the same ratio with A B

Therefore A B are the least of those which have the same ratio with them

Q E D

PROPOSITION 22

The least numbers of those which have the same ratio with them are prime to one another

Let A, B be the least numbers of those which have the same ratio with them,

I say that A, B are prime to one another

For, if they are not prime to one another some number will measure them

Let some number measure them and let it be C

And as many times as C measures A , so many units let there be in D

and as many times as C measures B so many units let there be in E

Since C measures A according to the units in D

therefore C by multiplying D has made A [VII Def 15]

For the same reason also

C by multiplying E has made B

Thus the number C by multiplying the two numbers D, E has made A, B ,

therefore as D is to E so is A to B , [VII 17]

therefore D, E are in the same ratio with A, B being less than they which is impossible

Therefore no number will measure the numbers A, B

Therefore A, B are prime to one another

Q E D

PROPOSITION 23

If two numbers be prime to one another the number which measures the one of them will be prime to the remaining number

Let A, B be two numbers prime to one another, and let any number C measure A ,

I say that C, B are also prime to one another

For if C, B are not prime to one another

some number will measure C, B

Let a number measure them and let it be D

Since D measures C and C measures A therefore D also measures A

But it also measures B

therefore D measures A, B which are prime to one another which is impossible [VII Def 12]

Therefore no number will measure the numbers C, B

Therefore C, B are prime to one another

Q E D

PROPOSITION 24

If two numbers be prime to any number their product also will be prime to the same

For let the two numbers A, B be prime to any number C and let A by multiplying B make D

I say that C, D are prime to one another


For if C, D are not prime to one another some number will measure C, D

Let a number measure them and let it be E

Now, since C, A are prime to one another

and a certain number E measures C ,
therefore A E are prime to one another [VII 23]

As many times then, as E measures D so many units let there be in F ,



therefore F also measures

D according to the units in E [VII 16]

Therefore E by multiplying F has made D [VII Def 15]

But further, A by multiplying B has also made D ,
therefore the product of E , F is equal to the product of A , B

But if the product of the extremes be equal to that of
the means the four numbers are proportional [VII 19]

therefore as E is to A , so is B to F

But A E are prime to one another

numbers which are prime to one another are also the least of those which have
the same ratio [VII 21]

and the least numbers of those which have the same ratio with them measure
those which have the same ratio the same number of times the greater the
greater and the less the less that is the antecedent the antecedent and the
consequent the consequent [VII 20]

therefore E measures B

But it also measures C ,
therefore E measures B , C which are prime to one another which is impossible
[VII Def 12]

Therefore no number will measure the numbers C D

Therefore C , D are prime to one another Q E D

PROPOSITION 25

If two numbers be prime to one another the product of one of them into itself will
be prime to the remaining one

Let A B be two numbers prime to one another

and let A by multiplying itself make C



I say that B C are prime to one another

For let D be made equal to A

Since A B are prime to one another and A is equal to D

therefore D B are also prime to one another

Therefore each of the two numbers D , A is prime to B


therefore the product of D A will also be prime to B [VII 24]

But the number which is the product of D A is C

Therefore C B are prime to one another Q E D

PROPOSITION 26

If two numbers be prime to two numbers both to each their products also will be
prime to one another



For let the two numbers A B be prime
to the two numbers C D both to each,
and let A by multiplying B make E and let
 C by multiplying D make F

I say that E F are prime to one another

For, since each of the numbers A B is prime to C

therefore the product of A B will also be prime to C [VII 24]

But the product of A , B is E

therefore E , C are prime to one another

For the same reason

E D are also prime to one another

Therefore each of the numbers C , D is prime to E

Therefore the product of C , D will also be prime to E [VII 24]

But the product of C , D is F

Therefore E , F are prime to one another

Q E D

PROPOSITION 27

If two numbers be prime to one another and each by multiplying itself make a certain number, the products will be prime to one another and if the original numbers by multiplying the products make certain numbers the latter will also be prime to one another [and this is always the case with the extremes]

Let A , B be two numbers prime to one another

let A by multiplying itself make C , and by multiplying C make D
and let B by multiplying itself make E , and by multiplying E make F ,

I say that both C , E and D F are prime to one another

For since A B are prime to one another and A by multiplying itself has made C

therefore C , B are prime to one another [VII 25]

Since then C , B are prime to one another

and B by multiplying itself has made E

therefore C E are prime to one another [id]

Again since A B are prime to one another

and B by multiplying itself has made E ,

therefore A E are prime to one another [id]

Since then the two numbers A C are prime to the two numbers B E , both to each

therefore also the product of A C is prime to the product of B , E [VII 26]

And the product of A C is D and the product of B , E is F

Therefore D F are prime to one another

Q E D

PROPOSITION 28

If two numbers be prime to one another the sum will also be prime to each of them and if the sum of two numbers be prime to any one of them, the original numbers will also be prime to one another

For let two numbers AB BC prime to one another be added

I say that the sum AC is also prime to each of the numbers AB BC


For if CA $1B$ are not prime to one another

some number will measure CA AB

Let a number measure them and let it be D

Since then D measures CA AB

therefore it will also measure the remainder BC



But it also measures BA ,
 therefore D measures AB, BC which are prime to one another which is impossible [VII Def 12]

Therefore no number will measure the numbers CA, AB , therefore CA, AB are prime to one another

For the same reason

AC, CB are also prime to one another

Therefore CA is prime to each of the numbers AB, BC

Again let CA, AB be prime to one another,

I say that AB, BC are also prime to one another

For if AB, BC are not prime to one another,

some number will measure AB, BC

Let a number measure them and let it be D

Now since D measures each of the numbers AB, BC it will also measure the whole CA

Put it also measures AB

therefore D measures CA, AB which are prime to one another

which is impossible

[VII Def 12]

Therefore no number will measure the numbers AB, BC

Therefore AB, BC are prime to one another

Q E D

PROPOSITION 29

Any prime number is prime to any number which it does not measure

Let A be a prime number and let it not measure B

I say that B, A are prime to one another

For if B, A are not prime to one another,

some number will measure them

_____A

_____B

Let C measure them

_____C

Since C measures B

and A does not measure B

therefore C is not the same with A

Now since C measures B, A

therefore it also measures A which is prime though it is not the same with it which is impossible

Therefore no number will measure B, A

Therefore A, B are prime to one another

Q E D

PROPOSITION 30

If two numbers by multiplying one another make some number and any prime number measure the product it will also measure one of the original numbers

For let the two numbers A, B by multiplying one another make C and let any prime number D measure C

A _____

I say that D measures one of the numbers A, B

B _____

For let it not measure A

C _____

Now D is prime

D _____

therefore A, D are prime to one another [VII 29]

E _____

And as many times as D measures C so many units let there be in E

Since then D measures C according to the units in E

therefore D by multiplying E has made C [VII Def 15]
 Further A by multiplying B has also made C ,
 therefore the product of D, E is equal to the product of A, B
 Therefore as D is to A , so is B to E [VII 19]
 But D, A are prime to one another,
 primes are also least [VII 21]
 and the least measure the numbers which have the same ratio the same number of times, the greater the greater and the less the less that is the antecedent the antecedent and the consequent the consequent, [VII 20]
 therefore D measures B
 Similarly we can also show that if D does not measure B , it will measure A
 Therefore D measures one of the numbers A, B Q E D

PROPOSITION 31

Any composite number is measured by some prime number

Let A be a composite number,

I say that A is measured by some prime number

For since A is composite

some number will measure it

Let a number measure it and let it be B

Now, if B is prime, what was enjoined will have been done

A _____

B _____

C _____

But if it is composite some number will measure it

Let a number measure it, and let it be C

Then since C measures B ,

and B measures A

therefore C also measures A

And if C is prime what was enjoined will have been done

But if it is composite some number will measure it

Thus if the investigation be continued in this way some prime number will be found which will measure the number before it which will also measure A

For if it is not found an infinite series of numbers will measure the number A each of which is less than the other

which is impossible in numbers

Therefore some prime number will be found which will measure the one before it which will also measure A

Therefore any composite number is measured by some prime number

Q E D

PROPOSITION 32

Any number either is prime or is measured by some prime number

Let A be a number

I say that A either is prime or is measured by some prime number

A _____

If now A is prime that which was enjoined will have been done

But if it is composite some prime number will measure it

[VII 31]

Therefore any number either is prime or is measured by some prime number

Q E D

PROPOSITION 33

Given as many numbers as we please, to find the least of those which have the same ratio with them

Let A, B, C be the given numbers, as many as we please,

thus it is required to find the least of those which have the same ratio with A, B, C

A, B, C are either prime to one another or not

Now, if A, B, C are prime to one another, they are the least of those which have the same ratio with them [VII 21]

But if not, let D the greatest common measure of A, B, C be taken [VII 3]

and, as many times as D measures the numbers A, B, C respectively, so many units let there be in the numbers E, F, G respectively

Therefore the numbers E, F, G measure the numbers A, B, C respectively according to the units in D [VII 16]

Therefore E, F, G measure A, B, C the same number of times

therefore E, F, G are in the same ratio with A, B, C [VII Def 20]

I say next that they are the least that are in that ratio

For if E, F, G are not the least of those which have the same ratio with A, B, C , there will be numbers less than E, F, G which are in the same ratio with A, B, C

Let them be H, K, L , therefore H measures A the same number of times that the numbers K, L measure the numbers B, C respectively

Now, as many times as H measures A , so many units let there be in M

therefore the numbers K, L also measure the numbers B, C respectively according to the units in M

And since H measures A according to the units in M

therefore M also measures A according to the units in H [VII 16]

For the same reason

M also measures the numbers B, C according to the units in the numbers K, L respectively,

Therefore M measures A, B, C

Now since H measures A according to the units in M

therefore H by multiplying M has made A [VII Def 15]

For the same reason also

E by multiplying D has made A

Therefore the product of E, D is equal to the product of H, M

Therefore as E is to H so is M to D

[VII 19]

But E is greater than H

therefore M is also greater than D

And it measures A, B, C

which is impossible for by hypothesis D is the greatest common measure of A, B, C

Therefore there cannot be any numbers less than E, F, G which are in the same ratio with A, B, C

Therefore E, F, G are the least of those which have the same ratio with A, B, C Q E D

PROPOSITION 34

Given two numbers to find the least number which they measure

Let A, B be the two given numbers,

thus it is required to find the least number which they measure

Now A, B are either prime to one another or not

First let A, B be prime to one another, and let A

by multiplying B make C ,

therefore also B by multiplying A has made C

Therefore A, B measure C

I say next that it is also the least number they measure

For if not, A, B will measure some number which is less than C

Let them measure D

Then as many times as A measures D so many units let there be in E , and as many times as B measures D , so many units let there be in F ,

therefore A by multiplying E has made D

and B by multiplying F has made D [VII Def 15]

therefore the product of A, E is equal to the product of B, F

Therefore as A is to B , so is F to E [VII 19]

But A, B are prime

primes are also least [VII 21]

and the least measure the numbers which have the same ratio the same number of times the greater the greater and the less the less [VII 20]

therefore B measures E as consequent consequent

And since A by multiplying B, E has made C, D

therefore as B is to E so is C to D [VII 17]

But B measures E ,

therefore C also measures D the greater the less which is impossible

Therefore A, B do not measure any number less than C

therefore C is the least that is measured by A, B

Next, let A, B not be prime to one another

and let F, E the least numbers of those which have the same ratio with A, B , be taken, [VII 33]

therefore the product of A, E is equal to the product of B, F [VII 19]

And let A by multiplying E make C



therefore also B by multiplying F has made C ,

therefore A, B measure C

I say next that it is also the least number that they measure

For, if not A, B will measure some number which is less than C

Let them measure D



And, as many times as A measures D , so many units let there be in G
and as many times as B measures D so many units let there be in H

Therefore A by multiplying G has made D ,

and B by multiplying H has made D

Therefore the product of A, G is equal to the product of B, H ,

therefore, as A is to B , so is H to G [VII 19]

But as A is to B , so is F to E

Therefore also as F is to E so is H to G

But F, E are least

and the least measure the numbers which have the same ratio the same number of times the greater the greater and the less the less [VII 20]

therefore E measures G

And since A by multiplying E G has made C, D

therefore as E is to G , so is C to D [VII 17]

But E measures G

therefore C also measures D , the greater the less

which is impossible

Therefore A, B will not measure any number which is less than C

Therefore C is the least that is measured by A, B Q E D

PROPOSITION 35


If two numbers measure any number the least number measured by them will also measure the same

For let the two numbers A, B measure any number CD

and let E be the least that they measure

I say that E also measures CD

For if E does not measure CD let E measuring DF leave CF less than itself



Now since A, B measure E

and E measures DF ,

therefore A, B will also measure DF

But they also measure the whole CD ,

therefore they will also measure the remainder CF which is less than E

which is impossible

Therefore E cannot fail to measure CD

therefore it measures it

Q E D

PROPOSITION 36

Given three numbers to find the least number which they measure

Let A, B, C be the three given numbers,

thus it is required to find the least number which they measure

A ———

B ———

C ———

D ———

E ———

Let D the least number measured by the two numbers A, B be taken [VII 34]

Then C either measures or does not measure D

First let it measure it

But A, B also measure D

therefore A, B, C measure D

I say next that it is also the least that they measure

For, if not, A, B, C will measure some number which is less than D

Let them measure E

Since A, B, C measure E ,

therefore also A, B measure E ;

Therefore the least number measured by A, B will also measure E [VII 35]

But D is the least number measured by A, B ,

therefore D will measure E , the greater the less
which is impossible

Therefore A, B, C will not measure any number which is less than D ,

therefore D is the least that A, B, C measure

Again let C not measure D ,

and let E the least number measured by C, D , be
taken [VII 34]

Since A, B measure D

and D measures E

therefore also A, B measure E

But C also measures E

therefore also A, B, C measure E

I say next that it is also the least that they measure

For if not A, B, C will measure some number which is less than E

Let them measure F

Since A, B, C measure F

therefore also A, B measure F ,

therefore the least number measured by A, B will also measure F [VII 35]

But D is the least number measured by A, B ,

therefore D measures F

But C also measures F ,

therefore D, C measure F

so that the least number measured by D, C will also measure F —

But E is the least number measured by C, D ,

therefore E measures F the greater the less
which is impossible

Therefore A, B, C will not measure any number which is less than E

Therefore E is the least that is measured by A, B, C Q. E. D.

PROPOSITION 37

If a number be measured by any number the number which is measured will have a part called by the same name as the measuring number

For let the number A be measured by any number B



I say that A has a part called by the same name as B

For as many times as B measures A so many units let
there be in C

Since B measures A according to the units in C ,
and the unit D also measures the number C according
to the units in it

therefore the unit D measures the number C the same number of times as B
measures A

Therefore alternately the unit D measures the number B the same number
of times as C measures A [VII 15]



therefore, whatever part the unit D is of the number B , the same part is C of A also

But the unit D is a part of the number B called by the same name as it,
therefore C is also a part of A called by the same name as B ,
so that A has a part C which is called by the same name as B Q E D

PROPOSITION 38

If a number have any part whatever, it will be measured by a number called by the same name as the part

For let the number A have any part whatever, B
and let C be a number called by the same name as the part B ,
A _____ I say that C measures A
_____B For since B is a part of A called by the same name
_____C as C ,
_____D and the unit D is also a part of C called by the same
name as it
therefore whatever part the unit D is of the number C
the same part is B of A also

therefore the unit D measures the number C the same number of times that B measures A

Therefore, alternately the unit D measures the number B the same number of times that C measures A [VII 15]

Therefore C measures A Q E D

PROPOSITION 39

To find the number which is the least that will have given parts

Let A, B, C be the given parts
thus it is required to find the number which is the least that will have the parts A, B, C

Let D, E, F be numbers called by the same name as the parts A, B, C
and let G the least number measured
by D, E, F , be taken [VII 36]
Therefore G has parts called by the
same name as D, E, F [VII 37]
But A, B, C are parts called by the
same name as D, E, F
therefore G has the parts A, B, C

I say next that it is also the least number that has
For if not there will be some number less than G which will have the parts A, B, C

Let it be H
Since H has the parts A, B, C
therefore H will be measured by numbers called by the same name as the parts A, B, C [VII 38]

But D, E, F are numbers called by the same name as the parts A, B, C ,
therefore H is measured by D, E, F

And it is less than G which is impossible
Therefore there will be no number less than G that will have the parts A, B, C Q E D

BOOK EIGHT


PROPOSITION 1

If there be as many numbers as we please in continued proportion and the extremes of them be prime to one another, the numbers are the least of those which have the same ratio with them

Let there be as many numbers as we please A, B, C, D , in continued proportion and let the extremes of them A, D be prime to one another

I say that A, B, C, D are the least of those which have the same ratio with them

For if not let E, F, G, H be less than A, B, C, D and in the same ratio with them



Now since A, B, C, D are in the same ratio with E, F, G, H and the multitude of the numbers A, B, C, D is equal to the multitude of the numbers E, F, G, H ,

therefore *ex aequali*
as A is to D so is F to H [VII 14]

But A, D are prime

primes are also least [VII 21]

and the least numbers measure those which have the same ratio the same number of times the greater the greater and the less the less, that is, the antecedent the antecedent and the consequent the consequent [VII 20]

Therefore A measures E , the greater the less
which is impossible

Therefore E, F, G, H which are less than A, B, C, D are not in the same ratio with them

Therefore A, B, C, D are the least of those which have the same ratio with them Q. E. D.

PROPOSITION 2

To find numbers in continued proportion as many as may be prescribed and the least that are in a given ratio

I let the ratio of A to B be the given ratio in least numbers, thus it is required to find numbers in continued proportion as many as may be prescribed and the least that are in the ratio of A to B

Let four be prescribed

let A by multiplying itself make C and by multiplying B let it make D

let B by multiplying itself make E ,

further let A by multiplying C, D, E make F, G, H

and let B by multiplying E make A

Now, since A by multiplying itself has made C ,

and by multiplying B has made D

therefore, as A is to B , so is C to D

[VII 17]

Again since A by multiplying B has made D

and B by multiplying itself has made E

therefore the numbers A, B by multiplying B have made the numbers D, E respectively

Therefore as A is to B so is D to E ,

[VII 18]

But as A is to B , so is C to D ,

therefore also as C is to D so is D to E

And since A by multiplying C D has made F, G ,

therefore as C is to D , so is F to G

[VII 17]

But as C is to D , so was A to B ,

therefore also as A is to B , so is F to G

Again since A by multiplying D, E has made G, H ,

therefore, as D is to E , so is G to H

[VII 17]

But as D is to E , so is A to B

Therefore also as A is to B , so is G to H

And since A, B by multiplying E have made H, K ,

therefore as A is to B , so is H to K

[VII 18]

But as A is to B so is F to G and G to H

Therefore also as F is to G so is G to H and H to K ,

therefore C, D, E and F, G, H, K are proportional in the ratio of A to B

I say next that they are the least numbers that are so

For since A, B are the least of those which have the same ratio with them, and the least of those which have the same ratio are prime to one another,

[VI 22]

therefore A, B are prime to one another

And the numbers A, B by multiplying themselves respectively have made the numbers C, E and by multiplying the numbers C, E respectively have made the numbers F, K

therefore C, E and F, K are prime to one another respectively

[VII 27]

But if there be as many numbers as we please in continued proportion and the extremes of them be prime to one another they are the least of those which have the same ratio with them

[VIII 1]

Therefore C, D, E and F, G, H, K are the least of those which have the same ratio with A, B


Q E D

PROISM From this it is manifest that, if three numbers in continued proportion be the least of those which have the same ratio with them the extremes of them are squares and if four numbers, cubes

PROPOSITION 3

If as many numbers as we please in continued proportion be the least of those which have the same ratio with them the extremes of them are prime to one another

Let as many numbers as we please A, B, C, D in continued proportion be the least of those which have the same ratio with them,



I say that the extremes of them A, D are prime to one another

For let two numbers E, F , the least that are in the ratio of A, B, C, D be taken, [VII 33]

then three others G, H, K with the same property,

and others more by one continually [VIII 2]

until the multitude taken becomes equal to the multitude of the numbers A, B, C, D

Let them be taken and let them be L, M, N, O

Now, since E, F are the least of those which have the same ratio with them they are prime to one another [VII 29]

And since the numbers E, F by multiplying themselves respectively have made the numbers G, K and by multiplying the numbers G, K respectively have made the numbers L, O [VIII 2 Por]

therefore both G, K and L, O are prime to one another [VII 27]

And since A, B, C, D are the least of those which have the same ratio with them

while L, M, N, O are the least that are in the same ratio with A, B, C, D and the multitude of the numbers A, B, C, D is equal to the multitude of the numbers L, M, N, O

therefore the numbers A, B, C, D are equal to the numbers L, M, N, O respectively

therefore A is equal to L and D to O

And L, O are prime to one another

Therefore A, D are also prime to one another

Q E D

PROPOSITION 4

Given as many ratios as we please in least numbers to find numbers in continued proportion which are the least in the given ratios

Let the given ratios in least numbers be that of A to B , that of C to D and that of E to F

thus it is required to find numbers in continued proportion which are the least that are in the ratio of A to B in the ratio of C to D and in the ratio of E to F


Let G the least number measured by B, C be taken [VII 34]

And as many times as B measures G so many times also let A measure H , and as many times as C measures G so many times also let D measure K

Now E either measures or does not measure K

First let it measure it

And as many times as E measures K , so many times let F measure L also
 Now since A measures H the same number of times that B measures G
 therefore as A is to B , so is H to G [VII Def 20, VII 13]



For the same reason also

as C is to D so is G to K

and further as E is to F , so is K to L ,

therefore H, G, K, L are continuously proportional in the ratio of A to B , in the ratio of C to D and in the ratio of E to F

I say next that they are also the least that have this property

For, if H, G, K, L are not the least numbers continuously proportional in the ratios of A to B , of C to D and of E to F , let them be N, O, M, P

Then since, as A is to B , so is N to O

while A, B are least

and the least numbers measure those which have the same ratio the same number of times the greater the greater and the less the less that is the antecedent the antecedent and the consequent the consequent,

therefore B measures O

[VII 20]

For the same reason

C also measures O ,

therefore B, C measure O

therefore the least number measured by B, C will also measure O [VII 35]


But G is the least number measured by B, C ,

therefore G measures O the greater the less

which is impossible

Therefore there will be no numbers less than H, G, K, L which are continuously in the ratio of A to B , of C to D and of E to F

Next let E not measure K



Let M the least number measured by E, K be taken

And as many times as K measures M so many times let H, G measure N, O respectively

and as many times as E measures M so many times let F measure P also
 Since H measures N the same number of times that G measures O ,
 therefore, as H is to G , so is N to O [VII 13 and Def 20]
 But as H is to G , so is A to B ,
 therefore also as A is to B , so is N to O
 For the same reason also
 as C is to D , so is O to M
 Again since E measures M the same number of times that F measures P ,
 therefore as L is to F so is M to P [VII 13 and Def 20]
 therefore N, O, M, P are continuously proportional in the ratios of A to B , of
 C to D and of E to F
 I say next that they are also the least that are in the ratios A, B, C, D, E, F
 For if not there will be some numbers less than N, O, M, P continuously
 proportional in the ratios A, B, C, D, E, F
 Let them be Q, R, S, T
 Now since as Q is to R , so is A to B
 while A, B are least
 and the least numbers measure those which have the same ratio with them the
 same number of times the antecedent the antecedent and the consequent the
 consequent, [VII 20]

 therefore B measures R
 For the same reason C also measures R ,
 therefore B, C measure R
 Therefore the least number measured by B, C will also measure R [VII 35]
 But G is the least number measured by B, C ,
 therefore G measures R
 And as G is to R so is A to S [VII 13]
 therefore A also measures S
 But E also measures S
 therefore E, A measure S
 Therefore the least number measured by E, A will also measure S [VII 35]
 But M is the least number measured by E, A ,
 therefore M measures S the greater the less
 which is impossible

Therefore there will not be any numbers less than N, O, M, P continuously
 proportional in the ratios of A to B of C to D and of E to F ,
 therefore N, O, M, P are the least numbers continuously proportional in the
 ratios A, B, C, D, E, F Q E D

PROPOSITION 5

Plane numbers have to one another the ratio compounded of the ratios of their sides

Let A, B be plane numbers and let the numbers C, D be the sides of A and
 E, F of B

I say that A has to B the ratio compounded of the ratios of the sides

For the ratios being given which C has to E and D to F , let the least num-
 bers G, H, K that are continuously in the ratios C, E, D, F be taken so that
 as C is to E so is G to H

and as D is to F so is H to K [VIII 4]

And let D by multiplying E make L

Now since D by multiplying C has made A , and by multiplying E has made L ,

therefore as C is to E , so is A to L [VI 17]

But as C is to E so is G to H ,

therefore also as G is to H , so is A to L

Again since E by multiplying D has made L , and further by multiplying F has made B

therefore, as D is to F so is L to B [VII 17]

But as D is to F , so is H to K ,

therefore also as H is to K , so is L to B

But it was also proved that

as G is to H , so is A to L ,

therefore *ex aequali*,

as G is to K so is A to B [VII 14]

But G has to K the ratio compounded of the ratios of the sides

therefore A also has to B the ratio compounded of the ratios of the sides

Q E D

PROPOSITION 6

If there be as many numbers as we please in continued proportion, and the first do not measure the second neither will any other measure any other

Let there be as many numbers as we please A, B, C, D, E in continued proportion and let A not measure B ,

I say that neither will any other measure any other

Now it is manifest that A, B, C, D, E do not measure one another in order for A does not even measure B

I say then that neither will any other measure any other

For if possible let A measure C

And however many A, B, C are let as many numbers F, G, H the least of those which have

the same ratio with A, B, C be taken [VII 33]

Now, since F, G, H are in the same ratio with A, B, C , and the multitude of the numbers A, B, C is equal to the multitude of the numbers F, G, H

therefore, *ex aequali* as A is to C so is F to H [VII 14]

And since, as A is to B so is F to G

while A does not measure B ,

therefore neither does F measure G , [VII Def 20]

therefore F is not an unit for the unit measures any number

Now F, H are prime to one another [VIII 3]

And as F is to H so is A to C

therefore neither does A measure C

Similarly we can prove that neither will any other measure any other

Q E D

PROPOSITION 7


If there be as many numbers as we please in continued proportion, and the first measure the last it will measure the second also

Let there be as many numbers as we please, A, B, C, D , in continued proportion, and let A measure D ,
I say that A also measures B

For if A does not measure B , neither will any other of the numbers measure any other [VIII 6]

But A measures D

Therefore A also measures B




PROPOSITION 8

If between two numbers there fall numbers in continued proportion with them then however many numbers fall between them in continued proportion so many will also fall in continued proportion between the numbers which have the same ratio with the original numbers

Let the numbers C, D fall between the two numbers A, B in continued proportion with them, and let E be made in the same ratio to F as A is to B ,

I say that as many numbers as have fallen between A, B in continued proportion so many will also fall between E, F in continued proportion

For as many as A, B, C, D are in multitude let so many numbers G, H, K, L the least of those which have the same ratio with A, C, D, B , be taken [VII 33]
therefore the extremes of them G, L are prime to one another [VIII 3]



Now since A, C, D, B are in the same ratio with G, H, K, L and the multitude of the numbers A, C, D, B is equal to the multitude of the numbers G, H, K, L

therefore *ex aequali* as A is to B so is G to L [VII 14]

But, as A is to B so is E to F

therefore also as G is to L so is E to F

But G, L are prime,

primes are also least [VII 21]

and the least numbers measure those which have the same ratio the same number of times the greater the greater and the less the less that is the antecedent the antecedent and the consequent the consequent [VII 20]

Therefore G measures E the same number of times as L measures F

Next as many times as G measures E so many times let H, K also measure M, N respectively

therefore G, H, K, L measure E, M, N, F the same number of times

Therefore G, H, K, L are in the same ratio with E, M, N, F [VII Def 20]

But G, H, K, L are in the same ratio with A, C, D, B

therefore A, C, D, B are also in the same ratio with E, M, N, F

But A, C, D, B are in continued proportion

therefore E, M, N, F are also in continued proportion

Therefore, as many numbers as have fallen between A, B in continued proportion with them, so many numbers have also fallen between E, F in continued proportion

[VIII 2]

PROPOSITION 9

If two numbers be prime to one another and numbers fall between them in continued proportion, then however many numbers fall between them in continued proportion so many will also fall between each of them and an unit in continued proportion

Let A, B be two numbers prime to one another and let C, D fall between them in continued proportion


and let the unit E be set out,

I say that as many numbers as fall between A, B in continued proportion so many will also fall between either of the numbers A, B and the unit in continued proportion

For let two numbers F, G , the least that are in the ratio of A, C, D, B be taken

three numbers H, K, L with the same property, and others more by one continually until their multitude is equal to the multitude of A, C, D, B

[VIII 2]



Let them be taken and let them be M, N, O, P

It is now manifest that F by multiplying itself has made H and by multiplying H has made M while G by multiplying itself has made L and by multiplying L has made P

[VIII 2, Por]

And since M, N, O, P are the least of those which have the same ratio with F, G

and A, C, D, B are also the least of those which have the same ratio with F, G

[VIII 1]

while the multitude of the numbers M, N, O, P is equal to the multitude of the numbers A, C, D, B

therefore M, N, O, P are equal to A, C, D, B respectively,

therefore M is equal to A , and P to B

Now since F by multiplying itself has made H

therefore F measures H according to the units in F

But the unit E also measures F according to the units in it therefore the unit E measures the number F the same number of times as F measures H

Therefore as the unit I is to the number F so is I to H [VII Def 20]

Again, since F by multiplying H has made M ,

therefore H measures M according to the units in F

But the unit L also measures the number F according to the units in it, therefore the unit E measures the number F the same number of times as H measures M

Therefore, as the unit E is to the number F , so is H to M

But it was also proved that as the unit L is to the number F so is F to H , therefore also as the unit E is to the number F , so is F to H , and H to M

But M is equal to A

therefore, as the unit E is to the number F , so is F to H and H to A

For the same reason also

as the unit L is to the number G so is G to L and L to B

Therefore as many numbers as have fallen between A, B in continued proportion so many numbers also have fallen between each of the numbers A, B and the unit E in continued proportion Q E D

PROPOSITION 10

If numbers fall between each of two numbers and an unit in continued proportion however many numbers fall between each of them and an unit in continued proportion so many also will fall between the numbers themselves in continued proportion

For let the numbers D, E and F, G respectively fall between the two numbers A, B and the unit C in continued proportion

I say that as many numbers as have fallen between each of the numbers A, B and the unit C in continued proportion so many numbers will also fall between A, B in continued proportion

For let D by multiplying I make H and let the numbers D, F by multiplying H make K, L respectively

Now since as the unit C is to the number D so is D to E

therefore the unit C measures the number D the same number of times as D measures E [VII Def 20]

But the unit C measures the number D according to the units in D


therefore the number D also measures E according to the units in D , therefore D by multiplying itself has made E

Again since as C is to the number D so is E to I therefore the unit C measures the number D the same number of times as E measures A

But the unit C measures the number D according to the units in D , therefore E also measures A according to the units in D therefore D by multiplying E has made A

For the same reason also

F by multiplying itself has made G and by multiplying G has made B



And since D by multiplying itself has made E and by multiplying F has made H ,

therefore as D is to F , so is E to H [VII 17]

For the same reason also

as D is to F , so is H to G [VII 18]

Therefore also as E is to H so is H to G

Again, since D by multiplying the numbers E H has made A Λ respectively,

therefore as E is to H , so is A to Λ [VII 17]

But, as E is to H , so is D to F ,

therefore also, as D is to F so is A to K

Again since the numbers D F by multiplying H have made Λ , L respectively,

therefore, as D is to F , so is Λ to L [VII 18]

But as D is to F , so is A to Λ

therefore also as A is to K , so is K to L

Further, since F by multiplying the numbers H G has made L , B respectively

therefore as H is to G , so is L to B , [VII 17]

But, as H is to G so is D to F ,

therefore also as D is to F , so is L to B

But it was also proved that,

as D is to F so is A to Λ and Λ to L

therefore also as A is to K , so is K to L and L to B

Therefore A Λ , L , B are in continued proportion

Therefore as many numbers as fall between each of the numbers A , B and the unit C in continued proportion so many also will fall between A , B in continued proportion

Q E D

PROPOSITION 11

Between two square numbers there is one mean proportional number and the square has to the square the ratio duplicate of that which the side has to the side

Let A , B be square numbers

and let C be the side of A and D of B

I say that between A B there is one mean proportional number and A has to B the ratio duplicate of that which C has to D

A _____

For let C by multiplying D make E

B _____

Now since A is a square and C is its side

C _____ D _____

therefore C by multiplying itself has made A

E _____

For the same reason also

D by multiplying itself has made B

Since then C by multiplying the numbers C D has made A E respectively

therefore as C is to D , so is A to E [VII 17]

For the same reason also

as C is to D so is E to B

[VII 18]

Therefore also as A is to E so is E to B

Therefore between A B there is one mean proportional number

I say next that A also has to B the ratio duplicate of that which C has to D

For since A E B are three numbers in proportion

therefore A has to B the ratio duplicate of that which A has to E [v Def 9]

But as A is to E , so is C to D

Therefore A has to B the ratio duplicate of that which the side C has to D
Q E D

PROPOSITION 12


Between two cube numbers there are two mean proportional numbers, and the cube has to the cube the ratio triplicate of that which the side has to the side

Let A, B be cube numbers,

and let C be the side of A , and D of B ,

I say that between A, B there are two mean proportional numbers and A has to B the ratio triplicate of that which C has to D

For let C by multiplying itself make E and by multiplying D let it make F
let D by multiplying itself make G



and let the numbers C, D by multiplying F make H, K respectively

Now since A is a cube and C its side

and C by multiplying itself has made E ,

therefore C by multiplying itself has made E and by multiplying E has made A

For the same reason also

D by multiplying itself has made G and by multiplying G has made B

And, since C by multiplying the numbers C, D has made E, F respectively

therefore, as C is to D so is E to F [vii 17]

For the same reason also,

as C is to D so is F to G [vii 18]

Again since C by multiplying the numbers E, F has made A, H respectively

therefore as E is to F , so is A to H [vii 17]

But as E is to F so is C to D

Therefore also as C is to D , so is A to H

Again since the numbers C, D by multiplying F have made H, K respectively,

therefore as C is to D , so is H to K [vii 18]

Again since D by multiplying each of the numbers F, G has made K, B respectively

therefore as F is to G , so is K to B [vii 17]

But, as F is to G , so is C to D

therefore also as C is to D , so is A to H , H to K and K to B

Therefore H, K are two mean proportionals between A, B

I say next that A also has to B the ratio triplicate of that which C has to D

For since A, H, K, B are four numbers in proportion, therefore A has to B the ratio triplicate of that which A has to H [v Def 10]

But, as A is to H so is C to D ,

therefore A also has to B the ratio triplicate of that which C has to D
Q E D

PROPOSITION 13

If there be as many numbers as we please in continued proportion, and each by multiplying itself make some number, the products will be proportional and, if the original numbers by multiplying the products make certain numbers the latter will also be proportional

Let there be as many numbers as we please, A, B, C , in continued proportion so that as A is to B so is B to C , let A, B, C by multiplying themselves make D, E, F , and by multiplying D, E, F let them make G, H, K

I say that D, E, F and G, H, K are in continued proportion

A _____	G _____
B _____	H _____
C _____	K _____
D _____	
E _____	M _____
F _____	N _____
L _____	P _____
O _____	Q _____

For let A by multiplying B make L

and let the numbers A, B by multiplying L make M, N respectively

And again let B by multiplying C make O ,

and let the numbers B, C by multiplying O make P, Q respectively

Then in manner similar to the foregoing we can prove that

D, L, E and G, M, N, H are continuously proportional in the ratio of A to B and further E, O, F and H, P, Q, K are continuously proportional in the ratio of B to C

Now, as A is to B , so is B to C ,

therefore D, L, E are also in the same ratio with E, O, F ,

and further G, M, N, H in the same ratio with H, P, Q, K

And the multitude of D, L, E is equal to the multitude of E, O, F and that of G, M, N, H to that of H, P, Q, K ,

therefore *ex aequali*

as D is to E so is E to F

and as G is to H , so is H to K [VII 14]

Q, E, D

PROPOSITION 14

If a square measure a square the side will also measure the side and if the side measure the side the square will also measure the square

Let A, B be square numbers let C, D be their sides and let A measure B

I say that C also measures D

A _____

B _____

C _____ D _____

E _____

For let C by multiplying D make E ,

therefore A, E, B are continuously proportional in the ratio of C to D [VIII 11]

And since A, E, B are continuously proportional and A measures B

therefore A also measures E

[VIII 7]

therefore A has to B the ratio duplicate of that which A has to E [v Def 9]

But, as A is to E , so is C to D

Therefore A has to B the ratio duplicate of that which the side C has to D
Q E D

PROPOSITION 12

Between two cube numbers there are two mean proportional numbers, and the cube has to the cube the ratio triplicate of that which the side has to the side

Let A, B be cube numbers,

and let C be the side of A , and D of B ,

I say that between A, B there are two mean proportional numbers and A has to B the ratio triplicate of that which C has to D

For let C by multiplying

itself make E and by multi


plying D let it make F ,

let D by multiplying itself

make G

and let the numbers C, D by

multiplying F make H, K respectively



Now, since A is a cube and C its side

and C by multiplying itself has made E ,

therefore C by multiplying itself has made E and by multiplying E has made A

For the same reason also

D by multiplying itself has made G and by multiplying G has made B

And since C by multiplying the numbers C, D has made E, F respectively
therefore, as C is to D so is E to F [vii 17]

For the same reason also

as C is to D so is F to G [vii 18]

Again since C by multiplying the numbers E, F has made A, H respectively
therefore as E is to F , so is A to H [vii 17]

But as E is to F , so is C to D

Therefore also as C is to D , so is A to H

Again since the numbers C, D by multiplying F have made H, K respectively

therefore, as C is to D , so is H to K [vii 18]

Again since D by multiplying each of the numbers F, G has made K, B respectively

therefore, as F is to G , so is K to B [vii 17]

But as F is to G so is C to D ,

therefore also as C is to D so is A to H H to K , and K to B

Therefore H, K are two mean proportionals between A, B

I say next that A also has to B the ratio triplicate of that which C has to D

For since A, H, K, B are four numbers in proportion

therefore A has to B the ratio triplicate of that which A has to H [v Def 10]

But, as A is to H so is C to D

therefore A also has to B the ratio triplicate of that which C has to D
Q E D

PROPOSITION 13

If there be as many numbers as we please in continued proportion, and each by multiplying itself make some number, the products will be proportional and if the original numbers by multiplying the products make certain numbers the latter will also be proportional

Let there be as many numbers as we please, A, B, C , in continued proportion, so that as A is to B , so is B to C ,
let A, B, C by multiplying themselves make D, E, F , and by multiplying D, E, F let them make G, H, K

I say that D, E, F and G, H, K are in continued proportion

A ———	G —————
B ———	H —————
C ———	K —————
D ———	
E ———	M —————
F ———	N —————
L ———	P —————
O ———	Q —————

For let A by multiplying B make L

and let the numbers A, B by multiplying L make M, N respectively

And again let B by multiplying C make O ,

and let the numbers B, C by multiplying O make P, Q respectively

Then in manner similar to the foregoing we can prove that

D, L, E and G, M, N, H are continuously proportional in the ratio of A to B
and further E, O, F and H, P, Q, K are continuously proportional in the ratio of B to C

Now, as A is to B , so is B to C ,

therefore D, L, E are also in the same ratio with E, O, F ,

and further G, M, N, H in the same ratio with H, P, Q, K

And the multitude of D, L, E is equal to the multitude of E, O, F and that of G, M, N, H to that of H, P, Q, K ,

therefore *ex aequali*

as D is to E so is E to F

and,

as G is to H so is H to K

[VII 14]

Q, E, D

PROPOSITION 14

If a square measure a square the side will also measure the side and if the side measure the side the square will also measure the square

Let A, B be square numbers, let C, D be their sides, and let A measure B ,

I say that C also measures D

A ———

B —————

— C ——— D ———

E ———

For let C by multiplying D make E

therefore A, E, B are continuously proportional in the ratio of C to D

[VIII 11]

And since A, E, B are continuously proportional and A measures B ,

therefore A also measures E

[VIII 7]

And as A is to E , so is C to D ,

therefore also C measures D

[VII Def 20]

Again, let C measure D ,

I say that A also measures B

For, with the same construction we can in a similar manner prove that A , E , B are continuously proportional in the ratio of C to D

And since as C is to D , so is A to E ,

and C measures D

therefore A also measures E

[VII Def 20]

And A , E , B are continuously proportional,

therefore A also measures B

Therefore etc

Q E D

PROPOSITION 15

If a cube number measure a cube number the side will also measure the side and, if the side measure the side the cube will also measure the cube

For let the cube number A measure the cube B ,

and let C be the side of A and D of B ,

I say that C measures D

For let C by multiplying itself make E

and let D by multiplying itself make G

further let C by multiplying D make F

and let C , D by multiplying F make H , K respectively

Now it is manifest that E , F , G and A , H , K , B are continuously proportional in the ratio of C to D

[VIII 11 12]

And since A , H , K , B are continuously proportional,

and A measures B ,

therefore it also measures H

[VIII 7]

And as A is to H so is C to D

therefore C also measures D

[VII Def 20]

Next let C measure D ,

I say that A will also measure B

For with the same construction, we can prove in a similar manner that A , H , K , B are continuously proportional in the ratio of C to D

And since C measures D

and as C is to D so is A to H ,

therefore A also measures H

[VII Def 20]

so that A measures B also

Q E D

PROPOSITION 16

If a square number do not measure a square number, neither will the side measure the side and if the side do not measure the side, neither will the square measure the square

Let A , B be square numbers and let C , D be their sides and let A not measure B ,

A _____ I say that neither does C measure D
 B _____ For if C measures D A will also measure B [VIII 14]
 C _____ But A does not measure B
 D _____ therefore neither will C measure D
 Again let C not measure D ,
 I say that neither will A measure B
 For if A measures B C will also measure D [VIII 14]
 But C does not measure D
 therefore neither will A measure B Q E D

PROPOSITION 17

If a cube number do not measure a cube number neither will the side measure the side and if the side do not measure the side, neither will the cube measure the cube

For let the cube number A not measure the cube number B ,
 and let C be the side of A and D of B ,
 I say that C will not measure D
 For if C measures D , A will also measure B [VIII 15]
 But A does not measure B ,
 therefore neither does C measure D

Again let C not measure D ,
 I say that neither will A measure B
 For if A measures B C will also measure D [VIII 15]
 But C does not measure D
 therefore neither will A measure B Q E D

PROPOSITION 18

Between two similar plane numbers there is one mean proportional number and the plane number has to the plane number the ratio duplicate of that which the corresponding side has to the corresponding side

Let A , B be two similar plane numbers and let the numbers C , D be the sides of A and E , F of B

A _____	C _____
B _____	D _____
G _____	E _____
	F _____

Now since similar plane numbers are those which have their sides proportional [VII Def 21]

therefore as C is to D so is E to F

I say then that between A , B there is one mean proportional number and A has to B the ratio duplicate of that which C has to E or D to F , that is of that which the corresponding side has to the corresponding side

Now since as C is to D so is E to F

therefore alternately as C is to E so is D to F [VII 13]

And since A is plane and C , D are its sides

therefore D by multiplying C has made A

For the same reason also

E by multiplying *F* has made *B*

Now let *D* by multiplying *L* make *G*

Then since *D* by multiplying *C* has made *A*, and by multiplying *E* has made *G*

therefore, as *C* is to *E*, so is *A* to *G* [VII 17]

But, as *C* is to *E*, so is *D* to *F*,

therefore also as *D* is to *F*, so is *A* to *G*

Again since *E* by multiplying *D* has made *G*, and by multiplying *F* has made *B*,

therefore as *D* is to *F* so is *G* to *B* [VII 17]

But it was also proved that

as *D* is to *F*, so is *A* to *G*,

therefore also as *A* is to *G* so is *G* to *B*

Therefore *A*, *G*, *B* are in continued proportion

Therefore between *A*, *B* there is one mean proportional number

I say next that *A* also has to *B* the ratio duplicate of that which the corresponding side has to the corresponding side that is of that which *C* has to *E* or *D* to *F*

For since *A*, *G*, *B* are in continued proportion

A has to *B* the ratio duplicate of that which it has to *G* [V Def 9]

And as *A* is to *G* so is *C* to *E* and so is *D* to *F*

Therefore *A* also has to *B* the ratio duplicate of that which *C* has to *E* or *D* to *F* Q E D

PROPOSITION 19

Between two similar solid numbers there fall two mean proportional numbers and the solid number has to the similar solid number the ratio triplicate of that which the corresponding side has to the corresponding side


Let *A*, *B* be two similar solid numbers and let *C*, *D*, *E* be the sides of *A* and *F*, *G*, *H* of *B*

Now since similar solid numbers are those which have their sides proportional [VII Def 21]

therefore as *C* is to *D* so is *F* to *G*

and as *D* is to *E* so is *G* to *H*

I say that between *A*, *B* there fall two mean proportional numbers, and *A* has to *B* the ratio triplicate of that which *C* has to *F*, *D* to *G*, and also *E* to *H*



For let *C* by multiplying *D* make *K* and let *F* by multiplying *G* make *L*

Now since *C*, *D* are in the same ratio with *F*, *G*

and *K* is the product of *C*, *D*, and *L* the product of *F*, *G*, *K*, *L* are similar plane numbers, [VII Def 21]

therefore between A, L there is one mean proportional number [VIII 18]

Let it be M

Therefore M is the product of D, F , as was proved in the theorem preceding this [VIII 18]

Now, since D by multiplying C has made K , and by multiplying F has made M ,

therefore as C is to F so is K to M [VII 17]

But, as K is to M so is M to L

Therefore K, M, L are continuously proportional in the ratio of C to F

And since as C is to D so is F to G

alternately therefore, as C is to F , so is D to G [VII 13]

For the same reason also

as D is to G so is E to H

Therefore K, M, L are continuously proportional in the ratio of C to F in the ratio of D to G and also in the ratio of E to H

Next let E, H by multiplying M make N, O respectively

Now, since A is a solid number and C, D, E are its sides

therefore E by multiplying the product of C, D has made A

But the product of C, D is K ,

therefore E by multiplying K has made A

For the same reason also

H by multiplying L has made B

Now since E by multiplying K has made A and further also by multiplying M has made N ,

therefore as K is to M , so is A to N [VII 17]

But, as K is to M so is C to F, D to G and also E to H ,

therefore also as C is to F, D to G , and E to H , so is A to N

Again, since E, H by multiplying M have made N, O respectively

therefore as E is to H , so is N to O [VII 18]

But, as E is to H so is C to F and D to G ,

therefore also, as C is to F, D to G and E to H so is A to N and N to O

Again since H by multiplying M has made O and further also by multiplying L has made B ,

therefore as M is to L so is O to B [VII 17]

But as M is to L so is C to F, D to G and E to H

Therefore also as C is to F, D to G and E to H so not only is O to B but also A to N and N to O

Therefore A, N, O, B are continuously proportional in the aforesaid ratios of the sides

I say that A also has to B the ratio triplicate of that which the corresponding side has to the corresponding side that is of the ratio which the number C has to F or D to G and also L to H

For since A, N, O, B are four numbers in continued proportion therefore A has to B the ratio triplicate of that which A has to N [v Def 10]

But as A is to N so it was proved that C is to F, D to G and also E to H

Therefore A also has to B the ratio triplicate of that which the corresponding side has to the corresponding side that is of the ratio which the number C has to F, D to G and also E to H

PROPOSITION 20

If one mean proportional number fall between two numbers the numbers will be similar plane numbers

For let one mean proportional number C fall between the two numbers A, B ,

I say that A, B are similar plane numbers

Let D, E , the least numbers of those which have the same ratio with A, C be taken, [VII 33]

therefore D measures A the same number of times that E measures C [VII 20]


Now, as many times as D measures A so many units let there be in F ,

therefore F by multiplying D has made A

so that A is plane and D, F are its sides

Again since D, E are the least of the numbers which have the same ratio with C, B

therefore D measures C the same number of times that E measures B [VII 20]



As many times then as E measures B so many units let there be in G

therefore G measures B according to the units in G ,

therefore G by multiplying E has made B

Therefore B is plane and E, G are its sides

Therefore A, B are plane numbers

I say next that they are also similar

For since F by multiplying D has made A and by multiplying E has made C

therefore as D is to E so is A to C that is C to B [VII 17]

Again since E by multiplying F, G has made C, B respectively,

therefore as F is to G , so is C to B [VII 17]

But as C is to B so is D to E

therefore also as D is to E so is F to G

And alternately as D is to F so is E to G [VII 13]

Therefore A, B are similar plane numbers for their sides are proportional
Q E D

PROPOSITION 21

If two mean proportional numbers fall between two numbers the numbers are similar solid numbers

For let two mean proportional numbers C, D fall between the two numbers A, B ,

I say that A, B are similar solid numbers


For let three numbers E, F, G the least of those which have the same ratio with A, C, D be taken [VII 33 or VIII 2]

therefore the extremes of them E, G are prime to one another [VIII 3]

Now, since one mean proportional number F has fallen between E, G ,
therefore E, G are similar plane numbers [VIII 20]

Let then H, K be the sides of E and L, M of G

Therefore it is manifest from the theorem before this that E, F, G are continuously proportional in the ratio of H to L and that of K to M



Now since E, F, G are the least of the numbers which have the same ratio with A, C, D ,
and the multitude of the numbers E, F, G is equal to the multitude of the numbers A, C, D ,

therefore *ex aequali* as E is to G so is A to D [VII 14]

But E, G are prime

primes are also least [VII 21]

and the least measure those which have the same ratio with them the same number of times the greater the greater and the less the less that is, the antecedent the antecedent and the consequent the consequent [VII 20]

therefore E measures A the same number of times that G measures D

Now, as many times as E measures A , so many units let there be in N

Therefore N by multiplying E has made A

But E is the product of H, K

therefore N by multiplying the product of H, K has made A

Therefore A is solid and H, K, N are its sides

Again since E, F, G are the least of the numbers which have the same ratio as C, D, B

therefore E measures C the same number of times that G measures B

Now as many times as E measures C so many units let there be in O

Therefore G measures B according to the units in O

therefore O by multiplying G has made B

But G is the product of L, M

therefore O by multiplying the product of L, M has made B

Therefore B is solid and L, M, O are its sides

therefore A, B are solid

I say that they are also similar

For since N, O by multiplying E have made A, C

therefore as N is to O so is A to C that is E to F [VII 18]

But as E is to F so is H to L and K to M

therefore also as H is to L so is K to M and N to O

And H, K, N are the sides of A and O, L, M the sides of B

Therefore A, B are similar solid numbers

Q E D

PROPOSITION 22

If three numbers be in continued proportion, and the first be square, the third will also be square

Let A, B, C be three numbers in continued proportion, and let A the first be square

I say that C the third is also square

For since between A, C there is one mean proportional number, B

therefore A, C are similar plane numbers [VIII 20]

But A is square,

therefore C is also square

Q E D

PROPOSITION 23

If four numbers be in continued proportion and the first be cube the fourth will also be cube

Let A, B, C, D be four numbers in continued proportion and let A be cube,

I say that D is also cube

For, since between A, D there are two mean proportional numbers B, C

therefore A, D are similar solid numbers

[VIII 21]

But A is cube

therefore D is also cube

Q E D

PROPOSITION 24

If two numbers have to one another the ratio which a square number has to a square number and the first be square the second will also be square

For let the two numbers A, B have to one another the ratio which the square number C has to the square number D and let A be square

I say that B is also square

For since C, D are square

C, D are similar plane numbers

Therefore one mean proportional number falls between C, D [VIII 18]

And as C is to D , so is A to B

therefore one mean proportional number falls between A, B also [VIII 8]

And A is square

therefore B is also square

[VIII 27]
Q E D

PROPOSITION 25

If two numbers have to one another the ratio which a cube number has to a cube number and the first be cube the second will also be cube

For let the two numbers A, B have to one another the ratio which the cube number C has to the cube number D and let A be cube


I say that B is also cube

For since C, D are cube

C, D are similar solid numbers

Therefore two mean proportional numbers fall between C, D [VIII 19]

And, as many numbers as fall between C, D in continued proportion, so many will also fall between those which have the same ratio with them, [VIII 8] so that two mean proportional numbers fall between A, B also



Let E, F so fall

Since, then, the four numbers A, E, F, B are in continued proportion and A is cube

therefore B is also cube

[VIII 23]


Q E D

PROPOSITION 26

Similar plane numbers have to one another the ratio which a square number has to a square number

Let A, B be similar plane numbers,

I say that A has to B the ratio which a square number has to a square number



For since A, B are similar plane numbers

therefore one mean proportional number falls between A, B [VIII 18]

Let it so fall and let it be C ,

and let D, E, F , the least numbers of those which have the same ratio with A, C, B , be taken [VII 33 or VIII 2]

therefore the extremes of them D, F are square [VIII 2 Por]

And since as D is to F , so is A to B ,

and D, F are square

therefore A has to B the ratio which a square number has to a square number


Q E D

PROPOSITION 27

Similar solid numbers have to one another the ratio which a cube number has to a cube number

Let A, B be similar solid numbers

I say that A has to B the ratio which a cube number has to a cube number



For since A, B are similar solid numbers

therefore two mean proportional numbers fall between A, B

[VIII 19]

Let C D so fall
 and let E F G H the least numbers of those which have the same ratio with
 A , C , D , B , and equal with them in multitude be taken, [VII 33 or VIII 2]
 therefore the extremes of them E , H are cube [VIII 2, Por]

And as E is to H , so is A to B ,
 therefore A also has to B the ratio which a cube number has to a cube number

—

— Q E D

BOOK NINE

PROPOSITION 1

If two similar plane numbers by multiplying one another make some number, the product will be square

Let A B be two similar plane numbers and let A by multiplying B make C ,
I say that C is square

A _____
For let A by multiplying itself make D

B _____
Therefore D is square

C _____
Since then A by multiplying itself has made

D _____
 D and by multiplying B has made C

therefore as A is to B , so is D to C [VII 17]

And since A B are similar plane numbers
therefore one mean proportional number falls between A B [VIII 18]

But if numbers fall between two numbers in continued proportion as many
as fall between them so many also fall between those which have the same
ratio, [VIII 8]

so that one mean proportional number falls between D C also

And D is square

therefore C is also square [VIII 22]

Q E D

PROPOSITION 2

If two numbers by multiplying one another make a square number they are similar plane numbers

Let A , B be two numbers and let A by multiplying B make the square number C

A _____
I say that A B are similar plane numbers

B _____
For let A by multiplying itself make D ,

C _____
therefore D is square

D _____
Now since A by multiplying itself has

made D and by multiplying B has made C

therefore as A is to B , so is D to C [VII 17]

And since D is square and C is so also

therefore D C are similar plane numbers

Therefore one mean proportional number falls between D C [VIII 18]

And as D is to C so is A to B

therefore one mean proportional number falls between A B also [VIII 8]

But if one mean proportional number fall between two numbers they are
similar plane numbers [VIII 20]

therefore A B are similar plane numbers

Q E D

PROPOSITION 3

If a cube number by multiplying itself make some number, the product will be cube

For let the cube number A by multiplying itself make B ,

I say that B is cube

For let C , the side of A , be taken and let C by multiplying itself make D

It is then manifest that C by multiplying D has made A

Now since C by multiplying itself has made D ,

therefore C measures D according to the units in itself

But further the unit also measures C according to the units in it



therefore as the unit is to C so is C to D [VII Def 20]

Again, since C by multiplying D has made A ,

therefore D measures A according to the units in C

But the unit also measures C according to the units in it,

therefore as the unit is to C so is D to A

But as the unit is to C so is C to D ,

therefore also as the unit is to C so is C to D and D to A

Therefore between the unit and the number A two mean proportional numbers C , D have fallen in continued proportion

Again since A by multiplying itself has made B

therefore A measures B according to the units in itself

But the unit also measures A according to the units in it,

therefore as the unit is to A , so is A to B [VII Def 20]

But between the unit and A two mean proportional numbers have fallen, therefore two mean proportional numbers will also fall between A B [VIII 8]

But if two mean proportional numbers fall between two numbers and the first be cube the second will also be cube [VIII 23]

And A is cube

therefore B is also cube

Q E D

PROPOSITION 4

If a cube number by multiplying a cube number make some number, the product will be cube

For let the cube number A by multiplying the cube number B make C ,

I say that C is cube

For let A by multiplying itself make D ,

therefore D is cube [IX 3]

And since A by multiplying itself has made D and by multiplying B has made C

therefore as A is to B so is D to C

[VII 17]

And, since A B are cube numbers

A B are similar solid numbers

Therefore two mean proportional numbers fall between A B [VIII 19]
so that two mean proportional numbers will fall between D C also [VIII 8]

And D is cube

therefore C is also cube

[VIII 23]
Q E D

PROPOSITION 5

If a cube number by multiplying any number make a cube number, the multiplied number will also be cube

For let the cube number A by multiplying any number B make the cube number C ,

I say that B is cube

A _____ For let A by multiplying itself make D ,
 B _____ therefore D is cube [IX 3]

C _____ Now since A by multiplying itself has
 D _____ made D and by multiplying B has made
 C ,

therefore as A is to B so is D to C [VII 17]

And since D , C are cube,

they are similar solid numbers

Therefore two mean proportional numbers fall between D , C [VIII 19]

And, as D is to C , so is A to B ,

therefore two mean proportional numbers fall between A , B also [VIII 8]

And A is cube,

therefore B is also cube [VIII 23]

PROPOSITION 6

If a number by multiplying itself make a cube number, it will itself also be cube

For let the number A by multiplying itself make the cube number B ,

I say that A is also cube

A _____ For let A by multiplying B make C

B _____ Since then, A by multiplying itself has made B , and
 C _____ by multiplying B has made C

therefore C is cube

And since A by multiplying itself has made B ,

therefore A measures B according to the units in itself

But the unit also measures A according to the units in it

Therefore as the unit is to A , so is A to B [VII Def 20]

And since A by multiplying B has made C ,

therefore B measures C according to the units in A

But the unit also measures A according to the units in it

Therefore as the unit is to A , so is B to C [VII Def 20]

But as the unit is to A so is A to B ,

therefore also as A is to B , so is B to C

And since B , C are cube

they are similar solid numbers

Therefore there are two mean proportional numbers between B , C [VIII 19]

And as B is to C , so is A to B

Therefore there are two mean proportional numbers between A , B also [VIII 8]

And B is cube

therefore A is also cube

[cf VIII 23]

Q E D

PROPOSITION 7

If a composite number by multiplying any number make some number, the product will be solid

For, let the composite number A by multiplying any number B make C ,

I say that C is solid

For since A is composite, it will be measured by some number

[vii Def 13]

Let it be measured by D
and as many times as D measures A ,
so many units let there be in E

A _____
B _____
C _____
D _____ E _____

Since then D measures A according to the units in E ,
therefore E by multiplying D has made A [vii Def 15]

And since A by multiplying B has made C ,
and A is the product of D E ,

therefore the product of D E by multiplying B has made C

Therefore C is solid and D , E , B are its sides Q E D

PROPOSITION 8

If as many numbers as we please beginning from an unit be in continued proportion the third from the unit will be square as will also those which successively leave out one the fourth will be cube as will also all those which leave out two and the seventh will be at once cube and square as will also those which leave out five

Let there be as many numbers as we please, A B , C D , E , F , beginning from an unit and in continued proportion

I say that B the third from the unit is square as are also all those which leave out one C the fourth is cube as are also all those which leave out two and F , the seventh is at once cube and square as are also all those which leave out five

A _____
B _____
C _____
D _____
E _____
F _____

For since as the unit is to A , so is A to B ,
therefore the unit measures the number A the same number of times that A measures B [vii Def 20]

But the unit measures the number A according to the units in it,
therefore A also measures B according to the units in A

Therefore A by multiplying itself has made B ,
therefore B is square

And since B C D are in continued proportion and B is square
therefore D is also square [viii 22]

For the same reason

F is also square

Similarly we can prove that all those which leave out one are square

I say next that C the fourth from the unit is cube as are also all those which leave out two

For since as the unit is to A so is B to C
therefore the unit measures the number A the same number of times that B measures C

But the unit measures the number 1 according to the units in 1,

therefore B also measures C according to the units in A

Therefore A by multiplying B has made C

Since then A by multiplying itself has made B and by multiplying B has made C ,

therefore C is cube

And since C, D, E, F are in continued proportion and C is cube,

therefore F is also cube [VIII 23]

But it was also proved square,

therefore the seventh from the unit is both cube and square

Similarly we can prove that all the numbers which leave out five are also both cube and square Q E D

PROPOSITION 9

If as many numbers as we please beginning from an unit be in continued proportion and the number after the unit be square, all the rest will also be square And if the number after the unit be cube, all the rest will also be cube

Let there be as many numbers as we please, A, B, C, D, E, F , beginning from an unit and in continued proportion and let A ,
the number after the unit be square,

I say that all the rest will also be square

Now it has been proved that B , the third from the unit is square as are also all those which leave out one

[IX 8]

I say that all the rest are also square

For since A, B, C are in continued proportion,

and A is square

therefore C is also square [VIII 22]

Again since B, C, D are in continued proportion

and B is square

D is also square [VIII 22]

Similarly we can prove that all the rest are also square

Next, let A be cube

I say that all the rest are also cube

Now it has been proved that C , the fourth from the unit is cube as also are all those which leave out two, [IX 8]

I say that all the rest are also cube

For, since as the unit is to A so is A to B ,
therefore the unit measures A the same number of times as A measures B

But the unit measures A according to the units in it

therefore A also measures B according to the units in itself,

therefore A by multiplying itself has made B

And A is cube

But if a cube number by multiplying itself make some number the product is cube [IX 3]

Therefore B is also cube

And since the four numbers A, B, C, D are in continued proportion

and A is cube

D also is cube

[VIII 23]

For the same reason

E is also cube and similarly all the rest are cube Q E D

PROPOSITION 10

If as many numbers as we please beginning from an unit be in continued proportion and the number after the unit be not square neither will any other be square except the third from the unit and all those which leave out one And if the number after the unit be not cube neither will any other be cube except the fourth from the unit and all those which leave out two

Let there be as many numbers as we please *A B C, D E F*, beginning from an unit and in continued proportion

and let *A* the number after the unit not be square,

I say that neither will any other be square except the third from the unit <and those which leave out one>

For if possible let *C* be square

But *B* is also square

[IX 8]

[therefore *B, C* have to one another the ratio which a square number has to a square number]

And as *B* is to *C* so is *A* to *B*

therefore *A B* have to one another the ratio which a square number has to a square number

[so that *A B* are similar plane numbers]

And *B* is square,

therefore *A* is also square

which is contrary to the hypothesis.

Therefore *C* is not square

Similarly we can prove that neither is any other of the numbers square except the third from the unit and those which leave out one

Next let *A* not be cube

I say that neither will any other be cube except the fourth from the unit and those which leave out two

For if possible let *D* be cube

Now *C* is also cube for it is fourth from the unit

[IX 8]

And as *C* is to *D* so is *B* to *C*

therefore *B* also has to *C* the ratio which a cube has to a cube

And *C* is cube

therefore *B* is also cube

[VIII 20]

And since as the unit is to *A* so is *A* to *B*

and the unit measures *A* according to the units in it

therefore *A* also measures *B* according to the units in itself

therefore *A* by multiplying itself has made the cube number *B*

But if a number by multiplying itself make a cube number it is also itself cube

[IX 6]

Therefore *A* is also cube

which is contrary to the hypothesis

Therefore *D* is not cube

Similarly we can prove that neither is any other of the numbers cube except the fourth from the unit and those which leave out two Q E D

PROPOSITION 11

If as many numbers as we please beginning from an unit be in continued proportion the less measures the greater according to some one of the numbers which have place among the proportional numbers

Let there be as many numbers as we please, B, C, D, E beginning from the unit A and in continued proportion

A ————— I say that B , the least of the numbers B, C, D, E
measures E according to some one of the numbers C, D

B ————— For since as the unit A is to B , so is D to E
 C ————— therefore the unit A measures the number B the same
 D ————— number of times as D measures E ,
 E ————— therefore alternately, the unit A measures D the same
number of times as B measures E [VII 15]

But the unit A measures D according to the units in it,

therefore B also measures E according to the units in D
so that B the less measures E the greater according to some number of those
which have place among the proportional numbers —

PORISM And it is manifest that whatever place the measuring number has
reckoned from the unit the same place also has the number according to which
it measures reckoned from the number measured in the direction of the num-
ber before it —

Q E D

PROPOSITION 12

If as many numbers as we please beginning from an unit be in continued proportion by however many prime numbers the last is measured the next to the unit will also be measured by the same

Let there be as many numbers as we please A, B, C, D beginning from an
unit and in continued proportion,

I say that by however many prime numbers D is measured A will also be
measured by the same

For let D be measured by any prime number E

A —————	F —————	I say that E measures A
B —————	G —————	For suppose it does not
C —————	H —————	now E is prime and any prime
D —————		number is prime to any which it
E —————		does not measure [VII 29]

therefore E, A are prime to one another

And since E measures D let it measure it according to F

therefore E by multiplying F has made D

Again since 1 measures D according to the units in C [IX 11 and Por]

therefore A by multiplying C has made D

But further E has also by multiplying F made D

therefore the product of A, C is equal to the product of E, F

Therefore as A is to E so is F to C [VII 19]

But A, E are prime

primes are also least [VII 21]

and the least measure those which have the same ratio the same number of
times the antecedent the antecedent and the consequent the consequent

[VII 20]

therefore E measures C

Let it measure it according to G ,

therefore E by multiplying G has made C

But, further, by the theorem before this

A has also by multiplying B made C [IX 11 and Por]

Therefore the product of A , B is equal to the product of E , G

Therefore as A is to E , so is G to B [VII 19]

But A , E are prime,

primes are also least, [VII 21]

and the least numbers measure those which have the same ratio with them the same number of times, the antecedent the antecedent and the consequent the consequent [VII 20]

therefore E measures B

Let it measure it according to H ,

therefore E by multiplying H has made B

But further, A has also by multiplying itself made B , [IX 8]

therefore the product of E H is equal to the square on A

Therefore as E is to A , so is A to H [VII 19]

But A , E are prime,

primes are also least [VII 21]

and the least measure those which have the same ratio the same number of times the antecedent the antecedent and the consequent the consequent, [VII 20]

therefore E measures A as antecedent antecedent

But again it also does not measure it

which is impossible

Therefore E , A are not prime to one another

Therefore they are composite to one another

But numbers composite to one another are measured by some number [VII Def 14]

And since E is by hypothesis prime

and the prime is not measured by any number other than itself,

therefore E measures A , E ,

so that E measures A

[But it also measures D

therefore E measures A , D]

Similarly we can prove that by however many prime numbers D is measured A will also be measured by the same Q E D

PROPOSITION 13

If as many numbers as we please beginning from an unit be in continued proportion and the number after the unit be prime the greatest will not be measured by any except those which have a place among the proportional numbers

Let there be as many numbers as we please A B C D beginning from an unit and in continued proportion and let A the number after the unit be prime

I say that D the greatest of them will not be measured by any other number except A B C

For, if possible let it be measured by E and let E not be the same with any of the numbers A B C

It is then manifest that E is not prime

For, if E is prime and measures D

it will also measure A [ix 12], which is prime, though it is not the same with it

A —————	E —————	which is impossible
B —————	F —————	Therefore E is not prime
C —————	G —————	Therefore it is composite
D —————	H —————	But any composite num
		ber is measured by some
		prime number, [vii 31]

therefore E is measured by some prime number

I say next that it will not be measured by any other prime except A

For, if E is measured by another

and E measures D

that other will also measure D

so that it will also measure A [ix 12] which is prime though it is not the same with it

which is impossible

Therefore A measures E

And since E measures D , let it measure it according to F

I say that F is not the same with any of the numbers A B , C

For if F is the same with one of the numbers A B C

and measures D according to E

therefore one of the numbers A , B C also measures D according to E

But one of the numbers A , B , C measures D according to some one of the numbers A B , C , [ix 11]

therefore E is also the same with one of the numbers A B , C

which is contrary to the hypothesis

Therefore F is not the same as any one of the numbers A B , C

Similarly we can prove that F is measured by A , by proving again that F is not prime

For if it is and measures D

it will also measure A [ix 12] which is prime though it is not the same with it

which is impossible

therefore F is not prime

Therefore it is composite

But any composite number is measured by some prime number [vii 31]

therefore F is measured by some prime number

I say next that it will not be measured by any other prime except A

For if any other prime number measures F

and F measures D

that other will also measure D

so that it will also measure A [ix 12] which is prime though it is not the same with it

which is impossible

Therefore A measures F

And since E measures D according to F

therefore E by multiplying F has made D

But further A has also by multiplying C made D

therefore the product of A C is equal to the product of E F [ix 11]

Therefore proportionally, as A is to E , so is F to C

[vii 19]

But A measures E ,

therefore F also measures C

Let it measure it according to G

Similarly then we can prove that G is not the same with any of the numbers

A , B , and that it is measured by A

And since F measures C according to G

therefore F by multiplying G has made C

But further A has also by multiplying B made C

[ix 11]

therefore the product of A , B is equal to the product of F , G

Therefore proportionally, as A is to F so is G to B

[vii 19]

But A measures F

therefore G also measures B

Let it measure it according to H

Similarly then we can prove that H is not the same with A

And since G measures B according to H

therefore G by multiplying H has made B

But further A has also by multiplying itself made B

[ix 8]

therefore the product of H , G is equal to the square on A

Therefore as H is to A , so is A to G

[vii 19]

But A measures G

therefore H also measures A which is prime though it is not the same with it
which is absurd

Therefore D the greatest will not be measured by any other number except

A , B , C

Q E D

PROPOSITION 14

If a number be the least that is measured by prime numbers it will not be measured by any other prime number except those originally measuring it

For let the number 1 be the least that is measured by the prime numbers

B , C , D

I say that A will not be measured by
any other prime number except B , C , D

A —————

B —

E —————

C —

For if possible let it be measured

F —————

D —

by the prime number E and let E not

be the same with any one of the numbers B , C , D

Now since E measures A let it measure it according to F ,

therefore E by multiplying F has made A

And A is measured by the prime numbers B , C , D

But if two numbers by multiplying one another make some number and any prime number measure the product it will also measure one of the original numbers

[vii 30]

therefore B , C , D will measure one of the numbers E , F

Now they will not measure E

for E is prime and not the same with any one of the numbers B , C , D

Therefore they will measure F which is less than A

which is impossible for A is by hypothesis the least number measured by B , C , D

Therefore no prime number will measure A except B , C , D

Q E D

PROPOSITION 15

If three numbers in continued proportion be the least of those which have the same ratio with them any two whatever added together will be prime to the remaining number

Let A, B, C , three numbers in continued proportion be the least of those which have the same ratio with them

A ————— B —————
 C —————
 D — E — F

Isay that any two of the numbers A, B, C whatever added together are prime to the remaining number namely A, B to C, B, C to A , and further, A, C to B

For let two numbers DE, EF , the least of those which have the same ratio with A, B, C , be taken [VIII 2]

It is then manifest that DE by multiplying itself has made A , and by multiplying EF has made B , and further EF by multiplying itself has made C [VIII 2]

Now since DE, EF are least,
 they are prime to one another [VI 22]

But, if two numbers be prime to one another
 their sum is also prime to each [VII 28]

therefore DF is also prime to each of the numbers DE, EF

But further, DE is also prime to EF ,
 therefore DF, DE are prime to EF

But, if two numbers be prime to any number
 their product is also prime to the other [VI 24]

so that the product of FD, DE is prime to EF

hence the product of FD, DE is also prime to the square on EF [VI 25]

But the product of FD, DE is the square on DE together with the product of DE, EF [II 3]

therefore the square on DE together with the product of DE, EF is prime to the square on EF

And the square on DE is A

the product of DE, EF is B ,

and the square on EF is C ,

therefore A, B added together are prime to C

Similarly we can prove that B, C added together are prime to A

I say next that A, C added together are also prime to B

For since DF is prime to each of the numbers DE, EF

the square on DF is also prime to the product of DE, EF [VII 24 25]

But the squares on DE, EF together with twice the product of DE, EF are equal to the square on DF [II 4]

therefore the squares on DE, EF together with twice the product of DE, EF are prime to the product of DE, EF

Separando the squares on DE, EF together with once the product of DE, EF are prime to the product of DE, EF

Therefore *separando* again the squares on DE, EF are prime to the product of DE, EF

And the square on DE is A

the product of DE, EF is B

and the square on EF is C

Therefore A, C added together are prime to B

Q E D

PROPOSITION 16

If two numbers be prime to one another, the second will not be to any other number as the first is to the second

For let the two numbers A, B be prime to one another

I say that B is not to any other number as A is to B

For if possible as A is to B , so let B be to C

Now A, B are prime

primes are also least

[VII 21]

and the least numbers measure those which have the same ratio the same number of times, the antecedent the antecedent and the consequent the consequent,

therefore A measures B as antecedent antecedent

But it also measures itself

therefore A measures A, B which are prime to one another

which is absurd

Therefore B will not be to C , as A is to B

Q E D

PROPOSITION 17

If there be as many numbers as we please in continued proportion and the extremes of them be prime to one another, the last will not be to any other number as the first to the second

For let there be as many numbers as we please A, B, C, D in continued proportion

and let the extremes of them, A, D , be prime to one another

I say that D is not to any other number as A is to B

For if possible as A is to B so let D be to E

therefore alternately as A is to D so is B to E

[VII 13]

But A, D are prime

primes are also least

[VII 21]

and the least numbers measure those which have the same ratio the same number of times the antecedent the antecedent and the consequent the consequent

[VII 20]

Therefore A measures B

And as A is to B so is B to C

Therefore B also measures C

so that A also measures C

And since as B is to C so is C to D

and B measures C

therefore C also measures D

But A measured C

so that A also measures D

But it also measures itself,

therefore A measures A, D which are prime to one another

which is impossible

Therefore D will not be to any other number as A is to B

Q E D

PROPOSITION 18

Given two numbers, to investigate whether it is possible to find a third proportional to them

Let A, B be the given two numbers, and let it be required to investigate whether it is possible to find a third proportional to them

Now A, B are either prime to one another or not

And if they are prime to one another, it has been proved that it is impossible to find a third proportional to them [IX 16]

Next let A, B not be prime to one another,

and let B by multiplying itself make C

Then A either measures C or does not measure it

First let it measure it according to D ,

therefore A by multiplying D has made C


But further, B has also by multiplying itself made C

therefore the product of A, D is equal to the square on B

Therefore as A is to B so is B to D [VII 19]

therefore a third proportional number D has been found to A, B

Next let A not measure C ,



I say that it is impossible to find a third proportional number to A, B

For if possible let D , such third proportional have been found

Therefore the product of A, D is equal to the square on B

But the square on B is C

therefore the product of A, D is equal to C

Hence A by multiplying D has made C

therefore A measures C according to D

But by hypothesis it also does not measure it

which is absurd

Therefore it is not possible to find a third proportional number to A, B when A does not measure C Q E D

PROPOSITION 19

Given three numbers to investigate when it is possible to find a fourth proportional to them

A _____

B _____

C _____

Let A, B, C be the given three numbers and let it be required to investigate when it is possible to find a fourth proportional to them

[The Greek text of this proposition is corrupt. However analogously to Proposition 18 the condition that a fourth proportional to A, B, C exists is that A measure the product of B and C]

PROPOSITION 20

Prime numbers are more than any assigned multitude of prime numbers

Let A, B, C be the assigned prime numbers

I say that there are more prime numbers than A, B, C

For let the least number measured by $A B C$ be taken,
and let it be DE

let the unit DF be added to DE

Then EF is either prime or not

First let it be prime

then the prime numbers $A B, C EF$ have
been found which are more than $A B C$

Next, let EF not be prime

therefore it is measured by some prime number [v.11 31]

Let it be measured by the prime number G

I say that G is not the same with any of the numbers A, B, C

For if possible let it be so

Now $A B C$ measure DE

therefore G also will measure DE

But it also measures EF

Therefore G being a number will measure the remainder the unit DF
which is absurd

Therefore G is not the same with any one of the numbers $A B, C$

And by hypothesis it is prime

Therefore the prime numbers $A B C G$ have been found which are more
than the assigned multitude of $A B C$ Q E D

PROPOSITION 21

If as many even numbers as we please be added together the whole is even

For let as many even numbers as we please $AB, BC CD DE$ be added together

I say that the whole AE is even

For since each of the numbers AB

$BC CD DE$ is even it has a half part A B C D E

[v.11 Def 6]

so that the whole AE also has a half part

But an even number is that which is divisible into two equal parts [id]
therefore AE is even Q E D

PROPOSITION 22

If as many odd numbers as we please be added together and their multitude be even, the whole will be even

For let as many odd numbers as we please $AB BC CD DE$ even in multitude be added together,

I say that the whole AE is even

For since each of the numbers
 $AB BC CD DE$ is odd if an unit
be subtracted from each each of
the remainders will be even

A B C D E

[v.11 Def 7]
[ix 21]

so that the sum of them will be even

But the multitude of the units is also even

Therefore the whole AE is also even

[ix 21]
Q E D

PROPOSITION 23

If as many odd numbers as we please be added together, and their multitude be odd the whole will also be odd

For let as many odd numbers as we please, AB, BC, CD , the multitude of which is odd be added together,

$A \quad B \quad C \quad D$

I say that the whole AD is also odd

Let the unit DE be subtracted from CD ,

therefore the remainder CE is even [VII Def 7]

But CA is also even, [IX 22]

therefore the whole AE is also even [IX 21]

And DE is an unit

Therefore AD is odd [VII Def 7]

Q E D

PROPOSITION 24

If from an even number an even number be subtracted, the remainder will be even

For from the even number AB let the even number BC be subtracted

$A \quad C \quad B$

I say that the remainder CA is even

For since AB is even it has a half part [VII Def 6]

For the same reason BC also has a half part

so that the remainder [CA also has a half part and] AC is therefore even

Q E D

PROPOSITION 25

If from an even number an odd number be subtracted, the remainder will be odd

For from the even number AB let the odd number BC be subtracted

$A \quad C \quad D \quad B$

I say that the remainder CA is odd

For let the unit CD be subtracted from BC ,

therefore DB is even [VII Def 7]

But AB is also even,

therefore the remainder AD is also even [IX 24]

And CD is an unit

therefore CA is odd [VII Def 7]

Q E D

PROPOSITION 26

If from an odd number an odd number be subtracted the remainder will be even

For from the odd number AB let the odd number BC be subtracted

$A \quad C \quad D \quad B$

I say that the remainder CA is even

For since AB is odd let the unit BD be subtracted

therefore the remainder AD is even [VII Def 7]

For the same reason CD is also even [VII Def 7]

so that the remainder CA is also even [IX 24]

Q E D

PROPOSITION 27

If from an odd number an even number be subtracted the remainder will be odd

For from the odd number AB let the even number BC be subtracted

I say that the remainder CA is odd
 Let the unit AD be subtracted,
 therefore DB is even [VII Def 7]
 But BC is also even,
 therefore the remainder CD is even [IX 24]
 Therefore CA is odd [VII Def 7]

$A \quad D \quad C \quad B$
 $Q \quad E \quad D$

PROPOSITION 28

If an odd number by multiplying an even number make some number, the product will be even

For let the odd number A by multiplying the even number B make C ,

I say that C is even

For since A by multiplying B has made C ,
 therefore C is made up of as many numbers equal to B
 as there are units in A [VII Def 15]

$A \quad \text{---}$
 $B \quad \text{---}$
 $C \quad \text{---}$

And B is even

therefore C is made up of even numbers

But if as many even numbers as we please be added together, the whole is even [IX 21]

Therefore C is even

$Q \quad E \quad D$

PROPOSITION 29

If an odd number by multiplying an odd number make some number, the product will be odd

For let the odd number A by multiplying the odd number B make C ,

I say that C is odd

For since A by multiplying B has made C
 therefore C is made up of as many numbers equal to
 B as there are units in A [VII Def 15]

$A \quad \text{---}$
 $B \quad \text{---}$
 $C \quad \text{---}$

And each of the numbers $A \quad B$ is odd

therefore C is made up of odd numbers the multitude of which is odd

Thus C is odd

[IX 25]

$Q \quad E \quad D$

PROPOSITION 30

If an odd number measure an even number, it will also measure the half of it

For let the odd number A measure the even number B ,

I say that it will also measure the half of it

For since 1 measures B

let it measure it according to C

I say that C is not odd

$A \quad \text{---}$
 $B \quad \text{---}$
 $C \quad \text{---}$

For if possible let it be so

Then since A measures B according to C

therefore A by multiplying C has made B

Therefore B is made up of odd numbers the multitude of which is odd

Therefore B is odd [IX 23]

which is absurd for by hypothesis it is even

Therefore C is not odd

therefore C is even

Thus A measures B an even number of times

For this reason then it also measures the half of it

Q E D

PROPOSITION 31

If an odd number be prime to any number, it will also be prime to the double of it

For let the odd number A be prime to any number B ,

and let C be double of B ,

A —————

I say that A is prime to C

B —————

For if they are not prime to one another
some number will measure them

C —————

Let a number measure them and let it be D

D —————

Now A is odd,

therefore D is also odd

And since D which is odd measures C

and C is even

therefore $\{D\}$ will measure the half of C also

[IX 30]

But B is half of C ,

therefore D measures B

But it also measures A

therefore D measures A B which are prime to one another
which is impossible

Therefore A cannot but be prime to C

Therefore A C are prime to one another

Q E D

PROPOSITION 32

Each of the numbers which are continually doubled beginning from a dyad is even times even only

For let as many numbers as we please B C D have been continually doubled beginning from the dyad A ,

A —

I say that B C D are even times even only

B —

Now that each of the numbers B C D is
even times even is manifest for it is
doubled from a dyad

C —

D —

I say that it is also even times even only

For let an unit be set out

Since then as many numbers as we please beginning from an unit are in continued proportion

and the number A after the unit is prime

therefore D the greatest of the numbers A B C D will not be measured by any other number except A B C

[IX 13]

And each of the numbers A B C is even

therefore D is even times even only

[VII Def 8]

Similarly we can prove that each of the numbers B C is even times even only

Q E D

PROPOSITION 33

If a number have its half odd it is even times odd only

For let the number A have its half odd

I say that A is even times odd only

Now that it is even times odd is manifest for the half of it being odd measures it an even number of times [vii Def 9]

I say next that it is also even times odd only

For if A is even times even also

it will be measured by an even number according to an even number

[vii Def 8]

so that the half of it will also be measured by an even number though it is odd

which is absurd

Therefore A is even times odd only

Q E D

PROPOSITION 34

If a number neither be one of those which are continually doubled from a dyad, nor have its half odd it is both even-times even and even-times odd

For let the number A neither be one of those doubled from a dyad nor have its half odd

I say that A is both even times even and even times odd

Now that A is even times even is manifest

for it has not its half odd

[vii Def 8]

I say next that it is also even times odd

For if we bisect A then bisect its half and do this continually we shall come upon some odd number which will measure A according to an even number

For if not we shall come upon a dyad

and A will be among those which are doubled from a dyad

which is contrary to the hypothesis

Thus A is even times odd

But it was also proved even times even

Therefore A is both even times even and even times odd

Q E D

PROPOSITION 35


If as many numbers as we please be in continued proportion and there be subtracted from the second and the last numbers equal to the first then as the excess of the second is to the first so will the excess of the last be to all those before it

Let there be as many numbers as we please in continued proportion A, BC, D, EF beginning from A as least

and let there be subtracted from BC and

EF the numbers BG, FH each equal to A

I say that as GC is to A so is EH to A, BC, D



For let FK be made equal to BC and FL equal to D

Then since FK is equal to BC

and of these the part FH is equal to the part BG

therefore the remainder HK is equal to the remainder GC

And since as EF is to D so is D to BC and BC to A

while D is equal to FL BC to FK and A to FH

therefore as EF is to FL so is LF to FK and FK to FH

Separando as EL is to LF so is LH to FK , and KH to PH [vii 11 13]

Therefore also as one of the antecedents is to one of the consequents so are all the antecedents to all the consequents [VII 12]

therefore as AH is to FH so are EL, LA, KH to LF, FK, HF

But AH is equal to CG, FH to A and LF, FA, HF to $D, BC, A,$

therefore as CG is to A , so is EH to D, BC, A

Therefore as the excess of the second is to the first so is the excess of the last to all those before it : Q E D

PROPOSITION 36

If as many numbers as we please beginning from an unit be set out continuously in double proportion until the sum of all becomes prime and if the sum multiplied into the last make some number the product will be perfect

For let as many numbers as we please A, B, C, D beginning from an unit be set out in double proportion until the sum of all becomes prime

let E be equal to the sum and let E by multiplying D make $FG,$

I say that FG is perfect

For however many A, B, C, D are in multitude let so many E, HA, L, M be taken in double proportion beginning from $E,$

therefore *ex aequali* as A is to D so is E to M [VII 14]

Therefore the product of F, D is equal to the product of A, M [VII 19]

And the product of E, D is FG


therefore the product of A, M is also FG

Therefore A by multiplying M has made FG

therefore M measures FG according to the units in A

And A is a dyad

therefore FG is double of M



But M, L, HK, E are continuously double of each other
therefore E, HA, L, M, FG are continuously proportional in double proportion

Now let there be subtracted from the second HA and the last FG the numbers HN, FO each equal to the first E

therefore as the excess of the second is to the first so is the excess of the last to all those before it [IX 35]

Therefore as NK is to E so is OG to M, L, HA, E

And NK is equal to E

therefore OG is also equal to M, L, HA, E

But FO is also equal to E

and E is equal to A, B, C, D and the unit

Therefore the whole FG is equal to E, HA, L, M and A, B, C, D and the unit

and it is measured by them

I say also that FG will not be measured by any other number except 4 B C , D , E HK , L M and the unit

For if possible let some number P measure FG
and let P not be the same with any of the numbers A , B , C D , E , HK L M

And as many times as P measures FG so many units let there be in Q
therefore Q by multiplying P has made FG

But further, E has also by multiplying D made FG
therefore as E is to Q so is P to D [VII 19]

And since A B C D are continuously proportional beginning from an unit
therefore D will not be measured by any other number except A B C [IX 13]

And by hypothesis P is not the same with any of the numbers A , B C ,
therefore P will not measure D

But as P is to D so is E to Q
therefore neither does E measure Q [VII Def 20]

And E is prime
and any prime number is prime to any number which it does not measure [VII 29]

Therefore E Q are prime to one another

But primes are also least [VII 21]
and the least numbers measure those which have the same ratio the same number of times the antecedent the antecedent and the consequent the consequent [VII 20]

and as E is to Q so is P to D

therefore E measures P the same number of times that Q measures D

But D is not measured by any other number except A B , C

therefore Q is the same with one of the numbers A , B , C

Let it be the same with B

And however many B C D are in multitude let so many E HK , L be taken beginning from E

Now E HK L are in the same ratio with B C D

therefore *ex æquali* as B is to D , so is E to L [VII 14]

Therefore the product of B L is equal to the product of D E [VII 19]

But the product of D E is equal to the product of Q P

therefore the product of Q P is also equal to the product of B L

Therefore as Q is to B so is L to P [VII 19]

And Q is the same with B

therefore L is also the same with P

which is impossible for by hypothesis P is not the same with any of the numbers set out

Therefore no number will measure FG except 4, B C , D E HK , L M and the unit

And FG was proved equal to 4 B C D E HK L M and the unit
and a perfect number is that which is equal to its own parts [VII Def 2^o]

therefore FG is perfect Q E D

BOOK TEN

DEFINITIONS I

1 Those magnitudes are said to be *commensurable* which are measured by the same measure and those *incommensurable* which cannot have any common measure

2 Straight lines are *commensurable in square* when the squares on them are measured by the same area, and *incommensurable in square* when the squares on them cannot possibly have any area as a common measure

3 With these hypotheses it is proved that there exist straight lines infinite in multitude which are commensurable and incommensurable respectively some in length only and others in square also with an assigned straight line Let then the assigned straight line be called *rational* and those straight lines which are commensurable with it whether in length and in square or in square only, *rational* but those which are incommensurable with it *irrational*

4 And let the square on the assigned straight line be called *rational* and those areas which are commensurable with it *rational* but those which are incommensurable with it *irrational* and the straight lines which produce them *irrational* that is in case the areas are squares the sides themselves but in case they are any other rectilineal figures the straight lines on which are described squares equal to them


BOOK X PROPOSITIONS

PROPOSITION 1

Two unequal magnitudes being set out if from the greater there be subtracted a magnitude greater than its half and from that which is left a magnitude greater than its half and if this process be repeated continually there will be left some magnitude which will be less than the lesser magnitude set out

Let AB C be two unequal magnitudes of which AB is the greater

I say that if from AB there be subtracted a magnitude greater than its half and from that which is left a magnitude greater than its half and if this process be repeated continually there will be left some magnitude which will be less than the magnitude C



For C if multiplied will sometime be greater than AB [cf v Def 4]

Let it be multiplied and let DE be a multiple of C and greater than AB
 let DE be divided into the parts DF FG GE equal to C
 from AB let there be subtracted BH greater than its half
 and from AH AK greater than its half

and let this process be repeated continually until the divisions in AB are equal in multitude with the divisions in DE

Let then AK , KH , HB be divisions which are equal in multitude with DF , FG , GE

Now since DE is greater than AB

and from DE there has been subtracted EG less than its half,

and from AB BH greater than its half

therefore the remainder GD is greater than the remainder HA

And since GD is greater than HA ,

and there has been subtracted, from GD the half GF ,

and from HA , HK greater than its half

therefore the remainder DF is greater than the remainder AK

But DF is equal to C

therefore C is also greater than AK

Therefore AK is less than C

Therefore there is left of the magnitude AB the magnitude AK which is less than the lesser magnitude set out namely C Q E D

And the theorem can be similarly proved even if the parts subtracted be halves

PROPOSITION 2

If when the less of two unequal magnitudes is continually subtracted in turn from the greater that which is left never measures the one before it the magnitudes will be incommensurable

For there being two unequal magnitudes AB , CD and AB being the less when the less is continually subtracted in turn from the greater, let that which is left over never measure the one before it

I say that the magnitudes AB , CD are incommensurable

For if they are commensurable some magnitude will measure them

Let a magnitude measure them if possible and let it be E ,

let AB measuring FD leave CF less than itself,

let CF measuring BG leave AG less than itself

and let this process be repeated continually until there is left some magnitude which is less than E

Suppose this done and let there be left AG less than E

Then since E measures AB

while AB measures DF

therefore E will also measure FD

But it measures the whole CD also


therefore it will also measure the remainder CF

But CF measures BG

therefore F also measures BG

But it measures the whole AB also

therefore it will also measure the remainder AG the greater the less which is impossible



Therefore no magnitude will measure the magnitudes AB CD ,
 therefore the magnitudes AB , CD are incommensurable [x Def 1]
 Therefore etc. Q E D

PROPOSITION 3

Given two commensurable magnitudes, to find their greatest common measure

Let the two given commensurable magnitudes be AB , CD of which AB is the less

thus it is required to find the greatest common measure of AB , CD

Now the magnitude AB either measures CD or it does not

If then it measures it—and it measures itself also— AB is a common measure of AB , CD


And it is manifest that it is also the greatest,

for a greater magnitude than the magnitude AB will not measure AB

Next, let AB not measure CD

Then if the less be continually subtracted in turn from the greater, that which is left over will sometime measure the one before it, because AB , CD are not incommensurable,

[cf x 2]



let AB , measuring ED leave EC less than itself
 let EC measuring FB leave AF less than itself
 and let AF measure CE

Since, then AF measures CE ,

while CE measures FB ,

therefore AF will also measure FB

But it measures itself also,

therefore AF will also measure the whole AB

But AB measures DE

therefore AF will also measure ED

But it measures CE also,

therefore it also measures the whole CD

Therefore AF is a common measure of AB CD

I say next that it is also the greatest

For if not there will be some magnitude greater than AF which will measure AB CD

Let it be G

Since then G measures AB

while AB measures ED

therefore G will also measure ED

But it measures the whole CD also

therefore G will also measure the remainder CE

But CE measures FB

therefore G will also measure FB

But it measures the whole AB also

and it will therefore measure the remainder AF the greater the less which is impossible

Therefore no magnitude greater than AF will measure AB CD
 therefore AF is the greatest common measure of AB CD

Therefore the greatest common measure of the two given commensurable magnitudes AB , CD has been found Q E D

POBISM From this it is manifest that if a magnitude measure two magnitudes it will also measure their greatest common measure

PROPOSITION 4

Given three commensurable magnitudes, to find their greatest common measure

Let A B C be the three given commensurable magnitudes, thus it is required to find the greatest common measure of A , B , C

Let the greatest common measure of the two magnitudes A B be taken and let it be D [\propto 3]
 then D either measures C or does not measure it

A —————

B —————

C —————

D ——— E — F —

First let it measure it

Since then D measures C

while it also measures A B

therefore D is a common measure of A B C

And it is manifest that it is also the greatest for a greater magnitude than the magnitude D does not measure A B

Next let D not measure C

I say first that C D are commensurable

For since 1 B C are commensurable

some magnitude will measure them

and this will of course measure A B also

so that it will also measure the greatest common measure of A B namely D [\propto 3 Por]

But it also measures C

so that the said magnitude will measure C D

therefore C D are commensurable

Now let their greatest common measure be taken and let it be E [\propto 3]

Since then E measures D

while D measures A B

therefore E will also measure A B

But it measures C also

therefore E measures A B C

therefore E is a common measure of A , B C

I say next that it is also the greatest

For if possible let there be some magnitude F greater than E and let it measure A B C

Now since F measures A B C

it will also measure A B

and will measure the greatest common measure of A , B [\propto 3 Por]

But the greatest common measure of A B is D

therefore F measures D

But it measures C also

therefore F measures C D

therefore F will also measure the greatest common measure of C D [\propto 3 Por]

But that is E ,

therefore F will measure E , the greater the less
which is impossible

Therefore no magnitude greater than the magnitude E will measure A, B, C ,
therefore E is the greatest common measure of A, B, C if D do not measure C ,
and if it measure it D is itself the greatest common measure

Therefore the greatest common measure of the three given commensurable
magnitudes has been found

PORISM From this it is manifest that if a magnitude measure three magni-
tudes it will also measure their greatest common measure

Similarly too, with more magnitudes the greatest common measure can be
found and the porism can be extended Q E D

PROPOSITION 5


*Commensurable magnitudes have to one another the ratio which a number has to a
number*

Let A, B be commensurable magnitudes,

I say that A has to B the ratio which a number has to a number

For, since A, B are commensurable some magnitude will measure them

Let it measure them and let it be C



And as many times as C meas-
ures A , so many units let there be
in D ,

and as many times as C meas-
ures B , so many units let there
be in E

Since then C measures A according to the units in D

while the unit also measures D according to the units in it

therefore the unit measures the number D the same number of times as the
magnitude C measures A

therefore as C , is to A so is the unit to D , [vii Def 20]

therefore, inversely as A is to C , so is D to the unit [cf v 7 Por]

Again since C measures B according to the units in E

while the unit also measures E according to the units in it

therefore the unit measures E the same number of times as C measures B ,

therefore as C is to B so is the unit to E

But it was also proved that

as A is to C , so is D to the unit

therefore *ex aequali*

as A is to B so is the number D to E [v 22]

Therefore the commensurable magnitudes A, B have to one another the
ratio which the number D has to the number E Q E D


PROPOSITION 6

*If two magnitudes have to one another the ratio which a number has to a number,
the magnitudes will be commensurable*

For let the two magnitudes A, B have to one another the ratio which the
number D has to the number E

I say that the magnitudes A, B are commensurable

For let A be divided into as many equal parts as there are units in D
 and let C be equal to one of them,
 and let F be made up of as
 many magnitudes equal to C
 as there are units in E



Since then there are in A
 as many magnitudes equal to
 C as there are units in D

whatever part the unit is of D , the same part is C of A also
 therefore as C is to A so is the unit to D [VII Def 20]

But the unit measures the number D ,
 therefore C also measures A

And since as C is to A , so is the unit to D
 therefore inversely as A is to C , so is the number D to the unit
 [cf v 7 Por]

Again since there are in F as many magnitudes equal to C as there are units
 in E

therefore as C is to F so is the unit to E [VII Def 20]

But it was also proved that

as A is to C so is D to the unit

therefore *ex aequali* as A is to F so is D to E [v 2^o]

But as D is to E so is A to B

therefore also as A is to B so is it to F also [v 11]

Therefore A has the same ratio to each of the magnitudes B , F ,

therefore B is equal to F [v 9]

But C measures F

therefore it measures B also

Further it measures A also

therefore C measures A B

Therefore A is commensurable with B

Therefore etc

POINISM From this it is manifest that if there be two numbers as D E and
 a straight line as A it is possible to make a straight line $\{F\}$ such that the
 given straight line is to it as the number D is to the number E

And if a mean proportional be also taken between A F as B ,
 as A is to F so will the square on A be to the square on B that is as the first
 is to the third so is the figure on the first to that which is similar and similarly
 described on the second [vi 19 Por]

But as A is to F so is the number D to the number E
 therefore it has been contrived that as the number D is to the number E so
 also is the figure on the straight line A to the figure on the straight line B

Q E D

PROPOSITION 7

*Incommensurable magnitudes have not to one another the ratio which a number has
 to a number*

Let A , B be incommensurable magnitudes

I say that A has not to B the ratio which a number has to a number

For, if A has to B the ratio which a number has to a number, A will be com

measurable with B

[x 6]

But it is not,

$\frac{A}{B}$ therefore A has not to B the ratio which a number has to a number

Therefore etc

Q E D

PROPOSITION 8

If two magnitudes have not to one another the ratio which a number has to a number, the magnitudes will be incommensurable

For let the two magnitudes A, B not have to one another the ratio which a number has to a number,

I say that the magnitudes A, B are incommensurable

$\frac{A}{B}$ For if they are commensurable A will have to B the ratio which a number has to a number

[x 5]

But it has not,

therefore the magnitudes A, B are incommensurable

Therefore etc

Q E D

PROPOSITION 9

The squares on straight lines commensurable in length have to one another the ratio which a square number has to a square number and squares which have to one another the ratio which a square number has to a square number will also have their sides commensurable in length. But the squares on straight lines incommensurable in length have not to one another the ratio which a square number has to a square number, and squares which have not to one another the ratio which a square number has to a square number will not have their sides commensurable in length either

For let A, B be commensurable in length

$\frac{A}{C}$
 $\frac{B}{D}$

I say that the square on A has to the square on B the ratio which a square number has to a square number

For since A is commensurable in length with B ,

therefore A has to B the ratio which a number has to a number [x 5]

Let it have to it the ratio which C has to D

Since then, as A is to B , so is C to D

while the ratio of the square on A to the square on B is duplicate of the ratio of A to B ,

for similar figures are in the duplicate ratio of their corresponding sides

[vi 20 Por]

and the ratio of the square on C to the square on D is duplicate of the ratio of C to D

for between two square numbers there is one mean proportional number and the square number has to the square number the ratio duplicate of that which the side has to the side [viii 11]

therefore also as the square on A is to the square on B so is the square on C to the square on D

Next as the square on A is to the square on B , so let the square on C be to the square on D ,

I say that A is commensurable in length with B

For since as the square on A is to the square on B , so is the square on C to the square on D ,
while the ratio of the square on A to the square on B is duplicate of the ratio of A to B ,
and the ratio of the square on C to the square on D is duplicate of the ratio of C to D ,

therefore also as A is to B , so is C to D

Therefore A has to B the ratio which the number C has to the number D
therefore A is commensurable in length with B [x 6]

Next let A be incommensurable in length with B ,

I say that the square on A has not to the square on B the ratio which a square number has to a square number

For if the square on A has to the square on B the ratio which a square number has to a square number, A will be commensurable with B

But it is not

therefore the square on A has not to the square on B the ratio which a square number has to a square number

Again let the square on A not have to the square on B the ratio which a square number has to a square number,

I say that A is incommensurable in length with B

For, if A is commensurable with B the square on A will have to the square on B the ratio which a square number has to a square number

But it has not,

therefore A is not commensurable in length with B

Therefore etc

PORISM And it is manifest from what has been proved that straight lines commensurable in length are always commensurable in square also, but those commensurable in square are not always commensurable in length also

[**LEMMA** It has been proved in the arithmetical books that similar plane numbers have to one another the ratio which a square number has to a square number [viii 26]

and that if two numbers have to one another the ratio which a square number has to a square number they are similar plane numbers [Converse of viii 26]

And it is manifest from these propositions that numbers which are not similar plane numbers that is those which have not their sides proportional have not to one another the ratio which a square number has to a square number

For if they have they will be similar plane numbers which is contrary to the hypothesis

Therefore numbers which are not similar plane numbers have not to one another the ratio which a square number has to a square number]

PROPOSITION 10

To find two straight lines incommensurable the one in length only and the other in square also with an assigned straight line

Let A be the assigned straight line,
thus it is required to find two straight lines incommensurable the one in length only and the other in square also with A

Let two numbers B C be set out which have not to one another the ratio

which a square number has to a square number that is which are not similar plane numbers,

and let it be contrived that,

A _____ as B is to C , so is the square on A to the square on D
 D _____ —for we have learnt how to do thus— [x 6 Por]
 E _____ therefore the square on A is commensurable with the
 B _____ square on D [x 6]
 C _____

And since B has not to C the ratio which a square number has to a square number,
 therefore neither has the square on A to the square on D the ratio which a square number has to a square number

therefore A is incommensurable in length with D [x 9]

Let E be taken a mean proportional between A D ,
 therefore as A is to D , so is the square on A to the square on E [v Def 9]

But A is incommensurable in length with D ,
 therefore the square on A is also incommensurable with the square on E [x 11]

therefore A is incommensurable in square with E

Therefore two straight lines D E have been found incommensurable D in length only and E in square and of course in length also with the assigned straight line A Q E D

PROPOSITION 11

If four magnitudes be proportional, and the first be commensurable with the second the third will also be commensurable with the fourth and if the first be incommensurable with the second the third will also be incommensurable with the fourth

Let A , B , C D be four magnitudes in proportion so that as A is to B so is C to D

and let A be commensurable with B ,

A _____ B _____ I say that C will also be commensurable
 C _____ D _____ with D

For since A is commensurable with B

therefore A has to B the ratio which a number has to a number [x 5]

And as A is to B so is C to D ,

therefore C also has to D the ratio which a number has to a number,

therefore C is commensurable with D [x 6]

Next let A be incommensurable with B

I say that C will also be incommensurable with D

For since A is incommensurable with B

therefore A has not to B the ratio which a number has to a number [x 7]

And as A is to B so is C to D

therefore neither has C to D the ratio which a number has to a number

therefore C is incommensurable with D [x 8]

Therefore etc

Q E D

PROPOSITION 12

Magnitudes commensurable with the same magnitude are commensurable with one another also

For let each of the magnitudes A B be commensurable with C

I say that A is also commensurable with B

For since A is commensurable with C ,
therefore A has to C the ratio
which a number has to a
number [x 5]

Let it have the ratio which
 D has to E

Again since C is commen-
surable with B ,

therefore C has to B the ratio which a number has to a number [x 5]

Let it have the ratio which F has to G

And given any number of ratios we please namely the ratio which D has to
 E and that which F has to G

let the numbers H, K, L be taken continuously in the given ratios, [cf viii 4]

so that as D is to E , so is H to K ,

and, as F is to G , so is K to L

Since then, as A is to C , so is D to E ,

while as D is to E so is H to K ,

therefore also, as A is to C so is H to K [v 11]

Again since as C is to B so is F to G ,

while as F is to G so is K to L

therefore also as C is to B so is K to L [v 11]

But also as A is to C , so is H to K

therefore *ex aequali* as A is to B , so is H to L [v 27]

Therefore A has to B the ratio which a number has to a number,

therefore A is commensurable with B [x 6]

Therefore etc

Q E D

PROPOSITION 13

If two magnitudes be commensurable and the one of them be incommensurable with
any magnitude the remaining one will also be incommensurable with the same

Let A, B be two commensurable magnitudes and let one of them A be in
commensurable with any other magnitude C

I say that the remaining one B will also be incommensurable with C

For if B is commensurable with C

while A is also commensurable with B

A is also commensurable with C

[x 12]

But it is also incommensurable with it

which is impossible

Therefore B is not commensurable with C

therefore it is incommensurable with it

Therefore etc


Q E D

LEMMA

Given two unequal straight lines to find by what square the square on the greater is
greater than the square on the less

Let AB, C be the given two unequal straight lines and let AB be the greater
of them,

thus it is required to find by what square the square on AB is greater than the square on C



Let the semicircle ADB be described on AB
and let AD be fitted into it equal to C , [iv 11]
let DB be joined

It is then manifest that the angle ADB is right [iii 31]

and that the square on AB is greater than the square on AD , that is, C by the square on DB [i 47]

Similarly also if two straight lines be given, the straight line the square on which is equal to the sum of the squares on them is found in this manner

Let AD DB be the given two straight lines, and let it be required to find the straight line the square on which is equal to the sum of the squares on them

Let them be placed so as to contain a right angle that formed by AD DB , and let AB be joined

It is again manifest that the straight line the square on which is equal to the sum of the squares on AD , DB is AB [i 47]

Q E D

PROPOSITION 14

If four straight lines be proportional and the square on the first be greater than the square on the second by the square on a straight line commensurable with the first the square on the third will also be greater than the square on the fourth by the square on a straight line commensurable with the third

And if the square on the first be greater than the square on the second by the square on a straight line incommensurable with the first, the square on the third will also be greater than the square on the fourth by the square on a straight line incommensurable with the third

Let A B C D be four straight lines in proportion so that as A is to B so is C to D



and let the square on A be greater than the square on B by the square on E

and let the square on C be greater than the square on D by the square on F ,

I say that if A is commensurable with E , C is also commensurable with F

and, if A is incommensurable with E C is also incommensurable with F

For since as A is to B so is C to D

therefore also as the square on A is to the square on B so is the square on C to the square on D [vi 22]

But the squares on E B are equal to the square on A

and the squares on D F are equal to the square on C

Therefore as the squares on E B are to the square on B so are the squares on D F to the square on D

therefore *separando* as the square on E is to the square on B so is the square on F to the square on D , [v 17]

therefore also as E is to B so is F to D [vi 22]

therefore inversely as B is to E so is D to F

But as A is to B , so also is C to D ,

therefore *ex aequali*, as A is to E , so is C to F [v 22]

Therefore if A is commensurable with E C is also commensurable with F ,
and if A is incommensurable with E , C is also incommensurable with F [x 11]

Therefore etc


Q E D

PROPOSITION 15

If two commensurable magnitudes be added together, the whole will also be commensurable with each of them and, if the whole be commensurable with one of them the original magnitudes will also be commensurable

For let the two commensurable magnitudes AB , BC be added together

I say that the whole AC is also commensurable with each of the magnitudes AB
 BC



For since AB , BC are commensurable,
some magnitude will measure them

Let it measure them and let it be D

Since then D measures AB , BC , it will also measure the whole AC

But it measures AB BC also,

therefore D measures AB BC AC

therefore AC is commensurable with each of the magnitudes AB , BC

[x Def 1]

Next, let AC be commensurable with AB ,

I say that AB , BC are also commensurable

For since AC , AB are commensurable some magnitude will measure them

Let it measure them and let it be D

Since then D measures CA AB it will also measure the remainder BC

But it measures AB also

therefore D will measure AB BC ,

therefore AB BC are commensurable

[x Def 1]

Therefore etc

Q E D

PROPOSITION 16

If two incommensurable magnitudes be added together the whole will also be incommensurable with each of them and if the whole be incommensurable with one of them the original magnitudes will also be incommensurable

For let the two incommensurable magnitudes AB BC be added together I say that the whole AC is also incommensurable with each of the magnitudes AB BC

For if CA AB are not incommensurable some magnitude will measure them

Let it measure them if possible and let it be D

Since then D measures CA AB

therefore it will also measure the remainder BC

But it measures AB also

therefore D measures AB BC

Therefore AB BC are commensurable

but they were also by hypothesis incommensurable
which is impossible




Therefore no magnitude will measure CA, AB ,
 therefore CA, AB are incommensurable [\propto Def 1]
 Similarly we can prove that AC, CB are also incommensurable
 Therefore AC is incommensurable with each of the magnitudes AB, BC
 Next let AC be incommensurable with one of the magnitudes AB, BC
 First let it be incommensurable with AB ,
 I say that AB, BC are also incommensurable
 For, if they are commensurable some magnitude will measure them
 Let it measure them, and let it be D
 Since then, D measures AB, BC ,
 therefore it will also measure the whole AC
 But it measures AB also
 therefore D measures CA, AB
 Therefore CA, AB are commensurable,
 but they were also by hypothesis incommensurable
 which is impossible
 Therefore no magnitude will measure AB, BC
 therefore AB, BC are incommensurable [\propto Def 1]
 Therefore etc Q E D

LEMMA

If to any straight line there be applied a parallelogram deficient by a square figure the applied parallelogram is equal to the rectangle contained by the segments of the straight line resulting from the application

For let there be applied to the straight line AB the parallelogram AD deficient by the square figure DB ,



I say that AD is equal to the rectangle contained by AC, CB

This is indeed at once manifest

for since DB is a square

DC is equal to CB

and AD is the rectangle AC, CD , that is the rectangle AC, CB

Therefore etc

Q E D

PROPOSITION 17

If there be two unequal straight lines and to the greater there be applied a parallelogram equal to the fourth part of the square on the less and deficient by a square figure and if it divide it into parts which are commensurable in length then the square on the greater will be greater than the square on the less by the square on a straight line commensurable with the greater

And if the square on the greater be greater than the square on the less by the square on a straight line commensurable with the greater and if there be applied to the greater a parallelogram equal to the fourth part of the square on the less and deficient by a square figure, it will divide it into parts which are commensurable in length

Let A, BC be two unequal straight lines of which BC is the greater and let there be applied to BC a parallelogram equal to the fourth part of the square on the less A that is equal to the square on the half of A and deficient by a square figure Let this be the rectangle BD, DC [cf Lemma]

and let BD be commensurable in length with DC ,


I say that the square on BC is greater than the square on A by the square on a straight line commensurable with BC

For let BC be bisected at the point E

and let EF be made equal to DE

Therefore the remainder DC is equal to BF

And since the straight line BC has been cut into equal parts at E and into unequal parts at D therefore the rectangle contained by BD , DC , together with the square on ED , is equal to the square on EC ,



[11 5]

And the same is true of their quadruples, therefore four times the rectangle BD , DC , together with four times the square on DE is equal to four times the square on EC

But the square on A is equal to four times the rectangle BD , DC and the square on DF is equal to four times the square on DE , for DF is double of DE

And the square on BC is equal to four times the square on EC , for again BC is double of CE

Therefore the squares on A , DF are equal to the square on BC , so that the square on BC is greater than the square on A by the square on DF

It is to be proved that BC is also commensurable with DF

Since BD is commensurable in length with DC

therefore BC is also commensurable in length with CD [x 15]

But CD is commensurable in length with CD , BF for CD is equal to BF [x 6]

Therefore BC is also commensurable in length with BF , CD [x 12]

so that BC is also commensurable in length with the remainder FD [x 15] therefore the square on BC is greater than the square on A by the square on a straight line commensurable with BC

Next let the square on BC be greater than the square on A by the square on a straight line commensurable with BC

let a parallelogram be applied to BC equal to the fourth part of the square on A and deficient by a square figure and let it be the rectangle BD , DC

It is to be proved that BD is commensurable in length with DC

With the same construction we can prove similarly that the square on BC is greater than the square on A by the square on FD

But the square on BC is greater than the square on A by the square on a straight line commensurable with BC

Therefore BC is commensurable in length with FD

so that BC is also commensurable in length with the remainder the sum of BF , DC [x 15]

But the sum of BF , DC is commensurable with DC [x 6]

so that BC is also commensurable in length with CD [x 12]

and therefore *separando* BD is commensurable in length with DC [x 15]

Therefore etc Q E D

PROPOSITION 18

If there be two unequal straight lines and to the greater there be applied a parallelogram equal to the fourth part of the square on the less and deficient by a square

figure, and if it divide it into parts which are incommensurable the square on the greater will be greater than the square on the less by the square on a straight line incommensurable with the greater

And if the square on the greater be greater than the square on the less by the square on a straight line incommensurable with the greater, and if there be applied to the greater a parallelogram equal to the fourth part of the square on the less and deficient by a square figure it divides it into parts which are incommensurable

Let A, BC be two unequal straight lines of which BC is the greater, and to BC let there be applied a parallelogram equal to the fourth part of the square on the less A , and deficient by a square figure Let this be the rectangle BD, DC , [cf Lemma before $\propto 17$] and let BD be incommensurable in length with DC , I say that the square on BC is greater than the square on A by the square on a straight line incommensurable with BC

For with the same construction as before, we can prove similarly that the square on BC is greater than the square on A by the square on FD

It is to be proved that BC is incommensurable in length with DF

Since BD is incommensurable in length with DC , therefore BC is also incommensurable in length with CD [$\propto 16$]

But DC is commensurable with the sum of BF, DC , [$\propto 6$]

therefore BC is also incommensurable with the sum of BF, DC [$\propto 13$]

so that BC is also incommensurable in length with the remainder FD [$\propto 16$]

And the square on BC is greater than the square on A by the square on FD , therefore the square on BC is greater than the square on A by the square on a straight line incommensurable with BC

Again, let the square on BC be greater than the square on A by the square on a straight line incommensurable with BC and let there be applied to BC a parallelogram equal to the fourth part of the square on A and deficient by a square figure Let this be the rectangle BD, DC

It is to be proved that BD is incommensurable in length with DC

For with the same construction, we can prove similarly that the square on BC is greater than the square on A by the square on FD

But the square on BC is greater than the square on A by the square on a straight line incommensurable with BC ,

therefore BC is incommensurable in length with FD , so that BC is also incommensurable with the remainder the sum of BF, DC [$\propto 16$]

But the sum of BF, DC is commensurable in length with DC , [$\propto 6$]

therefore BC is also incommensurable in length with DC [$\propto 13$]

so that, *separando* BD is also incommensurable in length with DC [$\propto 16$]

Therefore etc Q E D

LEMMA

Since it has been proved that straight lines commensurable in length are always commensurable in square also while those commensurable in square are not always commensurable in length also but can of course be either commensurable or incommensurable in length it is manifest that if any straight line be commensurable in length with a given rational straight line, it is called

rational and commensurable with the other not only in length but in square also, since straight lines commensurable in length are always commensurable in square also

But, if any straight line be commensurable in square with a given rational straight line then, if it is also commensurable in length with it it is called in this case also rational and commensurable with it both in length and in square but if again any straight line being commensurable in square with a given rational straight line be incommensurable in length with it, it is called in this case also rational but commensurable in square only

PROPOSITION 19

The rectangle contained by rational straight lines commensurable in length is rational

For let the rectangle AC be contained by the rational straight lines AB , BC commensurable in length,

I say that AC is rational

For on AB let the square AD be described,

therefore AD is rational [x Def 4]

And since AB is commensurable in length with BC ,

while AB is equal to BD ,

therefore BD is commensurable in length with BC

And as BD is to BC , so is DA to AC [VI 1]

Therefore DA is commensurable with AC [x 11]


But DA is rational

therefore AC is also rational

[x Def 4]

Therefore etc

Q E D



PROPOSITION 20

If a rational area be applied to a rational straight line it produces as breadth a straight line rational and commensurable in length with the straight line to which it is applied

For let the rational area AC be applied to AB , a straight line once more rational in any of the aforesaid ways producing BC as breadth,

I say that BC is rational and commensurable in length with BA

For on AB let the square AD be described

therefore AD is rational [x Def 4]

But AC is also rational

therefore DA is commensurable with AC

And as DA is to AC so is DB to BC

Therefore DB is also commensurable with BC , [VI 1]

and DB is equal to BA ,


therefore AB is also commensurable with BC

But AB is rational

therefore BC is also rational and commensurable in length with AB

Therefore etc

Q E D




PROPOSITION 21

The rectangle contained by rational straight lines commensurable in square only is irrational and the side of the square equal to it is irrational Let the latter be called medial

For let the rectangle AC be contained by the rational straight lines AB , BC commensurable in square only,

I say that AC is irrational, and the side of the square equal to it is irrational, and let the latter be called *medial*



For on AB let the square AD be described,

therefore AD is rational

[x Def 4]

And since AB is incommensurable in length with BC

for by hypothesis they are commensurable in square only,

while $4B$ is equal to BD

therefore DB is also incommensurable in length with BC

And, as DB is to BC , so is AD to AC

[vi 1]

therefore DA is incommensurable with AC

[x 11]

But DA is rational,

therefore AC is irrational

so that the side of the square equal to AC is also irrational

[x Def 4]

And let the latter be called *medial*

Q E D

LEMMA


If there be two straight lines then, as the first is to the second so is the square on the first to the rectangle contained by the two straight lines

Let FE , EG be two straight lines

I say that, as FE is to EG so is the square on FE to the rectangle FE , EG

For on FE let the square DF be described

and let GD be completed



Since then, as FE is to EG so is FD to DG ,

[vi 1]

and FD is the square on FE

and DG the rectangle DE EG that is the rectangle FE EG

therefore as FE is to EG so is the square on FE to the rectangle FE , EG

Similarly also as the rectangle GE , EF is to the square on EF that is as GD is to FD , so is GE to EF

Q E D

PROPOSITION 22

The square on a medial straight line if applied to a rational straight line produces as breadth a straight line rational and incommensurable in length with that to which it is applied



Let A be medial and CB rational

and let a rectangular area BD equal to the square on A be applied to BC producing CD as breadth

I say that CD is rational and incommensurable in length with CB

For since A is medial the square on it is equal to a rectangular area contained by rational straight lines commensurable in square only

[x 21]

Let the square on it be equal to GF

But the square on it is also equal to BD

therefore BD is equal to GF



But it is also equiangular with it,
and in equal and equiangular parallelograms the sides about the equal angles
are reciprocally proportional, [vi 14]

therefore, proportionally, as BC is to EG , so is EF to CD

Therefore also as the square on BC is to the square on EG , so is the square
on EF to the square on CD [vi 22]

But the square on CB is commensurable with the square on EG , for each of
these straight lines is rational

therefore the square on EF is also commensurable with the square on CD [x 11]

But the square on EF is rational,

therefore the square on CD is also rational, [x Def 4]

therefore CD is rational

And since EF is incommensurable in length with EG ,

for they are commensurable in square only,

and as EF is to EG so is the square on EF to the rectangle FE, EG , [Lemma]

therefore the square on EF is incommensurable with the rectangle FE, EG [x 11]

But the square on CD is commensurable with the square on EF , for the
straight lines are rational in square

and the rectangle DC, CB is commensurable with the rectangle FE, EG for
they are equal to the square on A ,

therefore the square on CD is also incommensurable with the rectangle DC
 CB [x 13]

But as the square on CD is to the rectangle DC, CB , so is DC to CB , [Lemma]

therefore DC is incommensurable in length with CB [x 11]

Therefore CD is rational and incommensurable in length with CB

Q E D

PROPOSITION 23

A straight line commensurable with a medial straight line is medial

Let A be medial and let B be commensurable with A ,

I say that B is also medial

For let a rational straight line CD be set out
and to CD let the rectangular area CE equal to the
square on A be applied producing ED as breadth,
therefore ED is rational and incommensurable in
length with CD [x 22]

And let the rectangular area CF equal to the square
on B be applied to CD producing DF as breadth

Since, then A is commensurable with B
the square on A is also commensurable with the
square on B


But EC is equal to the square on A

and CF is equal to the square on B ,

therefore EC is commensurable with CF

And as EC is to CF so is ED to DF

therefore ED is commensurable in length with DF



[vi 1]

[x 11]

But ED is rational and incommensurable in length with DC
 therefore DF is also rational [x Def 3] and incommensurable in length with DC [x 13]

Therefore CD, DF are rational and commensurable in square only

But the straight line the square on which is equal to the rectangle contained by rational straight lines commensurable in square only is medial, [x 21]
 therefore the side of the square equal to the rectangle CD, DF is medial

And B is the side of the square equal to the rectangle CD, DF ,

therefore B is medial Q E D

PORISM From this it is manifest that an area commensurable with a medial area is medial

[And in the same way as was explained in the case of rationals [Lemma following x 18] it follows as regards medials, that a straight line commensurable in length with a medial straight line is called *medial* and commensurable with it not only in length but in square also since in general, straight lines commensurable in length are always commensurable in square also

But if any straight line be commensurable in square with a medial straight line then, if it is also commensurable in length with it, the straight lines are called in this case too medial and commensurable in length and in square but if in square only, they are called medial straight lines commensurable in square only]

PROPOSITION 24

The rectangle contained by medial straight lines commensurable in length is medial

For let the rectangle AC be contained by the medial straight lines AB, BC which are commensurable in length

I say that AC is medial

For on AB let the square AD be described

therefore AD is medial

And, since AB is commensurable in length with BC

while AB is equal to BD ,

therefore DB is also commensurable in length with BC ,

so that DA is also commensurable with AC [vi 1 x 11]

But DA is medial

therefore AC is also medial

[x 23 Por.]

Q E D

PROPOSITION 25

The rectangle contained by medial straight lines commensurable in square only is either rational or medial

For let the rectangle AC be contained by the medial straight lines AB, BC which are commensurable in square only

I say that AC is either rational or medial

For on AB, BC let the squares AD, BE be described

therefore each of the squares AD, BE is medial

Let a rational straight line FG be set out

to FG let there be applied the rectangular parallelogram GH equal to AD producing FH as breadth

to HM let there be applied the rectangular parallelogram MA equal to AC , producing HA as breadth



and further to AN let there be similarly applied AL equal to BE producing KL as breadth,

therefore FH HA AL are in a straight line

Since then each of the squares AD , BE
is medial

and AD is equal to GH , and BE to NL
therefore each of the rectangles GH , NL is
also medial

And they are applied to the rational
straight line FG

therefore each of the straight lines FH KL
is rational and incommensurable in length
with FG [x 22]

And since AD is commensurable with BE

therefore GH is also commensurable with NL

And as GH is to NL so is GH to KL

therefore FH is commensurable in length with KL

Therefore FH KL are rational straight lines commensurable in length,

therefore the rectangle FH KL is rational

And since DB is equal to BA and OB to BC ,

therefore as DB is to BC so is AB to BO

But as DB is to BC so is DA to AC

and as AB is to BO so is AC to CO ,

therefore as DA is to AC so is AC to CO

But AD is equal to GH AC to MA and CO to NL ,

therefore as GH is to MA , so is MA to NL

therefore also as FH is to HA so is HA to AL

therefore the rectangle FH KL is equal to the square on HA

But the rectangle FH KL is rational

therefore the square on HA is also rational

Therefore HA is rational

And if it is commensurable in length with FG

HN is rational

but if it is incommensurable in length with FG

HA HM are rational straight lines commensurable in square only and there-
fore HN is medial


Therefore HN is either rational or medial

But HA is equal to AC

therefore AC is either rational or medial

Therefore etc

Q E D



PROPOSITION 26

A medial area does not exceed a medial area by a rational area

For if possible let the medial area AB exceed the medial area AC by the rational area DB

and let a rational straight line EF be set out
to EF let there be applied the rectangular parallelogram FH equal to AB pro-
ducing EH as breadth

and let the rectangle FG equal to AC be subtracted

therefore the remainder BD is equal to the remainder ΛH

But DB is rational

therefore ΛH is also rational

Since then each of the rectangles AB , AC is medial

and AB is equal to FH and AC to FG

therefore each of the rectangles FH , FG is also medial

And they are applied to the rational straight line EF ,

therefore each of the straight lines HE EG is rational and incommensurable in length with EF [x 22]

And, since [DB is rational and is equal to ΛH ,

therefore] KH is [also] rational,

and it is applied to the rational straight line EF

therefore GH is rational and commensurable in length with EF [x 20]

But EG is also rational and is incommensurable in length with EF

therefore EG is incommensurable in length with GH [x 13]

And as EG is to GH so is the square on EG to the rectangle EG , GH , therefore the square on EG is incommensurable with the rectangle EG , GH [x 11]

But the squares on EG , GH are commensurable with the square on EG for both are rational

and twice the rectangle EG , GH is commensurable with the rectangle EG , GH for it is double of it, [x 6]

therefore the squares on EG , GH are incommensurable with twice the rectangle EG , GH [x 13]

therefore also the sum of the squares on EG , GH and twice the rectangle EG , GH , that is the square on EH [II 4] is incommensurable with the squares on EG , GH [x 16]

But the squares on EG , GH are rational

therefore the square on EH is irrational [x Def 4]

Therefore EH is irrational

But it is also rational

which is impossible

Therefore etc

Q E D

PROPOSITION 27

To find medial straight lines commensurable in square only which contain a rational rectangle

Let two rational straight lines A , B commensurable in square only be set out,

let C be taken a mean proportional between A , B [VI 13]


and let it be contrived that

as A is to B , so is C to D [VI 12]

Then since A , B are rational and commensurable in square only

the rectangle A , B that is the square on C

[VI 17] is medial [x 21]



Therefore C is medial [x 21]
 And since as A is to B , so is C to D ,
 and A, B are commensurable in square only — —
 therefore C, D are also commensurable in square only [x 11]
 And C is medial,
 therefore D is also medial [x 23, addition]
 Therefore C, D are medial and commensurable in square only
 I say that they also contain a rational rectangle
 For since as A is to B so is C to D
 therefore alternately as A is to C so is B to D [v 16]
 But as A is to C so is C to B ,
 therefore also as C is to B so is B to D
 therefore the rectangle $C D$ is equal to the square on B
 But the square on B is rational
 therefore the rectangle $C D$ is also rational
 Therefore medial straight lines commensurable in square only have been
 found which contain a rational rectangle Q E D

PROPOSITION 28

To find medial straight lines commensurable in square only which contain a medial rectangle

Let the rational straight lines A, B, C commensurable in square only be set out

let D be taken a mean proportional between A, B , [v 13]
 and let it be contrived that
 as B is to C so is D to E [vi 12]

Since A, B are rational straight
 lines commensurable in square only
 therefore the rectangle $A B$ that
 is the square on D [vi 17] is medi
 al [x 21]



Therefore D is medial [x 21]
 And since B, C are commensurable in square only
 and as B is to C so is D to E
 therefore D, E are also commensurable in square only [x 11]
 But D is medial

therefore E is also medial [x 23 addition]
 Therefore D, E are medial straight lines commensurable in square only

I say next that they also contain a medial rectangle

For since as B is to C so is D to E
 therefore alternately as B is to D so is C to E [v 16]
 But as B is to D so is D to A

therefore also as D is to A so is C to E
 therefore the rectangle $A C$ is equal to the rectangle $D E$ [vi 16]
 But the rectangle $A C$ is medial [x 21]


therefore the rectangle $D E$ is also medial

Therefore medial straight lines commensurable in square only have been
 found which contain a medial rectangle Q E D

LEMMA 1

To find two square numbers such that their sum is also square

Let two numbers AB , BC be set out, and let them be either both even or both odd



Then since whether an even number is subtracted from an even number, or an odd number from an odd number the remainder is even [ix 24 26]

therefore the remainder AC is even

Let AC be bisected at D

Let AB , BC also be either similar plane numbers or square numbers which are themselves also similar plane numbers

Now the product of AB , BC together with the square on CD is equal to the square on BD [ii 6]

And the product of AB , BC is square inasmuch as it was proved that if two similar plane numbers by multiplying one another make some number the product is square [ix 1]

Therefore two square numbers, the product of AB , BC , and the square on CD have been found which when added together make the square on BD

And it is manifest that two square numbers, the square on BD and the square on CD , have again been found such that their difference, the product of AB , BC is a square, whenever AB , BC are similar plane numbers


But when they are not similar plane numbers two square numbers the square on BD and the square on DC , have been found such that their difference the product of AB , BC is not square Q E D

LEMMA 2

To find two square numbers such that their sum is not square

For let the product of AB , BC as we said be square and CA even

and let CA be bisected by D



It is then manifest that the square product of AB , BC together with the square on CD is equal to the square on BD

[See Lemma 1]

Let the unit DE be subtracted, therefore the product of AB , BC together with the square on CE is less than the square on BD

I say then that the square product of AB , BC together with the square on CE will not be square

For if it is square it is either equal to the square on BE or less than the square on BE but cannot any more be greater lest the unit be divided

First if possible, let the product of AB , BC together with the square on CE be equal to the square on BE

and let GA be double of the unit DE

Since then the whole AC is double of the whole CD

and in them AG is double of DE

therefore the remainder GC is also double of the remainder EC ,

therefore GC is bisected by E

Therefore the product of GB , BC together with the square on CE is equal to the square on BE [11 6]

But the product of AB , BC together with the square on CE is also, by hypothesis equal to the square on BE ,
therefore the product of GB , BC together with the square on CE is equal to the product of AB , BC together with the square on CE

And, if the common square on CE be subtracted,
it follows that AB is equal to GB
which is absurd

Therefore the product of AB , BC together with the square on CE is not equal to the square on BE

I say next that neither is it less than the square on BE

For, if possible, let it be equal to the square on BF
and let HA be double of DF

Now it will again follow that HC is double of CF
so that CH has also been bisected at F

and for this reason the product of HB , BC together with the square on FC is equal to the square on BF [11 6]

But, by hypothesis the product of AB , BC together with the square on CE is also equal to the square on BF

Thus the product of HB , BC together with the square on CF will also be equal to the product of AB , BC together with the square on CE
which is absurd

Therefore the product of AB , BC together with the square on CE is not less than the square on BE

And it was proved that neither is it equal to the square on BE

Therefore the product of AB , BC together with the square on CE is not square
Q E D

PROPOSITION 29

To find two rational straight lines commensurable in square only and such that the square on the greater is greater than the square on the less by the square on a straight line commensurable in length with the greater

For let there be set out any rational straight line AB and two square numbers CD , DE such that their difference CE is not square, [Lemma 1]

let there be described on AB the semicircle AFB

and let it be contrived that

as DC is to CE so is the square on BA to the square on AF [x 6 Por]

Let FB be joined

Since as the square on BA is to the square on AF ,
so is DC to CE


therefore the square on BA has to the square on AF
the ratio which the number DC has to the number CE ,

therefore the square on BA is commensurable with the square on AF [x 6]

But the square on AB is rational [x Def 4]

therefore the square on AF is also rational

therefore AF is also rational [11]



And since DC has not to CE the ratio which a square number has to a square number
neither has the square on BA to the square on AF the ratio which a square number has to a square number

therefore AB is incommensurable in length with AF [x 9]

Therefore BA, AF are rational straight lines commensurable in square only
And since as DC is to CE , so is the square on BA to the square on AF ,
therefore, *convertendo* as CD is to DE , so is the square on AB to the square on BF [v 19 Por, III 31 i 47]

But CD has to DE the ratio which a square number has to a square number
therefore also the square on AB has to the square on BF the ratio which a square number has to a square number,

therefore AB is commensurable in length with BF [x 9]

And the square on AB is equal to the squares on AF, FB
therefore the square on AB is greater than the square on AF by the square on BF commensurable with AB

Therefore there have been found two rational straight lines BA, AF commensurable in square only and such that the square on the greater AB is greater than the square on the less AF by the square on BF commensurable in length with AB Q E D

PROPOSITION 30

To find two rational straight lines commensurable in square only and such that the square on the greater is greater than the square on the less by the square on a straight line incommensurable in length with the greater

Let there be set out a rational straight line AB ,

and two square numbers CE, ED such that their sum CD is not square [Lemma 2]

let there be described on AB the semicircle AFB ,

let it be contrived that

as DC is to CE , so is the square on BA to the square on AF [x 6, Por]

and let FB be joined

Then in a similar manner to the preceding we can prove that BA, AF are rational straight lines commensurable in square only

And since as DC is to CE , so is the square on BA to the square on AF
therefore *convertendo* as CD is to DE so is the square on AB to the square on BF [v 19 Por III 31 i 47]


But CD has not to DE the ratio which a square number has to a square number

therefore neither has the square on AB to the square on BF the ratio which a square number has to a square number,

therefore AB is incommensurable in length with BF [x 9]

And the square on AB is greater than the square on AF by the square on FB incommensurable with AB

Therefore AB, AF are rational straight lines commensurable in square only and the square on AB is greater than the square on AF by the square on FB incommensurable in length with AB Q E D



PROPOSITION 31

To find two medial straight lines commensurable in square only, containing a rational rectangle and such that the square on the greater is greater than the square on the less by the square on a straight line commensurable in length with the greater

Let there be set out two rational straight lines A B commensurable in square only and such that the square on A being the greater is greater than the square on B the less by the square on a straight line commensurable in length with A [x 29]

And let the square on C be equal to the rectangle A B

Now the rectangle A B is medial [x 21]

therefore the square on C is also medial,

therefore C is also medial [x 21]

Let the rectangle C D be equal to the square on B

Now the square on B is rational

therefore the rectangle C D is also rational

And since as A is to B so is the rectangle A B to the square on B

while the square on C is equal to the rectangle A B ,

and the rectangle C D is equal to the square on B ,

therefore as A is to B so is the square on C to the rectangle C D ,

But as the square on C is to the rectangle C D so is C to D ,

therefore also as A is to B so is C to D

But A is commensurable with B in square only

therefore C is also commensurable with D in square only [x 11]

And C is medial

therefore D is also medial [x 23 addition]

And since as A is to B so is C to D

and the square on A is greater than the square on B by the square on a straight line commensurable with A ,

therefore also the square on C is greater than the square on D by the square on a straight line commensurable with C [x 14]

Therefore two medial straight lines C D commensurable in square only and containing a rational rectangle have been found and the square on C is greater than the square on D by the square on a straight line commensurable in length with C

Similarly also it can be proved that the square on C exceeds the square on D by the square on a straight line incommensurable with C when the square on A is greater than the square on B by the square on a straight line incommensurable with A [x 30]


PROPOSITION 32

To find two medial straight lines commensurable in square only containing a medial rectangle and such that the square on the greater is greater than the square on the less by the square on a straight line commensurable with the greater

Let there be set out three rational straight lines A B C commensurable in square only and such that the square on A is greater than the square on C by the square on a straight line commensurable with A [x 29]

and let the square on D be equal to the rectangle A B

Therefore the square on D is medial



therefore D is also medial

[x 21]

Let the rectangle D, E be equal to the rectangle B, C

A _____

D _____

B _____

E _____

C _____

Then since, as the rectangle A, B is to the rectangle B, C , so is A to C ,

while the square on D is equal to the rectangle A, B ,

and the rectangle D, E is equal to the rectangle B, C ,

therefore, as A is to C , so is the square on D to the rectangle D, E

But as the square on D is to the rectangle D, E , so is D to E ,

therefore also as A is to C , so is D to E

But A is commensurable with C in square only,

therefore D is also commensurable with E in square only [x 11]

But D is medial,

therefore E is also medial

[x 23 addition]

And, since, as A is to C , so is D to E ,

while the square on A is greater than the square on C by the square on a straight line commensurable with A ,

therefore also the square on D will be greater than the square on E by the square on a straight line commensurable with D [x 14]

I say next that the rectangle D, E is also medial

For, since the rectangle B, C is equal to the rectangle D, E , while the rectangle B, C is medial [x 21]


therefore the rectangle D, E is also medial

Therefore two medial straight lines D, E , commensurable in square only and containing a medial rectangle have been found such that the square on the greater is greater than the square on the less by the square on a straight line commensurable with the greater

Similarly again it can be proved that the square on D is greater than the square on E by the square on a straight line incommensurable with D , when the square on A is greater than the square on C by the square on a straight line incommensurable with A [x 30]

LEMMA

Let ABC be a right angled triangle having the angle A right and let the perpendicular AD be drawn,



I say that the rectangle CB, BD is equal to the square on BA ,

the rectangle BC, CD equal to the square on CA ,

the rectangle BD, DC equal to the square on AD

and further, the rectangle BC, AD equal to the rectangle BA, AC

And first that the rectangle CB, BD is equal to the square on BA

For since in a right angled triangle AD has been drawn from the right angle perpendicular to the base therefore the triangles ABD, ADC are similar both to the whole ABC and to one another [vi 8]

And since the triangle ABC is similar to the triangle ABD , therefore as CB is to BA so is BA to BD , [vi 4]

therefore the rectangle CB BD is equal to the square on AB [VI 14]

For the same reason the rectangle BC CD is also equal to the square on AC

And since if in a right angled triangle a perpendicular be drawn from the right angle to the base the perpendicular so drawn is a mean proportional between the segments of the base [VI 8 Por]

therefore as BD is to DA , so is AD to DC ,

therefore the rectangle BD DC is equal to the square on AD [VI 17]

I say that the rectangle BC AD is also equal to the rectangle BA AC

For since as we said ABC is similar to ABD ,

therefore as BC is to CA so is BA to AD [VI 4]

Therefore the rectangle BC AD is equal to the rectangle BA AC [VI 16]

Q E D


PROPOSITION 33

To find two straight lines incommensurable in square which make the sum of the squares on them rational but the rectangle contained by them medial

Let there be set out two rational straight lines AB BC commensurable in square only and such that the square on the greater AB is greater than the square on the less BC by the square on a straight line incommensurable with AB [X 30]

let BC be bisected at D

let there be applied to AB a parallelogram equal to the square on either of the straight lines BD DC and deficient by a square figure and let it be the rectangle AE EB , [VI 28]



let the semicircle AFB be described on AB ,

let EF be drawn at right angles to AB ,

and let AF FB be joined

Then since AB BC are unequal straight lines and the square on AB is greater than the square on BC by the square on a straight line incommensurable with AB while there has been applied to AB a parallelogram equal to the fourth part of the square on BC that is to the square on half of it and deficient by a square figure making the rectangle AE EB ,

therefore AE is incommensurable with EB [X 18]

And as AE is to EB so is the rectangle BA AE to the rectangle AB BE

while the rectangle BA AE is equal to the square on AF

and the rectangle AB BE to the square on BF

therefore the square on AF is incommensurable with the square on BF ,

therefore AF BF are incommensurable in square

And since AB is rational

therefore the square on AB is also rational,

so that the sum of the squares on AF BF is also rational [I 4]

And since again the rectangle AE EB is equal to the square on EF

and by hypothesis the rectangle AE EB is also equal to the square on BD

therefore FE is equal to BD

therefore BC is double of FE

so that the rectangle AB BC is also commensurable with the rectangle AB EF

But the rectangle AB BC is medial, [x 21]
 therefore the rectangle AB , EF is also medial [x 23 Por]
 But the rectangle AB , EF is equal to the rectangle AF FB , [Lemma]
 therefore the rectangle AF FB is also medial

But it was also proved that the sum of the squares on these straight lines is rational

Therefore two straight lines AF , FB incommensurable in square have been found which make the sum of the squares on them rational but the rectangle contained by them medial Q E D

PROPOSITION 34


To find two straight lines incommensurable in square which make the sum of the squares on them medial but the rectangle contained by them rational

Let there be set out two medial straight lines AB BC , commensurable in square only such that the rectangle which they contain is rational and the square on AB is greater than the square on BC by the square on a straight line incommensurable with AB , [x 31, ad fin]

let the semicircle ADB be described on AB ,

let BC be bisected at E

let there be applied to AB a parallelogram equal to the square on BE and deficient by a square figure, namely the rectangle AF FB [v 123]



therefore AF is incommensurable in length with FB [x 18]

Let FD be drawn from F at right angles to AB ,
 and let AD DB be joined

Since AF is incommensurable in length with FB
 therefore the rectangle BA , AF is also incommensurable with the rectangle AB , BF [x 11]

But the rectangle BA AF is equal to the square on AD and the rectangle AB , BF to the square on DB ,

therefore the square on AD is also incommensurable with the square on DB

And since the square on AB is medial

therefore the sum of the squares on AD DB is also medial [III 31 + 47]

And since BC is double of DF ,

therefore the rectangle AB BC is also double of the rectangle AB FD

But the rectangle AB BC is rational

therefore the rectangle AB FD is also rational [x 6]

But the rectangle AB FD is equal to the rectangle AD DB [Lemma]

so that the rectangle AD DB is also rational

Therefore two straight lines AD DB incommensurable in square have been found which make the sum of the squares on them medial but the rectangle contained by them rational Q E D

PROPOSITION 35

To find two straight lines incommensurable in square which make the sum of the squares on them medial and the rectangle contained by them medial and moreover incommensurable with the sum of the squares on them

Let there be set out two medial straight lines AB , BC commensurable in square only containing a medial rectangle and such that the square on AB is greater than the square on BC by the square on a straight line incommensurable with AB , [x 32, *ad fin*]

let the semicircle ADB be described on AB ,
and let the rest of the construction be as above

Then, since AF is incommensurable in length with FB [x 18]
 AD is also incommensurable in square with DB [x 11]

And since the square on AB is medial
therefore the sum of the squares on AD , DB is also medial [III 31 i 47]

And since the rectangle AF , FB is equal to the square on each of the straight lines BE , DF ,

therefore BE is equal to DF ,⁽¹⁾
therefore BC is double of FD

so that the rectangle AB , BC is also double of the rectangle AB , FD

But the rectangle AB , BC is medial,

therefore the rectangle AB , FD is also medial [x 32 *Por*]

And it is equal to the rectangle AD , DB , [Lemma after x 3^o]

therefore the rectangle AD , DB is also medial

And since AB is incommensurable in length with BC ,
while CB is commensurable with BE ,

therefore AB is also incommensurable in length with BE [x 13]

so that the square on AB is also incommensurable with the rectangle AB , BE [x 11]

But the squares on AD , DB are equal to the square on AB , [i 47]
and the rectangle AB , FD , that is the rectangle AD , DB , is equal to the rectangle AB , BE

therefore the sum of the squares on AD , DB is incommensurable with the rectangle AD , DB

Therefore two straight lines AD , DB incommensurable in square have been found which make the sum of the squares on them medial and the rectangle contained by them medial and moreover incommensurable with the sum of the squares on them

Q E D


PROPOSITION 36

If two rational straight lines commensurable in square only be added together, the whole is irrational and let it be called binomial

For let two rational straight lines AB , BC commensurable in square only be added together

I say that the whole AC is irrational

For since AB is incommensurable in length with BC —for they are commensurable in square only—
and as AB is to BC , so is the rectangle AB , BC to the square on BC ,
therefore the rectangle AB , BC is incommensurable with the square on BC [x 11]



But twice the rectangle AB, BC is commensurable with the rectangle AB, BC [x 6], and the squares on AB, BC are commensurable with the square on BC —for AB, BC are rational straight lines commensurable in square only—

[x 15]

therefore twice the rectangle AB, BC is incommensurable with the squares on AB, BC

[x 13]

And, *componendo* twice the rectangle AB, BC together with the squares on AB, BC , that is, the square on AC [II 4] is incommensurable with the sum of the squares on AB, BC

[x 16]

But the sum of the squares on AB, BC is rational

therefore the square on AC is irrational

so that AC is also irrational

[x Def 4]

And let it be called *binomial*

Q E D

PROPOSITION 37

If two medial straight lines commensurable in square only and containing a rational rectangle be added together the whole is irrational, and let it be called a first binomial straight line

For let two medial straight lines AB, BC commensurable in square only and containing a rational rectangle be added together

I say that the whole AC is irrational

For since AB is incommensurable in length with BC

therefore the squares on AB, BC are also incommensurable with twice the rectangle AB, BC ,

[cf x 36, II 9-20]

and *componendo* the squares on AB, BC together with twice the rectangle AB, BC , that is the square on AC [II 4] is incommensurable with the rectangle AB, BC

[x 16]

But the rectangle AB, BC is rational for by hypothesis AB, BC are straight lines containing a rational rectangle

therefore the square on AC is irrational,

therefore AC is irrational

[x Def 4]

And let it be called a first binomial straight line

Q E D

PROPOSITION 38

If two medial straight lines commensurable in square only and containing a medial rectangle be added together the whole is irrational and let it be called a second binomial straight line

For let two medial straight lines AB, BC commensurable in square only and containing a medial rectangle be added together

A B C

I say that AC is irrational

For let a rational straight line DE be set out and let the parallelogram DF equal to the square on AC be applied to DE producing DG as breadth


[I 44]

Then since the square on AC is equal to

the squares on AB, BC and twice the rectangle AB, BC

[II 4]

let EH equal to the squares on AB, BC be applied to DE



therefore the remainder HF is equal to twice the rectangle AB, BC

And, since each of the straight lines AB, BC is medial

therefore the squares on AB, BC are also medial

But by hypothesis, twice the rectangle AB, BC is also medial

And EH is equal to the squares on AB, BC ,

while FH is equal to twice the rectangle AB, BC ,

therefore each of the rectangles EH, HF is medial

And they are applied to the rational straight line DE ,

therefore each of the straight lines DH, HG is rational and incommensurable in length with DE [x 2]

Since then AB is incommensurable in length with BC ,

and as AB is to BC so is the square on AB to the rectangle AB, BC ,

therefore the square on AB is incommensurable with the rectangle AB, BC [x 11]

But the sum of the squares on AB, BC is commensurable with the square on AB [x 15]

and twice the rectangle AB, BC is commensurable with the rectangle AB, BC [x 6]

Therefore the sum of the squares on AB, BC is incommensurable with twice the rectangle AB, BC [x 13]

But EH is equal to the squares on AB, BC ,

and FH is equal to twice the rectangle AB, BC

Therefore EH is incommensurable with FH

so that DH is also incommensurable in length with HG [vi 1, x 11]

Therefore DH, HG are rational straight lines commensurable in square only, so that DG is irrational [x 36]

But DE is rational,

and the rectangle contained by an irrational and a rational straight line is irrational [cf x 20]

therefore the area DF is irrational

and the side of the square equal to it is irrational [x Def 4]

But AC is the side of the square equal to DF

therefore AC is irrational


And let it be called a second binomial straight line Q E D

PROPOSITION 39

If two straight lines incommensurable in square which make the sum of the squares on them rational but the rectangle contained by them medial, be added together, the whole straight line is irrational and let it be called major

I or let two straight lines AB, BC incommensurable in square and fulfilling the given conditions [x 33] be added together

I say that AC is irrational



For, since the rectangle AB, BC is medial

twice the rectangle AB, BC is also medial [x 6 and 23 Por]

But the sum of the squares on AB, BC is rational

therefore twice the rectangle AB, BC is incommensurable with the sum of the squares on AB, BC

so that the squares on AB, BC together with twice the rectangle AB, BC that

is, the square on AC , is also incommensurable with the sum of the squares on AB, BC , [x 16]

therefore the square on AC is irrational

so that AC is also irrational

[x Def 4]

And let it be called *major*

Q E D

PROPOSITION 40

If two straight lines incommensurable in square which make the sum of the squares on them medial but the rectangle contained by them rational be added together, the whole straight line is irrational and let it be called the side of a rational plus a medial area

For let two straight lines AB, BC incommensurable in square and fulfilling the given conditions [x 34] be added together, I say that AC is irrational

For, since the sum of the squares on AB, BC is medial while twice the rectangle AB, BC is rational

therefore the sum of the squares on AB, BC is incommensurable with twice the rectangle AB, BC ,

so that the square on AC is also incommensurable with twice the rectangle AB, BC [x 16]

But twice the rectangle AB, BC is rational,

therefore the square on AC is irrational

Therefore AC is irrational

[x Def 4]

And let it be called the *side of a rational plus a medial area*

Q E D

PROPOSITION 41

If two straight lines incommensurable in square which make the sum of the squares on them medial and the rectangle contained by them medial and also incommensurable with the sum of the squares on them be added together, the whole straight line is irrational and let it be called the side of the sum of two medial areas

For let two straight lines AB, BC incommensurable in square and satisfying the given conditions [x 35] be added together

I say that AC is irrational

Let a rational straight line DE be set out and let there be applied to DE the rectangle DF equal to the squares on AB, BC , and the rectangle GH equal to twice the rectangle AB, BC

therefore the whole DH is equal to the square on AC [II 4]

Now since the sum of the squares on AB, BC is medial, and is equal to DF ,


therefore DF is also medial

And it is applied to the rational straight line DE therefore DG is rational and incommensurable in length with DE [x 22]

For the same reason GK is also rational and incommensurable in length with GF that is DE

And since the squares on AB, BC are incommensurable with twice the rectangle AB, BC

DF is incommensurable with GH



so that DG is also incommensurable with GK [vi 1, x 11]

And they are rational

therefore DG GK are rational straight lines commensurable in square only,

therefore DA is irrational and what is called binomial [x 36]

But DE is rational

therefore DH is irrational and the side of the square which is equal to it is irrational [x Def 4]

But AC is the side of the square equal to HD ,
therefore AC is irrational

And let it be called the *side of the sum of two medial areas* Q E D

LEMMA

And that the aforesaid irrational straight lines are divided only in one way into the straight lines of which they are the sum and which produce the types in question we will now prove after premising the following lemma

Let the straight line AB be set out let the whole be cut into unequal parts at each of the points C D

and let AC be supposed greater than DB

I say that the squares on AC CB are greater than the squares on AD DB

For let AB be bisected at E

Then since AC is greater than DB ,

let DC be subtracted from each

therefore the remainder AD is greater than the remainder CB

But AE is equal to EB

therefore DE is less than EC

therefore the points C D are not equidistant from the point of bisection

And since the rectangle AC , CB together with the square on EC is equal to the square on EB [ii 5]

and further the rectangle AD DB together with the square on DE is equal to the square on EB [id]

therefore the rectangle AC CB together with the square on EC is equal to the rectangle AD DB together with the square on DE

And of these the square on DE is less than the square on EC ,
therefore the remainder the rectangle AC CB is also less than the rectangle AD DB

so that twice the rectangle AC CB is also less than twice the rectangle AD , DB

Therefore also the remainder the sum of the squares on AC , CB , is greater than the sum of the squares on AD DB Q E D

PROPOSITION 42


A binomial straight line is divided into its terms at one point only

Let AB be a binomial straight line divided into its terms at C ,

therefore AC CB are rational straight lines

commensurable in square only

I say that AB is not divided at another point into two rational straight lines commensurable in square only



For if possible let it be divided at D also so that AD DB are also rational straight lines commensurable in square only

It is then manifest that AC is not the same with DB

For, if possible let it be so

Then AD will also be the same as CB ,

and as AC is to CB , so will BD be to DA ,

thus AB will be divided at D also in the same way as by the division at C which is contrary to the hypothesis

Therefore AC is not the same with DB

For this reason also the points C , D are not equidistant from the point of bisection

Therefore that by which the squares on AC , CB differ from the squares on AD DB is also that by which twice the rectangle AD DB differs from twice the rectangle AC , CB ,

because both the squares on AC CB together with twice the rectangle AC , CB and the squares on AD DB together with twice the rectangle AD , DB are equal to the square on AB [II 4]

But the squares on AC , CB differ from the squares on AD , DB by a rational area

for both are rational

therefore twice the rectangle AD , DB also differs from twice the rectangle AC , CB by a rational area though they are medial [x 21]

which is absurd for a medial area does not exceed a medial by a rational area [x 26]


Therefore a binomial straight line is not divided at different points

therefore it is divided at one point only Q E D

PROPOSITION 43

A first binomial straight line is divided at one point only

Let AB be a first binomial straight line divided at C , so that AC CB are medial straight lines commensurable in square only and containing a rational rectangle; [x 37]



I say that AB is not so divided at another point

For if possible let it be divided at D also, so that AD , DB are also medial straight lines commensurable in square only and containing a rational rectangle

Since then that by which twice the rectangle AD DB differs from twice the rectangle AC CB is that by which the squares on AC CB differ from the squares on AD DB

while twice the rectangle AD DB differs from twice the rectangle AC , CB by a rational area—for both are rational—

therefore the squares on AC CB also differ from the squares on AD DB by a rational area though they are medial

which is absurd

[x 26]

Therefore a first binomial straight line is not divided into its terms at different points

therefore it is so divided at one point only

Q E D


PROPOSITION 44

A second bimedral straight line is divided at one point only

Let AB be a second bimedral straight line divided at C , so that AC , CB are medial straight lines commensurable in square only and containing a medial rectangle, [x 28]

it is then manifest that C is not at the point of bisection, because the segments are not commensurable in length

I say that AB is not so divided at another point



For if possible let it be divided at D also so that AC is not the same with DB , but AC is supposed greater

it is then clear that the squares on AD , DB are also as we proved above [Lemma] less than the squares on AC , CB

and suppose that AD , DB are medial straight lines commensurable in square only and containing a medial rectangle

Now let a rational straight line LF be set out

let there be applied to EF the rectangular parallelogram EA equal to the square on AB ,

and let EG , equal to the squares on AC , CB , be subtracted
therefore the remainder HK is equal to twice the rectangle AC , CB [x 4]

Again let there be subtracted EL equal to the squares on AD , DB , which were proved less than the squares on AC , CB [Lemma]

therefore the remainder MA is also equal to twice the rectangle AD , DB

Now since the squares on AC , CB are medial

therefore EG is medial

And it is applied to the rational straight line EF ,

therefore EH is rational and incommensurable in length with EF [x 27]

For the same reason

HN is also rational and incommensurable in length with EF

And since AC , CB are medial straight lines commensurable in square only

therefore AC is incommensurable in length with CB

But as AC is to CB so is the square on AC to the rectangle AC , CB ,
therefore the square on AC is incommensurable with the rectangle AC , CB [x 11]

But the squares on AC , CB are commensurable with the square on AC , for
 AC , CB are commensurable in square [x 15]

And twice the rectangle AC , CB is commensurable with the rectangle AC , CB [x 6]

Therefore the squares on AC , CB are also incommensurable with twice the rectangle AC , CB [x 13]

But EG is equal to the squares on AC CB ,
 and HA is equal to twice the rectangle AC , CB ,
 therefore EG is incommensurable with HA ,

so that EH is also incommensurable in length with HN [VI 1, x 11]

And they are rational

therefore EH HN are rational straight lines commensurable in square only

But if two rational straight lines commensurable in square only be added together the whole is the irrational which is called binomial [x 36]

Therefore EN is a binomial straight line divided at H [x 36]

In the same way EM , MN will also be proved to be rational straight lines commensurable in square only,

and EN will be a binomial straight line divided at different points H and M

And EH is not the same with MN

For the squares on AC CB are greater than the squares on AD , DB

But the squares on AD , DB are greater than twice the rectangle AD DB
 therefore also the squares on AC CB , that is, EG are much greater than twice the rectangle AD , DB that is MA

so that EH is also greater than MN


Therefore EH is not the same with MN

Q E D

PROPOSITION 45

A major straight line is divided at one and the same point only

Let AB be a major straight line divided at C , so that AC CB are incommensurable in square and make the sum of the squares on AC CB rational but the rectangle AC CB medial,



I say that AB is not so divided at another point

For if possible let it be divided at D also so that AD DB are also incommensurable in square and make the sum of the squares on AD DB rational but the rectangle contained by them medial

Then since that by which the squares on AC CB differ from the squares on AD DB is also that by which twice the rectangle AD DB differs from twice the rectangle AC CB

while the squares on AC CB exceed the squares on AD DB by a rational area—for both are rational—

therefore twice the rectangle AD DB also exceeds twice the rectangle AC CB by a rational area though they are medial

which is impossible

[x 26]


Therefore a major straight line is not divided at different points

therefore it is only divided at one and the same point Q E D

PROPOSITION 46

The side of a rational plus a medial area is divided at one point only

Let AB be the side of a rational plus a medial area divided at C so that AC , CB are incommensurable in square and make the sum of the squares on AC CB medial but twice the rectangle AC CB rational [x 40]



I say that AB is not so divided at another point

For if possible let it be divided at D also so that AD DB are also incommensurable in square and make the sum of the squares on AD DB medial but twice the rectangle AD DB rational

measurable in square and make the sum of the squares on AD DB medial but twice the rectangle AD , DB rational

Since then that by which twice the rectangle AC CB differs from twice the rectangle AD DB is also that by which the squares on AD DB differ from the squares on AC CB

while twice the rectangle AC CB exceeds twice the rectangle AD DB by a rational area,

therefore the squares on AD , DB also exceed the squares on AC , CB by a rational area though they are medial

which is impossible

[x 26]

Therefore the side of a rational plus a medial area is not divided at different points,

therefore it is divided at one point only

Q E D

PROPOSITION 47

The side of the sum of two medial areas is divided at one point only

Let AB be divided at C so that AC CB are incommensurable in square and make the sum of the squares on AC CB medial and the rectangle AC , CB medial and also incommensurable with the sum of the squares on them

I say that AB is not divided at another point so as to fulfil the given conditions

For if possible let it be divided at D so that again AC is of course not the same as BD but AC is supposed greater

let a rational straight line EF be set out

and let there be applied to EF the rectangle EG equal to the squares on AC CB

and the rectangle HA equal to twice the rectangle AC CB

therefore the whole EA is equal to the square on AB

[ii 4]

Again let EL equal to the squares on AD DB be applied to EF , therefore the remainder twice the rectangle AD DB is equal to the remainder MA

And since by hypothesis the sum of the squares on AC CB is medial therefore EG is also medial

And it is applied to the rational straight line EF

therefore HE is rational and incommensurable in length with EF

[x 27]

For the same reason

HA is also rational and incommensurable in length with EF

And since the sum of the squares on AC CB is incommensurable with twice the rectangle AC CB

therefore EG is also incommensurable with GN

so that EH is also incommensurable with HN


[xi 1 xi 11]

And they are rational

therefore EH HN are rational straight lines commensurable in square only,

therefore EN is a binomial straight line divided at H

[x 36]



Similarly we can prove that it is also divided at M

And EH is not the same with MN ,

therefore a binomial has been divided at different points

which is absurd [x 42]

Therefore a side of the sum of two medial areas is not divided at different points

therefore it is divided at one point only Q E D

DEFINITIONS II

1 Given a rational straight line and a binomial divided into its terms such that the square on the greater term is greater than the square on the lesser by the square on a straight line commensurable in length with the greater, then if the greater term be commensurable in length with the rational straight line set out, let the whole be called a *first binomial* straight line

2 but if the lesser term be commensurable in length with the rational straight line set out let the whole be called a *second binomial*

3 and if neither of the terms be commensurable in length with the rational straight line set out let the whole be called a *third binomial*


4 Again if the square on the greater term be greater than the square on the lesser by the square on a straight line incommensurable in length with the greater then if the greater term be commensurable in length with the rational straight line set out let the whole be called a *fourth binomial*,

5 if the lesser a *fifth binomial*,

6 and if neither a *sixth binomial*

PROPOSITION 48

To find the first binomial straight line

Let two numbers AC CB be set out such that the sum of them AB has to BC the ratio which a square number has to a square number but has not to CA the ratio which a square number has to a square number, [Lemma 1 after x 28]

 let any rational straight line D be set out and let EF be commensurable in length with D

Therefore EF is also rational

Let it be contrived that

as the number BA is to AC so is the square on EF to the square on FG [x 6 Por]

But AB has to AC the ratio which a number has to a number therefore the square on EF also has to the square on FG the ratio which a number has to a number

so that the square on EF is commensurable with the square on FG [x 6]

And EF is rational

therefore FG is also rational

And since BA has not to AC the ratio which a square number has to a square number

neither therefore has the square on EF to the square on FG the ratio which a square number has to a square number

therefore EF is incommensurable in length with FG [x 9]

Therefore EF , FG are rational straight lines commensurable in square only,
therefore EG is binomial [x 36]

I say that it is also a first binomial straight line

For since as the number BA is to AC , so is the square on EF to the square on FG ,

while BA is greater than AC ,

therefore the square on EF is also greater than the square on FG

Let then the squares on FG , H be equal to the square on EF

Now since, as BA is to AC , so is the square on EF to the square on FG ,
therefore, *convertendo*

as AB is to BC , so is the square on EF to the square on H [x 19 Por]

But AB has to BC the ratio which a square number has to a square number
therefore the square on EF also has to the square on H the ratio which a square number has to a square number

Therefore EF is commensurable in length with H [x 9]
therefore the square on EF is greater than the square on FG by the square on a straight line commensurable with EF

And EF , FG are rational and EF is commensurable in length with D

Therefore EF is a first binomial straight line

Q E D

PROPOSITION 49

To find the second binomial straight line

Let two numbers AC , CB be set out such that the sum of them AB has to BC the ratio which a square number has to a square number but has not to AC the ratio which a square number has to a square number,
let a rational straight line D be set out and let EF be commensurable in length with D

therefore EF is rational

Let it be contrived then that

as the number CA is to AB so also is the square on EF to the square on FG

[x 6 Por]

therefore the square on EF is commensurable with the square on FG [x 6]

Therefore FG is also rational

Now since the number CA has not to AB the ratio which a square number has to a square number neither has the square on EF to the square on FG the ratio which a square number has to a square number

Therefore EF is incommensurable in length with FG , [x 9]
therefore EF , FG are rational straight lines commensurable in square only

therefore EG is binomial

[x 36]

It is next to be proved that it is also a second binomial straight line

For since inversely as the number BA is to AC , so is the square on GF to the square on FE ,

while BA is greater than AC

therefore the square on GF is greater than the square on FE

Let the squares on EF , H be equal to the square on GF ,

therefore *convertendo* as AB is to BC so is the square on FG to the square on H

[x 19 Por]

But AB has to BC the ratio which a square number has to a square number,

therefore the square on FG also has to the square on H the ratio which a square number has to a square number

Therefore FG is commensurable in length with H , [x 9]
so that the square on FG is greater than the square on FE by the square on a straight line commensurable with FG

And FG FE are rational straight lines commensurable in square only and EF , the lesser term is commensurable in length with the rational straight line D set out

Therefore EG is a second binomial straight line


Q E D

PROPOSITION 50

To find the third binomial straight line

Let two numbers AC , CB be set out such that the sum of them AB has to BC the ratio which a square number has to a square number but has not to AC the ratio which a square number has to a square number

Let any other number D , not square be set out also, and let it not have to either of the numbers BA , AC the ratio which a square number has to a square number



Let any rational straight line E be set out and let it be contrived that as D is to AB so is the square on E to the square on FG , [x 6, Por]

therefore the square on E is commensurable with the square on FG [x 6]

And E is rational,

therefore FG is also rational

And since D has not to AB the ratio which a square number has to a square number

neither has the square on E to the square on FG the ratio which a square number has to a square number

therefore E is incommensurable in length with FG [x 9]

Next let it be contrived that as the number BA is to AC so is the square on FG to the square on GH [x 6 Por]

therefore the square on FG is commensurable with the square on GH [x 6]

But FG is rational

therefore GH is also rational

And since BA has not to $1C$ the ratio which a square number has to a square number

neither has the square on FG to the square on HG the ratio which a square number has to a square number

therefore FG is incommensurable in length with GH [x 9]

Therefore FG GH are rational straight lines commensurable in square only

therefore FH is binomial [x 36]

I say next that it is also a third binomial straight line

For since as D is to AB so is the square on E to the square on FG

and as BA is to AC so is the square on FG to the square on GH

therefore *ex aequali*, as D is to AC so is the square on E to the square on GH

[x 22]

But D has not to AC the ratio which a square number has to a square number,
therefore neither has the square on E to the square on GH the ratio which a square number has to a square number,

therefore E is incommensurable in length with GH [x 9]

And since, as BA is to AC , so is the square on FG to the square on GH ,

therefore the square on FG is greater than the square on GH

Let then the squares on GH , A be equal to the square on FG ,

therefore *convertendo* as AB is to BC , so is the square on FG to the square on K [v 19, Por]

But AB has to BC the ratio which a square number has to a square number,
therefore the square on FG also has to the square on K the ratio which a square number has to a square number,

therefore FG is commensurable in length with A [x 9]

Therefore the square on FG is greater than the square on GH by the square on a straight line commensurable with FG

And FG , GH are rational straight lines commensurable in square only, and neither of them is commensurable in length with E

Therefore FH is a third binomial straight line Q E D

PROPOSITION 51

To find the fourth binomial straight line

Let two numbers AC , CB be set out such that AB neither has to BC , nor yet to AC the ratio which a square number has to a square number

Let a rational straight line D be set out

and let EF be commensurable in length with D ,

therefore EF is also rational

Let it be contrived that, as the number BA is to AC so is the square on EF to the square on FG [x 6, Por]

therefore the square on EF is commensurable with the square on FG , [x 6]

therefore FG is also rational

Now since BA has not to AC the ratio which a square number has to a square number

neither has the square on EF to the square on FG the ratio which a square number has to a square number

therefore EF is incommensurable in length with FG [x 9]

Therefore EF , FG are rational straight lines commensurable in square only so that EG is binomial

I say next that it is also a fourth binomial straight line

For since as BA is to AC so is the square on EF to the square on FG


therefore the square on EF is greater than the square on FG

Let then the squares on FG , H be equal to the square on EF ,

therefore *convertendo* as the number AB is to BC , so is the square on EF to the square on H [v 19 Por]

But AB has not to BC the ratio which a square number has to a square number

therefore neither has the square on EF to the square on H the ratio which a square number has to a square number



Therefore EF is incommensurable in length with H , [x 9]
 therefore the square on EF is greater than the square on GF by the square on
 a straight line incommensurable with GF

And EF , FG are rational straight lines commensurable in square only, and
 EF is commensurable in length with D

Therefore EG is a fourth binomial straight line Q E D

PROPOSITION 52

To find the fifth binomial straight line

Let two numbers AC , CB be set out such that AB has not to either of them
 the ratio which a square number has to a square number,

let any rational straight line D be set out

and let EF be commensurable with D ,

therefore EF is rational

Let it be contrived that, as CA is to AB , so is the square on EF
 to the square on FG [x 6, Por]

But CA has not to AB the ratio which a square number
 has to a square number,

therefore neither has the square on EF to the square on FG the
 ratio which a square number has to a square number

Therefore EF , FG are rational straight lines commensurable
 in square only, [x 9]

therefore EG is binomial [x 36]

I say next that it is also a fifth binomial straight line

For since, as CA is to AB so is the square on EF to the square on FG
 inversely, as BA is to AC so is the square on FG to the square on FE ,

therefore the square on GF is greater than the square on FE

Let then the squares on EF , H be equal to the square on GF ,
 therefore, *convertendo* as the number AB is to BC , so is the square on GF to
 the square on H [v 19 Por]

But AB has not to BC the ratio which a square number has to a square
 number

therefore neither has the square on FG to the square on H the ratio which a
 square number has to a square number

Therefore FG is incommensurable in length with H [x 9]
 so that the square on FG is greater than the square on FE by the square on a
 straight line incommensurable with FG

And GF , FE are rational straight lines commensurable in square only and
 the lesser term EF is commensurable in length with the rational straight line
 D set out

Therefore EG is a fifth binomial straight line Q E D

PROPOSITION 53

To find the sixth binomial straight line

Let two numbers AC , CB be set out such that AB has not to either of them
 the ratio which a square number has to a square number

and let there also be another number D which is not square and which has not
 to either of the numbers BA , AC the ratio which a square number has to a
 square number

Let any rational straight line E be set out,
and let it be contrived that, as D is to AB so is the square on E to the square on FG [x 6 Por]

therefore the square on E is commensurable with the square on FG [x 6]

And E is rational

therefore FG is also rational

Now since D has not to AB the ratio which a square number has to a square number

neither has the square on E to the square on FG the ratio which a square number has to a square number,
therefore E is incommensurable in length with FG [x 9]

Again let it be contrived that as BA is to AC , so is the square on FG to the square on GH [x 6, Por]

Therefore the square on FG is commensurable with the square on GH [x 6]

Therefore the square on GH is rational

therefore GH is rational

And since BA has not to AC the ratio which a square number has to a square number,

neither has the square on FG to the square on GH the ratio which a square number has to a square number,

therefore FG is incommensurable in length with GH [x 9]

Therefore FG, GH are rational straight lines commensurable in square only
therefore FH is binomial [x 10]

It is next to be proved that it is also a sixth binomial straight line

For since as D is to AB so is the square on E to the square on FG
and also as BA is to AC , so is the square on FG to the square on GH
therefore *ex aequali* as D is to AC so is the square on E to the square on GH [x 22]

But D has not to AC the ratio which a square number has to a square number

therefore neither has the square on E to the square on GH the ratio which a square number has to a square number

therefore E is incommensurable in length with GH [x 9]

But it was also proved incommensurable with FG

therefore each of the straight lines FG, GH is incommensurable in length with E

And since as BA is to AC so is the square on FG to the square on GH
therefore the square on FG is greater than the square on GH

Let then the squares on CH, A be equal to the square on FG

therefore *convertendo* as AB is to BC , so is the square on FG to the square on A [x 19 Por]

But AB has not to BC the ratio which a square number has to a square number

so that neither has the square on FG to the square on A the ratio which a square number has to a square number

Therefore FG is incommensurable in length with A [x 9]
therefore the square on FG is greater than the square on GH by the square on a straight line incommensurable with FG

And FG GH are rational straight lines commensurable in square only and neither of them is commensurable in length with the rational straight line E set out

Therefore FH is a sixth binomial straight line

Q E D

LEMMA

Let there be two squares AB , BC , and let them be placed so that DB is in a straight line with BE ,

therefore FB is also in a straight line with BG

Let the parallelogram AC be completed

I say that AC is a square that DG is a mean proportional between AB , BC and further that DC is a mean proportional between AC , CB

For, since DB is equal to BF and BE to BG ,

therefore the whole DE is equal to the whole FG

But DE is equal to each of the straight lines AH , KC ,

and FG is equal to each of the straight lines AK , HC [I. 34],

therefore each of the straight lines AH , KC is also equal to each of the straight lines AK , HC

Therefore the parallelogram AC is equilateral

And it is also rectangular

therefore AC is a square

And since as FB is to BG so is DB to BE

while as FB is to BG so is AB to DG ,

and as DB is to BE , so is DG to BC

[VI. 1]

therefore also as AB is to DG so is DG to BC

[V. 11]

Therefore DG is a mean proportional between AB BC

I say next that DC is also a mean proportional between AC CB

For since as AD is to DK so is KG to GC —

for they are equal respectively—

and *componendo* as AK is to AD so is AC to CG

[V. 18]

while as AK is to AD so is AC to CD

and as KC is to CG so is DC to CB

[VI. 1]

therefore also as AC is to DC so is DC to BC

[V. 11]

Therefore DC is a mean proportional between AC CB

Being what it was proposed to prove

PROPOSITION 54

If an area be contained by a rational straight line and the first binomial, the side of the area is the irrational straight line which is called binomial

For let the area AC be contained by the rational straight line AB and the first binomial AD ,

I say that the side of the area AC is the irrational straight line which is called binomial



For since AD is a first binomial straight line let it be divided into its terms at E

and let AE be the greater term

It is then manifest that AE ED are rational straight lines commensurable in square only

the square on AE is greater than the square on ED by the square on a straight line commensurable with AE ,
 and AE is commensurable in length with the rational straight line AB set out
 1 2 13 [x Def 11 1]

Let ED be bisected at the point F



Then since the square on AE is greater than the square on ED by the square on a straight line commensurable with AE

therefore, if there be applied to the greater AE a parallelogram equal to the fourth part of the square on the less that is to the square on EF , and deficient by a square figure it divides it into commensurable parts [x 17]

Let then the rectangle AG GE equal to the square on EF be applied to AE
 therefore AG is commensurable in length with FG

Let GH EK FL be drawn from G E F parallel to either of the straight lines AB CD

let the square SN be constructed equal to the parallelogram AH and the square NQ equal to GA [ii 14]

and let them be placed so that MN is in a straight line with NO ,

therefore RN is also in a straight line with NP

And let the parallelogram SQ be completed

therefore SQ is a square [Lemma]

Now since the rectangle AG GE is equal to the square on EF

therefore as AG is to EF so is FE to EG , [vi 14]

therefore also as AH is to EL so is EL to KG [vi 14]

therefore EL is a mean proportional between AH GA

But AH is equal to SN and GA to NQ

therefore EL is a mean proportional between SN NQ

But MR is also a mean proportional between the same SN NQ [Lemma]

therefore EL is equal to MR

so that it is also equal to PO

But AH GA are also equal to SN NQ

therefore the whole AC is equal to the whole SQ that is to the square on MO

therefore MO is the side of AC

I say next that MO is binomial

For since AC is commensurable with GE

therefore AF is also commensurable with each of the straight lines AG GE [x 15]

But AE is also by hypothesis commensurable with AB

therefore AG GE are also commensurable with AB [x 15]

And AB is rational

therefore each of the straight lines AG GE is also rational

therefore each of the rectangles AH GA is rational, [x 19]

and AH is commensurable with GK

But AH is equal to SN , and GA to AQ ,
therefore SN, NQ that is, the squares on MN, NO are rational and commensurable

And since AE is incommensurable in length with ED
while AE is commensurable with AG , and DE is commensurable with EF
therefore AG is also incommensurable with EF [x 13]
so that AH is also incommensurable with EL [vi 1, x 11]

But AH is equal to SN , and EL to MR ,
therefore SN is also incommensurable with MR

But, as SN is to MR , so is PN to NR , [vi 1]
therefore PN is incommensurable with NR [x 11]

But PN is equal to MN , and NR to NO ,
therefore MN is incommensurable with NO

And the square on MN is commensurable with the square on NO
and each is rational,

therefore MN, NO are rational straight lines commensurable in square only
Therefore MO is binomial [x 36] and the side ' of AC Q E D

PROPOSITION 55

If an area be contained by a rational straight line and the second binomial the 'side' of the area is the irrational straight line which is called a first bimedial

For let the area $ABCD$ be contained by the rational straight line AB and the second binomial AD ,


I say that the side of the area AC is a first bimedial straight line

For, since AD is a second binomial straight line let it be divided into its terms at E so that AE is the greater term
therefore AE, ED are rational straight lines commensurable in square only, the square on AE is greater than the square on ED by the square on a straight line commensurable with AE
and the lesser term ED is commensurable in length with AB [x Def II 2]

Let ED be bisected at F
and let there be applied to AE the rectangle $AG GE$ equal to the square on EF and deficient by a square figure

therefore AG is commensurable in length with GE [x 17]

Through $G E F$ let $GH EK FL$ be drawn parallel to $AB CD$
let the square SN be constructed equal to the parallelogram AH and the square NQ equal to GA



and let them be placed so that MN is in a straight line with NO

therefore RN is also in a straight line with NP

Let the square SQ be completed

It is then manifest from what was proved before that MR is a mean proportional between SN , NQ and is equal to EL , and that MO is the "side" of the area AC

It is now to be proved that MO is a first bimedial straight line

Since AE is commensurable in length with ED

while ED is commensurable with AB

therefore AE is incommensurable with AB [x 13]

And since AG is commensurable with EG ,

AE is also commensurable with each of the straight lines AG GE [x 15]

But AE is incommensurable in length with AB ,

therefore AG GE are also incommensurable with AB [x 13]

Therefore BA AG and BA , GE are pairs of rational straight lines commensurable in square only

so that each of the rectangles AH , GA is medial [x 21]

Hence each of the squares SN NQ is medial

Therefore MN NO are also medial

And since AG is commensurable in length with GE

AH is also commensurable with GA , [xi 1, x 11]

that is SN is commensurable with NQ

that is the square on MN with the square on NO

And since AE is incommensurable in length with ED ,

while AE is commensurable with AG ,

and ED is commensurable with EF

therefore AG is incommensurable with EF , [x 13]

so that AH is also incommensurable with EL ,

that is SN is incommensurable with MR ,

that is PN with NR , [vi 1, x 11]

that is MN is incommensurable in length with NO

But MA NO were proved to be both medial and commensurable in square therefore MA NO are medial straight lines commensurable in square only

I say next that they also contain a rational rectangle

For since DE is by hypothesis commensurable with each of the straight lines AB EF

therefore EF is also commensurable with EA [x 1^o]

And each of them is rational,

therefore EL that is MR is rational [x 10]

and MR is the rectangle MN NO

But if two medial straight lines commensurable in square only and containing a rational rectangle be added together the whole is irrational and is called a first bimedial straight line [x 3^d]

Therefore MO is a first bimedial straight line Q E D

PROPOSITION 56



If an area be contained by a rational straight line and the third binomial the "side" of the area is the irrational straight line called a second bimedial

For let the area $ABCD$ be contained by the rational straight line AB and the third binomial AD divided into its terms at E of which terms AE is the greater

I say that the "side" of the area AC is the irrational straight line called a second bimedial

For let the same construction be made as before

Now, since AD is a third binomial straight line,



therefore AE , ED are rational straight lines commensurable in square only, the square on AE is greater than the square on ED by the square on a straight line commensurable with AE , and neither of the terms AE , ED is commensurable in length with AB

[x Deff 11 3]

Then in manner similar to the foregoing, we shall prove that MO is the 'side' of the area AC

and MN NO are medial straight lines commensurable in square only, so that MO is bimedial

It is next to be proved that it is also a second bimedial straight line

Since DE is incommensurable in length with AB that is with EK

and DE is commensurable with EF

therefore EF is incommensurable in length with EK [x 13]

And they are rational,

therefore FE EK are rational straight lines commensurable in square only

Therefore EL that is MR is medial

[x 21]

And it is contained by MN , NO

therefore the rectangle MN NO is medial

Therefore MO is a second bimedial straight line

[x 38]

Q E D

PROPOSITION 57

If an area be contained by a rational straight line and the fourth binomial the 'side' of the area is the irrational straight line called major

For let the area AC be contained by the rational straight line AB and the fourth binomial AD divided into its terms at E , of which terms let AE be the greater

I say that the side of the area AC is the irrational straight line called major

For since AD is a fourth binomial straight line

therefore AE ED are rational straight lines commensurable in square only the square on AE is greater than the square on ED by the square on a straight line incommensurable with AE

and AE is commensurable in length with AB [x Deff 11 4]


Let DE be bisected at F

and let there be applied to AE a parallelogram the rectangle AG , GE equal to the square on EF

therefore AG is incommensurable in length with GE [x 18]

Let GH EK FL be drawn parallel to AB

and let the rest of the construction be as before
it is then manifest that MO is the 'side' of the area AC



It is next to be proved that MO is the irrational straight line called major
Since AG is incommensurable with EG
 AH is also incommensurable with GK that is SN with NQ [vi 1, x 11]
therefore MN NO are incommensurable in square

And since AE is commensurable with AB
 AH is rational [x 10]

and it is equal to the squares on MN NO

therefore the sum of the squares on MN NO is also rational

And, since DE is incommensurable in length with AB that is with EK
while DE is commensurable with EF ,

therefore EF is incommensurable in length with EK [x 13]

Therefore EA LF are rational straight lines commensurable in square only
therefore LE that is, MR , is medial [x 21]

And it is contained by MN NO ,

therefore the rectangle MN NO is medial

And the {sum} of the squares on MN NO is rational

and MN NO are incommensurable in square

But if two straight lines incommensurable in square and making the sum of the squares on them rational but the rectangle contained by them medial be added together the whole is irrational and is called major [x 39]

Therefore MO is the irrational straight line called major and is the 'side' of the area AC Q E D

PROPOSITION 58

If an area be contained by a rational straight line and the fifth binomial the "side" of the area is the irrational straight line called the side of a rational plus a medial area

For let the area AC be contained by the rational straight line AB and the fifth binomial AD divided into its terms at E so that AE is the greater term,
I say that the 'side' of the area AC is the irrational straight line called the side of a rational plus a medial area

For let the same construction be made as before shown,

it is then manifest that MO is the 'side' of the area AC



It is then to be proved that MO is the side of a rational plus a medial area
For since AG is incommensurable with GE [x 15]

therefore EH is also commensurable with HE [vi 1, x 11]

that is the square on MA with the square on NO ,

therefore MA NO are incommensurable in square

And since AD is a fifth binomial straight line and ED the lesser segment therefore ED is commensurable in length with AB [X. Def. II. 5]



But AE is incommensurable with ED ,

therefore AB is also incommensurable in length with AE [X. 13]

Therefore AA that is the sum of the squares on MN NO , is medial [X. 21]

And, since DE is commensurable in length with AB , that is, with FA ,

while DE is commensurable with EF ,

therefore EF is also commensurable with EA [X. 12]

And EK is rational,

therefore EL that is, MR , that is the rectangle MN NO , is also rational [X. 19]

Therefore MN , NO are straight lines incommensurable in square which make the sum of the squares on them medial but the rectangle contained by them rational

Therefore MO is the side of a rational plus a medial area [X. 40] and is the "side of the area AC "



PROPOSITION 59

If an area be contained by a rational straight line and the sixth binomial, the "side" of the area is the irrational straight line called the side of the sum of two medial areas

For let the area $ABCD$ be contained by the rational straight line AB and the sixth binomial AD , divided into its terms at E so that AE is the greater term

I say that the "side" of AC is the side of the sum of two medial areas

Let the same construction be made as before shown



It is then manifest that MO is the "side" of AC and that MA is incommensurable in square with NO

Now since EA is incommensurable in length with AB

therefore EA AB are rational straight lines commensurable in square only therefore AA that is the sum of the squares on MN NO is medial [X. 21]

Again since ED is incommensurable in length with AB ,

therefore FE is also incommensurable with EK , [x 13]
 therefore FE , EK are rational straight lines commensurable in square only,
 therefore EL that is MR , that is the rectangle MN , NO is medial [x 9]
 And since AE is incommensurable with EF

AK is also incommensurable with EL [vi 1 x 11]

But AK is the sum of the squares on MN , NO

and EL is the rectangle MN , NO -

therefore the sum of the squares on MN , NO is incommensurable with the rectangle MN , NO


And each of them is medial, and MN , NO are incommensurable in square

Therefore MO is the side of the sum of two medial areas [x 41] and is the side of AC Q E D

LEMMA

If a straight line be cut into unequal parts, the squares on the unequal parts are greater than twice the rectangle contained by the unequal parts

Let AB be a straight line and let it be cut into unequal parts at C and let AC be the greater



I say that the squares on AC , CB are greater than twice the rectangle AC , CB
 For let AB be bisected at D

Since then, a straight line has been cut into equal parts at D , and into unequal parts at C

therefore the rectangle AC , CB together with the square on CD is equal to the square on AD [II 5]

so that the rectangle AC , CB is less than the square on AD ,
 therefore twice the rectangle AC , CB is less than double of the square on AD

But the squares on AC , CB are double of the squares on AD , DC , [II 9]
 therefore the squares on AC , CB are greater than twice the rectangle AC , CB Q E D

PROPOSITION 60

The square on the binomial straight line applied to a rational straight line produces as breadth the first binomial

Let AB be a binomial straight line divided into its terms at C so that AC is the greater term

let a rational straight line DF be set out
 and let $DEFG$ equal to the square on AB be applied to DE producing DG as its breadth


I say that DG is a first binomial straight line

For let there be applied to DE the rectangle DH equal to the square on AC and KL equal to the square on BC

therefore the remainder twice the rectangle AC , CB is equal to MF

Let MG be bisected at N and let NO be drawn parallel [to ML or GF]

Therefore each of the rectangles MO , NO is equal to once the rectangle AC , CB



Now, since AB is a binomial divided into its terms at C

therefore AC, CB are rational straight lines commensurable in square only, [x 36]
 therefore the squares on AC, CB are rational and commensurable with one another

so that the sum of the squares on AC, CB is also rational [x 15]

And it is equal to DL ,

therefore DL is rational

And it is applied to the rational straight line DE ,

therefore DM is rational and commensurable in length with DE [x 20]

Again since AC, CB are rational straight lines commensurable in square only

therefore twice the rectangle AC, CB that is MF is medial [x 21]

And it is applied to the rational straight line ML ,

therefore MG is also rational and incommensurable in length with ML , that is DE [x 23]

But MD is also rational and is commensurable in length with DE ,

therefore DV is incommensurable in length with MG [x 13]

And they are rational

therefore DM, MG are rational straight lines commensurable in square only

therefore DG is binomial [x 36]

It is next to be proved that it is also a first binomial straight line

Since the rectangle AC, CB is a mean proportional between the squares on AC, CB [cf Lemma after x 53]

therefore MO is also a mean proportional between DH, KL

Therefore, as DH is to MO so is MO to KL

that is as DA is to MN so is MA to NK [vi 1]

therefore the rectangle DK, KV is equal to the square on MA [vi 17]

And since the square on AC is commensurable with the square on CB

DH is also commensurable with KL

so that DA is also commensurable with KM [vi 1 x 11]

And, since the squares on AC, CB are greater than twice the rectangle AC, CB , [Lemma]

therefore DL is also greater than MF ,

so that DM is also greater than MG [vi 1]

And the rectangle DA, KM is equal to the square on MN that is to the fourth part of the square on MG

and DA is commensurable with KM

But if there be two unequal straight lines and to the greater there be applied a parallelogram equal to the fourth part of the square on the less and deficient by a square figure and if it divide it into commensurable parts the square on the greater is greater than the square on the less by the square on a straight line commensurable with the greater [x 17]

therefore the square on DM is greater than the square on MG by the square on a straight line commensurable with DM

And DM, MG are rational

and DM , which is the greater term is commensurable in length with the rational straight line DE set out

Therefore DG is a first binomial straight line


[x Def II 1]

Q E D

PROPOSITION 61

The square on the first binomial straight line applied to a rational straight line produces as breadth the second binomial

Let AB be a first binomial straight line divided into its medials at C , of which medials AC is the greater, let a rational straight line DE be set out, and let there be applied to DE the parallelogram DF equal to the square on AB , producing DG as its breadth



I say that DG is a second binomial straight line. For let the same construction as before be made. Then, since AB is a first binomial divided at C , therefore AC CB are medial straight lines commensurable in square only and containing a rational rectangle, [x 34]
so that the squares on AC , CB are also medial [x 21]

Therefore DL is medial [x 15 and 13, Por.] And it has been applied to the rational straight line DE , therefore MD is rational and incommensurable in length with DE [x 20]

Again, since twice the rectangle AC CB is rational, MF is also rational.

And it is applied to the rational straight line ML , therefore MG is also rational and commensurable in length with ML , that is DE [x 20]

therefore DM is incommensurable in length with MG [x 13]

And they are rational, therefore DM MG are rational straight lines commensurable in square only, therefore DG is binomial [x 36]

It is next to be proved that it is also a second binomial straight line. For, since the squares on AC CB are greater than twice the rectangle AC , CB

therefore DL is also greater than MF , so that DM is also greater than MG [vi 1]

And, since the square on AC is commensurable with the square on CB , DH is also commensurable with KL

so that DA is also commensurable with AM [vi 1 x 11]

And the rectangle DA AM is equal to the square on MN , therefore the square on DM is greater than the square on MG by the square on a straight line commensurable with DM [x 17]

And MG is commensurable in length with DE . Therefore DG is a second binomial straight line [x Def. 11]


PROPOSITION 62

The square on the second binomial straight line applied to a rational straight line produces as breadth the third binomial

Let AB be a second binomial straight line divided into its medials at C , so that AC is the greater segment, let DE be any rational straight line and to DE let there be applied the parallelogram DF equal to the square on AB and producing DG as its breadth,

I say that DG is a third binomial straight line

Let the same construction be made as before shown



Then, since AB is a second binomial divided at C ,

therefore AC , CB are medial straight lines commensurable in square only and containing a medial rectangle [x 35]

so that the sum of the squares on AC , CB is also medial [x 15 and 23 Por]

And it is equal to DL

therefore DL is also medial

And it is applied to the rational straight line DE

therefore MD is also rational and incommensurable in length with DE [x 22]

For the same reason

MG is also rational and incommensurable in length with ML that is, with DE , therefore each of the straight lines DM , MG is rational and incommensurable in length with DE

And since AC is incommensurable in length with CB ,

and as AC is to CB so is the square on AC to the rectangle AC , CB , therefore the square on AC is also incommensurable with the rectangle AC , CB [x 11]

Hence the sum of the squares on AC , CB is incommensurable with twice the rectangle AC , CB , [x 12 13]

that is, DL is incommensurable with MF ,

so that DM is also incommensurable with MG [x 1, x 11]

And they are rational,

therefore DG is binomial

[x 36]

It is to be proved that it is also a third binomial straight line

In manner similar to the foregoing we may conclude that DM is greater than MG

and that DK is commensurable with KM

And the rectangle DK , KM is equal to the square on MN , therefore the square on DM is greater than the square on MG by the square on a straight line commensurable with DM

And neither of the straight lines DM , MG is commensurable in length with DE

Therefore DG is a third binomial straight line

[x Def II 3]

Q E D

PROPOSITION 63

The square on the major straight line applied to a rational straight line produces as breadth the fourth binomial

Let AB be a major straight line divided at C so that AC is greater than CB

let DE be a rational straight line

and to DE let there be applied the parallelogram DF equal to the square on AB and producing DG as its breadth

I say that DG is a fourth binomial straight line

Let the same construction be made as before shown

Then since AB is a major straight line divided at C ,

AC , CB are straight lines incommensurable in square which make the sum of the squares on them rational but the rectangle contained by them medial

[x 39]

Since, then the sum of the squares on AC , CB is rational therefore DL is rational therefore DM is also rational and commensurable in length with DE

[x 20]

Again since twice the rectangle AC , CB , that is MF is medial, and it is applied to the rational straight line ML therefore MG is also rational and incommensurable in length with DE ,

[x 22]

therefore DM is also incommensurable in length with MG

[x 13]

Therefore DM , MG are rational straight lines commensurable in square only

therefore DG is binomial

[x 36]

It is to be proved that it is also a fourth binomial straight line

In manner similar to the foregoing we can prove that DM is greater than MG

and that the rectangle DK , AM is equal to the square on MN

Since then the square on LC is incommensurable with the square on CB , therefore DH is also incommensurable with KL ,

so that DK is also incommensurable with AM

[vi 1 x 11]

But if there be two unequal straight lines and to the greater there be applied a parallelogram equal to the fourth part of the square on the less and deficient by a square figure and if it divide it into incommensurable parts then the square on the greater will be greater than the square on the less by the square on a straight line incommensurable in length with the greater

[x 18]

therefore the square on DM is greater than the square on MG by the square on a straight line incommensurable with DM

And DM , MG are rational straight lines commensurable in square only and DM is commensurable with the rational straight line DE set out

Therefore DG is a fourth binomial straight line

[x Def II 4]

Q E D

PROPOSITION 64

The square on the side of a rational plus a medial area applied to a rational straight line produces as breadth the fifth binomial



Let AB be the side of a rational plus a medial area divided into its straight lines at C so that AC is the greater, let a rational straight line DE be set out and let there be applied to DE the parallelogram DF equal to the square on AB producing DG as its breadth,

I say that DG is a fifth binomial straight line

Let the same construction as before be made

Since then AB is the side of a rational plus a medial area divided at C

therefore AC , CB are straight lines incommensurable in square which make



the sum of the squares on them medial but the rectangle contained by them rational [x 40]

Since, then the sum of the squares on AC CB is medial

therefore DL is medial

so that DM is rational and incommensurable in length with DE [x 22]

Again, since twice the rectangle AC , CB , that is MF is rational

therefore MG is rational and commensurable with DE [x 20]

Therefore DM is incommensurable with MG [x 13]

therefore DM , MG are rational straight lines commensurable in square only

therefore DG is binomial [x 36]

I say next that it is also a fifth binomial straight line

For it can be proved similarly that the rectangle DA AM is equal to the square on MA

and that DA is incommensurable in length with AM ,

therefore the square on DM is greater than the square on MG by the square on a straight line incommensurable with DM [x 18]

And DM , MG are commensurable in square only, and the less MG , is commensurable in length with DE

Therefore DG is a fifth binomial

Q E D

PROPOSITION 65

The square on the side of the sum of two medial areas applied to a rational straight line produces as breadth the sixth binomial

Let AB be the side of the sum of two medial areas, divided at C

let DE be a rational straight line

and let there be applied to DE the parallelogram DF equal to the square on AB producing DG as its breadth,


I say that DG is a sixth binomial straight line

For let the same construction be made as before

Then since AB is the side of the sum of two medial areas divided at C

therefore AC CB are straight lines incommensurable in square which make the sum of the squares on them medial the rectangle contained by them medial and moreover the sum of the squares on them incommensurable with the rectangle contained by them [x 41]

so that in accordance with what was before proved each of the rectangles DL MF is medial



And they are applied to the rational straight line DE therefore each of the straight lines DM MG is rational and incommensurable in length with DE [x 22]

And since the sum of the squares on AC , CB is incommensurable with twice the rectangle AC CB

therefore DL is incommensurable with MF

Therefore DM is also incommensurable with MG [vi 1 x 11]

therefore DM MG are rational straight lines commensurable in square only therefore DG is binomial [x 36]

I say next that it is also a sixth binomial straight line

Similarly again we can prove that the rectangle DA , AM is equal to the square on MN ,

and that DA is incommensurable in length with KM ,
and for the same reason the square on DM is greater than the square on MG by the square on a straight line incommensurable in length with DM .

And neither of the straight lines DM , MG is commensurable in length with the rational straight line DE set out.

Therefore DG is a sixth binomial straight line.

Q E D

PROPOSITION 66

A straight line commensurable in length with a binomial straight line is itself also binomial and the same in order.

Let AB be binomial and let CD be commensurable in length with AB ,

I say that CD is binomial and the same

in order with AB .

For since AB is binomial

let it be divided into its terms at E

and let AE be the greater term,

therefore AE , EB are rational straight lines commensurable in square only. [v 36]

Let it be contrived that

as AB is to CD so is AE to CF , [vi 17]

therefore also the remainder EB is to the remainder FD as AB is to CD . [v 19]

But AB is commensurable in length with CD

therefore AE is also commensurable with CF , and EB with FD . [x 11]

And AE , EB are rational

therefore CF , FD are also rational

And as AE is to CF so is EB to FD . [v 11]

Therefore alternately as AE is to EB so is CF to FD . [v 16]

But AE , EB are commensurable in square only

therefore CF , FD are also commensurable in square only. [x 11]

And they are rational

therefore CD is binomial. [x 36]

I say next that it is the same in order with AB .

For the square on AE is greater than the square on EB either by the square on a straight line commensurable with AE or by the square on a straight line incommensurable with it.

If then the square on AE is greater than the square on EB by the square on a straight line commensurable with AE

the square on CF will also be greater than the square on FD by the square on a straight line commensurable with CF . [x 14]

And if AE is commensurable with the rational straight line set out, CF will also be commensurable with it. [x 17]

and for this reason each of the straight lines AB , CD is a first binomial that is the same in order. [x Def II 1]

But if EB is commensurable with the rational straight line set out FD is also commensurable with it. [x 17]

and for this reason again CD will be the same in order with AB .

for each of them will be a second binomial. [x Def II 2]

But, if neither of the straight lines AE , EB is commensurable with the rational straight line set out neither of the straight lines CF , FD will be commensurable with it [x 13]

and each of the straight lines AB , CD is a third binomial [x Deff II 3]

But, if the square on AE is greater than the square on EB by the square on a straight line incommensurable with AE , the square on CF is also greater than the square on FD by the square on a straight line incommensurable with CF [x 14]

And if AE is commensurable with the rational straight line set out, CF is also commensurable with it,

and each of the straight lines AB , CD is a fourth binomial [x Deff II 4]

But if EB is so commensurable, so is FD also and each of the straight lines AB , CD will be a fifth binomial [x Deff II 5]


But if neither of the straight lines AE , EB is so commensurable, neither of the straight lines CF , FD is commensurable with the rational straight line set out

and each of the straight lines AB , CD will be a sixth binomial [x Deff II 6]

Hence a straight line commensurable in length with a binomial straight line is binomial and the same in order Q E D

PROPOSITION 67

A straight line commensurable in length with a binomial straight line is itself also binomial and the same in order

Let AB be binomial, and let CD be commensurable in length with AB ,
 I say that CD is binomial and the same in order with AB
 For, since AB is binomial,

let it be divided into its medials at E ,
 therefore AE , EB are medial straight lines commensurable in square only [x 37 38]

And let it be contrived that
 as AB is to CD so is AE to CF ,
 therefore also the remainder EB is to the remainder FD as AB is to CD [v 19]

But AB is commensurable in length with CD ,
 therefore AE , EB are also commensurable with CF , FD respectively [x 11]

But AE , EB are medial,
 therefore CF , FD are also medial [x 23]

And since, as AE is to EB so is CF to FD [v 11]

and AE , EB are commensurable in square only,
 CF , FD are also commensurable in square only [x 11]

But they were also proved medial
 therefore CD is binomial

I say next that it is also the same in order with AB
 For since as AE is to EB so is CF to FD

therefore also as the square on AE is to the rectangle AE , EB , so is the square on CF to the rectangle CF , FD ,

therefore alternately

as the square on AE is to the square on CF so is the rectangle AE , EB to the rectangle CF , FD [v 16]

But the square on AE is commensurable with the square on CF ,

therefore the rectangle AE, EB is also commensurable with the rectangle CF, FD

If therefore the rectangle AE, EB is rational

the rectangle CF, FD is also rational

{and for this reason CD is a first bimedial}, [x 3,]

but if medial medial [x 23, Por]

and each of the straight lines AB, CD is a second bimedial [x 38]

And for this reason CD will be the same in order with AB Q E D

PROPOSITION 68

A straight line commensurable with a major straight line is itself also major

Let AB be major and let CD be commensurable with AB ,

I say that CD is major

Let AB be divided at E ,

therefore AE, EB are straight lines incommensurable in square which make the sum of the squares on them rational, but the rectangle contained by them medial [x 39]

Let the same construction be made as before

Then since as AB is to CD so is AE to CF and EB to FD ,

therefore also as AE is to CF , so is EB to FD [v 11]

But AB is commensurable with CD ,

therefore AE, EB are also commensurable with CF, FD respectively [x 11]

And since as AE is to CF so is EB to FD ,

alternately also

as AE is to EB , so is CF to FD , [v 16]

therefore also *componendo*

as AB is to BE , so is CD to DF , [v 18]

therefore also as the square on AB is to the square on BE , so is the square on CD to the square on DF [vi 20]

Similarly we can prove that as the square on AB is to the square on AE so also is the square on CD to the square on CF

Therefore also as the square on AB is to the squares on AE, EB so is the square on CD to the squares on CF, FD ,

therefore also alternately

as the square on AB is to the square on CD so are the squares on AE, EB to the squares on CF, FD [v 16]

But the square on AB is commensurable with the square on CD , therefore the squares on AE, EB are also commensurable with the squares on CF, FD

And the squares on AE, EB together are rational,

therefore the squares on CF, FD together are rational

Similarly also twice the rectangle AE, EB is commensurable with twice the rectangle CF, FD

And twice the rectangle AE, EB is medial,

therefore twice the rectangle CF, FD is also medial [x 23 Por]

Therefore CF, FD are straight lines incommensurable in square which make at the same time the sum of the squares on them rational but the rectangle contained by them medial therefore the whole CD is the irrational straight line called major [x 39]

Therefore a straight line commensurable with the major straight line is major

Q E D

PROPOSITION 69

A straight line commensurable with the side of a rational plus a medial area is itself also the side of a rational plus a medial area

Let AB be the side of a rational plus a medial area and let CD be commensurable with AB

it is to be proved that CD is also the side of a rational plus a medial area

Let AB be divided into its straight lines at E ,
therefore AE EB are straight lines incommensurable in square
which make the sum of the squares on them medial but the
rectangle contained by them rational [x 40]

Let the same construction be made as before

We can then prove similarly that

CF FD are incommensurable in square
and the sum of the squares on AE , EB is commensurable with the sum of the
squares on CF FD

and the rectangle AE , EB with the rectangle CF , FD ,
so that the sum of the squares on CF FD is also medial, and the rectangle CF
 FD rational

Therefore CD is the side of a rational plus a medial area Q E D

PROPOSITION 70

A straight line commensurable with the side of the sum of two medial areas is the side of the sum of two medial areas

Let AB be the side of the sum of two medial areas and CD commensurable with AB

it is to be proved that CD is also the side of the sum of two medial areas

For since AB is the side of the sum of two medial areas
let it be divided into its straight lines at E ,
therefore AE EB are straight lines incommensurable in square
which make the sum of the squares on them medial the rectangle
contained by them medial and furthermore the sum of the
squares on AE , EB incommensurable with the rectangle AE EB [x 41]

Let the same construction be made as before

We can then prove similarly that

CF FD are also incommensurable in square
the sum of the squares on AE EB is commensurable with the sum of the
squares on CF FD

and the rectangle AE EB with the rectangle CF FD
so that the sum of the squares on CF FD is also medial
the rectangle CF FD is medial

and moreover the sum of the squares on CF FD is incommensurable with the
rectangle CF FD

Therefore CD is the side of the sum of two medial areas Q E D

PROPOSITION 71

If a rational and a medial area be added together, four irrational straight lines arise namely a binomial or a first bimedial or a major or a side of a rational plus a medial area

Let AB be rational and CD medial

I say that the 'side' of the area AD is a binomial or a first bimedial or a major or a side of a rational plus a medial area

For AB is either greater or less than CD

First let it be greater

let a rational straight line EF be set out

let there be applied to EF the rectangle EG equal to AB , producing EH as breadth

and let HI equal to DC be applied to EF producing HK as breadth

Then since AB is rational and is equal to EG

therefore EG is also rational

And it has been applied to EF producing EH as breadth

therefore EH is rational and commensurable in length with EF [x 20]

Again since CD is medial and is equal to HI

therefore HI is also medial

And it is applied to the rational straight line EF producing HK as breadth, therefore HK is rational and incommensurable in length with EF [x 22]

And since CD is medial

while AB is rational

therefore AB is incommensurable with CD ,

so that EG is also incommensurable with HI

But as EG is to HI so is EH to HK [vi 1]

therefore EH is also incommensurable in length with HK [x 11]

And both are rational

therefore EH HK are rational straight lines commensurable in square only

therefore EK is a binomial straight line divided at H [x 36]

And since AB is greater than CD

while AB is equal to EG and CD to HI

therefore EG is also greater than HI

therefore EH is also greater than HK


The square then on EH is greater than the square on HK either by the square on a straight line commensurable in length with EH or by the square on a straight line incommensurable with it

First let the square on it be greater by the square on a straight line commensurable with itself

Now the greater straight line HE is commensurable in length with the rational straight line EF set out

therefore EK is a first binomial

[x Def 11 1]



But EF is rational

and if an area be contained by a rational straight line and the first binomial the side of the square equal to the area is binomial [x 54]

Therefore the "side" of EI is binomial

so that the "side" of AD is also binomial

Next let the square on EH be greater than the square on HA by the square on a straight line incommensurable with EH

Now the greater straight line EH is commensurable in length with the rational straight line EF set out

therefore EK is a fourth binomial [x Def II 4]

But EF is rational

and if an area be contained by a rational straight line and the fourth binomial, the 'side' of the area is the irrational straight line called major [x 57]

Therefore the 'side' of the area EI is major,

so that the "side" of the area AD is also major

Next, let AB be less than CD ,

therefore EG is also less than HI ,

so that EH is also less than HA

Now the square on HK is greater than the square on EH either by the square on a straight line commensurable with HA or by the square on a straight line incommensurable with it

First let the square on it be greater by the square on a straight line commensurable in length with itself

Now the lesser straight line EH is commensurable in length with the rational straight line EF set out,

therefore EK is a second binomial [x Def II 2]

But EF is rational

and if an area be contained by a rational straight line and the second binomial the side of the square equal to it is a first binomial, [x 55]

therefore the 'side' of the area EI is a first binomial

so that the 'side' of AD is also a first binomial

Next let the square on HA be greater than the square on HE by the square on a straight line incommensurable with HA

Now the lesser straight line EH is commensurable with the rational straight line EF set out,

therefore EK is a fifth binomial [x Def II 5]

But EF is rational

and, if an area be contained by a rational straight line and the fifth binomial the side of the square equal to the area is a side of a rational plus a medial area [x 58]

Therefore the 'side' of the area EI is a side of a rational plus a medial area so that the 'side' of the area AD is also a side of a rational plus a medial area

Therefore etc



Q E D

PROPOSITION 72

If two medial areas incommensurable with one another be added together, the remaining two irrational straight lines arise namely either a second binomial or a side of the sum of two medial areas

For let two medial areas AB , CD incommensurable with one another be added together,

I say that the "side" of the area AD is either a second binomial or a side of the sum of two medial areas



For AB is either greater or less than CD

First if it so chance let AB be greater than CD

Let the rational straight line EF be set out

and to EF let there be applied the rectangle EG equal to AB and producing EH as breadth and the rectangle HI equal to CD and producing HK as breadth.

Now since each of the areas AB , CD is medial

therefore each of the areas EG , HI is also medial

And they are applied to the rational straight line FE producing EH , HK as breadth

therefore each of the straight lines EH , HK is rational and incommensurable in length with EF [x 11]

And, since AB is incommensurable with CD ,

and AB is equal to EG , and CD to HI ,

therefore EG is also incommensurable with HI

But as EG is to HI , so is EH to HK [vi 1]

therefore EH is incommensurable in length with HK [x 11]

Therefore EH , HK are rational straight lines commensurable in square only,

therefore EK is binomial [x 36]

But the square on EH is greater than the square on HK either by the square on a straight line incommensurable with EH or by the square on a straight line commensurable with it

First let the square on it be greater by the square on a straight line commensurable in length with itself

Now neither of the straight lines EH , HK is commensurable in length with the rational straight line EF set out

therefore EK is a third binomial [x Def 11 3]

But FF is rational,

and if an area be contained by a rational straight line and the third binomial the "side" of the area is a second binomial [x 56]

therefore the "side" of EF that is of AD is a second binomial

Next let the square on EH be greater than the square on HK by the square on a straight line incommensurable in length with EH

Now each of the straight lines EH , HK is incommensurable in length with EF ,

therefore EK is a sixth binomial [x Def 11 6]

But, if an area be contained by a rational straight line and the sixth bi

nomial the 'side' of the area is the side of the sum of two medial areas [x 59] so that the "side" of the area AD is also the side of the sum of two medial areas

Therefore etc

Q E D

The binomial straight line and the irrational straight lines after it are neither the same with the medial nor with one another

For the square on a medial if applied to a rational straight line produces as breadth a straight line rational and incommensurable in length with that to which it is applied [x 22]

But the square on the binomial if applied to a rational straight line produces as breadth the first binomial [x 60]

The square on the first binomial if applied to a rational straight line produces as breadth the second binomial [x 61]

The square on the second binomial if applied to a rational straight line produces as breadth the third binomial [x 62]

The square on the major if applied to a rational straight line, produces as breadth the fourth binomial [x 63]

The square on the side of a rational plus a medial area, if applied to a rational straight line produces as breadth the fifth binomial [x 64]

The square on the side of the sum of two medial areas if applied to a rational straight line produces as breadth the sixth binomial [x 65]

And the said breadths differ both from the first and from one another from the first because it is rational and from one another because they are not the same in order

so that the irrational straight lines themselves also differ from one another

PROPOSITION 73

If from a rational straight line there be subtracted a rational straight line commensurable with the whole in square only the remainder is irrational and let it be called an apotome

For from the rational straight line AB let the rational straight line BC commensurable with the whole in square only be subtracted,

I say that the remainder AC is the irrational straight line called apotome

For, since AB is incommensurable in length with BC , and as AB is to BC so is the square on AB to the rectangle $AB BC$, therefore the square on AB is incommensurable with the rectangle $AB BC$ [x 11]

But the squares on $AB BC$ are commensurable with the square on AB [x 1.]

and twice the rectangle $AB BC$ is commensurable with the rectangle $AB BC$ [x 6]

And inasmuch as the squares on AB, BC are equal to twice the rectangle AB, BC together with the square on CA [ii 7]

therefore the squares on $AB BC$ are also incommensurable with the remainder the square on AC [x 13 16]

But the squares on $AB BC$ are rational therefore AC is irrational [x Def 4]


And let it be called an apotome Q E D

PROPOSITION 74

If from a medial straight line there be subtracted a medial straight line which is commensurable with the whole in square only, and which contains with the whole a rational rectangle the remainder is irrational. And let it be called a first apotome of a medial straight line.

For from the medial straight line AB let there be subtracted the medial straight line BC which is commensurable with AB in square only and with AB makes the rectangle AB, BC rational,

I say that the remainder AC is irrational and let it be called a first apotome of a medial straight line.



For since AB, BC are medial

the squares on AB, BC are also medial

But twice the rectangle AB, BC is rational, therefore the squares on AB, BC are incommensurable with twice the rectangle AB, BC ,

therefore twice the rectangle AB, BC is also incommensurable with the remainder the square on AC , [cf II 17]

since if the whole is incommensurable with one of the magnitudes, the original magnitudes will also be incommensurable [x 1b]

But twice the rectangle AB, BC is rational

therefore the square on AC is irrational,

therefore AC is irrational

[x Def 4]

And let it be called a first apotome of a medial straight line


Q E D

PROPOSITION 75

If from a medial straight line there be subtracted a medial straight line which is commensurable with the whole in square only and which contains with the whole a medial rectangle the remainder is irrational and let it be called a second apotome of a medial straight line.

For from the medial straight line AB let there be subtracted the medial straight line CB which is commensurable with the whole AB in square only and such that the rectangle AB, BC which it contains with the whole AB , is medial [x 28]

I say that the remainder AC is irrational and let it be called a second apotome of a medial straight line.



For let a rational straight line DI be set out, let DE , equal to the squares on AB, BC , be applied to DI , producing DG as breadth,

and let DH equal to twice the rectangle $AB \ BC$ be applied to DI producing DF as breadth,

therefore the remainder FE is equal to the square on AC [II 7]

Now, since the squares on AB, BC are medial and commensurable

therefore DE is also medial [X 15 and 23, Por.]

And it is applied to the rational straight line DI producing DG as breadth, therefore DG is rational and incommensurable in length with DI [X 22]

Again since the rectangle AB, BC is medial,

therefore twice the rectangle AB, BC is also medial [X 23, Por.]

And it is equal to DH ,

therefore DH is also medial

And it has been applied to the rational straight line DI , producing DF as breadth

therefore DF is rational and incommensurable in length with DI [X 22]

And, since AB, BC are commensurable in square only,

therefore AB is incommensurable in length with BC

therefore the square on AB is also incommensurable with the rectangle $AB \ BC$ [X 11]

But the squares on AB, BC are commensurable with the square on AB ,

[X 15]

and twice the rectangle $AB \ BC$ is commensurable with the rectangle $AB \ BC$,

[X 6]

therefore twice the rectangle AB, BC is incommensurable with the squares on AB, BC [X 13]

But DE is equal to the squares on AB, BC ,

and DH to twice the rectangle AB, BC ,

therefore DE is incommensurable with DH

But as DE is to DH so is GD to DF ,

[VI 11]

therefore GD is incommensurable with DF [X 11]

And both are rational,

therefore $GD \ DF$ are rational straight lines commensurable in square only

therefore FG is an apotome [X 73]

But DI is rational

and the rectangle contained by a rational and an irrational straight line is irrational, [deduction from X 20]

and its 'side' is irrational

And AC is the 'side' of FE

therefore AC is irrational

And let it be called a second apotome of a medial straight line Q.E.D.

PROPOSITION 76

If from a straight line there be subtracted a straight line which is incommensurable in square with the whole and which with the whole makes the squares on them added together rational but the rectangle contained by them medial the remainder is irrational and let it be called minor

For from the straight line AB let there be subtracted the straight line BC

$\overbrace{A \quad C \quad B}$ which is incommensurable in square with the whole and fulfils the given conditions [X 33]

I say that the remainder AC is the irrational straight line called minor

For, since the sum of the squares on AB , BC is rational while twice the rectangle AB BC is medial
 therefore the squares on AB , BC are incommensurable with twice the rectangle AB BC ,
 and *conuertendo*, the squares on AB , BC are incommensurable with the remainder the square on AC [II 7 & 16]

But the squares on AB , BC are rational
 therefore the square on AC is irrational,
 therefore AC is irrational

And let it be called *minor*

Q E D

PROPOSITION 77

If from a straight line there be subtracted a straight line which is incommensurable in square with the whole and which with the whole makes the sum of the squares on them medial, but twice the rectangle contained by them rational, the remainder is irrational and let it be called that which produces with a rational area a medial whole

For from the straight line AB let there be subtracted the straight line BC which is incommensurable in square with AB and fulfils the given conditions, [x 34]

I say that the remainder AC is the irrational straight line aforesaid

For, since the sum of the squares on AB BC is medial

while twice the rectangle AB , BC is rational

therefore the squares on AB BC are incommensurable with twice the rectangle AB , BC

therefore the remainder also the square on AC is incommensurable with twice the rectangle AB , BC [II 7 & 16]

And twice the rectangle AB BC is rational

therefore the square on AC is irrational,

therefore AC is irrational

And let it be called *that which produces with a rational area a medial whole*

Q E D


PROPOSITION 78

If from a straight line there be subtracted a straight line which is incommensurable in square with the whole and which with the whole makes the sum of the squares on them medial twice the rectangle contained by them medial and further, the squares on them incommensurable with twice the rectangle contained by them the remainder is irrational and let it be called that which produces with a medial area a medial whole

For from the straight line AB let there be subtracted the straight line BC incommensurable in square with AB and fulfilling the given conditions [x 30]

I say that the remainder AC is the irrational straight line called *that which produces with a medial area a medial whole*

For let a rational straight line DI be set out to DI let there be applied DE equal to the squares on AB BC , producing DG as breadth



and let DH equal to twice the rectangle AB, BC be subtracted

Therefore the remainder FE is equal to the square on AC [II 7]
so that AC is the side of FE

Now since the sum of the squares on AB, BC is medial and is equal to DE ,
therefore DE is medial

And it is applied to the rational straight line DI producing DG as breadth,
therefore DG is rational and incommensurable in length with DI [X 22]

Again since twice the rectangle AB, BC is medial and is equal to DH
therefore DH is medial

And it is applied to the rational straight line DI , producing DF as breadth
therefore DF is also rational and incommensurable in length with DI [X 22]

And since the squares on AB, BC are incommensurable with twice the rectangle AB, BC ,

therefore DE is also incommensurable with DH

But as DE is to DH , so also is DG to DF [VI 1]

therefore DG is incommensurable with DF [X 11]

And both are rational,

therefore GD, DF are rational straight lines commensurable in square only

Therefore FG is an apotome [X 73]

And FH is rational,

but the rectangle contained by a rational straight line and an apotome is irrational [deduction from X 20]

and its 'side' is irrational

And AC is the 'side' of FE

therefore AC is irrational

And let it be called that which produces with a medial area a medial whole

Q E D

PROPOSITION 79

To an apotome only one rational straight line can be annexed which is commensurable with the whole in square only

Let AB be an apotome and BC an annex to it,

$A \quad B \quad C \quad D$ therefore AC, CB are rational straight lines
commensurable in square only [X 73]

I say that no other rational straight line can be annexed to AB which is commensurable with the whole in square only

For if possible let BD be so annexed

therefore AD, DB are also rational straight lines commensurable in square only [X 73]

Now since the excess of the squares on AD, DB over twice the rectangle AD, DB is also the excess of the squares on AC, CB over twice the rectangle AC, CB ,

for both exceed by the same the square on AB [II 7]

therefore alternately the excess of the squares on AD, DB over the squares on AC, CB is the excess of twice the rectangle AD, DB over twice the rectangle AC, CB

But the squares on AD, DB exceed the squares on AC, CB by a rational area
for both are rational

therefore twice the rectangle AD, DB also exceeds twice the rectangle AC, CB by a rational area

which is impossible

for both are medial [x 21] and a medial area does not exceed a medial by a rational area [x 96]

Therefore no other rational straight line can be annexed to AB which is commensurable with the whole in square only

Therefore only one rational straight line can be annexed to an apotome which is commensurable with the whole in square only Q E D

PROPOSITION 80

To a first apotome of a medial straight line only one medial straight line can be annexed which is commensurable with the whole in square only and which contains with the whole a rational rectangle

For let AB be a first apotome of a medial straight line and let BC be an annex to AB , therefore AC CB are medial straight lines commensurable in square only and such that the rectangle AC CB which they contain is rational, [x 74]

I say that no other medial straight line can be annexed to AB which is commensurable with the whole in square only and which contains with the whole a rational area

For if possible let DB also be so annexed therefore AD DB are medial straight lines commensurable in square only and such that the rectangle AD DB which they contain is rational [x 74]

Now since the excess of the squares on AD DB over twice the rectangle AD DB is also the excess of the squares on AC CB over twice the rectangle AC CB ,

for they exceed by the same the square on AB , [ii 7] therefore alternately the excess of the squares on AD DB over the squares on AC CB is also the excess of twice the rectangle AD DB over twice the rectangle AC CB

But twice the rectangle AD DB exceeds twice the rectangle AC CB by a rational area

for both are rational

Therefore the squares on AD DB also exceed the squares on AC CB by a rational area

which is impossible

for both are medial [x 15 and 23 1 or] and a medial area does not exceed a medial by a rational area [x 96]

Therefore etc

Q E D

PROPOSITION 81

To a second apotome of a medial straight line only one medial straight line can be annexed which is commensurable with the whole in square only and which contains with the whole a medial rectangle

Let AB be a second apotome of a medial straight line and BC an annex to AB , therefore AC CB are medial straight lines commensurable in square only and such that the rectangle AC CB which they contain is medial [x 73]

I say that no other medial straight line can be annexed to AB which is com

measurable with the whole in square only and which contains with the whole a medial rectangle

For, if possible let BD also be so annexed,

therefore AD , DB are also medial straight lines commensurable in square only and such that the rectangle AD , DB which they contain is medial [x 75]

Let a rational straight line EF be set out
let EG equal to the squares on AC , CB be applied to EF , producing EM as breadth
and let HG equal to twice the rectangle AC , CB be subtracted producing HM as breadth

therefore the remainder EL is equal to the square on AB , [ii 7]

so that AB is the "side" of EL

Again let EI equal to the squares on AD , DB be applied to EF , producing EN as breadth

But EL is also equal to the square on AB ,
therefore the remainder HI is equal to twice the rectangle AD , DB [ii 7]

Now since AC , CB are medial straight lines
therefore the squares on AC , CB are also medial

And they are equal to EG

therefore EG is also medial [x 15 and 23 Por.]

And it is applied to the rational straight line EF , producing EM as breadth
therefore EM is rational and incommensurable in length with EF [x 22]

Again since the rectangle AC , CB is medial

twice the rectangle AC , CB is also medial [x 23 Por.]

And it is equal to HG ,

therefore HG is also medial

And it is applied to the rational straight line EF producing HM as breadth,
therefore HM is also rational and incommensurable in length with EF [x 22]

And since AC , CB are commensurable in square only

therefore AC is incommensurable in length with CB

But as AC is to CB , so is the square on AC to the rectangle AC , CB ,
therefore the square on AC is incommensurable with the rectangle AC , CB [x 11]

But the squares on AC , CB are commensurable with the square on AC ,
while twice the rectangle AC , CB is commensurable with the rectangle AC , CB [x 6]

therefore the squares on AC , CB are incommensurable with twice the rectangle AC , CB [x 13]

And EG is equal to the squares on AC , CB ,

while HG is equal to twice the rectangle AC , CB

therefore EG is incommensurable with HG


But, as EG is to HG so is EM to HM [vi 1]

therefore EM is incommensurable in length with HM [x 11]

And both are rational

therefore EM , HM are rational straight lines commensurable in square only

therefore EH is an apotome and HM an annex to it [x 73]



Similarly we can prove that HN is also an annex to it,
therefore to an apotome different straight lines are annexed which are com-
mensurable with the wholes in square only
which is impossible [x 19]

Therefore etc Q E D

PROPOSITION 82

To a minor straight line only one straight line can be annexed which is incommensurable in square with the whole and which makes with the whole, the sum of the squares on them rational but twice the rectangle contained by them medial

Let AB be the minor straight line and let BC be an annex to AB ,
therefore AC , CB are straight lines incommensurable in square which make
the sum of the squares on them rational, but twice A B C D
the rectangle contained by them medial [x 76]

I say that no other straight line can be annexed to AB fulfilling the same conditions

For, if possible let BD be so annexed,
therefore AD , DB are also straight lines incommensurable in square which fulfil the aforesaid conditions [x 16]

Now since the excess of the squares on AD , DB over the squares on AC , CB
is also the excess of twice the rectangle AD , DB over twice the rectangle AC , CB ,

while the squares on AD , DB exceed the squares on AC , CB by a rational area,
for both are rational

therefore twice the rectangle AD , DB also exceeds twice the rectangle AC , CB
by a rational area

which is impossible for both are medial [x 26]

Therefore to a minor straight line only one straight line can be annexed
which is incommensurable in square with the whole and which makes the
squares on them added together rational but twice the rectangle contained by
them medial Q E D

PROPOSITION 83

To a straight line which produces with a rational area a medial whole only one straight line can be annexed which is incommensurable in square with the whole straight line and which with the whole straight line makes the sum of the squares on them medial but twice the rectangle contained by them rational

Let AB be the straight line which produces with a rational area a medial whole

and let BC be an annex to AB ,
therefore AC , CB are straight lines incommensurable in square which fulfil the given conditions A B C D

I say that no other straight line can be annexed to AB which fulfils the same conditions [x 17]

For if possible let BD be so annexed
therefore AD , DB are also straight lines incommensurable in square which fulfil the given conditions [x 77]

Since then as in the preceding cases
the excess of the squares on AD , DB over the squares on AC , CB is also the

excess of twice the rectangle AD DB over twice the rectangle AC , CB , while twice the rectangle AD , DB exceeds twice the rectangle AC , CB by a rational area

for both are rational

therefore the squares on AD DB also exceed the squares on AC , CB by a rational area

which is impossible for both are medial [X 26]

Therefore no other straight line can be annexed to AB which is incommensurable in square with the whole and which with the whole fulfils the aforesaid conditions,

therefore only one straight line can be so annexed

Q E D

PROPOSITION 84


To a straight line which produces with a medial area a medial whole only one straight line can be annexed which is incommensurable in square with the whole straight line and which with the whole straight line makes the sum of the squares on them medial and twice the rectangle contained by them both medial and also incommensurable with the sum of the squares on them

Let AB be the straight line which produces with a medial area a medial whole

and BC an annex to it,

therefore AC , CB are straight lines incommensurable in square which fulfil the aforesaid conditions [X 78]

A B C D



I say that no other straight line can be annexed to AB which fulfils the aforesaid conditions

For if possible let BD be so annexed

so that AD DB are also straight lines incommensurable in square which make the squares on AD DB added together medial twice the rectangle AD DB medial and also the squares

on AD DB incommensurable with twice the rectangle AD DB [X 78]

Let a rational straight line EF be set out
let EG equal to the squares on AC CB be applied to EF producing EM as breadth

and let HG equal to twice the rectangle AC CB be applied to EF producing HM as breadth

therefore the remainder the square on AB [II 7] is equal to EL

therefore AB is the side of EL

Again let EI equal to the squares on AD DB be applied to EI producing EN as breadth

But the square on AB is also equal to EL ,
therefore the remainder twice the rectangle AD DB [II 7] is equal to HI

Now since the sum of the squares on AC CB is medial and is equal to EG
therefore EG is also medial

And it is applied to the rational straight line EF producing EM as breadth
therefore EM is rational and incommensurable in length with EF [X 22]

Again, since twice the rectangle AC , CB is medial and is equal to HG
therefore HG is also medial

And it is applied to the rational straight line EF producing HM as breadth
therefore HM is rational and incommensurable in length with EF [x 22]

And, since the squares on AC , CB are incommensurable with twice the rectangle AC CB

EG is also incommensurable with HG ,

therefore EM is also incommensurable in length with MH [vi 1 & 11]

And both are rational,

therefore EM , MH are rational straight lines commensurable in square only
therefore EH is an apotome and HM an annex to it [x 13]

Similarly we can prove that EH is again an apotome and HN an annex to it

Therefore to an apotome different rational straight lines are annexed which
are commensurable with the wholes in square only

which was proved impossible [x 9]

Therefore no other straight line can be so annexed to AB

Therefore to AB only one straight line can be annexed which is incommensurable in square with the whole and which with the whole makes the squares on them added together medial twice the rectangle contained by them medial, and also the squares on them incommensurable with twice the rectangle contained by them

Q E D

DEFINITIONS III

1 Given a rational straight line and an apotome if the square on the whole be greater than the square on the annex by the square on a straight line commensurable in length with the whole and the whole be commensurable in length with the rational straight line set out let the apotome be called a *first apotome*

2 But if the annex be commensurable in length with the rational straight line set out and the square on the whole be greater than that on the annex by the square on a straight line commensurable with the whole let the apotome be called a *second apotome*

3 But if neither be commensurable in length with the rational straight line set out and the square on the whole be greater than the square on the annex by the square on a straight line commensurable with the whole, let the apotome be called a *third apotome*

4 Again if the square on the whole be greater than the square on the annex by the square on a straight line incommensurable with the whole then if the whole be commensurable in length with the rational straight line set out let the apotome be called a *fourth apotome*

5 if the annex be so commensurable a *fifth*

6 and if neither a *sixth*

PROPOSITION 85


To find the first apotome

Let a rational straight line A be set out

and let BG be commensurable in length with A ,

therefore BG is also rational

Let two square numbers DE , EF be set out, and let their difference FD not be square,



therefore neither has ED to DF the ratio which a square number has to a square number

Let it be contrived that

as ED is to DF , so is the square on BG to the square on GC , [x 6 Por] therefore the square on BG is commensurable with the square on GC [x 6]

But the square on BG is rational,

therefore the square on GC is also rational,

therefore GC is also rational

And since ED has not to DF the ratio which a square number has to a square number,

therefore neither has the square on BG to the square on GC the ratio which a square number has to a square number,

therefore BG is incommensurable in length with GC [x 9]

And both are rational,

therefore BG , GC are rational straight lines commensurable in square only

therefore BC is an apotome [x 73]

I say next that it is also a first apotome

For let the square on H be that by which the square on BG is greater than the square on GC

Now since as ED is to FD so is the square on BG to the square on GC , therefore also, *convertendo* [v 19, Por]

as DE is to EF , so is the square on GB to the square on H

But DE has to EF the ratio which a square number has to a square number, for each is square,

therefore the square on GB also has to the square on H the ratio which a square number has to a square number

therefore BG is commensurable in length with H [x 9]

And the square on BG is greater than the square on GC by the square on H , therefore the square on BG is greater than the square on GC by the square on a straight line commensurable in length with BG

And the whole BG is commensurable in length with the rational straight line A set out

Therefore BC is a first apotome [x Defi III 1]

Therefore the first apotome BC has been found

(Being) that which it was required to find Q E D

PROPOSITION 86

To find the second apotome

Let a rational straight line A be set out and GC commensurable in length with A ,

therefore GC is rational

Let two square numbers DE , EF be set out and let their difference FD not be square

Now let it be contrived that as FD is to DE so is the square on CG to the square on GB

Therefore the square on CG is commensurable with the square on GB

But the square on CG is rational,
 therefore the square on GB is also rational,
 therefore GB is rational

And, since the square on GC has not to the
 square on GB the ratio which a square num-
 ber has to a square number
 CG is incommensurable in length with GB

[x 9]

And both are rational,
 therefore CG , GB are rational straight lines
 commensurable in square only,

therefore BC is an apotome

[x 73]

I say next that it is also a second apotome

For let the square on H be that by which the square on BG is greater than
 the square on GC

Since then as the square on BG is to the square on GC , so is the number ED
 to the number DF

therefore *convertendo*

as the square on BG is to the square on H , so is DE to EF [v 19 Por]

And each of the numbers DE , EF is square
 therefore the square on BG has to the square on H the ratio which a square
 number has to a square number

therefore BG is commensurable in length with H [x 9]

And the square on BG is greater than the square on GC by the square on H ,
 therefore the square on BG is greater than the square on GC by the square on a
 straight line commensurable in length with BG

And CG the annex is commensurable with the rational straight line A set
 out

Therefore BC is a second apotome

[x Def III 2]

Therefore the second apotome BC has been found

Q E D

PROPOSITION 87

To find the third apotome

Let a rational straight line A be set out
 let three numbers E BC CD be set out
 which have not to one another the ratio
 which a square number has to a square num-
 ber but let CB have to BD the ratio which a
 square number has to a square number

Let it be contrived that as E is to BC so is
 the square on A to the square on FG
 and as BC is to CD , so is the square on FG
 to the square on GH



[x 6 Por]

Since then, as E is to BC so is the square on A to the square on FG ,
 therefore the square on A is commensurable with the square on FG [x 6]

But the square on A is rational

therefore the square on FG is also rational

therefore FG is rational



And since E has not to BC the ratio which a square number has to a square number,
therefore neither has the square on A to the square on FG the ratio which a square number has to a square number,

therefore A is incommensurable in length with FG [x 9]

Again, since as BC is to CD so is the square on FG to the square on GH
therefore the square on FG is commensurable with the square on GH [x 6]

But the square on FG is rational,

therefore the square on GH is also rational,

therefore GH is rational

And since BC has not to CD the ratio which a square number has to a square number,

therefore neither has the square on FG to the square on GH the ratio which a square number has to a square number

therefore FG is incommensurable in length with GH [x 9]

And both are rational

therefore FG, GH are rational straight lines commensurable in square only,

therefore FH is an apotome [x 73]

I say next that it is also a third apotome

For since, as E is to BC so is the square on A to the square on FG

and, as BC is to CD so is the square on FG to the square on GH

therefore *ex aequali*, as E is to CD so is the square on A to the square on HG [v 22]

But E has not to CD the ratio which a square number has to a square number
therefore neither has the square on A to the square on GH the ratio which a square number has to a square number,

therefore A is incommensurable in length with GH [x 9]

Therefore neither of the straight lines FG, GH is commensurable in length with the rational straight line A set out

Now let the square on A be that by which the square on FG is greater than the square on GH

Since then as BC is to CD so is the square on FG to the square on GH
therefore *convertendo* as BC is to BD so is the square on FG to the square on A [v 19 Por]

But BC has to BD the ratio which a square number has to a square number,
therefore the square on FG also has to the square on A the ratio which a square number has to a square number

Therefore FG is commensurable in length with A [x 9]
and the square on FG is greater than the square on GH by the square on a straight line commensurable with FG

And neither of the straight lines FG, GH is commensurable in length with the rational straight line A set out

therefore FH is a third apotome [x Def III 3]

Therefore the third apotome FH has been found

Q E D


PROPOSITION 88

To find the fourth apotome

Let a rational straight line A be set out and BG commensurable in length with it

therefore BG is also rational

Let two numbers DF , FE be set out such that the whole DE has not to either of the numbers DF , FE the ratio which a square number has to a square number



Let it be contrived that as DE is to EF , so is the square on BG to the square on GC , [x 6 Por]

therefore the square on BG is commensurable with the square on GC [x 6]

But the square on BG is rational

therefore the square on GC is also rational,

therefore GC is rational

Now since DE has not to EF the ratio which a square number has to a square number,

therefore neither has the square on BG to the square on GC the ratio which a square number has to a square number,

therefore BG is incommensurable in length with GC [x 9]

And both are rational,

therefore BG , GC are rational straight lines commensurable in square only, therefore BC is an apotome [x 73]

Now let the square on H be that by which the square on BG is greater than the square on GC

Since then as DE is to EF so is the square on BG to the square on GC , therefore also *convertendo* as ED is to DF so is the square on GB to the square on H [v 10 Por]

But ED has not to DF the ratio which a square number has to a square number

therefore neither has the square on GB to the square on H the ratio which a square number has to a square number

therefore BG is incommensurable in length with H [x 9]

And the square on BG is greater than the square on GC by the square on H , therefore the square on BG is greater than the square on GC by the square on a straight line incommensurable with BG

And the whole BG is commensurable in length with the rational straight line A set out

Therefore BC is a fourth apotome [x Def m 4]

Therefore the fourth apotome has been found Q E D

PROPOSITION 89

To find the fifth apotome

Let a rational straight line A be set out and let CG be commensurable in length with A

therefore CG is rational

Let two numbers DF , FE be set out such that DE again has not to either of the numbers DF , FE the ratio which a square number has to a square number,

and let it be contrived that as FE is to ED so is the square on CG to the square on GB

Therefore the square on GB is also rational, [x 6]
therefore GB is also rational

Now since as DE is to EF , so is the square on BG to the square on GC

while DE has not to EF the ratio which a square number has to a square number

therefore neither has the square on BG to the square on GC the ratio which a square number has to a square number

therefore BG is incommensurable in length with GC [x 9]

And both are rational,
therefore BG GC are rational straight lines commensurable in square only,

therefore BC is an apotome [x 73]

I say next that it is also a fifth apotome

For let the square on H be that by which the square on BG is greater than the square on GC

Since then as the square on BG is to the square on GC , so is DE to EF ,
therefore, *convertendo* as ED is to DF so is the square on BG to the square on H [v 19 Por]

But ED has not to DF the ratio which a square number has to a square number,
therefore neither has the square on BG to the square on H the ratio which a square number has to a square number

therefore BG is incommensurable in length with H [x. 9]

And the square on BG is greater than the square on GC by the square on H ,
therefore the square on GB is greater than the square on GC by the square on a straight line incommensurable in length with GB

And the annex CG is commensurable in length with the rational straight line A set out


therefore BC is a fifth apotome [x Deff III 5]

Therefore the fifth apotome BC has been found q.e.d.

PROPOSITION 90

To find the sixth apotome

Let a rational straight line A be set out and three numbers E BC , CD not having to one another the ratio which a square number has to a square number and further let CB also not have to BD the ratio which a square number has to a square number



Let it be contrived that as E is to BC , so is the square on A to the square on FG ,
and as BC is to CD so is the square on FG to the square on GH [x 6 Por]

Now since as E is to BC so is the square on A to the square on FG

therefore the square on A is commensurable with the square on FG [x 6]

But the square on A is rational

therefore the square on FG is also rational

therefore FG is also rational

And, since E has not to BC the ratio which a square number has to a square number,

therefore neither has the square on A to the square on FG the ratio which a square number has to a square number

therefore A is incommensurable in length with FG [x 9]

Again since as BC is to CD so is the square on FG to the square on GH therefore the square on FG is commensurable with the square on GH [x 6]

But the square on FG is rational

therefore the square on GH is also rational,

therefore GH is also rational

And since BC has not to CD the ratio which a square number has to a square number,

therefore neither has the square on FG to the square on GH the ratio which a square number has to a square number

therefore FG is incommensurable in length with GH [x 9]

And both are rational

therefore FG , GH are rational straight lines commensurable in square only,

therefore FH is an apotome [x 13]

I say next that it is also a sixth apotome

For since as E is to BC , so is the square on A to the square on FG

and as BC is to CD so is the square on FG to the square on GH ,

therefore *ex æquali* as E is to CD , so is the square on A to the square on GH [v 22]

But E has not to CD the ratio which a square number has to a square number

therefore neither has the square on A to the square on GH the ratio which a square number has to a square number,

therefore A is incommensurable in length with GH , [x 9]

therefore neither of the straight lines FG GH is commensurable in length with the rational straight line A

Now let the square on A be that by which the square on FG is greater than the square on GH

Since then as BC is to CD so is the square on FG to the square on GH

therefore *conuertendo* as CB is to BD , so is the square on FG to the square on A [v 19, For]

But CB has not to BD the ratio which a square number has to a square number,

therefore neither has the square on FG to the square on A the ratio which a square number has to a square number

therefore FG is incommensurable in length with A [x 9]

And the square on FG is greater than the square on GH by the square on A , therefore the square on FG is greater than the square on GH by the square on a straight line incommensurable in length with FG

And neither of the straight lines FG GH is commensurable with the rational straight line A set out

Therefore FH is a sixth apotome

Therefore the sixth apotome FH has been found

[x Def III 6]
Q E D

PROPOSITION 91

If an area be contained by a rational straight line and a first apotome the "side" of the area is an apotome

For let the area AB be contained by the rational straight line AC and the first apotome AD


I say that the 'side' of the area AB is an apotome

For since AD is a first apotome let DG be its annex,
therefore AG GD are rational straight lines commensurable in square only [x 73]

And the whole AG is commensurable with the rational straight line AC set out

and the square on AG is greater than the square on GD by the square on a straight line commensurable in length with AG [x Def III 1]

if therefore there be applied to AG a parallelogram equal to the fourth part of the square on DG and deficient by a square figure it divides it into commensurable parts [x 17]



Let DG be bisected at E
let there be applied to AG a parallelogram equal to the square on EG and deficient by a square figure
and let it be the rectangle AF FG
therefore AF is commensurable with FG

And through the points E F G let EH FI GK be drawn parallel to AC

Now since AF is commensurable in length with FG
therefore AG is also commensurable in length with each of the straight lines AF FG [x 15]

But AG is commensurable with AC ,
therefore each of the straight lines AF FG is commensurable in length with AC [x 12]

And AC is rational

therefore each of the straight lines AF FG is also rational

so that each of the rectangles AI FA is also rational [x 19]

Now since DE is commensurable in length with EG

therefore DG is also commensurable in length with each of the straight lines DE EG [x 15]

But DG is rational and incommensurable in length with AC

therefore each of the straight lines DE , EG is also rational and incommensurable in length with AC [x 13]

therefore each of the rectangles DH EK is medial [x 21]

Now let the square LM be made equal to AI and let there be subtracted the square NO having a common angle with it the angle LPN , and equal to FA

therefore the squares LM NO are about the same diameter [vi 26]

Let PR be their diameter and let the figure be drawn

Since then the rectangle contained by AF , FG is equal to the square on EG
 therefore, as AF is to EG , so is EG to FG {vi 17}

But, as AF is to LG so is AI to EK
 and as EG is to FG so is EK to AF , {vi 11}
 therefore EK is a mean proportional between AI , AF {v 11}

But MN is also a mean proportional between LM , NO , as was before proved
 {Lemma after v 53}

and AI is equal to the square LM , and AF to AO ,
 therefore MA is also equal to EK

But EK is equal to DH and MN to LO ,
 therefore DA is equal to the gnomon UVW and NO
 But AA is also equal to the squares LM , NO

therefore the remainder AB is equal to ST

But ST is the square on LN ,
 therefore the square on LN is equal to AB ,
 therefore LN is the "side" of AB

I say next that LN is an apotome

For since each of the rectangles AI , FA is rational,
 and they are equal to LM , NO

therefore each of the squares LM , NO that is, the squares on LP , PN respectively is also rational

therefore each of the straight lines LP , PN is also rational

Again since DH is medial and is equal to LO

therefore LO is also medial

Since then LO is medial

while NO is rational

therefore LO is incommensurable with NO

But as LO is to NO so is LP to PN {vi 11}

therefore LP is incommensurable in length with PN {x 11}

And both are rational

therefore LP , PN are rational straight lines commensurable in square only
 therefore LN is an apotome {x 73}

And it is the "side" of the area AB

therefore the "side" of the area AB is an apotome

Therefore etc

Q E D

PROPOSITION 92

If an area be contained by a rational straight line and a second apotome, the "side" of the area is a first apotome of a medial straight line

For let the area AB be contained by the rational straight line AC and the second apotome AD


I say that the "side" of the area AB is a first apotome of a medial straight line

For let DG be the annex to AD

therefore AG , GD are rational straight lines commensurable in square only
 {x 73}

and the annex DG is commensurable with the rational straight line AC set out,
 while the square on the whole AG is greater than the square on the annex GD
 by the square on a straight line commensurable in length with AG
 {x Def m 1}

Since, then the square on AG is greater than the square on GD by the square on a straight line commensurable with AG ,



therefore, if there be applied to AG a parallelogram equal to the fourth part of the square on GD and deficient by a square figure it divides it into commensurable parts [x 17]

Let then DG be bisected at E , let there be applied to AG a parallelogram equal to the square on EG and deficient by a square figure, and let it be the rectangle AF, FG , therefore AF is commensurable in length with FG

Therefore AG is also commensurable in length with each of the straight lines AF, FG [x 15]

But AG is rational and incommensurable in length with AC , therefore each of the straight lines AF, FG is also rational and incommensurable in length with AC , [x 13]

therefore each of the rectangles AI, FK is medial [x 21]

Again since DE is commensurable with EG therefore DG is also commensurable with each of the straight lines DE, EG [x 15]

But DG is commensurable in length with AC

Therefore each of the rectangles DH, EK is rational [x 19]

Let then the square LM be constructed equal to AI , and let there be subtracted NO equal to FK and being about the same angle with LM namely the angle LPM ,

therefore the squares LM, NO are about the same diameter [vi 26]

Let PR be their diameter and let the figure be drawn

Since then AI, FK are medial and are equal to the squares on LP, PN ,

the squares on LP, PN are also medial

therefore LP, PN are also medial straight lines commensurable in square only

And since the rectangle AF, FG is equal to the square on EG ,

therefore, as AF is to EG so is EG to FG , [vi 17]

while, as AF is to EG so is AI to EK

and as EG is to FG so is EK to FK [vi 11]

therefore EK is a mean proportional between AI, FK [v 11]

But MN is also a mean proportional between the squares LM, NO

and AI is equal to LM and FK to NO ,

therefore MN is also equal to EK

But DH is equal to EK and LO equal to MN

therefore the whole DA is equal to the gnomon UVW and AO

Since then the whole AK is equal to LM, AO

and, in these DK is equal to the gnomon UVW and NO

therefore the remainder AB is equal to TS

But TS is the square on LN

therefore the square on LN is equal to the area AB ,

therefore LN is the "side" of the area AB

I say that LN is a first apotome of a medial straight line

For, since EK is rational and is equal to LO

therefore LO that is the rectangle LP, PN , is rational

But NO was proved medial,

therefore LO is incommensurable with NO

But, as LO is to NO , so is LP to PN , [VI 1]

therefore LP, PN are incommensurable in length [X 11]

Therefore LP, PN are medial straight lines commensurable in square only which contain a rational rectangle,

therefore LN is a first apotome of a medial straight line [X 74]

And it is the 'side' of the area AB

Therefore the "side" of the area AB is a first apotome of a medial straight line
Q E D

PROPOSITION 93

If an area be contained by a rational straight line and a third apotome the 'side' of the area is a second apotome of a medial straight line

For let the area AB be contained by the rational straight line AC and the third apotome AD

I say that the "side" of the area AB is a second apotome of a medial straight line

For let DG be the annex to AD

therefore AG, GD are rational straight lines commensurable in square only, and neither of the straight lines AG, GD is commensurable in length with the rational straight line AC set out

while the square on the whole AG is greater than the square on the annex DG by the square on a straight line commensurable with AG [X Def III 3]

Since, then, the square on AG is greater than the square on DG by the square on a straight line commensurable with AG ,

therefore if there be applied to AG a parallelogram equal to the fourth part of the square on DG and deficient by a square figure it will divide it into commensurable parts [X 17]

Let then DG be bisected at E

let there be applied to AG a parallelogram equal to the square on EG and deficient by a square figure,


and let it be the rectangle AF, FG

Let EH, FI, GK be drawn through the points E, F, G parallel to AC

Therefore AF, FG are commensurable

rational
therefore AI is also commensurable with FA [VI 1 X 11]

And, since AF, FG are commensurable in length, therefore AG is also commensurable in length with each of the straight lines AF, FG [X 13]



But AG is rational and incommensurable in length with AC ,

so that AF FG are so also

[x 13]

Therefore each of the rectangles AI FK is medial

[x 21]

Again since DE is commensurable in length with EG

therefore DG is also commensurable in length with each of the straight lines DE EG

[x 15]

But GD is rational and incommensurable in length with AC

therefore each of the straight lines DE , EG is also rational and incommensurable in length with AC

[x 13]

therefore each of the rectangles DIH EK is medial

[x 21]

And since AG GD are commensurable in square only

therefore AG is incommensurable in length with GD

But AG is commensurable in length with AF and DG with EG

therefore AF is incommensurable in length with EG

[x 13]

But as AF is to EG so is AI to EK

[vi 1]

therefore AI is incommensurable with EK

[x 11]

Now let the square LU be constructed equal to AI

and let there be subtracted AO equal to FK and being about the same angle with LM

therefore LM NO are about the same diameter

[vi 26]

Let PR be their diameter and let the figure be drawn

Now since the rectangle AF FG is equal to the square on EG

therefore as AF is to EG so is EG to FG

[vi 17]

But as AF is to EG , so is AI to EK

and as EG is to FG so is EK to FK ,

[vi 1]

therefore also as AI is to EK so is EK to FK

[v 11]

therefore EK is a mean proportional between AI FK

But MA is also a mean proportional between the squares LM NO

and AI is equal to LM and FK to NO

therefore EK is also equal to MN

But MA is equal to LO and EK equal to DH

therefore the whole DH is also equal to the gnomon UVW and AO

But AK is also equal to LM NO

therefore the remainder AB is equal to ST that is to the square on LN

therefore LN is the side of the area AB

I say that LN is a second apotome of a medial straight line

For since AI FK were proved medial and are equal to the squares on LP PN

therefore each of the squares on LP PN is also medial

therefore each of the straight lines LP PN is medial

And since AI is commensurable with FK

[vi 1 x 11]

therefore the square on LP is also commensurable with the square on PN

Again since AI was proved incommensurable with EK

therefore LM is also incommensurable with MA

that is the square on LP with the rectangle LP PN

so that LP is also incommensurable in length with PN ,

[vi 1, x 11]

therefore LP PN are medial straight lines commensurable in square only

I say next that they also contain a medial rectangle

For since EK was proved medial and is equal to the rectangle LP , PN

therefore the rectangle LP , PN is also medial
 so that LP PN are medial straight lines commensurable in square only which contain a medial rectangle

Therefore LN is a second apotome of a medial straight line, [x 75]
 and it is the side of the area AB

Therefore the 'side' of the area AB is a second apotome of a medial straight line
Q E D

PROPOSITION 94

If an area be contained by a rational straight line and a fourth apotome, the 'side of the area is minor

For let the area AB be contained by the rational straight line AC and the fourth apotome AD ,

I say that the side of the area AB is minor


For let DG be the annex to AD

therefore AG GD are rational straight lines commensurable in square only, AG is commensurable in length with the rational straight line AC set out and the square on the whole AG is greater than the square on the annex DG by the square on a straight line incommensurable in length with AG

[x Def iii 4]

Since then the square on AG is greater than the square on GD by the square on a straight line incommensurable in length with AG

therefore if there be applied to AG a parallelogram equal to the fourth part of the square on DG and deficient by a square figure it will divide it into incommensurable parts [x 18]



Let then DG be bisected at E
 let there be applied to AG a parallelogram equal to the square on EG and deficient by a square figure

and let it be the rectangle AF FG
 therefore AF is incommensurable in length with FG

Let EH FI GK be drawn through E F G parallel to AC BD

Since then AG is rational and commensurable in length with AC ,
 therefore the whole AK is rational [x 10]

Again since DG is incommensurable in length with AC and both are rational

therefore DK is medial [x 9]

Again since AF is incommensurable in length with FG

therefore AI is also incommensurable with FA [vi 1 x 11]

Now let the square LM be constructed equal to AI
 and let there be subtracted VO equal to FA and about the same angle the angle LPM

Therefore the squares LM AO are about the same diameter [vi 9]

Let PR be their diameter and let the figure be drawn

Since then, the rectangle AF FG is equal to the square on EG ,
therefore, proportionally, as AF is to EG so is EG to FG [vi 17]

But as AF is to EG , so is AI to EK ,
and as EG is to FG so is EK to FA , [vi 1]

therefore EK is a mean proportional between AI FA [v 11]

But MN is also a mean proportional between the squares LM , NO ,
and AI is equal to LM and FA to NO ,
therefore EK is also equal to MN

But DH is equal to EK , and LO is equal to MN ,
therefore the whole DA is equal to the gnomon UVW and NO

Since then the whole AA is equal to the squares LM , NO ,
and in these DA is equal to the gnomon UVW and the square NO
therefore the remainder AB is equal to ST , that is to the square on LN ,
therefore LN is the "side" of the area AB

I say that LN is the irrational straight line called minor

For, since AK is rational and is equal to the squares on LP , PN ,
therefore the sum of the squares on LP , PN is rational

Again since DA is medial,

and DA is equal to twice the rectangle LP , PN ,
therefore twice the rectangle LP PN is medial

And since AI was proved incommensurable with FA
therefore the square on LP is also incommensurable with the square on PN
Therefore LP , PN are straight lines incommensurable in square which make
the sum of the squares on them rational but twice the rectangle contained by
them medial

Therefore LN is the irrational straight line called minor, [x 76]
and it is the "side" of the area AB

Therefore the "side" of the area AB is minor Q E D

PROPOSITION 95

If an area be contained by a rational straight line and a fifth apotome the "side" of the area is a straight line which produces with a rational area a medial whole

For let the area AB be contained by the rational straight line AC and the fifth apotome AD

I say that the "side" of the area AB is a straight line which produces with a rational area a medial whole

For let DG be the annex to AD ,
therefore AG GD are rational straight lines commensurable in square only
the annex GD is commensurable in length with the rational straight line AC
set out

and the square on the whole AG is greater than the square on the annex DG by
the square on a straight line incommensurable with AG [x Def iii 5]

Therefore, if there be applied to AG a parallelogram equal to the fourth part
of the square on DG and deficient by a square figure it will divide it into in
commensurable parts [x 18]

Let then DG be bisected at the point E ,
let there be applied to AG a parallelogram equal to the square on EG and deficient by a square figure and let it be the rectangle AF , FG ,
therefore AF is incommensurable in length with FG

Now since AG is incommensurable in length with CA , and both are rational
therefore AK is medial [x 21]

Again since DG is rational and commensurable in length with AC

DA is rational [x 10]

Now let the square LM be constructed equal to AI and let the square NO equal to FA and about the same angle the angle LPM be subtracted

therefore the squares LM , NO are about the same diameter [vi 26]

Let PR be their diameter and let the figure be drawn

Similarly then we can prove that LN is the 'side' of the area AB

I say that LN is the straight line which produces with a rational area a medial whole

For since AK was proved medial and is equal to the squares on LP , PN ,
therefore the sum of the squares on LP , PN is medial

Again, since DA is rational and is equal to twice the rectangle LP , PN ,
the latter is itself also rational

And since AI is incommensurable with FA ,
therefore the square on LP is also incommensurable with the square on PN ,
therefore LP , PN are straight lines incommensurable in square which make
the sum of the squares on them medial but twice the rectangle contained by
them rational

Therefore the remainder LN is the irrational straight line called that which
produces with a rational area a medial whole [x 77]

and it is the 'side' of the area AB

Therefore the 'side' of the area AB is a straight line which produces with a
rational area a medial whole Q E D

PROPOSITION 96

If an area be contained by a rational straight line and a sixth apotome the "side of the area is a straight line which produces with a medial area a medial whole

For let the area AB be contained by the rational straight line AC and the sixth apotome AD

I say that the 'side' of the area AB is a straight line which produces with a medial area a medial whole


For let DG be the annex to AD

therefore AG , GD are rational straight lines commensurable in square only
neither of them is commensurable in length with the rational straight line AC
set out


and the square on the whole AG is greater than the square on the annex DG by
the square on a straight line incommensurable in length with AG

[x Def III 6]

Since then the square on AG is greater than the square on GD by the square
on a straight line incommensurable in length with AG ,



therefore if there be applied to $4G$ a parallelogram equal to the fourth part of the square on DG and deficient by a square figure it will divide it into incommensurable parts [x 18]



Let then DG be bisected at E
let there be applied to AG a parallelogram equal to the square on EG and deficient by a square figure, and let it be the rectangle AF, FG , therefore AF is incommensurable in length with FG

But as AF is to FG so is AI to FK [vi 1]
therefore AI is incommensurable with FK [x 11]

And since AG, AC are rational straight lines commensurable in square only

AK is medial [x 21]

Again since AC, DG are rational straight lines and incommensurable in length

DK is also medial [x 21]

Now, since AG, GD are commensurable in square only
therefore AG is incommensurable in length with GD

But as $4G$ is to GD so is AK to KD , [vi 1]
therefore AK is incommensurable with KD [x 11]

Now let the square LM be constructed equal to AI ,
and let NO equal to FK and about the same angle be subtracted,
therefore the squares LM, NO are about the same diameter [vi 26]

Let PR be their diameter and let the figure be drawn

Then in manner similar to the above we can prove that LN is the 'side' of the area AB

I say that LV is a straight line which produces with a medial area a medial whole

For since AK was proved medial and is equal to the squares on LP, PN
therefore the sum of the squares on LP, PN is medial

Again since DK was proved medial and is equal to twice the rectangle LP, PN

twice the rectangle LP, PN is also medial

And since AK was proved incommensurable with DK
the squares on LP, PN are also incommensurable with twice the rectangle LP, PN

And since AI is incommensurable with FK
therefore the square on LP is also incommensurable with the square on PN
therefore LP, PN are straight lines incommensurable in square which make the sum of the squares on them medial twice the rectangle contained by them medial and further the squares on them incommensurable with twice the rectangle contained by them

Therefore LV is the irrational straight line called that which produces with a medial area a medial whole [x 7^o]

and it is the "side" of the area AB

Therefore the "side" of the area is a straight line which produces with a medial area a medial whole Q E D

PROPOSITION 97


The square on an apotome applied to a rational straight line produces as breadth a first apotome

Let AB be an apotome and CD rational,
and to CD let there be applied CE equal to the square on AB and producing CF as breadth

I say that CF is a first apotome

For let BG be the annex to AB ,
therefore AG , GB are rational
straight lines commensurable
in square only [x 73]

To CD let there be applied
 CH equal to the square on AG
and KL equal to the square on
 BG



Therefore the whole CL is equal to the squares on AG , GB , and, in these, CE is equal to the square on AB ,
therefore the remainder FL is equal to twice the rectangle AG , GB [II 7]

Let FM be bisected at the point N ,
and let NO be drawn through N parallel to CD
therefore each of the rectangles FO , LN is equal to the rectangle AG , GB
Now since the squares on AG , GB are rational
and DM is equal to the squares on AG , GB ,
therefore DM is rational

And it has been applied to the rational straight line CD , producing CM as breadth

therefore CM is rational and commensurable in length with CD [x 90]

Again since twice the rectangle AG , GB is medial and FL is equal to twice the rectangle AG , GB ,

therefore FL is medial

And it is applied to the rational straight line CD producing FM as breadth
therefore FM is rational and incommensurable in length with CD [x 22]

And since the squares on AG , GB are rational
while twice the rectangle AG , GB is medial,
therefore the squares on AG , GB are incommensurable with twice the rectangle AG , GB

And CL is equal to the squares on AG , GB ,
and FL to twice the rectangle AG , GB ,
therefore DM is incommensurable with FL

But, as DM is to FL so is CM to FM ,
therefore CM is incommensurable in length with FM [VI 1]
[x 11]

And both are rational
therefore CM , MF are rational straight lines commensurable in square only
therefore CF is an apotome [x 73]

I say next that it is also a first apotome

For since the rectangle AG, GB is a mean proportional between the squares on AG, GB

and CH is equal to the square on AG ,

KL equal to the square on GB ,

and NL equal to the rectangle AG, GB ,

therefore NL is also a mean proportional between CH, KL ,

therefore as CH is to NL , so is NL to KL

But as CH is to NL , so is CA to NM ,

and, as NL is to KL so is NM to KM , [vi 1]

therefore the rectangle CK, KM is equal to the square on NM [vi 17] that is, to the fourth part of the square on FM

And since the square on AG is commensurable with the square on GB ,

CH is also commensurable with KL

But, as CH is to KL so is CA to KM [vi 1]

therefore CA is commensurable with KM [x 11]

Since, then, CM, MF are two unequal straight lines

and to CM there has been applied the rectangle CK, KM equal to the fourth part of the square on FM and deficient by a square figure

while CA is commensurable with KM

therefore the square on CM is greater than the square on MF by the square on a straight line commensurable in length with CM [x 17]

And CM is commensurable in length with the rational straight line CD set out,

therefore CF is a first apotome [x Deff III 1]

Therefore etc

Q E D

PROPOSITION 98


The square on a first apotome of a medial straight line applied to a rational straight line produces as breadth a second apotome

Let AB be a first apotome of a medial straight line and CD a rational straight line

and to CD let there be applied CE equal to the square on AB , producing CF as breadth

I say that CF is a second apotome

For let BG be the annex to AB ,
therefore AG, GB are medial straight lines commensurable in square only which contain a rational rectangle [x 74]



To CD let there be applied CH equal to the square on AG , producing CA as breadth, and KL equal to the square on GB , producing KM as breadth
therefore the whole CL is equal to the squares on AG, GB ,

therefore CL is also medial [x 15 and 23 Por]

And it is applied to the rational straight line CD producing CM as breadth
therefore CM is rational and incommensurable in length with CD [x 22]

Now since CL is equal to the squares on AG GB

and in these the square on AB is equal to CE

therefore the remainder twice the rectangle AG GB , is equal to FL [x 7]

But twice the rectangle AG GB is rational

therefore FL is rational

And it is applied to the rational straight line FE producing FM as breadth
therefore FM is also rational and commensurable in length with CD [x 20]

Now since the sum of the squares on AG GB that is CL is medial while
twice the rectangle AG GB that is FL is rational

therefore CL is incommensurable with FL

But as CL is to FL so is CM to FM , [vi 1]

therefore CM is incommensurable in length with FM [x 11]

And both are rational,

therefore CM MF are rational straight lines commensurable in square only,

therefore CF is an apotome [x 73]

I say next that it is also a second apotome

For let FM be bisected at N

and let NO be drawn through N parallel to CD ,

therefore each of the rectangles FO NL is equal to the rectangle AG GB

Now since the rectangle AG GB is a mean proportional between the squares
on AG GB

and the square on AG is equal to CH

the rectangle AG , GB to NL

and the square on GB to KL

therefore NL is also a mean proportional between CH KL

therefore as CH is to NL so is NL to KL

But as CH is to VL so is CK to NM

and as VL is to KL so is NM to MK [vi 1]

therefore as CK is to NM so is NM to KM [v 11]

therefore the rectangle CK KM is equal to the square on NM [vi 17] that is
to the fourth part of the square on FM

Since then CM MF are two unequal straight lines and the rectangle CK ,
 KM equal to the fourth part of the square on MF and deficient by a square
figure has been applied to the greater CM and divides it into commensurable
parts

therefore the square on CM is greater than the square on MF by the square on
a straight line commensurable in length with CM [x 11]

And the annex FM is commensurable in length with the rational straight
line CD set out

therefore CF is a second apotome [x Def m 2]

Therefore etc

Q E D


PROPOSITION 99

The square on a second apotome of a medial straight line applied to a rational
straight line produces as breadth a third apotome

Let AB be a second apotome of a medial straight line and CD rational
and to CD let there be applied CE equal to the square on AB producing CF as
breadth,

I say that CF is a third apotome

For let BG be the annex to AB ,
therefore AG GB are medial straight lines commensurable in square only
which contain a medial rectangle [x 75]



Let CH equal to the square
on AG be applied to CD , pro-
ducing CA as breadth
and let KL equal to the square
on BG be applied to KH pro-
ducing KM as breadth,
therefore the whole CL is equal
to the squares on AG , GB ,

therefore CL is also medial [x 15 and 23, Por.]

And it is applied to the rational straight line CD producing CM as breadth
therefore CM is rational and incommensurable in length with CD [x 22]

Now, since the whole CL is equal to the squares on AG GB and in these
 CE is equal to the square on AB

therefore the remainder LF is equal to twice the rectangle AG GB [ii 7]

Let then FM be bisected at the point N

and let NO be drawn parallel to CD ,

therefore each of the rectangles FO NL is equal to the rectangle AG , GB

But the rectangle AG , GB is medial,

therefore FL is also medial

And it is applied to the rational straight line EF producing FM as breadth
therefore FM is also rational and incommensurable in length with CD [x 22]

And since AG , GB are commensurable in square only,

therefore AG is incommensurable in length with GB

therefore the square on AG is also incommensurable with the rectangle AG
 GB [vi 1 x 11]

But the squares on AG GB are commensurable with the square on AG ,

and twice the rectangle AG GB with the rectangle AG , GB ,

therefore the squares on AG GB are incommensurable with twice the rectangle
 AG GB [x 13]

But CL is equal to the squares on AG GB

and FL is equal to twice the rectangle AG GB ,

therefore CL is also incommensurable with FL

But as CL is to FL so is CM to FM , [vi 1]

therefore CM is incommensurable in length with FM [x 11]

And both are rational,

therefore CM , MF are rational straight lines commensurable in square only

therefore CF is an apotome [v 73]

I say next that it is also a third apotome

For since the square on AG is commensurable with the square on GB

therefore CH is also commensurable with KL

so that CA is also commensurable with KM [vi 1 x 11]

And since the rectangle AG GB is a mean proportional between the squares
on AG GB

and CH is equal to the square on AG

KL equal to the square on GB

and NL equal to the rectangle AG GB

therefore NL is also a mean proportional between CH ; KL ,
 therefore as CH is to NL so is NL to KL

But, as CH is to NL so is CK to NM

and, as NL is to KL , so is NM to KM , [vi 1]

therefore as CK is to NM , so is NM to KM , [v 11]

therefore the rectangle CK KM is equal to [the square on NM , that is to] the fourth part of the square on FM

Since then CM MF are two unequal straight lines and a parallelogram equal to the fourth part of the square on FM and deficient by a square figure has been applied to CM and divides it into commensurable parts - -
 therefore the square on CM is greater than the square on MF by the square on a straight line commensurable with CM [x 17]

And neither of the straight lines CM MF is commensurable in length with the rational straight line CD set out

therefore CF is a third apotome [x Def iii 3]
 Q E D

Therefore etc

PROPOSITION 100


The square on a minor straight line applied to a rational straight line produces as breadth a fourth apotome

Let AB be a minor and CD a rational straight line and to the rational straight line CD let CE be applied equal to the square on AB and producing CF as breadth

I say that CF is a fourth apotome

For let BG be the annex to AB

therefore AG GB are straight lines incommensurable in square which make the sum of the squares on AG GB rational but twice the rectangle AG GB medial [x 76]



To CD let there be applied CH equal to the square on AG and producing CK as breadth

and KL equal to the square on BG producing KM as breadth,

therefore the whole CL is equal to the squares on AG , BG

And the sum of the squares on AG GB is rational

therefore CL is also rational

And it is applied to the rational straight line CD producing CM as breadth, therefore CM is also rational and commensurable in length with CD [x 20]

And since the whole CL is equal to the squares on AG GB and in these, CE is equal to the square on AB

therefore the remainder FL is equal to twice the rectangle AG GB [ii 7]

Let then FM be bisected at the point N

and let AO be drawn through A parallel to either of the straight lines CD ML , therefore each of the rectangles FO NL is equal to the rectangle AG GB

And since twice the rectangle AG GB is medial and is equal to FL

therefore FL is also medial

And it is applied to the rational straight line FE producing FM as breadth

therefore FM is rational and incommensurable in length with CD [x 22]

And since the sum of the squares on AG GB is rational

while twice the rectangle AG GB is medial,

the squares on AG GB are incommensurable with twice the rectangle AG GB

But GL is equal to the squares on AG , GB ,

and FL equal to twice the rectangle AG GB ,

therefore CL is incommensurable with FL

But as CL is to FL so is CM to MF ,

[vi 1]

therefore CM is incommensurable in length with MF

[x 11]

And both are rational,

therefore CM , MF are rational straight lines commensurable in square only

therefore CF is an apotome

[x 73]

I say that it is also a fourth apotome

For, since AG , GB are incommensurable in square

therefore the square on AG is also incommensurable with the square on GB

And CH is equal to the square on AG ,

and KL equal to the square on GB ,

therefore CH is incommensurable with KL

But as CH is to KL , so is CK to KM ,

[vi 1]

therefore CK is incommensurable in length with KM

[x 11]

And since the rectangle AG , GB is a mean proportional between the squares on AG , GB ,

and the square on AG is equal to CH ,

the square on GB to KL

and the rectangle AG GB to NL

therefore NL is a mean proportional between CH , KL

therefore as CH is to NL , so is NL to KL

But as CH is to NL so is CK to NM

and as NL is to KL so is NM to KM ,

[vi 1]

therefore as CK is to MN so is MN to KM

[v 11]

therefore the rectangle CK , KM is equal to the square on MN [vi 17] that is to the fourth part of the square on FM

Since then CM MF are two unequal straight lines and the rectangle CK , KM equal to the fourth part of the square on MF and deficient by a square figure has been applied to CM and divides it into incommensurable parts, therefore the square on CM is greater than the square on MF by the square on a straight line incommensurable with CM

[x 18]

And the whole CM is commensurable in length with the rational straight line CD set out,

therefore CF is a fourth apotome

[x Deff III 4]

Therefore etc

Q E D

PROPOSITION 101

The square on the straight line which produces with a rational area a medial whole, if applied to a rational straight line produces as breadth a fifth apotome

Let AB be the straight line which produces with a rational area a medial whole and CD a rational straight line and to CD let CE be applied equal to the square on AB and producing CF as breadth

I say that CF is a fifth apotome

For let BG be the annex to AB

therefore AG, GB are straight lines incommensurable in square which make the sum of the squares on them medial but twice the rectangle contained by them rational [x 7]

To CD let there be applied CH equal to the square on AG and KL equal to the square on GB ,

therefore the whole CL is equal to the squares on AG, GB

But the sum of the squares on AG, GB together is medial, therefore CL is medial

And it is applied to the rational straight line CD , producing CM as breadth therefore CM is rational and incommensurable with CD [x 2]

And, since the whole CL is equal to the squares on AG, GB , and in these CE is equal to the square on AB ,

therefore the remainder FL is equal to twice the rectangle AG, GB [ii 4]

Let then FM be bisected at N ,

and through N let NO be drawn parallel to either of the straight lines CD, ML , therefore each of the rectangles FO, NL is equal to the rectangle AG, GB

And, since twice the rectangle AG, GB is rational and equal to FL therefore FL is rational

And it is applied to the rational straight line EF producing FM as breadth therefore FM is rational and commensurable in length with CD [x 20]

Now since CL is medial and FL rational,

therefore CL is incommensurable with FL

But as CL is to FL so is CM to MF [vi 1]

therefore CM is incommensurable in length with MF [x 11]

And both are rational,

therefore CM, MF are rational straight lines commensurable in square only therefore CF is an apotome [x 73]

I say next that it is also a fifth apotome

For we can prove similarly that the rectangle CK, KM is equal to the square on NM that is to the fourth part of the square on FM

And since the square on AG is incommensurable with the square on GB ,

while the square on AG is equal to CH ,

and the square on GB to KL

therefore CH is incommensurable with KL

But as CH is to KL so is CK to KM , [vi 1]

therefore CK is incommensurable in length with KM [x 11]


Since then CM, MF are two unequal straight lines

and a parallelogram equal to the fourth part of the square on FM and deficient by a square figure has been applied to CM , and divides it into incommensurable parts

therefore the square on CM is greater than the square on MF by the square on a straight line incommensurable with CM [x 18]

And the annex FM is commensurable with the rational straight line CD set out therefore CF is a fifth apotome [x Def III 5]

Q E D




PROPOSITION 102

The square on the straight line which produces with a medial area a medial whole if applied to a rational straight line produces as breadth a sixth apotome

Let AB be the straight line which produces with a medial area a medial whole and CD a rational straight line, and to CD let CE be applied equal to the square on AB and producing CF as breadth

I say that CF is a sixth apotome

For let BG be the annex to AB ,



therefore AG GB are straight lines incommensurable in square which make the sum of the squares on them medial, twice the rectangle AG GB medial and the squares on AG GB incommensurable with twice the rectangle AG GB

[x 78]

Now to CD let there be applied CH equal to the square on AG and producing CL as breadth

and KL equal to the square on BG ,

therefore the whole CL is equal to the squares on AG GB

therefore CL is also medial

And it is applied to the rational straight line CD producing CM as breadth, therefore CM is rational and incommensurable in length with CD [x 22]

Since now CL is equal to the squares on AG GB ,

and in these CE is equal to the square on AB

therefore the remainder FL is equal to twice the rectangle AG GB [ii 7]

And twice the rectangle AG GB is medial,

therefore FL is also medial

And it is applied to the rational straight line FE producing FM as breadth, therefore FM is rational and incommensurable in length with CD [x 22]

And since the squares on AG GB are incommensurable with twice the rectangle AG GB

and CL is equal to the squares on AG GB

and FL equal to twice the rectangle AG GB

therefore CL is incommensurable with FL

But as CL is to FL so is CM to MF [vi 1]

therefore CM is incommensurable in length with MF [x 11]

And both are rational

Therefore CM MF are rational straight lines commensurable in square only

therefore CF is an apotome [x 73]

I say next that it is also a sixth apotome

For since FL is equal to twice the rectangle AG GB

let FM be bisected at N

and let NO be drawn through N parallel to CD

therefore each of the rectangles FO NL is equal to the rectangle AG GB

And since AG GB are incommensurable in square

therefore the square on AG is incommensurable with the square on GB
 But CH is equal to the square on AG ,

and KL is equal to the square on GB ,

therefore CH is incommensurable with KL

But as CH is to KL so is CA to KM [vi 1]

therefore CA is incommensurable with KM [x 11]

And since the rectangle AG GB is a mean proportional between the squares on AG , GB

and CH is equal to the square on AG ,

KL equal to the square on GB ,

and NL equal to the rectangle AG GB

therefore NL is also a mean proportional between CH , KL ,

therefore as CH is to NL so is NL to KL

And for the same reason as before the square on CM is greater than the square on MF by the square on a straight line incommensurable with CM [x 18]

And neither of them is commensurable with the rational straight line CD set out

therefore CF is a sixth apotome [v Def iii 6]

Q E D


PROPOSITION 103

A straight line commensurable in length with an apotome is an apotome and the same in order

Let AB be an apotome

and let CD be commensurable in length with AB

I say that CD is also an apotome and the same in order with AB



For since AB is an apotome let BE be the annex to it

therefore AE EB are rational straight lines commensurable in square only [x 7]

Let it be contrived that the ratio of BE to DF is the same as the ratio of AE to CD [x 17]

therefore also as one is to one so are all to all [v 12]

therefore also as the whole AF is to the whole CF so is AB to CD

But AB is commensurable in length with CD

Therefore AE is also commensurable with CF and BE with DF [x 11]

And AE EB are rational straight lines commensurable in square only
 therefore CF FD are also rational straight lines commensurable in square only [x 13]

Now since as AE is to CF so is BE to DF

alternately therefore as AE is to EB so is CF to FD [v 16]

And the square on AE is greater than the square on EB either by the square on a straight line commensurable with AE or by the square on a straight line incommensurable with it

If then the square on AE is greater than the square on EB by the square on a straight line commensurable with AE the square on CF will also be greater than the square on FD by the square on a straight line commensurable with CF [x 14]

And, if AE is commensurable in length with the rational straight line set out,

CF is so also [x 12]

if BE then DF also [id]

and if neither of the straight lines AF , EB , then neither of the straight lines CF , FD [x 13]

But if the square on AE is greater than the square on FB by the square on a straight line incommensurable with AE ,
the square on CF will also be greater than the square on FD by the square on a straight line incommensurable with CF [x 14]

And if AE is commensurable in length with the rational straight line set out,

CF is so also

if BF then DF also [x 12]

and if neither of the straight lines AE , EB , then neither of the straight lines CF , FD [x 13]

Therefore CD is an apotome and the same in order with AB Q E D

PROPOSITION 104

A straight line commensurable with an apotome of a medial straight line is an apotome of a medial straight line and the same in order

Let AB be an apotome of a medial straight line

and let CD be commensurable in length with AB ,

I say that CD is also an apotome of a medial straight line and the same in order with AB

For, since AB is an apotome of a medial straight line let EB be the annex to it

Therefore AE EB are medial straight lines commensurable in square only [x 74 75]

Let it be contrived that as AB is to CD , so is BE to DF [vi 12]

therefore AE is also commensurable with CF and BE with DF [v 12 x 11]

But AE EB are medial straight lines commensurable in square only

therefore CF FD are also medial straight lines [x 23] commensurable in square only [x 13]

therefore CD is an apotome of a medial straight line [x 74 75]

I say next that it is also the same in order with AB

Since as AE is to EB so is CF to FD

therefore also as the square on AE is to the rectangle AE EB so is the square on CF to the rectangle CF FD

But the square on AE is commensurable with the square on CF

therefore the rectangle AE , EB is also commensurable with the rectangle CF FD [v 16 x 11]

Therefore, if the rectangle AE EB is rational the rectangle CF FD will also be rational [x Def 4]

and if the rectangle AE , EB is medial the rectangle CF FD is also medial [x 23 Por]

Therefore CD is an apotome of a medial straight line and the same in order with AB [x 74 75]

Q E D


PROPOSITION 105

A straight line commensurable with a minor straight line is minor

Let AB be a minor straight line and CD commensurable with AB ,

I say that CD is also minor

Let the same construction be made as before,
then since AE EB are incommensurable in square



[x 76]

therefore CF FD are also incommensurable in square [x 13]

Now since as AE is to EB so is CF to FD [v 12 v 16]

therefore also as the square on AE is to the square on EB so is the square on CF to the square on FD [vi 27]

Therefore *componendo* as the squares on AE EB are to the square on EB so are the squares on CF FD to the square on FD [v 18]

But the square on BE is commensurable with the square on DF
therefore the sum of the squares on AE , EB is also commensurable with the sum of the squares on CF FD [v 16 x 11]

But the sum of the squares on AE EB is rational, [x 6]

therefore the sum of the squares on CF FD is also rational [x Def 4]

Again since as the square on AE is to the rectangle AE , EB so is the square on CF to the rectangle CF FD

while the square on AE is commensurable with the square on CF
therefore the rectangle AE EB is also commensurable with the rectangle CF , FD

But the rectangle AE EB is medial [x 76]

therefore the rectangle CF FD is also medial, [x 23 Por 1]
therefore CF FD are straight lines incommensurable in square which make the sum of the squares on them rational but the rectangle contained by them medial

Therefore CD is minor [x 6]


Q E D

PROPOSITION 106

A straight line commensurable with that which produces with a rational area a medial whole is a straight line which produces with a rational area a medial whole

Let AB be a straight line which produces with a rational area a medial whole
and CD commensurable with AB

I say that CD is also a straight line which produces with a rational area a medial whole



For let BE be the annex to AB

therefore AL LB are straight lines incommensurable in square which make the sum of the squares on AL FB medial but the rectangle contained by them rational [x 17]

Let the same construction be made

Then we can prove in manner similar to the foregoing that CF FD are in the same ratio as AF EB

the sum of the squares on AE EB is commensurable with the sum of the squares on CF FD

and the rectangle AE EB with the rectangle CF FD

so that CF , FD are also straight lines incommensurable in square which make the sum of the squares on CF , FD medial but the rectangle contained by them rational

Therefore CD is a straight line which produces with a rational area a medial whole

[x 77]

Q E D

PROPOSITION 107

A straight line commensurable with that which produces with a medial area a medial whole is itself also a straight line which produces with a medial area a medial whole

Let AB be a straight line which produces with a medial area a medial whole and let CD be commensurable with AB

I say that CD is also a straight line which produces with a medial area a medial whole

For let BE be the annex to AB , and let the same construction be made

therefore AE , EB are straight lines incommensurable in square which make the sum of the squares on them medial the rectangle contained by them medial and further the sum of the squares on them incommensurable with the rectangle contained by them

[x 78]

Now, as was proved, AE , EB are commensurable with CF , FD , the sum of the squares on AE , EB with the sum of the squares on CF , FD , and the rectangle AE , EB with the rectangle CF , FD ,

therefore CF , FD are also straight lines incommensurable in square which make the sum of the squares on them medial the rectangle contained by them medial and further, the sum of the squares on them incommensurable with the rectangle contained by them

Therefore CD is a straight line which produces with a medial area a medial whole


[x 78]

Q E D

PROPOSITION 108

If from a rational area a medial area be subtracted the 'side' of the remaining area becomes one of two irrational straight lines either an apotome or a minor straight line

For from the rational area BC let the medial area BD be subtracted




I say that the side of the remainder EC becomes one of two irrational straight lines either an apotome or a minor straight line

For let a rational straight line FG be set out

to FG let there be applied the rectangular parallelogram GH equal to BC

and let GK equal to DB be subtracted therefore the remainder EC is equal to LH

Since then BC is rational and BD medial



while BC is equal to GH , and BD to GK ,
therefore GH is rational and GK medial

And they are applied to the rational straight line FG ,
therefore FH is rational and commensurable in length with FG , [x 70]
while FK is rational and incommensurable in length with FG , [x 27]
therefore FH is incommensurable in length with FK [x 13]

Therefore FH , FK are rational straight lines commensurable in square only,
therefore KH is an apotome [x 73] and AK the annex to it

Now the square on HF is greater than the square on FK by the square on a straight line either commensurable with HF or not commensurable

First let the square on it be greater by the square on a straight line commensurable with it

Now the whole HF is commensurable in length with the rational straight line FG set out,

therefore AH is a first apotome [x Def III 1]

But the side of the rectangle contained by a rational straight line and a first apotome is an apotome [x 91]

Therefore the side of LH that is of EC is an apotome

But if the square on HF is greater than the square on FK by the square on a straight line incommensurable with HF
while the whole FH is commensurable in length with the rational straight line FG set out

AH is a fourth apotome [x Def III 4]

But the side of the rectangle contained by a rational straight line and a fourth apotome is minor [x 94]

Q E D

PROPOSITION 109

If from a medial area a rational area be subtracted there arise two other irrational straight lines either a first apotome of a medial straight line or a straight line which produces with a rational area a medial whole


For from the medial area BC let the rational area BD be subtracted

I say that the side of the remainder EC becomes one of two irrational straight lines either a first apotome of a medial straight line or a straight line which produces with a rational area a medial whole

For let a rational straight line FG be set out

and let the areas be similarly applied

It follows then that FH is rational and incommensurable in length with FG
while FK is rational and commensurable in length with FG
therefore FH , FK are rational straight lines commensurable in square only [x 13]



therefore AH is an apotome and AK the annex to it [x 73]

Now the square on HF is greater than the square on FK either by the square on a straight line commensurable with HF or by the square on a straight line incommensurable with it

If then the square on HF is greater than the square on FA by the square on a straight line commensurable with HF , while the annex FA is commensurable in length with the rational straight line FG set out

KH is a second apotome

[x Def III 2]

But FG is rational, so that the 'side' of LH , that is of EC , is a first apotome of a medial straight line

[x 92]

But, if the square on HF is greater than the square on FA by the square on a straight line incommensurable with HF , while the annex FA is commensurable in length with the rational straight line FG set out,

LH is a fifth apotome,

[x Def III 5]

so that the 'side' of EC is a straight line which produces with a rational area a medial whole


[x 93]

Q E D

PROPOSITION 110

If from a medial area there be subtracted a medial area incommensurable with the whole the two remaining irrational straight lines arise, either a second apotome of a medial straight line or a straight line which produces with a medial area a medial whole

For as in the foregoing figures let there be subtracted from the medial area BC the medial area BD incommensurable with the whole



I say that the 'side' of EC is one of two irrational straight lines either a second apotome of a medial straight line or a straight line which produces with a medial area a medial whole

For since each of the rectangles BC , BD is medial

and BC is incommensurable with BD

it follows that each of the straight lines FH , FA will be rational and uncommensurable

[x 22]

surable in length with FG

And since BC is incommensurable with BD

that is GH with GK

HF is also incommensurable with FK

[vi 1 x 11]

therefore FH , FA are rational straight lines commensurable in square only

therefore LH is an apotome

[x 73]

If then the square on FH is greater than the square on FK by the square on a straight line commensurable with FH

while neither of the straight lines FH , FA is commensurable in length with the rational straight line FG set out

LH is a third apotome

[x Def III 3]

But AL is rational

and the rectangle contained by a rational straight line and a third apotome is irrational,

and the "side" of it is irrational and is called a second apotome of a medial straight line, [x 93]
 so that the 'side' of LH , that is, of EC , is a second apotome of a medial straight line

But if the square on FH is greater than the square on FK by the square on a straight line incommensurable with FH
 while neither of the straight lines HF FK is commensurable in length with FG

KH is a sixth apotome [x Def iii 6]

But the "side" of the rectangle contained by a rational straight line and a sixth apotome is a straight line which produces with a medial area a medial whole [x 96]

Therefore the "side" of LH that is of EC is a straight line which produces with a medial area a medial whole Q E D

PROPOSITION 111

The apotome is not the same with the binomial straight line

Let AB be an apotome

I say that AB is not the same with the binomial straight line

For if possible let it be so,
 let a rational straight line DC be set out and to CD
 let there be applied the rectangle CE equal to the
 square on AB and producing DE as breadth

Then since AB is an apotome

DE is a first apotome [x 97]

Let EF be the annex to it

therefore DF FE are rational straight lines commensurable in square only

the square on DF is greater than the square on FE by
 the square on a straight line commensurable with DF

and DF is commensurable in length with the rational straight line DC set out [x Def iii 1]

Again since AB is binomial

therefore DE is a first binomial straight line [x 60]

Let it be divided into its terms at G

and let DG be the greater term

therefore DG GE are rational straight lines commensurable in square only
 the square on DG is greater than the square on GE by the square on a straight
 line commensurable with DG and the greater term DG is commensurable in
 length with the rational straight line DC set out [x Def iii 1]

Therefore DF is also commensurable in length with DG [x 12]

therefore the remainder GF is also commensurable in length with DF [x 15]


But DF is incommensurable in length with EF

therefore FG is also incommensurable in length with EF [x 13]

Therefore GF IF are rational straight lines commensurable in square only
 therefore EG is an apotome [x 73]

But it is also rational

which is impossible



Therefore the apotome is not the same with the binomial straight line

Q E D

The apotome and the irrational straight lines following it are neither the same with the medial straight line nor with one another

For the square on a medial straight line if applied to a rational straight line produces as breadth a straight line rational and incommensurable in length with that to which it is applied [x 22]

while the square on an apotome, if applied to a rational straight line produces as breadth a first apotome, [x 97]

the square on a first apotome of a medial straight line if applied to a rational straight line produces as breadth a second apotome [x 98]

the square on a second apotome of a medial straight line if applied to a rational straight line produces as breadth a third apotome, [x 99]

the square on a minor straight line if applied to a rational straight line, produces as breadth a fourth apotome [x 100]

the square on the straight line which produces with a rational area a medial whole if applied to a rational straight line produces as breadth a fifth apotome [x 101]

and the square on the straight line which produces with a medial area a medial whole, if applied to a rational straight line produces as breadth a sixth apotome [x 102]

Since then the said breadths differ from the first and from one another from the first because it is rational, and from one another since they are not the same in order

it is clear that the irrational straight lines themselves also differ from one another

And, since the apotome has been proved not to be the same as the binomial straight line [x 111]

but if applied to a rational straight line the straight lines following the apotome produce as breadths each according to its own order, apotomes and those following the binomial straight line themselves also according to their order produce the binomials as breadths

therefore those following the apotome are different and those following the binomial straight line are different so that there are, in order, thirteen irrational straight lines in all

Medial

Binomial

First bimedial

Second bimedial

Major

Side of a rational plus a medial area

Side of the sum of two medial areas

Apotome

First apotome of a medial straight line

Second apotome of a medial straight line

Minor

Producing with a rational area a medial whole

Producing with a medial area a medial whole


PROPOSITION 112

The square on a rational straight line applied to the binomial straight line produces as breadth an apotome the terms of which are commensurable with the terms of the binomial and moreover in the same ratio and further the apotome so arising will have the same order as the binomial straight line

Let A be a rational straight line

let BC be a binomial and let DC be its greater term,

let the rectangle BC , EF be equal to the square on A ,



I say that EF is an apotome the terms of which are commensurable with CD , DB and in the same ratio and further EF will have the same order as BC

For again let the rectangle BD , G be equal to the square on A

Since then the rectangle BC , EF is equal to the rectangle BD , G

therefore as CB is to BD so is G to EF [vi 16]

But CB is greater than BD

therefore G is also greater than EF [v 16 v 14]

Let EH be equal to G

therefore as CB is to BD so is HE to EF

therefore *separando* as CD is to BD so is HF to FE [v 17]

Let it be contrived that as HF is to FE so is FA to AE

therefore also the whole HA is to the whole KE as FA is to AE

for as one of the antecedents is to one of the consequents so are all the antecedents to all the consequents [v 12]

But as FA is to AE so is CD to DB [v 11]

therefore also as HA is to KE so is CD to DB [id]

But the square on CD is commensurable with the square on DB [x 36]

therefore the square on HA is also commensurable with the square on AE [x 22 x 11]

And as the square on HA is to the square on AE so is HA to AE , since the three straight lines HA , AF , AE are proportional [v Def 9]

Therefore HA is commensurable in length with AE

so that HE is also commensurable in length with EA [x 13]

Now since the square on 1 is equal to the rectangle EH , BD

while the square on A is rational

therefore the rectangle EH , BD is also rational

And it is applied to the rational straight line BD

therefore EH is rational and commensurable in length with BD [x 20]
so that EA being commensurable with it is also rational and commensurable in length with BD

Since then as CD is to DB so is FA to AE

while CD , DB are straight lines commensurable in square only,

therefore FA , AE are also commensurable in square only [x 11]

But AE is rational,

therefore FA is also rational

Therefore FA KE are rational straight lines commensurable in square only
therefore EF is an apotome [x 73]

Now the square on CD is greater than the square on DB either by the square on a straight line commensurable with CD or by the square on a straight line incommensurable with it

If then the square on CD is greater than the square on DB by the square on a straight line commensurable with CD the square on FA is also greater than the square on KE by the square on a straight line commensurable with FA

[x 14]

And if CD is commensurable in length with the rational straight line set out,
so also is FK [x 11 12]

if BD is so commensurable

so also is KE

[x 12]

but, if neither of the straight lines CD DB is so commensurable
neither of the straight lines FK KE is so

But if the square on CD is greater than the square on DB by the square on a straight line incommensurable with CD

the square on FA is also greater than the square on KE by the square on a straight line incommensurable with FK

[x 14]

And, if CD is commensurable with the rational straight line set out,
so also is FA ,

if BD is so commensurable,

so also is KE ,

but, if neither of the straight lines CD DB is so commensurable,
neither of the straight lines FK KE is so

so that FE is an apotome the terms of which FK KE are commensurable with the terms CD DB of the binomial straight line and in the same ratio and it has the same order as BC

Q E D

PROPOSITION 113

The square on a rational straight line if applied to an apotome produces as breadth the binomial straight line the terms of which are commensurable with the terms of the apotome and in the same ratio and further the binomial so arising has the same order as the apotome

Let A be a rational straight line and BD an apotome and let the rectangle BD KH be equal to the square on A so that the square on the rational straight line A when applied to the apotome BD produces KH as breadth


I say that AH is a binomial straight line the terms of which are commensurable with the terms of BD and in the same ratio and further AH has the same order as BD

For let DC be the annex to BD

therefore BC CD are rational straight lines commensurable in square only [x 73]

Let the rectangle BC G be also equal to the square on A

But the square on A is rational



therefore the rectangle BC, G is also rational

And it has been applied to the rational straight line BC

therefore G is rational and commensurable in length with BC [x 20]

Since now the rectangle BC, G is equal to the rectangle BD, KH ,

therefore, proportionally, as CB is to BD , so is KH to G [vi 16]

But BC is greater than BD ,

therefore KH is also greater than G [v 16 v 14]

Let KE be made equal to G

therefore KE is commensurable in length with BC

And since as CB is to BD so is KA to KE

therefore *convertendo* as BC is to CD so is KH to HE [v 19 Por]

Let it be contrived that, as KH is to HE so is HF to FE

therefore also the remainder KF is to FH as KH is to HE that is as BC is to CD [v 19]

But BC, CD are commensurable in square only,

therefore KF, FH are also commensurable in square only [x 11]

And since as KH is to HE so is KF to FH

while as KH is to HE so is HF to FE

therefore also as KF is to FH so is HF to FE [v 11]

so that also as the first is to the third so is the square on the first to the square on the second [v Def 9]

therefore also as KF is to FE so is the square on KF to the square on FH

But the square on KF is commensurable with the square on FH

for KF, FH are commensurable in square,

therefore KF is also commensurable in length with FE , [x 11]

so that KF is also commensurable in length with KE [x 13]

But KE is rational and commensurable in length with BC ,

therefore KF is also rational and commensurable in length with BC [x 12]

And since as BC is to CD so is KF to FH

alternately as BC is to KF so is DC to FH [v 16]

But BC is commensurable with KF

therefore FH is also commensurable in length with CD [x 11]

But BC, CD are rational straight lines commensurable in square only

therefore KF, FH are also rational straight lines [x Def 3] commensurable in square only

therefore KH is binomial [x 36]

If now the square on BC is greater than the square on CD by the square on a straight line commensurable with BC

the square on KF will also be greater than the square on FH by the square on a straight line commensurable with KF [x 14]

And, if BC is commensurable in length with the rational straight line set out so also is KF

if CD is commensurable in length with the rational straight line set out,

so also is FH

but if neither of the straight lines BC, CD

then neither of the straight lines KF, FH

But, if the square on BC is greater than the square on CD by the square on a straight line incommensurable with BC

the square on KF is also greater than the square on FH by the square on a

straight line incommensurable with KF

[τ 14]

And if BC is commensurable with the rational straight line set out

so also is ΔF

if CD is so commensurable

so also is ΓH

but if neither of the straight lines BC , CD ,


then neither of the straight lines ΔF , ΓH

Therefore KH is a binomial straight line the terms of which ΔF , ΓH are commensurable with the terms BC , CD of the apotome and in the same ratio and further ΔH has the same order as BD Q E D

PROPOSITION 114

If an area be contained by an apotome and the binomial straight line the terms of which are commensurable with the terms of the apotome and in the same ratio the side of the area is rational

For let an area the rectangle AB , CD be contained by the apotome AB and the binomial straight line CD



and let CE be the greater term of the latter let the terms CE , ED of the binomial straight line be commensurable with the terms AF , FB of the apotome and in the same ratio

and let the side of the rectangle AB , CD be G

I say that G is rational

For let a rational straight line H be set out and to CD let there be applied a rectangle equal to the square on H and producing KL as breadth

Therefore KL is an apotome

Let its terms be KM , ML commensurable with the terms CE , ED of the binomial straight line and in the same ratio [x 112]

But CE , ED are also commensurable with AF , FB and in the same ratio therefore as AF is to FB so is KM to ML

Therefore alternately as AF is to KM so is BF to LM

therefore also the remainder AB is to the remainder KL as AF is to KM [γ 19]

But AF is commensurable with KM [τ 12]

therefore AB is also commensurable with KL [x 11]

And as AB is to KL so is the rectangle CD , AB to the rectangle CD , KL [vi 1]

therefore the rectangle CD , AB is also commensurable with the rectangle CD , KL [τ 11]

But the rectangle CD , KL is equal to the square on H

therefore the rectangle CD , AB is commensurable with the square on H

But the square on G is equal to the rectangle CD , AB

therefore the square on G is commensurable with the square on H

But the square on H is rational

therefore the square on G is also rational

therefore G is rational

And it is the side of the rectangle CD , AB

Therefore etc

POINSETT And it is made manifest to us by this also that it is possible for a rational area to be contained by irrational straight lines : Q E D


PROPOSITION 115

From a medial straight line there arise irrational straight lines infinite in number, and none of them is the same as any of the preceding

Let A be a medial straight line

I say that from A there arise irrational straight lines infinite in number, and none of them is the same as any of the preceding

Let a rational straight line B be set out and let the square on C be equal to the rectangle $B A$,



therefore C is irrational [x Def 4]

for that which is contained by an irrational and a rational straight line is irrational [deduction from x 20]

And it is not the same with any of the preceding for the square on none of the preceding if applied to a rational straight line produces as breadth a medial straight line

Again let the square on D be equal to the rectangle B, C , therefore the square on D is irrational [deduction from x 20]

Therefore D is irrational [x Def 4] and it is not the same with any of the preceding for the square on none of the preceding if applied to a rational straight line produces C as breadth

Similarly if this arrangement proceeds *ad infinitum*, it is manifest that from the medial straight line there arise irrational straight lines infinite in number and none is the same with any of the preceding Q E D

BOOK ELEVEN

DEFINITIONS

- 1 A *solid* is that which has length breadth and depth
- 2 An extremity of a solid is a surface
- 3 A *straight line* is at right angles to a plane when it makes right angles with all the straight lines which meet it and are in the plane
- 4 A plane is at right angles to a plane when the straight lines drawn in one of the planes at right angles to the common section of the planes are at right angles to the remaining plane
- 5 The inclination of a straight line to a plane is assuming a perpendicular drawn from the extremity of the straight line which is elevated above the plane to the plane and a straight line joined from the point thus arising to the extremity of the straight line which is in the plane the angle contained by the straight line so drawn and the straight line standing up
- 6 The inclination of a plane to a plane is the acute angle contained by the straight lines drawn at right angles to the common section at the same point one in each of the planes
- 7 A plane is said to be *similarly inclined* to a plane as another is to another when the said angles of the inclinations are equal to one another
- 8 *Parallel planes* are those which do not meet
- 9 *Similar solid figures* are those contained by similar planes equal in multitude
- 10 *Equal and similar solid figures* are those contained by similar planes equal in multitude and in magnitude
- 11 A *solid angle* is the inclination constituted by more than two lines which meet one another and are not in the same surface towards all the lines
Otherwise A *solid angle* is that which is contained by more than two plane angles which are not in the same plane and are constructed to one point
- 12 A *pyramid* is a solid figure contained by planes which is constructed from one plane to one point
- 13 A *prism* is a solid figure contained by planes two of which namely those which are opposite are equal similar and parallel while the rest are parallelograms
- 14 When the diameter of a semicircle remaining fixed the semicircle is carried round and restored again to the same position from which it began to be moved the figure so comprehended is a *sphere*
- 15 The *axis of the sphere* is the straight line which remains fixed and about which the semicircle is turned
- 16 The *centre of the sphere* is the same as that of the semicircle
- 17 A *diameter of the sphere* is any straight line drawn through the centre and

terminated in both directions by the surface of the sphere

18 When one side of those about the right angle in a right angled triangle remaining fixed the triangle is carried round and restored again to the same position from which it began to be moved the figure so comprehended is a *cone*

And if the straight line which remains fixed be equal to the remaining side about the right angle which is carried round, the cone will be *right-angled*, if less *obtuse angled*, and if greater *acute-angled*

19 The *axis of the cone* is the straight line which remains fixed and about which the triangle is turned

20 And the *base* is the circle described by the straight line which is carried round

21 When one side of those about the right angle in a rectangular parallelogram remaining fixed, the parallelogram is carried round and restored again to the same position from which it began to be moved the figure so comprehended is a *cylinder*

22 The *axis of the cylinder* is the straight line which remains fixed and about which the parallelogram is turned

23 And the *bases* are the circles described by the two sides opposite to one another which are carried round

24 *Similar cones and cylinders* are those in which the axes and the diameters of the bases are proportional

25 A *cube* is a solid figure contained by six equal squares

26 An *octahedron* is a solid figure contained by eight equal and equilateral triangles

27 An *icosahedron* is a solid figure contained by twenty equal and equilateral triangles

28 A *dodecahedron* is a solid figure contained by twelve equal equilateral and equiangular pentagons

BOOK VI PROPOSITIONS

PROPOSITION I

A part of a straight line cannot be in the plane of reference and a part in a plane more elevated

I or if possible let a part *AB* of the straight line *ABC* be in the plane of reference and a part *BC* in a plane more elevated


There will then be in the plane of reference some straight line continuous with *AB* in a straight line

Let it be *BD*

therefore *AB* is a common segment of the two straight lines *ABC ABD*

which is impossible inasmuch as if we describe a circle with centre *B* and distance *AB* the diameters will cut off unequal circumferences of the circle

Therefore a part of a straight line cannot be in the plane of reference and a part in a plane more elevated



Q E D

PROPOSITION 2

If two straight lines cut one another, they are in one plane and every triangle is in one plane

For let the two straight lines AB , CD cut one another at the point E ,
I say that AB , CD are in one plane and every triangle is in one plane

For let points F , G be taken at random on EC , EB ,

let CB , FG be joined

and let FH , GK be drawn across

I say first that the triangle ECB is in one plane

For if part of the triangle ECB either FHC or GBK is in the plane of reference and the rest in another

a part also of one of the straight lines EC , EB will be in the plane of reference and a part in another

But if the part $FCBG$ of the triangle ECB be in the plane of reference and the rest in another

a part also of both the straight lines EC , EB will be in the plane of reference and a part in another

which was proved absurd [XI 1]

Therefore the triangle ECB is in one plane

But in whatever plane the triangle ECB is in that plane also is each of the straight lines EC , EB

and in whatever plane each of the straight lines EC , EB is in that plane are AB , CD also [XI 1]

Therefore the straight lines AB , CD are in one plane

and every triangle is in one plane

Q E D

PROPOSITION 3

If two planes cut one another their common section is a straight line

For let the two planes AB , BC cut one another

and let the line DB be their common section

I say that the line DB is a straight line

For if not from D to B let the straight line DEB be joined in the plane AB ,

and in the plane BC the straight line DFB

Then the two straight lines DEB , DFB will have the same extremities and will clearly enclose an area

which is absurd

Therefore DEB , DFB are not straight lines



Similarly we can prove that neither will there be any other straight line joined from D to B except DB the common section of the planes AB , BC

Therefore etc

Q E D

PROPOSITION 4

If a straight line be set up at right angles to two straight lines which cut one another, at their common point of section it will also be at right angles to the plane through them



For let a straight line EF be set up at right angles to the two straight lines AB , CD , which cut one another at the point E , from E ,


I say that EF is also at right angles to the plane through AB , CD

For let AE , EB , CE , ED be cut off equal to one another,

and let any straight line GEH be drawn across through E at random,

let AD , CB be joined,

and further, let FA , FG , FD , FC , FH , FB be joined from the point F taken at random < on EF >



Now since the two straight lines AE , ED are equal to the two straight lines CE , EB , and contain equal angles [1 15]

therefore the base AD is equal to the base CB ,

and the triangle AED will be equal to the triangle CEB , [1 4]

so that the angle DAE is also equal to the angle EBC

But the angle AGE is also equal to the angle BEH [1 15]

therefore AGE , BEH are two triangles which have two angles equal to two angles respectively and one side equal to one side namely that adjacent to the equal angles that is to say AE to EB ,

therefore they will also have the remaining sides equal to the remaining sides [1 26]

Therefore GE is equal to EH and AG to BH

And since AE is equal to EB

while FE is common and at right angles

therefore the base FA is equal to the base FB [1 4]

For the same reason

FC is also equal to FD

And since AD is equal to CB

and FA is also equal to FB

the two sides FA , AD are equal to the two sides FB , BC respectively,

and the base FD was proved equal to the base FC

therefore the angle FAD is also equal to the angle FBC [1 8]

And since again AG was proved equal to BH

and further FA also equal to FB

the two sides FA , AG are equal to the two sides FB , BH

And the angle FAG was proved equal to the angle FBH ,

therefore the base FG is equal to the base FH [1 4]

Now since again GE was proved equal to EH

and EF is common

the two sides GE , EF are equal to the two sides HE , EF ,

and the base FG is equal to the base FH

therefore the angle GFE is equal to the angle HEF [1 8]

Therefore each of the angles CEF , HFE is right

Therefore FE is at right angles to GH drawn at random through E

Similarly we can prove that FE will also make right angles with all the straight lines which meet it and are in the plane of reference

But a straight line is at right angles to a plane when it makes right angles

with all the straight lines which meet it and are in that same plane, [XI Def 3]
 therefore FE is at right angles to the plane of reference
 But the plane of reference is the plane through the straight lines AB CD
 Therefore FE is at right angles to the plane through AB , CD
 Therefore etc Q E D

PROPOSITION 5

If a straight line be set up at right angles to three straight lines which meet one another, at their common point of section the three straight lines are in one plane

For let a straight line AB be set up at right angles to the three straight lines BC , BD , BE , at their point of meeting at B ,

I say that BC BD , BE are in one plane

For suppose they are not but, if possible, let BD , BE be in the plane of reference and BC in one more elevated

let the plane through AB , BC be produced,
 it will thus make a common section in the plane of reference, a straight line [XI 3]

Let it make BF

Therefore the three straight lines AB , BC , BF are in one plane, namely that drawn through AB BC

Now, since AB is at right angles to each of the straight lines BD , BE
 therefore AB is also at right angles to the plane through BD , BE [XI 4]

But the plane through BD , BE is the plane of reference,
 therefore AB is at right angles to the plane of reference

Thus AB will also make right angles with all the straight lines which meet it and are in the plane of reference [XI Def 3]

But BF which is in the plane of reference meets it,
 therefore the angle ABF is right

But by hypothesis the angle ABC is also right,
 therefore the angle ABF is equal to the angle ABC

And they are in one plane
 which is impossible

Therefore the straight line BC is not in a more elevated plane,
 therefore the three straight lines BC , BD BE are in one plane

Therefore if a straight line be set up at right angles to three straight lines,
 at their point of meeting, the three straight lines are in one plane Q E D

PROPOSITION 6

If two straight lines be at right angles to the same plane the straight lines will be parallel

For let the two straight lines AB CD be at right angles to the plane of reference

I say that AB is parallel to CD

For let them meet the plane of reference at the points B , D

let the straight line BD be joined

let DE be drawn in the plane of reference at right angles to BD ,

let DE be made equal to AB

and let BE , AE , AD be joined

Now, since AB is at right angles to the plane of reference it will also make right angles with all the straight lines which meet it and are in the plane of reference [XI Def 3]

But each of the straight lines BD , BE is in the plane of reference and meets AB ,

therefore each of the angles ABD ABE is right

For the same reason

each of the angles CDB , CDE is also right

And since AB is equal to DE

and BD is common,

the two sides AB , BD are equal to the two sides

ED , DB ,

and they include right angles,

therefore the base AD is equal to the base BE [I 4]

And since AB is equal to DE ,

while AD is also equal to BE ,

the two sides AB , BE are equal to the two sides ED , DA ,

and AE is their common base,

therefore the angle ABE is equal to the angle EDA [I 8]

But the angle ABE is right

therefore the angle EDA is also right,

therefore ED is at right angles to DA

But it is also at right angles to each of the straight lines BD , DC , therefore ED is set up at right angles to the three straight lines BD DA DC at their point of meeting

therefore the three straight lines BD DA , DC are in one plane [XI 5]

But, in whatever plane DB DA are in that plane is AB also

for every triangle is in one plane, [XI 2]

therefore the straight lines AB BD , DC are in one plane

And each of the angles ABD BDC is right

therefore AB is parallel to CD [I 18]

Therefore etc

Q E D

PROPOSITION 7

If two straight lines be parallel and points be taken at random on each of them the straight line joining the points is in the same plane with the parallel straight lines

Let AB CD be two parallel straight lines

and let points E , F be taken at random on them respectively

I say that the straight line joining the points E , F is in the same plane with the parallel straight lines

For suppose it is not, but if possible let it be in a more elevated plane as EGF



and let a plane be drawn through LGF

it will then make as section in the plane of reference a straight line [XI 3]

Let it make it as EF ,

therefore the two straight lines EGF EF will enclose an area

which is impossible



Therefore the straight line joined from E to F is not in a plane more elevated, therefore the straight line joined from E to F is in the plane through the parallel straight lines AB , CD

Therefore etc

Q E D

PROPOSITION 8

If two straight lines be parallel and one of them be at right angles to any plane, the remaining one will also be at right angles to the same plane

Let AB , CD be two parallel straight lines,

and let one of them, AB , be at right angles to the plane of reference,

I say that the remaining one, CD , will also be at right angles to the same plane

For let AB , CD meet the plane of reference at the points B , D ,

and let BD be joined,

therefore AB , CD , BD are in one plane [XI 7]

Let DE be drawn in the plane of reference, at right angles to BD ,

let DE be made equal to AB

and let BE , AE , AD be joined

Now since AB is at right angles to the plane of reference therefore AB is also at right angles to all the straight lines which meet it and are in the plane of reference [XI Def 3]

therefore each of the angles ABD , ABE is right

And since the straight line BD has fallen on the parallels AB , CD ,

therefore the angles ABD , CDB are equal to two right angles [I 29]

But the angle ABD is right

therefore the angle CDB is also right

therefore CD is at right angles to BD

And since AB is equal to DE

and BD is common

the two sides AB , BD are equal to the two sides ED , DB ,

and the angle ABD is equal to the angle EDB

for each is right

therefore the base AD is equal to the base BE

And, since AB is equal to DE ,

and BE to AD ,

the two sides AB , BE are equal to the two sides ED , DA respectively,

and AE is their common base,

therefore the angle ABE is equal to the angle EDA

But the angle ABE is right

therefore the angle EDA is also right

therefore ED is at right angles to AD

But it is also at right angles to DB

therefore ED is also at right angles to the plane through BD , DA [XI 4]

Therefore ED will also make right angles with all the straight lines which meet it and are in the plane through BD , DA

But DC is in the plane through BD , DA , inasmuch as AB , BD are in the

plane through $BD, DA,$

[xi 7]

and DC is also in the plane in which AB, BD are

Therefore ED is at right angles to $DC,$

so that CD is also at right angles to DE

But CD is also at right angles to BD

Therefore CD is set up at right angles to the two straight lines DE, DB which cut one another from the point of section at $D,$

so that CD is also at right angles to the plane through DE, DB [xi 4]

But the plane through DE, DB is the plane of reference,

therefore CD is at right angles to the plane of reference

Therefore etc

Q E D

PROPOSITION 9

Straight lines which are parallel to the same straight line and are not in the same plane with it are also parallel to one another

For let each of the straight lines AB, CD be parallel to EF , not being in the same plane with it

I say that AB is parallel to CD

For let a point G be taken at random on EF

and from it let there be drawn GH in the plane through EF, AB , at right angles to EF and GK in the plane through FE, CD again at right angles to EF

Now since EF is at right angles to each of the straight lines GH, GK

therefore EF is also at right angles to the plane through GH, GK [xi 4]

And EF is parallel to AB

therefore AB is also at right angles to the plane through HG, GK [xi 6]

For the same reason

CD is also at right angles to the plane through HG, GK , therefore each of the straight lines AB, CD is at right angles to the plane through HG, GK

But if two straight lines be at right angles to the same plane the straight lines are parallel [xi 6]

therefore AB is parallel to CD

Q E D

PROPOSITION 10

If two straight lines meeting one another be parallel to two straight lines meeting one another not in the same plane they will contain equal angles

For let the two straight lines AB, BC meeting one another be parallel to the two straight lines DE, EF meeting one another not in the same plane,

I say that the angle ABC is equal to the angle DEF

For let BA, BC, ED, EF be cut off equal to one another and let AD, CF, BE, AC, DF be joined

Now since BA is equal and parallel to ED

therefore AD is also equal and parallel to BE

[i 33]

For the same reason

CF is also equal and parallel to BE

Therefore each of the straight lines AD , CF is equal and parallel to BE

But straight lines which are parallel to the same straight line and are not in the same plane with it are parallel to one another, [xi 9]

therefore AD is parallel and equal to CF

And AC , DF join them,
therefore AC is also equal and parallel to DF [i 33]

Now, since the two sides AB , BC are equal to the two sides DE , EF ,

and the base AC is equal to the base DF
therefore the angle ABC is equal to the angle DEF [i 8]

Therefore etc

Q E D

PROPOSITION 11

From a given elevated point to draw a straight line perpendicular to a given plane

Let A be the given elevated point and the plane of reference the given plane, thus it is required to draw from the point A a straight line perpendicular to the plane of reference

Let any straight line BC be drawn at random in the plane of reference

and let AD be drawn from the point A
perpendicular to BC [i 12]

If then AD is also perpendicular to the plane of reference that which was enjoined will have been done

But if not let DE be drawn from the point D at right angles to BC and in the plane of reference, [i 11]

let AF be drawn from A perpendicular to DE [i 12]

and let GH be drawn through the point F parallel to BC [i 31]

Now since BC is at right angles to each of the straight lines DA , DE ,
therefore BC is also at right angles to the plane through ED , DA [xi 4]

And GH is parallel to it,
but if two straight lines be parallel and one of them be at right angles to any plane the remaining one will also be at right angles to the same plane [vi 8]
therefore GH is also at right angles to the plane through ED , DA

Therefore GH is also at right angles to all the straight lines which meet it and are in the plane through ED , DA [xi Def 3]

But AF meets it and is in the plane through ED , DA

therefore GH is at right angles to FA ,
so that FA is also at right angles to GH

But AF is also at right angles to DE

therefore AF is at right angles to each of the straight lines GH , DE

But if a straight line be set up at right angles to two straight lines which cut one another at the point of section it will also be at right angles to the plane through them [xi 4]

therefore FA is at right angles to the plane through ED , GH

But the plane through ED , GH is the plane of reference

therefore AF is at right angles to the plane of reference

Therefore from the given elevated point A the straight line AF has been drawn perpendicular to the plane of reference Q E F

PROPOSITION 12

To set up a straight line at right angles to a given plane from a given point in it

Let the plane of reference be the given plane,

and A the point in it,

thus it is required to set up from the point A a straight line at right angles to the plane of reference


Let any elevated point B be conceived

from B let BC be drawn perpendicular to

the plane of reference [XI 11]

and through the point A let AD be drawn

parallel to BC [I 31]



Then since AD CB are two parallel straight lines

while one of them BC is at right angles to the plane of reference

therefore the remaining one AD is also at right angles to the plane of reference [XI 8]

Therefore AD has been set up at right angles to the given plane from the point A in it Q E F

PROPOSITION 13

From the same point two straight lines cannot be set up at right angles to the same plane on the same side

For if possible from the same point A let the two straight lines AB , AC be set up at right angles to the plane of reference and on the same side

and let a plane be drawn through BA AC

it will then make as section through A in the plane of reference a straight line [XI 3]

Let it make DAE

therefore the straight lines AB AC ,

DAE are in one plane

And since CA is at right angles to the plane of reference it will also make right angles with all the straight lines which meet it and are in the plane of reference [XI Def 3]

But DAE meets it and is in the plane of reference

therefore the angle CAE is right

For the same reason

the angle BAE is also right

therefore the angle CAE is equal to the angle BAE

And they are in one plane

which is impossible

Therefore etc

Q E D


PROPOSITION 14

Planes to which the same straight line is at right angles will be parallel

For let any straight line AB be at right angles to each of the planes CD , EF ,

I say that the planes are parallel

For, if not they will meet when produced



Let them meet,
they will then make as common section,
a straight line [xi 3]

Let them make GH ,
let a point K be taken at random on GH ,
and let AK , BK be joined

Now, since AB is at right angles to
the plane EF

therefore AB is also at right angles to BK which is a straight line in the plane
 EF produced, [xi Def 3]

therefore the angle ABK is right

For the same reason

the angle BAK is also right

Thus in the triangle ABK the two angles ABK BAK are equal to two
right angles

which is impossible [i 17]

Therefore the planes CD EF will not meet when produced


therefore the planes CD EF are parallel [xi Def 8]

Therefore planes to which the same straight line is at right angles are par-
allel Q E D

PROPOSITION 15

If two straight lines meeting one another be parallel to two straight lines meeting
one another not being in the same plane, the planes through them are parallel

For let the two straight lines AB , BC meeting one another be parallel to the
two straight lines DE EF meeting one another
not being in the same plane, I say that the
planes produced through AB BC and DE , EF
will not meet one another



For let BG be drawn from the point B per-
pendicular to the plane through DE EF [xi 11],

and let it meet the plane at the point G ,
through G let GH be drawn parallel to ED and
 GK parallel to EF [i 31]

Now since BG is at right angles to the plane
through DE EF

therefore it will also make right angles with all the straight lines which meet it
and are in the plane through DE EF [xi Def 3]

But each of the straight lines GH GK meets it and is in the plane through
 DE EF

therefore each of the angles BGH , BGK is right

And since BA is parallel to GH , [xi 9]

therefore the angles GBA , BGH are equal to two right angles [i 29]

But the angle BGH is right

therefore the angle GBA is also right

therefore GB is at right angles to BA

For the same reason

GB is also at right angles to BC

Since then the straight line GB is set up at right angles to the two straight lines BA, BC which cut one another, therefore GB is also at right angles to the plane through BA, BC [xi 4]


But planes to which the same straight line is at right angles are parallel, [xi 14]
therefore the plane through AB, BC is parallel to the plane through DE, EF

Therefore if two straight lines meeting one another be parallel to two straight lines meeting one another not in the same plane, the planes through them are parallel Q E D

PROPOSITION 16

If two parallel planes be cut by any plane their common sections are parallel

For let the two parallel planes AB, CD be cut by the plane $EFGH$,
and let EF, GH be their common sections,
I say that EF is parallel to GH



For if not EF, GH will when produced meet either in the direction of FH or of EG

Let them be produced as in the direction of FH and let them first meet at K

Now since EFA is in the plane AB

therefore all the points on EFA are also in the plane AB [xi 1]

But A is one of the points on the straight line EFA ,

therefore K is in the plane AB

For the same reason

K is also in the plane CD

therefore the planes AB, CD will meet when produced

But they do not meet because they are by hypothesis parallel
therefore the straight lines EF, GH will not meet when produced in the direction of FH

Similarly we can prove that neither will the straight lines EF, GH meet when produced in the direction of EG

But straight lines which do not meet in either direction are parallel [i Def 23]


Therefore EF is parallel to GH

Therefore etc

Q E D

PROPOSITION 17

If two straight lines be cut by parallel planes they will be cut in the same ratios
For let the two straight lines AB CD be cut by the parallel planes GH , KL , MN at the points A , E , B and C , F , D ,



I say that, as the straight line AE is to EB so is CF to FD

For let AC BD , AD be joined,
let AD meet the plane KL at the point O ,
and let EO OF be joined

Now, since the two parallel planes KL MN are cut by the plane $EBDO$,
their common sections EO BD are parallel [XI 16]

For the same reason since the two parallel planes GH KL are cut by the plane $AOCF$,

their common sections AC , OF are parallel [id]

And, since the straight line EO has been drawn parallel to BD , one of the sides of the triangle ABD

therefore proportionally, as AE is to EB so is AO to OD [VI 2]

Again since the straight line OF has been drawn parallel to AC one of the sides of the triangle ADC

proportionally as AO is to OD , so is CF to FD [id]

But it was also proved that as AO is to OD so is AE to EB ,
therefore also as AE is to EB , so is CF to FD [v 11]

Therefore etc

Q E D

PROPOSITION 18

If a straight line be at right angles to any plane, all the planes through it will also be at right angles to the same plane

For let any straight line AB be at right angles to the plane of reference

I say that all the planes through AB are also at right angles to the plane of reference

For let the plane DE be drawn through AB
let CE be the common section of the plane DE
and the plane of reference

let a point F be taken at random on CE
and from F let FG be drawn in the plane DE
at right angles to CE [I 11]

Now since AB is at right angles to the plane of reference AB is also at right angles to all the straight lines which meet it and are in the plane of reference [XI Def 3]

so that it is also at right angles to CE

therefore the angle ABF is right

But the angle GFB is also right

therefore AB is parallel to FG [I 28]

But AB is at right angles to the plane of reference,

therefore FG is also at right angles to the plane of reference [xi 8]

Now a plane is at right angles to a plane when the straight lines drawn in one of the planes at right angles to the common section of the planes are at right angles to the remaining plane [xi Def 4]

And FG drawn in one of the planes DE at right angles to CE the common section of the planes was proved to be at right angles to the plane of reference therefore the plane DE is at right angles to the plane of reference

Similarly also it can be proved that all the planes through AB are at right angles to the plane of reference

Therefore etc

Q E D

PROPOSITION 19

If two planes which cut one another be at right angles to any plane their common section will also be at right angles to the same plane

For let the two planes AB BC be at right angles to the plane of reference and let BD be their common section

I say that BD is at right angles to the plane of reference

For suppose it is not and from the point D let DE be drawn in the plane AB at right angles to the straight line AD and DF in the plane BC at right angles to CD

Now since the plane AB is at right angles to the plane of reference and DE has been drawn in the plane AB at right angles to AD their common section

therefore DF is at right angles to the plane of reference

[xi Def 4]

Similarly we can prove that

DF is also at right angles to the plane of reference

Therefore from the same point D two straight lines have been set up at right angles to the plane of reference on the same side which is impossible

[xi 13]

Therefore no straight line except the common section DB of the planes AB BC can be set up from the point D at right angles to the plane of reference

Q E D

Therefore etc

PROPOSITION 20



If a solid angle be contained by three plane angles any two taken together in any manner are greater than the remaining one

For let the solid angle at A be contained by the three plane angles BAC CAD DAB

I say that any two of the angles BAC CAD DAB taken together in any manner are greater than the remaining one

If now the angles BAC CAD DAB are equal to one another it is manifest that any two are greater than the remaining one

But if not let BAC be greater and on the straight line AB and at the point A on it let the angle BAE be



constructed in the plane through BA AC , equal to the angle DAB

let AE be made equal to AD ,

and let BEC , drawn across through the point E cut the straight lines AB , AC at the points B C ,

let DB , DC be joined

Now since DA is equal to AE ,

and AB is common

two sides are equal to two sides,

and the angle DAB is equal to the angle BAE ,

therefore the base DB is equal to the base BE [I 4]

And, since the two sides BD , DC are greater than BC [I 20]

and of these DB was proved equal to BE

therefore the remainder DC is greater than the remainder EC

Now, since DA is equal to AE ,

and AC is common

and the base DC is greater than the base EC ,

therefore the angle DAC is greater than the angle EAC [I 25]

But the angle DAB was made equal to the angle BAE ,

therefore the angles DAB , DAC are greater than the angle BAC

Similarly we can prove that the remaining angles also taken together two and two, are greater than the remaining one

Therefore etc

Q E D

PROPOSITION 21

Any solid angle is contained by plane angles less than four right angles

Let the angle at A be a solid angle contained by the plane angles BAC , CAD DAB ,

I say that the angles BAC CAD DAB are less than four right angles

For let points B , C , D be taken at random on the straight lines AB , AC AD respectively,

and let BC , CD , DB be joined

Now since the solid angle at B is contained by the three plane angles CBA ABD CBD any two are greater than the remaining one

[XI 20]

therefore the angles CBA ABD are greater than the angle CBD

For the same reason


the angles BCA ACD are also greater than the angle BCD , and the angles CDA ADB are greater than the angle CDB

therefore the six angles CBA ABD BCA ACD CDA ADB are greater than the three angles CBD BCD CDB

But the three angles CBD BDC BCD are equal to two right angles [I 32] therefore the six angles CBA ABD BCA , ACD CDA ADB are greater than two right angles

And since the three angles of each of the triangles ABC ACD ADB are equal to two right angles

therefore the nine angles of the three triangles the angles CBA ACB , BAC , ACD CDA CAD ADB , DBA BAD are equal to six right angles



and of them the six angles $ABC, BCA, ACD, CDA, ADB, DBA$ are greater than two right angles,
therefore the remaining three angles BAC, CAD, DAB containing the solid angle are less than four right angles

Therefore etc

Q E D

PROPOSITION 22

If there be three plane angles of which two taken together in any manner are greater than the remaining one, and they are contained by equal straight lines, it is possible to construct a triangle out of the straight lines joining the extremities of the equal straight lines

Let there be three plane angles ABC, DEF, GHA , of which two taken together in any manner are greater than the remaining one, namely

the angles ABC, DEF greater than the angle GHA ,


the angles DEF, GHA greater than the angle ABC

and further, the angles GHA, ABC greater than the angle DEF ,

let the straight lines AB, BC, DE, EF, GH, HA be equal,

and let AC, DF, GA be joined,

I say that it is possible to construct a triangle out of straight lines equal to AC, DF, GA that is that any two of the straight lines AC, DF, GA are greater than the remaining one



Now if the angles ABC, DEF, GHA are equal to one another it is manifest that AC, DF, GA being equal also it is possible to construct a triangle out of straight lines equal to AC, DF, GA

But if not let them be unequal

and on the straight line HA and at the point H on it let the angle KHL be constructed equal to the angle ABC ,

let HL be made equal to one of the straight lines AB, BC, DE, EF, GH, HA

and let KL, GL be joined

Now since the two sides AB, BC are equal to the two sides KH, HL

and the angle at B is equal to the angle KHL


therefore the base AC is equal to the base KL

And since the angles ABC, GHA are greater than the angle DEF ,

while the angle ABC is equal to the angle KHL

therefore the angle GHL is greater than the angle DEF

And since the two sides GH, HL are equal to the two sides DE, EF



{ 4 }

and the angle GHL is greater than the angle DEF ,
therefore the base GL is greater than the base DF

[1 24]

But GK, KL are greater than GL

Therefore GK, KL are much greater than DF

But KL is equal to AC ,

therefore AC, GK are greater than the remaining straight line DF

Similarly we can prove that

AC, DF are greater than GK ,

and further DF, GK are greater than AC




Therefore it is possible to construct a triangle out of straight lines equal to
 AC, DF, GK

Q E D

PROPOSITION 23

To construct a solid angle out of three plane angles two of which taken together in any manner, are greater than the remaining one thus the three angles must be less than four right angles

Let the angles ABC, DEF, GHA be the three given plane angles and let two of these taken together in any manner be greater than the remaining one, while further the three are less than four right angles
thus it is required to construct a solid angle out of angles equal to the angles ABC, DEF, GHA



Let AB, BC, DE, EF, GH, HA be cut off equal to one another
and let AC, DF, GK be joined

it is therefore possible to construct a triangle out of straight lines equal to AC, DF, GK

[xi 22]

Let LMN be so constructed that AC is equal to LM, DF to MN and further, GK to NL ,
let the circle LMN be described about the triangle LMN

let its centre be taken and let it be O ,

let LO, MO, NO be joined

I say that $\angle B$ is greater than $\angle O$

For if not $\angle B$ is either equal to $\angle O$, or less

First let it be equal

Then since AB is equal to LO

while AB is equal to BC and OL to OM


the two sides AB, BC are equal to the two sides LO, OM respectively,

and by hypothesis the base AC is equal to the base LM ,

therefore the angle ABC is equal to the angle LOM

[1 8]

For the same reason



the angle DEF is also equal to the angle MON ,
 and further the angle GKH to the angle NOL ,
 therefore the three angles ABC, DEF, GHK are equal to the three angles LOM, MOV, NOL

But the three angles LOM, MON, NOL are equal to four right angles,
 therefore the angles ABC, DEF, GHK are equal to four right angles

But they are also, by hypothesis, less than four right angles
 which is absurd

Therefore AB is not equal to LO

I say next that neither is AB less than LO

For, if possible let it be so

and let OP be made equal to AB and OQ equal to BC ,
 and let PQ be joined

Then, since AB is equal to BC ,

OP is also equal to OQ ,

so that the remainder LP is equal to QV

Therefore LM is parallel to PQ

and LMO is equiangular with PQO ,

therefore, as OL is to LM , so is OP to PQ ,

and alternately as LO is to OP , so is LM to PQ

But LO is greater than OP ,

therefore LM is also greater than PQ

But LM was made equal to AC ,

therefore AC is also greater than PQ

Since then the two sides AB, BC are equal to the two sides PO, OQ ,

and the base AC is greater than the base PQ

therefore the angle ABC is greater than the angle POQ [I 25]

Similarly we can prove that

the angle DEF is also greater than the angle MON ,

and the angle GKH greater than the angle NOL

Therefore the three angles ABC, DEF, GHK are greater than the three angles LOM, MON, NOL

But by hypothesis the angles ABC, DEF, GHK are less than four right angles

therefore the angles LOM, MON, NOL are much less than four right angles

But they are also equal to four right angles

which is absurd

Therefore AB is not less than LO

And it was proved that neither is it equal,

therefore AB is greater than LO

Let then OR be set up from the point O at right angles to the plane of the circle LMN [XI 17]

and let the square on OR be equal to that area by which the square on AB is greater than the square on LO [Lemma]

let RL, RM, RN be joined

Then since RO is at right angles to the plane of the circle LMN ,
 therefore RO is also at right angles to each of the straight lines LO, MO, NO

And since LO is equal to OM

while OR is common and at right angles

therefore the base RL is equal to the base RM

[1 4]

For the same reason

RN is also equal to each of the straight lines RL RM ,

therefore the three straight lines RL RM , RN are equal to one another

Next, since by hypothesis the square on OR is equal to that area by which the square on AB is greater than the square on LO

therefore the square on AB is equal to the squares on LO OR

But the square on LR is equal to the squares on LO OR , for the angle LOR is right,

[1 47]

therefore the square on AB is equal to the square on RL ,

therefore AB is equal to RL

But each of the straight lines BC DE EF , GH HA is equal to AB ,

while each of the straight lines RM , RN is equal to RL ,

therefore each of the straight lines AB , BC , DE EF GH HA

is equal to each of the straight lines RL , RM RN

And since the two sides LR RM are equal to the two sides AB , BC ,

and the base LM is by hypothesis equal to the base AC ,

therefore the angle LRM is equal to the angle ABC

[1 8]

For the same reason

the angle MRN is also equal to the angle DEF ,

and the angle LRN to the angle GHA

Therefore, out of the three plane angles LRM , MRN , LRN , which are equal to the three given angles ABC , DEF GHA the solid angle at R has been constructed which is contained by the angles LRM MRN LRN Q E F

LEMMA

But how it is possible to take the square on OR equal to that area by which the square on AB is greater than the square on LO we can show as follows

Let the straight lines AB LO be set out,

and let AB be the greater,

let the semicircle ABC be described on AB

and into the semicircle ABC let AC be fitted equal to the straight line LO not being greater than the diameter AB

[iv 1]

let CB be joined

Since then the angle ACB is an angle in the semicircle ACB

therefore the angle ACB is right

[iii 31]

Therefore the square on AB is equal to the squares on AC CB

[1 47]

Hence the square on AB is greater than the square on LO by the square on CB

But AC is equal to LO

Therefore the square on AB is greater than the square on LO by the square on CB

If then we cut off OR equal to BC the square on AB will be greater than the square on LO by the square on OR


PROPOSITION 24

If a solid be contained by parallel planes the opposite planes are parallelogrammic

For let the solid $CDHG$ be contained by the parallel planes AC , GF AH DF BF , AE ,

I say that the opposite planes in it are equal and parallelogrammic

For since the two parallel planes BG CE are cut by the plane AC their common sections are parallel [xi 16]



Therefore AB is parallel to DC

Again since the two parallel planes BF AE are cut by the plane AC their common sections are parallel [xi 16]

Therefore BC is parallel to AD

But AB was also proved parallel to DC

therefore AC is a parallelogram

Similarly we can prove that each of the planes DF , FG , GB , BF , AE is a parallelogram

Let AH DF be joined

Then since AB is parallel to DC and BH to CF the two straight lines AB BH which meet one another are parallel to the two straight lines DC CF which meet one another not in the same plane, therefore they will contain equal angles, [xi 10]

therefore the angle ABH is equal to the angle DCF

And since the two sides AB BH are equal to the two sides DC CF , [i 33]

and the angle ABH is equal to the angle DCF

therefore the base AH is equal to the base DF

and the triangle ABH is equal to the triangle DCF [i 4]

And the parallelogram BG is double of the triangle ABH and the parallelogram CE double of the triangle DCF [i 34]

therefore the parallelogram BG is equal to the parallelogram CE

Similarly we can prove that

AC is also equal to GF

and AE to BF

Q E D

Therefore etc

PROPOSITION 25

If a parallelepipedal solid be cut by a plane which is parallel to the opposite planes then as the base is to the base so will the solid be to the solid

For let the parallelepipedal solid $ABCD$ be cut by the plane FG which is parallel to the opposite planes RA DH

I say that as the base $AFFV$ is to the base $EHCF$ so is the solid $ABFU$ to the solid $EHCD$

For let AH be produced in each direction

let any number of straight lines whatever AK KL be made equal to AE ,


and any number whatever HM MN equal to EH ,

and let the parallelograms IP KV HW MS and the solids LQ , KR , DM , NT be completed

Then since the straight lines LA KA AE are equal to one another

the parallelograms LP KV IF are also equal to one another,

KO , KB , AG are equal to one another,
and further, LV , KQ , AR are equal to one another, for they are opposite
[xi 24]



For the same reason

the parallelograms EC , HW , MS are also equal to one another,

HG , HI , IN are equal to one another

and further, DH , MI , NT are equal to one another

Therefore in the solids LQ , KR , AU three planes are equal to three planes

But the three planes are equal to the three opposite,

therefore the three solids LQ , KR , AU are equal to one another

For the same reason

the three solids ED , DM , MT are also equal to one another

Therefore whatever multiple the base LF is of the base AF , the same multiple also is the solid LU of the solid AU

For the same reason

whatever multiple the base NF is of the base FH the same multiple also is the solid NU of the solid HU

And if the base LF is equal to the base NF the solid LU is also equal to the solid NU

if the base LF exceeds the base NF the solid LU also exceeds the solid NU ,
and, if one falls short, the other falls short

Therefore there being four magnitudes the two bases AF , FH and the two solids AU , UH

equimultiples have been taken of the base AF and the solid AU namely the base LF and the solid LU ,

and equimultiples of the base FH and the solid HU , namely the base NF and the solid NU

and it has been proved that if the base LF exceeds the base NF the solid LU also exceeds the solid NU

if the bases are equal the solids are equal,

and if the base falls short the solid falls short

Therefore as the base AF is to the base FH , so is the solid AU to the solid UH

[v Def 5]

Q E D

PROPOSITION 26

On a given straight line and at a given point on it, to construct a solid angle equal to a given solid angle

Let AB be the given straight line, A the given point on it and the angle at D contained by the angles EDC , EDF , FDC , the given solid angle, thus it is required to construct on the straight line AB , and at the point A on it a solid angle equal to the solid angle at D

For let a point F be taken at random on DF

let FG be drawn from F perpendicular to the plane through ED DC , and let it meet the plane at G [xi 11]

let DG be joined, let there be constructed on the straight line AB and at the point A on it the angle BAL equal to the angle EDC , and the angle BAK equal to the angle EDG

[i 23]

let AK be made equal to DG let KH be set up from the point K at right angles to the plane through BA , AL , [xi 12]

let KH be made equal to GF

and let HA be joined

I say that the solid angle at A , contained by the angles BAL , BAH , HAL is equal to the solid angle at D contained by the angles EDC , EDF , FDC

For let AB DE be cut off equal to one another

and let HB KB FE GE be joined

Then since FG is at right angles to the plane of reference it will also make right angles with all the straight lines which meet it and are in the plane of reference [xi Def 3]

therefore each of the angles FGD FGE is right

For the same reason

each of the angles HKA HKB is also right

And since the two sides KA AB are equal to the two sides GD DE respectively

and they contain equal angles

therefore the base KB is equal to the base GE [i 4]

But KH is also equal to GF

and they contain right angles

therefore HB is also equal to FE [i 4]

Again since the two sides KA KH are equal to the two sides DG GF ,

and they contain right angles

therefore the base HA is equal to the base FD [i 4]

But AB is also equal to DE


therefore the two sides HA AB are equal to the two sides DF DE

And the base HB is equal to the base FE

therefore the angle BAH is equal to the angle EDF [i 8]

For the same reason

the angle HAL is also equal to the angle FDC




And the angle BAL is also equal to the angle EDC

Therefore on the straight line AB , and at the point A on it a solid angle has been constructed equal to the given solid angle at D Q E F

PROPOSITION 27

On a given straight line to describe a parallelepipedal solid similar and similarly situated to a given parallelepipedal solid

Let AB be the given straight line and CD the given parallelepipedal solid thus it is required to describe on the given straight line AB a parallelepipedal solid similar and similarly situated to the given parallelepipedal solid CD



For on the straight line AB and at the point A on it let the solid angle contained by the angles BAH HAK KAB , be constructed equal to the solid angle at C , so that the angle BAH is equal to the angle ECF the angle BAK equal to the

angle ECG , and the angle KAH to the angle GCF

and let it be contrived that

as EC is to CG , so is BA to AK ,

and as GC is to CF , so is KA to AH

[vi 12]

Therefore also *ex aequali*

as EC is to CF , so is BA to AH

[v 22]

Let the parallelogram HB and the solid AL be completed

Now since, as EC is to CG so is BA to AK ,

and the sides about the equal angles ECG BAK are thus proportional

therefore the parallelogram GE is similar to the parallelogram AB

For the same reason

the parallelogram AH is also similar to the parallelogram GF and further FE to HB ,

therefore three parallelograms of the solid CD are similar to three parallelograms of the solid AL

But the former three are both equal and similar to the three opposite parallelograms

and the latter three are both equal and similar to the three opposite parallelograms,

therefore the whole solid CD is similar to the whole solid AL [xi Def 9]

Therefore on the given straight line AB there has been described AL similar and similarly situated to the given parallelepipedal solid CD Q E F

PROPOSITION 28

If a parallelepipedal solid be cut by a plane through the diagonals of the opposite planes the solid will be bisected by the plane

For let the parallelepipedal solid AB be cut by the plane $CDEF$ through the diagonals CF , DE of opposite planes

I say that the solid AB will be bisected by the plane $CDEF$


For since the triangle CGF is equal to the triangle CFB ,

[i 34]

and ADE to DEH ,
 while the parallelogram CA is also equal to the parallelogram EB , for they are opposite and GE to CH ,
 therefore the prism contained by the two triangles CGF , ADE and the three parallelograms GE , AC , CE is also equal to the prism contained by the two triangles CFB , DEH and the three parallelograms CH , BE , CE , for they are contained by planes equal both in multitude and in magnitude [11 Def 10]

Hence the whole solid AB is bisected by the plane $CDEF$

Q E D



PROPOSITION 29

Parallelepipedal solids which are on the same base and of the same height and in which the extremities of the sides which stand up are on the same straight lines are equal to one another

Let CM , CN be parallelepipedal solids on the same base AB and of the same height

and let the extremities of their sides which stand up namely AG , AF , LM , LN , CD , CE , BH , BA , be on the same straight lines FN , DA

I say that the solid CM is equal to the solid CN

For since each of the figures CH , CA is a parallelogram CB is equal to each of the straight lines DH , EA [1 34]

hence DH is also equal to EA

Let EH be subtracted from each

therefore the remainder DE is equal to the remainder HK

Hence the triangle DCE is also equal to the triangle HBK [1 8 4]

and the parallelogram DG to the parallelogram HN [1 36]

For the same reason

the triangle AFG is also equal to the triangle MLN

But the parallelogram CF is equal to the parallelogram BM and CG to BN , for they are opposite


therefore the prism contained by the two triangles AFG , DCE and the three parallelograms AD , DG , CG is equal to the prism contained by the two triangles MLN , HBA and the three parallelograms BM , HN , BN

Let there be added to each the solid of which the parallelogram AB is the base and GEH its opposite

therefore the whole parallelepipedal solid CM is equal to the whole parallelepipedal solid CN

Therefore etc

Q E D



PROPOSITION 30

Parallelepipedal solids which are on the same base and of the same height and in which the extremities of the sides which stand up are not on the same straight lines are equal to one another

Let CM CN be parallelepipedal solids on the same base AB and of the same height, and let the extremities of their sides which stand up namely $AF, AG, LM, LN, CD, CE, BH, BK$, not be on the same straight lines,

I say that the solid CM is equal to the solid CN

For let NK DH be produced and meet one another at R

and further, let FM, GE be produced to P, Q

let AO, LP, CQ, BR be joined

Then the solid CM , of which the parallelogram $ACBL$ is the base, and $FDHM$ its opposite, is equal to the solid CP , of which the parallelogram $ACBL$ is the base and $OQRP$ its opposite for they are on the same base $ACBL$ and of the same height, and the extremities of their sides which stand up namely $AF, AO, LM, LP, CD, CQ, BH, BR$, are on the same straight lines FP, DR [XI 29]

But the solid CP , of which the parallelogram $ACBL$ is the base and $OQRP$ its opposite, is equal to the solid CN , of which the parallelogram $ACBL$ is the base and $GEKN$ its opposite, for they are again on the same base $ACBL$ and of the same height and the extremities of their sides which stand up namely $AG, AO, CE, CQ, LN, LP, BK, BR$ are on the same straight lines GQ, NR

Hence the solid CM is also equal to the solid CN

Therefore etc



Q E D

PROPOSITION 31

Parallelepipedal solids which are on equal bases and of the same height are equal to one another

Let the parallelepipedal solids AE CF , of the same height be on equal bases AB, CD

I say that the solid AE is equal to the solid CF



First let the sides which stand up $HK, BE, AG, LM, PQ, DF, CO, RS$, be at right angles to the bases AB, CD

let the straight line RT be produced in a straight line with CR on the straight line RT and at the point R on it let the angle TRU be constructed equal to the angle ALB , [I 23]

let RT be made equal to AL , and RU equal to LB ,

and let the base RW and the solid λU be completed

Now since the two sides TR RU are equal to the two sides AL LB ,

and they contain equal angles,

therefore the parallelogram RW is equal and similar to the parallelogram HL

Since again AL is equal to RT , and LM to RS

and they contain right angles

therefore the parallelogram RX is equal and similar to the parallelogram AM

For the same reason

LE is also equal and similar to SU ,

therefore three parallelograms of the solid AE are equal and similar to three parallelograms of the solid λU

But the former three are equal and similar to the three opposite and the latter three to the three opposite [XI 21]

therefore the whole parallelepipedal solid AE is equal to the whole parallelepipedal solid XU [XI Def 10]

Let DR , WU be drawn through and meet one another at Y ,

let aTb be drawn through T parallel to DY ,

let PD be produced to a

and let the solids λX , RI be completed

Then the solid λY of which the parallelogram RA is the base and Yc its opposite is equal to the solid XU of which the parallelogram RA is the base and UI its opposite

for they are on the same base RY and of the same height and the extremities of their sides which stand up namely RY RU , Tb TH , Sc Sd λc λI , are on the same straight lines YW eV [XI 20]

But the solid λU is equal to AE

therefore the solid λY is also equal to the solid AE

And since the parallelogram $RUWT$ is equal to the parallelogram YT for they are on the same base RT and in the same parallels RT , YW , [I 35]

while $RUWT$ is equal to CD since it is also equal to AB ,

therefore the parallelogram YT is also equal to CD

But DT is another parallelogram,

therefore as the base CD is to DT so is YT to DT [v 7]

And since the parallelepipedal solid CI has been cut by the plane RF which is parallel to opposite planes

as the base CD is to the base DT so is the solid CF to the solid RI [XI 25]

For the same reason

since the parallelepipedal solid YI has been cut by the plane RY which is parallel to opposite planes

as the base YT is to the base TD so is the solid YX to the solid RI [XI 25]

But as the base CD is to DT so is YT to DT

therefore also as the solid CF is to the solid RI so is the solid YX to RI [v 11]

Therefore each of the solids CF YX has to RI the same ratio



therefore the solid CF is equal to the solid YX [v 9]

But XY was proved equal to AE

therefore AE is also equal to CF

Next, let the sides standing up $AG, HK, BE, LM, CN, PQ, DF, RS$ not be at right angles to the bases AB, CD ,

I say again that the solid AE is equal to the solid CF



For from the points K, E, G, M, Q, F, N, S let $KO, ET, GU, MV, QW, FX, NY, SI$ be drawn perpendicular to the plane of reference and let them meet the plane at the points O, T, U, V, W, X, Y, I

and let $OT, OU, UV, TV, WX, WY, YI, IX$ be joined

Then the solid KV is equal to the solid QI ,
for they are on the equal bases KM, QS and of the same height and their sides which stand up are at right angles to their bases [First part of this Prop]

But the solid KV is equal to the solid AE

and QI to CF

for they are on the same base and of the same height while the extremities of their sides which stand up are not on the same straight lines [xi 30]

Therefore the solid AE is also equal to the solid CF

Therefore etc



Q E D

PROPOSITION 32

Parallelepipedal solids which are of the same height are to one another as their bases

Let AB, CD be parallelepipedal solids of the same height

I say that the parallelepipedal solids AB, CD are to one another as their bases that is that as the base AE is to the base CF so is the solid AB to the solid CD



For let FH equal to AE be applied to FG [I 45]
and on FH as base and with the same height as that of CD let the parallelepipedal solid GA be completed

Then the solid AB is equal to the solid GA
for they are on equal bases AE, FH and of the same height [xi 31]

And since the parallelepipedal solid CA is cut by the plane DG which is parallel to opposite planes

But the former three parallelograms are equal and similar to their opposites and the latter three to their opposites, [xi 24]
 therefore the whole solid AP is equal and similar to the whole solid CD [xi Def 10]

Let the parallelogram GK be completed,
 and on the parallelograms GK, KL as bases and with the same height as that of AB , let the solids EO, LQ be completed

Then since owing to the similarity of the solids AB, CD

as AE is to CF so is EG to FN , and EH to FR

while CF is equal to EK , FN to EL , and FR to EM

therefore as AE is to EK , so is GE to FI , and HE to EM

But, as AE is to EK so is AG to the parallelogram GK ,

as GE is to EL , so is GK to KL

and as HE is to EM , so is QE to KM , [vi 1]

therefore also as the parallelogram AG is to GK so is GK to KL and QE to KM

But, as AG is to GK , so is the solid AB to the solid EO ,

as GK is to KL so is the solid OE to the solid QL

and, as QE is to KM , so is the solid QL to the solid AP , [xi 32]

therefore also as the solid AB is to EO so is EO to QL and QL to AP

But if four magnitudes be continuously proportional, the first has to the fourth the ratio triplicate of that which it has to the second [v Def 10]

therefore the solid AB has to AP the ratio triplicate of that which AB has to EO

But as AB is to EO , so is the parallelogram AG to GK and the straight line AE to EK [vi 1]

hence the solid AB has also to AP the ratio triplicate of that which AE has to EK

But the solid AP is equal to the solid CD ,

and the straight line EK to CF

therefore the solid AB has also to the solid CD the ratio triplicate of that which the corresponding side of it AE has to the corresponding side CF

Therefore etc

Q E D

PROPOSITION 34
 From this it is manifest that if four straight lines be continuous
 proportional as the first is to the fourth so will a parallelepipedal solid on
 the first be to the similar and similarly described parallelepipedal solid on the
 second inasmuch as the first has to the fourth the ratio triplicate of that which
 it has to the second

PROPOSITION 34

In equal parallelepipedal solids the bases are reciprocally proportional to the heights and those parallelepipedal solids in which the bases are reciprocally proportional to the heights are equal

Let AB, CD be equal parallelepipedal solids,

I say that in the parallelepipedal solids AB, CD the bases are reciprocally proportional to the heights

that is as the base EHI is to the base NQ so is the height of the solid CD to the height of the solid AB

First, let the sides which stand up namely $AG, EF, LB, HA, CM, NO, PD$

QR be at right angles to their bases,

I say that, as the base EH is to the base NQ , so is CM to AG

If now the base EH is equal to the base NQ

while the solid AB is also equal to the solid CD ,

CM will also be equal to AG



For parallelepipedal solids of the same height are to one another as the bases, [xi 32]

and as the base EH is to NQ so will CM be to AG

and it is manifest that in the parallelepipedal solids AB , CD the bases are reciprocally proportional to the heights

Next, let the base EH not be equal to the base NQ ,

but let EH be greater



Now the solid AB is equal to the solid CD

therefore CM is also greater than AG

Let then CT be made equal to AG ,

and let the parallelepipedal solid VC be completed on NQ as base and with CT as height

Now since the solid AB is equal to the solid CD

and CV is outside them

while equals have to the same the same ratio

therefore as the solid AB is to the solid CV so is the solid CD to the solid CV [v 1]

But as the solid AB is to the solid CV so is the base EH to the base NQ

for the solids AB , CV are of equal height

and as the solid CD is to the solid CV so is the base MQ to the base TQ [xi 32]

[xi 23] and CM to CT [vi 1]

therefore also as the base EH is to the base NQ so is MC to CT

But CT is equal to AG

therefore also as the base EH is to the base NQ so is MC to AG

Therefore in the parallelepipedal solids AB , CD the bases are reciprocally proportional to the heights

Again in the parallelepipedal solids AB , CD let the bases be reciprocally proportional to the heights that is as the base EH is to the base NQ so let the height of the solid CD be to the height of the solid AB ,

I say that the solid AB is equal to the solid CD

Let the sides which stand up be again at right angles to the bases

Now if the base EH is equal to the base NQ

and as the base EH is to the base NQ so is the height of the solid CD to the height of the solid AB ,

therefore the height of the solid CD is also equal to the height of the solid AB

But parallelepipedal solids on equal bases and of the same height are equal to one another, [xi 31]

therefore the solid AB is equal to the solid CD

Next let the base EH not be equal to the base NQ ,

but let EH be greater,

therefore the height of the solid CD is also greater than the height of the solid AB ,

that is CM is greater than AG

Let CT be again made equal to AG

and let the solid CV be similarly completed

Since, as the base EH is to the base NQ so is MC to AG

while AG is equal to CT

therefore as the base EH is to the base NQ so is CM to CT

But as the base EH is to the base NQ , so is the solid AB to the solid CV ,

for the solids AB , CV are of equal height, [xi 32]

and, as CM is to CT , so is the base MQ to the base QT [xi 1]

and the solid CD to the solid CV [xi 25]

Therefore also as the solid AB is to the solid CV , so is the solid CD to the solid CV ,

therefore each of the solids AB CD has to CV the same ratio

Therefore the solid AB is equal to the solid CD [v 9]

Now let the sides which stand up FE , BL , GA , HK , ON , DP , MC , RQ not be at right angles to their bases



let perpendiculars be drawn from the points F G , B , K , O M , D , R to the planes through EH , NQ

and let them meet the planes at S T , U , V W X , Y , a

and let the solids FV Oa be completed,

I say that in this case too if the solids AB CD are equal the bases are reciprocally proportional to the heights that is as the base EH is to the base NQ , so is the height of the solid CD to the height of the solid AB

Since the solid AB is equal to the solid CD



while AB is equal to BT ,

for they are on the same base FA and of the same height

[xi 29 30]

and the solid CD is equal to DX

for they are again on the same base RO and of the same height, [id]

therefore the solid BT is also equal to the solid DX

[Therefore, as the base FK is to the base OR , so is the height of the solid DX to the height of the solid BT [Part 1]

But the base FA is equal to the base EH ,

and the base OR to the base NQ ,

therefore as the base EH is to the base NQ , so is the height of the solid DX to the height of the solid BT

But the solids DX , BT and the solids DC , BA have the same heights respectively,

therefore as the base EH is to the base AQ so is the height of the solid DC to the height of the solid AB

Therefore in the parallelepipedal solids AB , CD the bases are reciprocally proportional to the heights

Again in the parallelepipedal solids AB , CD let the bases be reciprocally proportional to the heights

that is as the base EH is to the base NQ , so let the height of the solid CD be to the height of the solid AB ,

I say that the solid AB is equal to the solid CD

For with the same construction

since as the base EH is to the base AQ so is the height of the solid CD to the height of the solid AB

while the base EH is equal to the base FK ,

and NQ to OR

therefore as the base FK is to the base OR so is the height of the solid CD to the height of the solid AB

But the solids AB , CD and BT , DX have the same heights respectively therefore as the base FK is to the base OR , so is the height of the solid DX to the height of the solid BT

Therefore in the parallelepipedal solids BT , DX the bases are reciprocally proportional to the heights

therefore the solid BT is equal to the solid DX [Part 1]

But BT is equal to BA

for they are on the same base FA and of the same height, [xi 29 30]

and the solid DX is equal to the solid DC [id]

Therefore the solid AB is also equal to the solid CD Q E D

PROPOSITION 35

If there be two equal plane angles and on their vertices there be set up elevated straight lines containing equal angles with the original straight lines respectively if on the elevated straight lines points be taken at random and perpendiculars be drawn from them to the planes in which the original angles are, and if from the points so arising in the planes straight lines be joined to the vertices of the original angles they will contain with the elevated straight lines equal angles


Let the angles BAC , EDF be two equal rectilineal angles and from the points A , D let the elevated straight lines AG , DM be set up containing with the original straight lines equal angles respectively, namely the angle MAE to the angle GAB and the angle MDF to the angle GAC ,

let points G, M be taken at random on AG, DM ,
 let GL, MN be drawn from the points G, M perpendicular to the planes through
 BA, AC and ED, DF , and let them meet the planes at L, N ,
 and let LA, ND be joined

I say that the angle GAL is equal to the angle MDN

Let AH be made equal
 to DM ,
 and let HA be drawn
 through the point H paral-
 lel to GL

But GL is perpendicular
 to the plane through BA ,
 AC ,
 therefore HK is also per-
 pendicular to the plane
 through BA, AC [x1 8]



From the points K, N let KC, NF KB, NE be drawn perpendicular to the
 straight lines AC, DF, AB, DE :

and let HC, CB MF, FE be joined

Since the square on HA is equal to the squares on HK, KA ,

and the squares on KC, CA are equal to the square on KA , [1 47]

therefore the square on HA is also equal to the squares on HK, KC, CA

But the square on HC is equal to the squares on HK, KC , [1 47]

therefore the square on HA is equal to the squares on HC, CA

Therefore the angle HCA is right [1 48]

For the same reason

the angle DFM is also right

Therefore the angle ACH is equal to the angle DFM

But the angle HAC is also equal to the angle MDF

Therefore MDF, HAC are two triangles which have two angles equal to two
 angles respectively and one side equal to one side namely that subtending
 one of the equal angles that is, HA equal to MD ,

therefore they will also have the remaining sides equal to the remaining sides
 respectively [1 26]

Therefore AC is equal to DF

Similarly we can prove that AB is also equal to DE

Since then AC is equal to DF and AB to DE

the two sides CA, AB are equal to the two sides FD, DE

But the angle CAB is also equal to the angle FDE

therefore the base BC is equal to the base EF , the triangle to the triangle, and
 the remaining angles to the remaining angles [1 4]

therefore the angle ACB is equal to the angle DFE

But the right angle ACK is also equal to the right angle DFN

therefore the remaining angle BCK is also equal to the remaining angle EFN

For the same reason

the angle CBK is also equal to the angle FEN

Therefore BCK, FEN are two triangles which have two angles equal to two
 angles respectively and one side equal to one side namely that adjacent to
 the equal angles that is BC equal to EF

therefore they will also have the remaining sides equal to the remaining sides [1 26]

Therefore CA is equal to FA

But AC is also equal to DF ,

therefore the two sides AC CA are equal to the two sides DF FA ,
and they contain right angles

Therefore the base AA is equal to the base DN [1 4]

And since AH is equal to DM

the square on AH is also equal to the square on DM

But the squares on AA AH are equal to the square on AM ,
for the angle AAH is right, [1 4]

and the squares on DN DM are equal to the square on DN ,
for the angle DNM is right, [1 47]

therefore the squares on AA AH are equal to the squares on DN DM ,

and of these the square on AA is equal to the square on DN

therefore the remaining square on AH is equal to the square on DM

therefore HK is equal to MN

And since the two sides HA AA are equal to the two sides MD DM respectively,

and the base AA was proved equal to the base MN ,
therefore the angle HAA is equal to the angle MDN [1 8]


Therefore etc

PROPOSITION. From this it is manifest that if there be two equal plane angles and if there be set up on them elevated straight lines which are equal and contain equal angles with the original straight lines respectively, the perpendiculars drawn from their extremities to the planes in which are the original angles are equal to one another Q E D

PROPOSITION 36

If three straight lines be proportional the parallelepipedal solid formed out of the three is equal to the parallelepipedal solid on the mean which is equilateral but equiangular with the aforesaid solid


Let A B C be three straight lines in proportion so that as A is to B so is B to C



A —————

B —————

C —————



I say that the solid formed out of A B C is equal to the solid on B which is equilateral but equiangular with the aforesaid solid

Let there be set out the solid angle at E contained by the angles DEG , GEF , FED ,

let each of the straight lines DE , GE , EF be made equal to B , and let the parallelepipedal solid EK be completed

let LM be made equal to A ,
and on the straight line LM , and at the point L on it, let there be constructed a solid angle equal to the solid angle at E namely that contained by NLO , OLM , MLN

let LO be made equal to B , and LN equal to C

Now since as A is to B so is B to C
while A is equal to LM B to each of the straight lines LO ED and C to LN ,
therefore as LM is to EF so is DE to LN

Thus the sides about the equal angles NLM , DEF are reciprocally proportional,

therefore the parallelogram MN is equal to the parallelogram DF [xi 14]

And since the angles DEF , NLM are two plane rectilineal angles and on them the elevated straight lines LO , EG are set up which are equal to one another and contain equal angles with the original straight lines respectively therefore the perpendiculars drawn from the points G O to the planes through NL , LM and DE EF are equal to one another, [xi 35 Por]

hence the solids LH EK are of the same height

But parallelepipedal solids on equal bases and of the same height are equal to one another [xi 31]

therefore the solid HL is equal to the solid EK

And LH is the solid formed out of A B , C and EK the solid on B
therefore the parallelepipedal solid formed out of A B , C is equal to the solid on B which is equilateral but equiangular with the aforesaid solid


Q E D

PROPOSITION 37

If four straight lines be proportional the parallelepipedal solids on them which are similar and similarly described will also be proportional and if the parallelepipedal solids on them which are similar and similarly described be proportional the straight lines will themselves also be proportional

Let AB , CD EF , GH be four straight lines in proportion so that as AB is to CD so is EF to GH ,

and let there be described on AB , CD , EF GH the similar and similarly situ



ated parallelepipedal solids KA LC , ME NG ,

I say that as KA is to LC so is ME to NG

For since the parallelepipedal solid KA is similar to LC ,
therefore KA has to LC the ratio triplicate of that which AB has to CD [xi 33]

For the same reason
 ME also has to NG the ratio triplicate of that which EF has to GH [id]
And as AB is to CD so is EF to GH

Therefore also as KA is to LC , so is ME to NG

Next as the solid KA is to the solid LC so let the solid ME be to the solid NG ,

I say that as the straight line AB is to CD so is EF to GH

For since again KA has to LC the ratio triplicate of that which AB has to CD [xi 33]

and ME also has to NG the ratio triplicate of that which EF has to GH [id]
and as KA is to LC , so is ME to NG

therefore also as AB is to CD so is EF to GH

Therefore etc

Q E D

PROPOSITION 38

If the sides of the opposite planes of a cube be bisected and planes be carried through the points of section the common section of the planes and the diameter of the cube bisect one another

For let the sides of the opposite planes CF AH of the cube AF be bisected at the points K L M N O Q P R and through the points of section let the planes KN OR be carried
let US be the common section of the planes and DG the diameter of the cube AF

I say that UT is equal to TS and DT to TG

For let DL LE BS SG be joined

Then since DO is parallel to PE

the alternate angles DOU UPE are equal to one another [i 29]

And since DO is equal to PE and
 OU to UP

and they contain equal angles
therefore the base DU is equal to the
base UE

the triangle DOU is equal to the tri-
angle PUE

and the remaining angles are equal to
the remaining angles [i 4]
therefore the angle ODU is equal to
the angle PUE


For this reason DLE is a straight
line [i 14]

For the same reason BSG is also a
straight line

and BS is equal to SG

Now since CA is equal and paral-
lel to DB ,

while CA is also equal and parallel to EG



therefore DB is also equal and parallel to EG

[vi 9]

And the straight lines DE BG join their extremities,

therefore DE is parallel to BG

[i 32]

Therefore the angle EDT is equal to the angle BGT ,

for they are alternate,

[i 29]

and the angle DTU is equal to the angle GTS

[i 15]

Therefore DTU , GTS are two triangles which have two angles equal to two angles and one side equal to one side namely that subtending one of the equal angles that is DU equal to GS

for they are the halves of DE BG

therefore they will also have the remaining sides equal to the remaining sides

[i 26]

Therefore DT is equal to TG and UT to TS

Therefore etc

Q E D

PROPOSITION 39



If there be two prisms of equal height and one have a parallelogram as base and the other a triangle and if the parallelogram be double of the triangle the prisms will be equal

Let $ABCDEF$ $GHLKMN$ be two prisms of equal height

let one have the parallelogram AF as base and the other the triangle GHA ,

and let the parallelogram AF be double of the triangle GHA

I say that the prism $ABCDEF$ is equal to the prism $GHLKMN$



For let the solids AO GP be completed

Since the parallelogram AF is double of the triangle GHA

while the parallelogram HK is also double of the triangle GHA

[i 34]

therefore the parallelogram AF is equal to the parallelogram HA

But parallelepipedal solids which are on equal bases and of the same height are equal to one another

[xi 31]

therefore the solid AO is equal to the solid GP

And the prism $ABCDEF$ is half of the solid AO

and the prism $GHLKMN$ is half of the solid GP

[xi 28]

therefore the prism $ABCDEF$ is equal to the prism $GHLKMN$

Therefore etc

Q E D

BOOK TWELVE

PROPOSITIONS

PROPOSITION 1



Similar polygons inscribed in circles are to one another as the squares on the diameters

Let ABC FGH be circles
let $ABCDE$ $FGHKL$ be similar polygons inscribed in them and let BM GN be diameters of the circles,

I say that as the square on BM is to the square on GN , so is the polygon $ABCDE$ to the polygon $FGHKL$

For let BE AM GL FN be joined

Now since the polygon $ABCDE$ is similar to the polygon $FGHKL$,



the angle BAE is equal to the angle GFL

and as BA is to AE so is GF to FL

[vi Def 1]

Thus BAF GFL are two triangles which have one angle equal to one angle namely the angle BAE to the angle GFL and the sides about the equal angles proportional

therefore the triangle ABE is equiangular with the triangle FGL

[vi 6]

Therefore the angle AEB is equal to the angle FLG

But the angle AEB is equal to the angle AMB

for they stand on the same circumference,

[iii 27]

and the angle FLG to the angle FNH

therefore the angle AMB is also equal to the angle FNH

But the right angle BAM is also equal to the right angle GFN

[iii 31]

therefore the remaining angle is equal to the remaining angle

[i 32]

Therefore the triangle ABM is equiangular with the triangle GFN

Therefore proportionally as BM is to GN so is BA to GF

[vi 4]

But the ratio of the square on BM to the square on GN is duplicate of the ratio of BM to GN and the ratio of the polygon $ABCDE$ to the polygon $FGHKL$ is duplicate of the ratio of BA to GF , [VI 20] therefore also as the square on BM is to the square on GN so is the polygon $ABCDE$ to the polygon $FGHKL$

Therefore etc


Q E D

PROPOSITION 2

Circles are to one another as the squares on the diameters

Let $ABCD$ $EFGH$ be circles and BD FH their diameters

I say that as the circle $ABCD$ is to the circle $EFGH$, so is the square on BD to the square on FH



For if the square on BD is not to the square on FH as the circle $ABCD$ is to the circle $EFGH$ then as the square on BD is to the square on FH so will the circle $ABCD$ be either to some less area than the circle $EFGH$ or to a greater

First let it be in that ratio to a less area S

Let the square $EFGH$ be inscribed in the circle $EFGH$ then the inscribed square is greater than the half of the circle $EFGH$ inasmuch as if through the points E F G H we draw tangents to the circle the square $EFGH$ is half the square circumscribed about the circle and the circle is less than the circumscribed square

hence the inscribed square $EFGH$ is greater than the half of the circle $EFGH$

Let the circumferences EF FG GH HE be bisected at the points K L M N

and let EK KF FL LG GM MH HN NE be joined

therefore each of the triangles EKF FLG GMH HNE is also greater than the half of the segment of the circle about it inasmuch as if through the points K L M N we draw tangents to the circle and complete the parallelograms on the straight lines EF FG GH HE each of the triangles EKF FLG GMH HNE will be half of the parallelogram about it

while the segment about it is less than the parallelogram hence each of the triangles EKF FLG GMH HNE is greater than the half of the segment of the circle about it

Thus by bisecting the remaining circumferences and joining straight lines

and by doing this continually we shall leave some segments of the circle which will be less than the excess by which the circle $EFGH$ exceeds the area S

For it was proved in the first theorem of the tenth book that if two unequal magnitudes be set out and if from the greater there be subtracted a magnitude greater than the half and from that which is left a greater than the half and if this be done continually there will be left some magnitude which will be less than the lesser magnitude set out

Let segments be left such as described and let the segments of the circle $EFGH$ on $EK KF FL LG GM MH HN NE$ be less than the excess by which the circle $EFGH$ exceeds the area S

Therefore the remainder the polygon $EAKFLGMHN$ is greater than the area S

Let there be inscribed also in the circle $ABCD$ the polygon $AOBPCQDR$ similar to the polygon $EAKFLGMHN$

therefore as the square on BD is to the square on FH so is the polygon $AOBPCQDR$ to the polygon $EAKFLGMHN$ [xii 1]

But as the square on BD is to the square on FH so also is the circle $ABCD$ to the area S

therefore also as the circle $ABCD$ is to the area S so is the polygon $AOBPCQDR$ to the polygon $EAKFLGMHN$ [v 11]

therefore alternately as the circle $ABCD$ is to the polygon inscribed in it so is the area S to the polygon $EAKFLGMHN$ [v 16]

But the circle $ABCD$ is greater than the polygon inscribed in it

therefore the area S is also greater than the polygon $EAKFLGMHN$

But it is also less

which is impossible

Therefore as the square on BD is to the square on FH so is not the circle $ABCD$ to any area less than the circle $EFGH$

Similarly we can prove that neither is the circle $EFGH$ to any area less than the circle $ABCD$ as the square on FH is to the square on BD

I say next that neither is the circle $ABCD$ to any area greater than the circle $EFGH$ as the square on BD is to the square on FH

For if possible let it be in that ratio to a greater area S

Therefore inversely as the square on FH is to the square on BD so is the area S to the circle $ABCD$

But as the area S is to the circle $ABCD$ so is the circle $EFGH$ to some area less than the circle $ABCD$

therefore also as the square on FH is to the square on BD so is the circle $EFGH$ to some area less than the circle $ABCD$ [v 11]

which was proved impossible

Therefore as the square on BD is to the square on FH so is not the circle $ABCD$ to any area greater than the circle $EFGH$

And it was proved that neither is it in that ratio to any area less than the circle $EFGH$

therefore as the square on BD is to the square on FH so is the circle $ABCD$ to the circle $EFGH$

Therefore etc

Q E D

LEMMA

I say that the area S being greater than the circle $EFGH$, as the area S is to the circle $ABCD$, so is the circle $EFGH$ to some area less than the circle $ABCD$

For let it be contrived that as the area S is to the circle $ABCD$ so is the circle $EFGH$ to the area T

I say that the area T is less than the circle $ABCD$

For since as the area S is to the circle $ABCD$, so is the circle $EFGH$ to the area T ,

therefore alternately, as the area S is to the circle $EFGH$, so is the circle $ABCD$ to the area T [v 16]

But the area S is greater than the circle $EFGH$,

therefore the circle $ABCD$ is also greater than the area T

Hence as the area S is to the circle $ABCD$ so is the circle $EFGH$ to some area less than the circle $ABCD$ Q E D


PROPOSITION 3

Any pyramid which has a triangular base is divided into two pyramids equal and similar to one another similar to the whole and having triangular bases and into two equal prisms and the two prisms are greater than the half of the whole pyramid

Let there be a pyramid of which the triangle ABC is the base and the point D the vertex

I say that the pyramid $ABCD$ is divided into two pyramids equal to one another having triangular bases and similar to the whole pyramid and into two equal prisms and the two prisms are greater than the half of the whole pyramid

For let AB BC CA AD DB DC be bisected at the points E F G H , K L and let HE , EG GH HA KL LH AF FG be joined



Since AE is equal to EB and AH to DH therefore EH is parallel to DB [vi 2]

For the same reason

HK is also parallel to AB

Therefore $HEBK$ is a parallelogram

therefore HK is equal to EB

[i 34]

But EB is equal to EA

therefore AE is also equal to HA

But AH is also equal to HD

therefore the two sides EA AH are equal to the two sides AH HD respectively

and the angle EAH is equal to the angle KHD ,

therefore the base EH is equal to the base KD

[i 4]

Therefore the triangle AEH is equal and similar to the triangle HKD

For the same reason

the triangle ANG is also equal and similar to the triangle HLD

Now since two straight lines EH HG meeting one another are parallel to two straight lines AD DL meeting one another and are not in the same plane they will contain equal angles [xi 10]

Therefore the angle EHG is equal to the angle ADL

And since the two straight lines EH, HG are equal to the two KD, DL respectively

and the angle EHG is equal to the angle ADL

therefore the base EG is equal to the base KL , [1 4]

therefore the triangle EHG is equal and similar to the triangle ADL

For the same reason

the triangle AEG is also equal and similar to the triangle HKL

Therefore the pyramid of which the triangle AEG is the base and the point H the vertex is equal and similar to the pyramid of which the triangle HKL is the base and the point D the vertex [xi Def 10]

And since HA has been drawn parallel to AB , one of the sides of the triangle ADB

the triangle ADB is equiangular to the triangle DHA [1 29]

and they have their sides proportional

therefore the triangle ADB is similar to the triangle DHK [xi Def 1]

For the same reason

the triangle DBC is also similar to the triangle DAL and the triangle ADC to the triangle DLH

Now since the two straight lines BA, AC meeting one another are parallel to the two straight lines KH, HL meeting one another not in the same plane, they will contain equal angles [xi 10]

Therefore the angle BAC is equal to the angle KHL

And as BA is to AC so is KH to HL

therefore the triangle ABC is similar to the triangle HKL

Therefore also the pyramid of which the triangle ABC is the base and the point D the vertex is similar to the pyramid of which the triangle HKL is the base and the point D the vertex

But the pyramid of which the triangle HKL is the base and the point D the vertex was proved similar to the pyramid of which the triangle AEG is the base and the point H the vertex

Therefore each of the pyramids $AEGH, HKLD$ is similar to the whole pyramid $ABCD$

Next since BF is equal to FC

the parallelogram $EBFC$ is double of the triangle GFC

And since if there be two prisms of equal height and one have a parallelogram as base and the other a triangle and if the parallelogram be double of the triangle the prisms are equal [xi 39]

therefore the prism contained by the two triangles BKF, FHG and the three parallelograms $FBFG, FBKH, HKFG$ is equal to the prism contained by the two triangles GFC, HKL and the three parallelograms $KFCL, LCGH, HKFG$

And it is manifest that each of the prisms namely that in which the parallelogram $EBFG$ is the base and the straight line HA is its opposite and that in which the triangle GFC is the base and the triangle HKL its opposite is greater than each of the pyramids of which the triangles LEG, HAK are the bases and the points H, D the vertices

inasmuch as if we join the straight lines IF, FA the prism in which the parallelogram $EBFC$ is the base and the straight line HA its opposite is greater than the pyramid of which the triangle FBF is the base and the point K the vertex

But the pyramid of which the triangle EBF is the base and the point K the vertex is equal to the pyramid of which the triangle AEG is the base and the point H the vertex

for they are contained by equal and similar planes

Hence also the prism in which the parallelogram $PBFG$ is the base and the straight line HA its opposite is greater than the pyramid of which the triangle AEG is the base and the point H the vertex

But the prism in which the parallelogram $EBFG$ is the base and the straight line HA its opposite is equal to the prism in which the triangle GFC is the base and the triangle HKL its opposite

and the pyramid of which the triangle AEG is the base and the point H the vertex is equal to the pyramid of which the triangle HKL is the base and the point D the vertex

Therefore the said two prisms are greater than the said two pyramids of which the triangles AEG HKL are the bases and the points H D the vertices

Therefore the whole pyramid of which the triangle ABC is the base and the point D the vertex has been divided into two pyramids equal to one another and into two equal prisms and the two prisms are greater than the half of the whole pyramid


Q E D

PROPOSITION 4

If there be two pyramids of the same height which have triangular bases and each of them be divided into two pyramids equal to one another and similar to the whole and into two equal prisms then as the base of the one pyramid is to the base of the other pyramid so will all the prisms in the one pyramid be to all the prisms being equal in multitude in the other pyramid

Let there be two pyramids of the same height which have the triangular bases ABC DEF and vertices the points G H and let each of them be divided into two pyramids equal to one another and similar to the whole and into two equal prisms [xii 3]

I say that as the base ABC is to the base DEF so are all the prisms in the pyramid $ABCG$ to all the prisms being equal in multitude in the pyramid $DEFH$



For since BO is equal to OC and AL to LC
therefore LO is parallel to AB
and the triangle ABC is similar to the triangle LOC

For the same reason

the triangle DEF is also similar to the triangle RVF

And since BC is double of CO and EF of FV ,

therefore as BC is to CO so is EF to FV

And on BC CO are described the similar and similarly situated rectilinear figures ABC , LOC ,

and on EF FV the similar and similarly situated figures DEF , RVF ,
therefore as the triangle ABC is to the triangle LOC so is the triangle DEF to the triangle RVF , [vi 22]

therefore alternately as the triangle ABC is to the triangle DEF , so is the triangle LOC to the triangle RVF [v 16]

But as the triangle LOC is to the triangle RVF so is the prism in which the triangle LOC is the base and PMN its opposite to the prism in which the triangle RVF is the base and STU its opposite [Lemma following]

therefore also as the triangle ABC is to the triangle DEF , so is the prism in which the triangle LOC is the base and PMN its opposite to the prism in which the triangle RVF is the base and STU its opposite

But as the said prisms are to one another so is the prism in which the parallelogram $KBOL$ is the base and the straight line PM its opposite to the prism in which the parallelogram $QEVR$ is the base and the straight line ST its opposite [xi 39 of xii 3]

Therefore also the two prisms that in which the parallelogram $KBOL$ is the base and PM its opposite and that in which the triangle LOC is the base and PMN its opposite are to the prisms in which $QEVR$ is the base and the straight line ST its opposite and in which the triangle RVF is the base and STU its opposite in the same ratio [v 12]

Therefore also as the base ABC is to the base DEF so are the said two prisms to the said two prisms

And similarly if the pyramids $PMNG$ $STUH$ be divided into two prisms and two pyramids

as the base PMN is to the base STU so will the two prisms in the pyramid $PMNG$ be to the two prisms in the pyramid $STUH$

But as the base PMN is to the base STU so is the base ABC to the base DEF

for the triangles PMN STU are equal to the triangles LOC RVF respectively

Therefore also as the base ABC is to the base DEF so are the four prisms to the four prisms

And similarly also if we divide the remaining pyramids into two pyramids and into two prisms then as the base ABC is to base the DEF so will all the prisms in the pyramid $ABCG$ be to all the prisms being equal in multitude in the pyramid $DEFH$ Q E D

LEMMA

But that as the triangle LOC is to the triangle RVF so is the prism in which the triangle LOC is the base and PMN its opposite to the prism in which the triangle RVF is the base and STU its opposite we must prove as follows

For in the same figure let perpendiculars be conceived drawn from G , H to

the planes ABC , DEF , these are of course equal because by hypothesis, the pyramids are of equal height

Now, since the two straight lines GC and the perpendicular from G are cut by the parallel planes ABC , PMN

they will be cut in the same ratios [xi 17]

And GC is bisected by the plane PMN at N , therefore the perpendicular from G to the plane ABC will also be bisected by the plane PMN

For the same reason the perpendicular from H to the plane DEF will also be bisected by the plane STU

And the perpendiculars from G H to the planes ABC , DEF are equal, therefore the perpendiculars from the triangles PMN , STU to the planes ABC , DEF are also equal

Therefore the prisms in which the triangles LOC , RVF are bases, and PMN , STU their opposites, are of equal height

Hence also the parallelepipedal solids described from the said prisms are of equal height and are to one another as their bases, [xi 32]


therefore their halves namely the said prisms, are to one another as the base LOC is to the base RVF Q E D

PROPOSITION 5

Pyramids which are of the same height and have triangular bases are to one another as the bases

Let there be pyramids of the same height of which the triangles ABC , DEF are the bases and the points G H the vertices,

I say that as the base ABC is to the base DEF , so is the pyramid $ABCG$ to the pyramid $DEFH$



For if the pyramid $ABCG$ is not to the pyramid $DEFH$ as the base ABC is to the base DEF , then as the base ABC is to the base DEF so will the pyramid $ABCG$ be either to some solid less than the pyramid $DEFH$ or to a greater

Let it first be in that ratio to a less solid W and let the pyramid $DEFH$ be divided into two pyramids equal to one another and similar to the whole and into two equal prisms

then the two prisms are greater than the half of the whole pyramid [xii 3]

Again let the pyramids arising from the division be similarly divided and let this be done continually until there are left over from the pyramid $DEFH$ some pyramids which are less than the excess by which the pyramid $DEFH$ exceeds the solid Π [x 1]

Let such be left and let them be for the sake of argument, $DQRS STUH$ therefore the remainders, the prisms in the pyramid $DEFH$, are greater than the solid Π

Let the pyramid $ABCG$ also be divided similarly, and a similar number of times with the pyramid $DEFH$

therefore as the base ABC is to the base DEF so are the prisms in the pyramid $ABCG$ to the prisms in the pyramid $DEFH$ [xii 4]

But as the base ABC is to the base DEF so also is the pyramid $ABCG$ to the solid Π

therefore also as the pyramid $ABCG$ is to the solid Π , so are the prisms in the pyramid $ABCG$ to the prisms in the pyramid $DEFH$, [v 11]

therefore alternately as the pyramid $ABCG$ is to the prisms in it so is the solid Π to the prisms in the pyramid $DEFH$ [v 16]

But the pyramid $ABCG$ is greater than the prisms in it, therefore the solid Π is also greater than the prisms in the pyramid $DEFH$

But it is also less

which is impossible

Therefore the prism $ABCG$ is not to any solid less than the pyramid $DEFH$ as the base ABC is to the base DEF

Similarly it can be proved that neither is the pyramid $DEFH$ to any solid less than the pyramid $ABCG$ as the base DEF is to the base ABC

I say next that neither is the pyramid $ABCG$ to any solid greater than the pyramid $DEFH$ as the base ABC is to the base DEF

For if possible let it be in that ratio to a greater solid Π therefore inversely as the base DEF is to the base ABC , so is the solid Π to the pyramid $ABCG$

But as the solid Π is to the solid $ABCG$ so is the pyramid $DEFH$ to some solid less than the pyramid $ABCG$ as was before proved [xii 2 Lemma]

therefore also as the base DEF is to the base ABC , so is the pyramid $DEFH$ to some solid less than the pyramid $ABCG$ [v 11]

which was proved absurd

Therefore the pyramid $ABCG$ is not to any solid greater than the pyramid $DEFH$ as the base ABC is to the base DEF

But it was proved that neither is it in that ratio to a less solid

Therefore as the base ABC is to the base DEF so is the pyramid $ABCG$ to the pyramid $DEFH$ Q E D

PROPOSITION 6

Pyramids which are of the same height and have polygonal bases are to one another as the bases

Let there be pyramids of the same height of which the polygons $ABCDE FGHAL$ are the bases and the points $M N$ the vertices

I say that as the base $ABCD$ is to the base $FGHAL$, so is the pyramid $ABCDEM$ to the pyramid $FGHALN$

For let AC AD FH , FA be joined

Since then $ABCM$, $ACDM$ are two pyramids which have triangular bases and equal height

they are to one another as the bases, [xii 5]

therefore as the base ABC is to the base ACD so is the pyramid $ABCM$ to the pyramid $ACDM$

And *componendo* as the base $ABCD$ is to the base ACD so is the pyramid $ABCDM$ to the pyramid $ACDM$ [v 18]

But also as the base ACD is to the base ADE so is the pyramid $ACDM$ to the pyramid $ADEM$ [xii 5]

Therefore *ex aequali* as the base $ABCD$ is to the base ADE , so is the pyramid $ABCDM$ to the pyramid $ADEM$ [v 22]

And again *componendo* as the base $ABCDE$ is to the base ADE , so is the pyramid $ABCDEM$ to the pyramid $ADEM$ [v 18]

Similarly also it can be proved that as the base $FGHKL$ is to the base FGH , so is the pyramid $FGHKLN$ to the pyramid $FGHN$

And since $ADEM$ $FGHN$ are two pyramids which have triangular bases and equal height

therefore as the base ADE is to the base FGH so is the pyramid $ADEM$ to the pyramid $FGHN$ [xii 5]

But as the base ADE is to the base $ABCDE$ so was the pyramid $ADEM$ to the pyramid $ABCDEM$

Therefore also *ex aequali* as the base $ABCDE$ is to the base FGH , so is the pyramid $ABCDEM$ to the pyramid $FGHN$ [v 22]

But further, as the base FGH is to the base $FGHKL$ so also was the pyramid $FGHN$ to the pyramid $FGHKLN$

Therefore also *ex aequali* as the base $ABCDE$ is to the base $FGHKL$ so is the pyramid $ABCDEM$ to the pyramid $FGHKLN$ [v 22]

Q E D

PROPOSITION 7

Any prism which has a triangular base is divided into three pyramids equal to one another which have triangular bases

Let there be a prism in which the triangle ABC is the base and DEF its opposite

I say that the prism $ABCDEF$ is divided into three pyramids equal to one another which have triangular bases

For let BD EC CD be joined

Since $ABED$ is a parallelogram and BD is its diameter


therefore the triangle ABD is equal to the triangle EBD [i 34]

therefore also the pyramid of which the triangle ABD is the base and the point C the vertex is equal to the pyramid of which the triangle DEB is the base and

the point C the vertex

[xii 5]

But the pyramid of which the triangle DEB is the base and the point C the vertex is the same with the pyramid of which the triangle EBC is the base and the point D the vertex



for they are contained by the same planes

Therefore the pyramid of which the triangle ABD is the base and the point C the vertex is also equal to the pyramid of which the triangle EBC is the base and the point D the vertex

Again since $FCBE$ is a parallelogram
and CE is its diameter

the triangle CEF is equal to the triangle CBE [i 34]

Therefore also the pyramid of which the triangle BCE is the base and the point D the vertex is equal to the pyramid of which the triangle ECF is the base and the point D the vertex [xii 5]

But the pyramid of which the triangle BCE is the base and the point D the vertex was proved equal to the pyramid of which the triangle ABD is the base and the point C the vertex

therefore also the pyramid of which the triangle CEF is the base and the point D the vertex is equal to the pyramid of which the triangle ABD is the base and the point C the vertex

therefore the prism $ABCDEF$ has been divided into three pyramids equal to one another which have triangular bases

And since the pyramid of which the triangle ABD is the base and the point C the vertex is the same with the pyramid of which the triangle CAB is the base and the point D the vertex

for they are contained by the same planes

while the pyramid of which the triangle ABD is the base and the point C the vertex was proved to be a third of the prism in which the triangle ABC is the base and DEF its opposite

therefore also the pyramid of which the triangle ABC is the base and the point D the vertex is a third of the prism which has the same base the triangle ABC , and DEF as its opposite

PROPOSITION From this it is manifest that any pyramid is a third part of the prism which has the same base with it and equal height Q E D

PROPOSITION 8

Similar pyramids which have triangular bases are in the triplicate ratio of their corresponding sides

Let there be similar and similarly situated pyramids of which the triangles ABC DEF are the bases and the points G H the vertices

I say that the pyramid $ABCG$ has to the pyramid $DEFH$ the ratio triplicate of that which BC has to EF

For let the parallelepipedal solids $BGML$ $EHQP$ be completed



Now since the pyramid $ABCG$ is similar to the pyramid $DEFH$

therefore the angle ABC is equal to the angle DEF

the angle GBC to the angle HLF

and the angle ABG to the angle DEH

and as AB is to DE , so is BC to EF , and BG to EH
 And since as AB is to DE , so is BC to EF
 and the sides are proportional about equal angles,



therefore the parallelogram BM is similar to the parallelogram EQ
 For the same reason

BN is also similar to ER and BA to EO

therefore the three parallelograms MB , BA , BN are similar to the three EQ , EO , ER

But the three parallelograms MB , BA , BN are equal and similar to their three opposites

and the three EQ , EO , ER are equal and similar to their three opposites

[xi 24]

Therefore the solids $BGML$, $EHQP$ are contained by similar planes equal in multitude

Therefore the solid $BGML$ is similar to the solid $EHQP$

But similar parallelepipedal solids are in the triplicate ratio of their corresponding sides

[xi 33]

Therefore the solid $BGML$ has to the solid $EHQP$ the ratio triplicate of that which the corresponding side BC has to the corresponding side EF

But as the solid $BGML$ is to the solid $EHQP$, so is the pyramid $ABCG$ to the pyramid $DEFH$,

inasmuch as the pyramid is a sixth part of the solid because the prism which is half of the parallelepipedal solid [xi 26] is also triple of the pyramid [xii 7]

Therefore the pyramid $ABCG$ also has to the pyramid $DEFH$ the ratio triplicate of that which BC has to EF

Q E D

PROPOSITION From this it is manifest that similar pyramids which have polygonal bases are also to one another in the triplicate ratio of their corresponding sides

For, if they are divided into the pyramids contained in them which have triangular bases by virtue of the fact that the similar polygons forming their bases are also divided into similar triangles equal in multitude and corresponding to the wholes

[vi 20]

then as the one pyramid which has a triangular base in the one complete pyramid is to the one pyramid which has a triangular base in the other complete pyramid so also will all the pyramids which have triangular bases contained in the one pyramid be to all the pyramids which have triangular bases contained in the other pyramid [v 12] that is the pyramid itself which has a polygonal base to the pyramid which has a polygonal base



But the pyramid which has a triangular base is to the pyramid which has a triangular base in the triplicate ratio of the corresponding sides
therefore also the pyramid which has a polygonal base has to the pyramid which has a similar base the ratio triplicate of that which the side has to the side

PROPOSITION 9

In equal pyramids which have triangular bases the bases are reciprocally proportional to the heights and those pyramids in which the bases are reciprocally proportional to the heights are equal

For let there be equal pyramids which have the triangular bases ABC DEF and vertices the points G H

I say that in the pyramids $ABCG$ $DEFH$ the bases are reciprocally proportional to the heights that is as the base ABC is to the base DEF so is the height of the pyramid $DEFH$ to the height of the pyramid $ABCG$



For let the parallelepipedal solids $BGML$ $EHQP$ be completed
Now since the pyramid $ABCG$ is equal to the pyramid $DEFH$,
and the solid $BGML$ is six times the pyramid $ABCG$
and the solid $EHQP$ six times the pyramid $DEFH$
therefore the solid $BGML$ is equal to the solid $EHQP$

But in equal parallelepipedal solids the bases are reciprocally proportional to the heights
therefore as the base BM is to the base EQ so is the height of the solid $EHQP$ to the height of the solid $BGML$ [xi 31]

But as the base BM is to IQ so is the triangle ABC to the triangle DEF [i 34]

Therefore also as the triangle ABC is to the triangle DEF so is the height of the solid $IHPQ$ to the height of the solid $BGML$ [vi 11]

But the height of the solid $IHPQ$ is the same with the height of the pyramid $DEFH$

and the height of the solid $BGML$ is the same with the height of the pyramid $ABCG$

therefore as the base ABC is to the base DEF so is the height of the pyramid $DEFH$ to the height of the pyramid $ABCG$

Therefore in the pyramids $ABCG$ $DEFH$ the bases are reciprocally proportional to the heights

Next in the pyramids $ABCG$, $DEFH$ let the bases be reciprocally proportional to the heights,
that is as the base ABC is to the base DEF so let the height of the pyramid $DEFH$ be to the height of the pyramid $ABCG$,

I say that the pyramid $ABCG$ is equal to the pyramid $DEFH$

For with the same construction
since as the base ABC is to the base DEF , so is the height of the pyramid $DEFH$ to the height of the pyramid $ABCG$

while as the base ABC is to the base DEF , so is the parallelogram BM to the parallelogram EQ

therefore also, as the parallelogram BM is to the parallelogram EQ , so is the height of the pyramid $DEFH$ to the height of the pyramid $ABCG$ [v 11]

But the height of the pyramid $DEFH$ is the same with the height of the parallelepiped $EHQP$

and the height of the pyramid $ABCG$ is the same with the height of the parallelepiped $BGML$,

therefore as the base BM is to the base EQ so is the height of the parallelepiped $EHQP$ to the height of the parallelepiped $BGML$

But those parallelepipedal solids in which the bases are reciprocally proportional to the heights are equal, [xi 34]

therefore the parallelepipedal solid $BGML$ is equal to the parallelepipedal solid $EHQP$

And the pyramid $ABCG$ is a sixth part of $BGML$ and the pyramid $DEFH$ a sixth part of the parallelepiped $EHQP$,

therefore the pyramid $ABCG$ is equal to the pyramid $DEFH$


Therefore etc

Q E D

PROPOSITION 10

Any cone is a third part of the cylinder which has the same base with it and equal height

For let a cone have the same base, namely the circle $ABCD$, with a cylinder and equal height,



I say that the cone is a third part of the cylinder that is that the cylinder is triple of the cone

For if the cylinder is not triple of the cone the cylinder will be either greater than triple or less than triple of the cone

First let it be greater than triple
and let the square $ABCD$ be inscribed in the circle $ABCD$, [iv 6]
then the square $ABCD$ is greater than the half of the circle $ABCD$

From the square $ABCD$ let there be set up a prism of equal height with the cylinder

Then the prism so set up is greater than the half of the cylinder
inasmuch as if we also circumscribe a square about the circle $ABCD$ [iv 7]
the square inscribed in the circle $ABCD$ is half of that circumscribed about it,
and the solids set up from them are parallelepipedal prisms of equal height

while parallelepipedal solids which are of the same height are to one another as their bases [xi 3^d]

therefore also the prism set up on the square $ABCD$ is half of the prism set up from the square circumscribed about the circle $ABCD$

[cf xi 28 or xii 6 and 7 For]

and the cylinder is less than the prism set up from the square circumscribed about the circle $ABCD$

therefore the prism set up from the square $ABCD$ and of equal height with the cylinder is greater than the half of the cylinder

Let the circumferences AB BC CD DA be bisected at the points E , F G H

and let AE FB BF FC CG GD DH HA be joined

then each of the triangles AEB BFC CGD DHA is greater than the half of that segment of the circle $ABCD$ which is about it as we proved before

[xii 2]

On each of the triangles AEB BFC CGD DHA let prisms be set up of equal height with the cylinder

then each of the prisms so set up is greater than the half part of that segment of the cylinder which is about it

inasmuch as if we draw through the points E F G H parallels to AB BC , CD DA complete the parallelograms on AB BC CD DA and set up from them parallelepipedal solids of equal height with the cylinder the prisms on the triangles AEB BFC CGD DHA are halves of the several solids set up

and the segments of the cylinder are less than the parallelepipedal solids set up hence also the prisms on the triangles AEB BFC CGD DHA are greater than the half of the segments of the cylinder about them

Thus bisecting the circumferences that are left joining straight lines setting up on each of the triangles prisms of equal height with the cylinder, and doing this continually

we shall leave some segments of the cylinder which will be less than the excess by which the cylinder exceeds the triple of the cone

[x 1]

Let such segments be left and let them be AL LB BF , FC CG , GD DH HA

therefore the remainder the prism of which the polygon $AEBFCGDH$ is the base and the height is the same as that of the cylinder is greater than triple of the cone

But the prism of which the polygon $AEBFCGDH$ is the base and the height the same as that of the cylinder is triple of the pyramid of which the polygon $AEBFCGDH$ is the base and the vertex is the same as that of the cone

[xii 7 For]

therefore also the pyramid of which the polygon $AEBFCGDH$ is the base and the vertex is the same as that of the cone is greater than the cone which has the circle $ABCD$ as base

But it is also less for it is enclosed by it

which is impossible

Therefore the cylinder is not greater than triple of the cone

I say next that neither is the cylinder less than triple of the cone

For if possible let the cylinder be less than triple of the cone

therefore inversely the cone is greater than a third part of the cylinder

Let the square $ABCD$ be inscribed in the circle $ABCD$,
therefore the square $ABCD$ is greater than the half of the circle $ABCD$

Now let there be set up from the square $ABCD$ a pyramid having the same vertex with the cone,
therefore the pyramid so set up is greater than the half part of the cone,
seeing that as we proved before, if we circumscribe a square about the circle,
the square $ABCD$ will be half of the square circumscribed about the circle,
and if we set up from the squares parallelepipedal solids of equal height with the cone which are also called prisms the solid set up from the square $ABCD$ will be half of that set up from the square circumscribed about the circle,
for they are to one another as their bases [xi 32]

Hence also the thirds of them are in that ratio
therefore also the pyramid of which the square $ABCD$ is the base is half of the pyramid set up from the square circumscribed about the circle

And the pyramid set up from the square about the circle is greater than the cone,

for it encloses it

Therefore the pyramid of which the square $ABCD$ is the base and the vertex is the same with that of the cone is greater than the half of the cone

Let the circumferences AB , BC , CD , DA be bisected at the points E , F , G , H ,

-and let AE , EB , BF , FC , CG , GD , DH , HA be joined
therefore also each of the triangles AEB , BFC , CGD , DHA is greater than the half part of that segment of the circle $ABCD$ which is about it

Now on each of the triangles AEB , BFC , CGD , DHA let pyramids be set up which have the same vertex as the cone,
therefore also each of the pyramids so set up is in the same manner, greater than the half part of that segment of the cone which is about it

Thus by bisecting the circumferences that are left, joining straight lines setting up on each of the triangles a pyramid which has the same vertex as the cone

and doing this continually,
we shall leave some segments of the cone which will be less than the excess by which the cone exceeds the third part of the cylinder [x 1]

Let such be left and let them be the segments on AE , EB , BF , FC , CG , GD , DH , HA ,
therefore the remainder the pyramid of which the polygon $AEBFCGDH$ is the base and the vertex the same with that of the cone is greater than a third part of the cylinder

But the pyramid of which the polygon $AEBFCGDH$ is the base and the vertex the same with that of the cone is a third part of the prism of which the polygon $AEBFCGDH$ is the base and the height is the same with that of the cylinder,

therefore the prism of which the polygon $AEBFCGDH$ is the base and the height is the same with that of the cylinder is greater than the cylinder of which the circle $ABCD$ is the base

But it is also less for it is enclosed by it

which is impossible

Therefore the cylinder is not less than triple of the cone

But it was proved that neither is it greater than triple,
therefore the cylinder is triple of the cone,
hence the cone is a third part of the cylinder
Therefore etc




Q E D

PROPOSITION 11

Cones and cylinders which are of the same height are to one another as their bases

Let there be cones and cylinders of the same height,
let the circles $ABCD$ $EFGH$ be their bases AL MN their axes and AC EG
the diameters of their bases

I say that as the circle $ABCD$ is to the circle $EFGH$ so is the cone AL to the cone EN



For if not then as the circle $ABCD$ is to the circle $EFGH$ so will the cone AL be either to some solid less than the cone EN or to a greater

First let it be in that ratio to a less solid O and let the solid X be equal to that by which the solid O is less than the cone EN

therefore the cone EN is equal to the solids O X

Let the square $EFGH$ be inscribed in the circle $EFGH$

therefore the square is greater than the half of the circle

Let there be set up from the square $EFGH$ a pyramid of equal height with the cone

therefore the pyramid so set up is greater than the half of the cone inasmuch as if we circumscribe a square about the circle and set up from it a pyramid of equal height with the cone the inscribed pyramid is half of the circumscribed pyramid

for they are to one another as their bases [XII 6]

while the cone is less than the circumscribed pyramid

Let the circumferences EF IG GH HE be bisected at the points P Q R S and let HP PE IQ QI FR RG GS SH be joined

Therefore each of the triangles HPF FQF FRG GSH is greater than the half of that segment of the circle which is about it

On each of the triangles HPF EQF FRG GSH let there be set up a pyramid of equal height with the cone

therefore also each of the pyramids so set up is greater than the half of that segment of the cone which is about it

Thus bisecting the circumferences which are left joining straight lines set

ting up on each of the triangles pyramids of equal height with the cone
and doing this continually
we shall leave some segments of the cone which will be less than the solid λ [x 1]

Let such be left, and let them be the segments on $HP, PE, EQ, QF, FR, RG, GS, SH$,

therefore the remainder the pyramid of which the polygon $HPEQFRGS$ is the base and the height the same with that of the cone, is greater than the solid O

Let there also be inscribed in the circle $ABCD$ the polygon $DTAUBVCW$ similar and similarly situated to the polygon $HPEQFRGS$,

and on it let a pyramid be set up of equal height with the cone AL

Since then, as the square on AC is to the square on EG , so is the polygon $DTAUBVCW$ to the polygon $HPEQFRGS$ [xii 1]

while as the square on AC is to the square on EG , so is the circle $ABCD$ to the circle $EFGH$, [xii 2]

therefore also as the circle $ABCD$ is to the circle $EFGH$ so is the polygon $DTAUBVCW$ to the polygon $HPEQFRGS$

But as the circle $ABCD$ is to the circle $EFGH$, so is the cone AL to the solid O ,

and as the polygon $DTAUBVCW$ is to the polygon $HPEQFRGS$, so is the pyramid of which the polygon $DTAUBVCW$ is the base and the point L the vertex to the pyramid of which the polygon $HPEQFRGS$ is the base and the point N the vertex [xii 6]

Therefore also as the cone AL is to the solid O , so is the pyramid of which the polygon $DTAUBVCW$ is the base and the point L the vertex to the pyramid of which the polygon $HPEQFRGS$ is the base and the point N the vertex [v 11]

therefore alternately as the cone AL is to the pyramid in it so is the solid O to the pyramid in the cone EN [v 16]

But the cone AL is greater than the pyramid in it
therefore the solid O is also greater than the pyramid in the cone EN

But it is also less

which is absurd

Therefore the cone AL is not to any solid less than the cone EN as the circle $ABCD$ is to the circle $EFGH$

Similarly we can prove that neither is the cone EN to any solid less than the cone AL as the circle $EFGH$ is to the circle $ABCD$

I say next that neither is the cone AL to any solid greater than the cone EN as the circle $ABCD$ is to the circle $EFGH$

For if possible let it be in that ratio to a greater solid O
therefore inversely, as the circle $EFGH$ is to the circle $ABCD$ so is the solid O to the cone AL

But as the solid O is to the cone AL so is the cone EN to some solid less than the cone AL

therefore also as the circle $EFGH$ is to the circle $ABCD$ so is the cone EN to some solid less than the cone AL

which was proved impossible

Therefore the cone AL is not to any solid greater than the cone EN as the circle $ABCD$ is to the circle $EFGH$

But it was proved that neither is it in this ratio to a less solid, therefore as the circle $ABCD$ is to the circle $EFGH$, so is the cone AL to the cone EN

But as the cone is to the cone so is the cylinder to the cylinder, for each is triple of each, [xii 10]

Therefore also as the circle $ABCD$ is to the circle $EFGH$ so are the cylinders on them which are of equal height

Therefore etc


Q E D

PROPOSITION 12

Similar cones and cylinders are to one another in the triplicate ratio of the diameters in their bases

Let there be similar cones and cylinders
let the circles $ABCD$, $EFGH$ be their bases BD FH the diameters of the bases, and KL MN the axes of the cones and cylinders,

I say that the cone of which the circle $ABCD$ is the base and the point L the vertex has to the cone of which the circle $EFGH$ is the base and the point N the vertex the ratio triplicate of that which BD has to FH



For if the cone $ABCDL$ has not to the cone $EFGHN$ the ratio triplicate of that which BD has to FH
the cone $ABCDL$ will have that triplicate ratio either to some solid less than the cone $EFGHN$ or to a greater

First let it have that triplicate ratio to a less solid O

Let the square $FFCH$ be inscribed in the circle $EFGH$ [iv 6]
therefore the square $FFCH$ is greater than the half of the circle $EFGH$

Now let there be set up on the square $FFCH$ a pyramid having the same vertex with the cone

therefore the pyramid so set up is greater than the half part of the cone

Let the circumferences FF GG HH LL be bisected at the points P Q R S
and let FP PP FQ QG GR RH HS SP be joined

Therefore each of the triangles FPP FQG GRH HSL is also greater than the half part of that segment of the circle $EFGH$ which is about it

Now on each of the triangles FPP FQG GRH HSL let a pyramid be set up having the same vertex with the cone,

therefore each of the pyramids so set up is also greater than the half part of that segment of the cone which is about it

Thus bisecting the circumferences so left joining straight lines, setting up on each of the triangles pyramids having the same vertex with the cone, and doing this continually,

we shall leave some segments of the cone which will be less than the excess by which the cone $EFGHN$ exceeds the solid O [x 1]

Let such be left and let them be the segments on $LP, PF, FQ, QG, GR, RH, HS, SE$

therefore the remainder, the pyramid of which the polygon $EPFQGRHS$ is the base and the point N the vertex is greater than the solid O

Let there be also inscribed in the circle $ABCD$ the polygon $ATBUCVDW$ similar and similarly situated to the polygon $EPFQGRHS$ and let there be set up on the polygon $ATBUCVDW$ a pyramid having the same vertex with the cone,

of the triangles containing the pyramid of which the polygon $ATBUCVDW$ is the base and the point L the vertex let LBT be one, and of the triangles containing the pyramid of which the polygon $EPFQGRHS$ is the base and the point N the vertex let NFP be one, and let AT, MP be joined

Now since the cone $ABCDL$ is similar to the cone $EFGHN$, therefore as BD is to FH , so is the axis AL to the axis MN [xi Def 24]

But, as BD is to FH , so is BA to FM ,

therefore also as BA is to FM so is KL to MN

And alternately as BA is to KL , so is FM to MN [v 16]

And the sides are proportional about equal angles, namely the angles BAL, FMV ,

therefore the triangle BAL is similar to the triangle FMN [vi 6]

Again, since as BA is to AT , so is FM to MP

and they are about equal angles namely the angles BAT, FMP , inasmuch as whatever part the angle BAT is of the four right angles at the centre A the same part also is the angle FMP of the four right angles at the centre M ,

since then the sides are proportional about equal angles

therefore the triangle BAT is similar to the triangle FMP [vi 6]

Again since it was proved that, as BK is to AL so is FM to MN

while BK is equal to KT and FM to PM

therefore as TA is to KL so is PM to MN

and the sides are proportional about equal angles namely the angles TAL, PMN for they are right,

therefore the triangle LKT is similar to the triangle NMP [vi 6]

And since owing to the similarity of the triangles LKB, NMF

as LB is to BA so is NF to FM

and owing to the similarity of the triangles BKT, FMP

as KB is to BT so is MF to FP

therefore *ex aequali* as LB is to BT , so is NF to FP [v 22]

Again since owing to the similarity of the triangles LTA, NPM

as LT is to TA so is NP to PM

and owing to the similarity of the triangles TAB, PMF ,

as KT is to TB so is MP to PF ,

therefore *ex aequali* as LT is to TB so is NP to PF [v 27]

But it was also proved that as TB is to BL , so is PF to FN

Therefore *ex aequali* as TL is to LB , so is PN to NF [v 27]

Therefore in the triangles LTB , NPF the sides are proportional,

therefore the triangles LTB , NPF are equiangular [vi 5]

hence they are also similar [vi Def 1]

Therefore the pyramid of which the triangle BKT is the base and the point L the vertex is also similar to the pyramid of which the triangle FMP is the base and the point N the vertex

for they are contained by similar planes equal in multitude [vi Def 9]

But similar pyramids which have triangular bases are to one another in the triplicate ratio of their corresponding sides [xii 8]

Therefore the pyramid $BA TL$ has to the pyramid $FMPN$ the ratio triplicate of that which BA has to FM

Similarly by joining straight lines from A , B , D , V , C , U to Λ , and from E , S , H , R , G , Q to M and setting up on each of the triangles pyramids which have the same vertex with the cones

we can prove that each of the similarly arranged pyramids will also have to each similarly arranged pyramid the ratio triplicate of that which the corresponding side BA has to the corresponding side FM that is which BD has to FH

And as one of the antecedents is to one of the consequents so are all the antecedents to all the consequents [v 17]

therefore also as the pyramid $BA TL$ is to the pyramid $FMPN$ so is the whole pyramid of which the polygon $ATBUCV DW$ is the base and the point L the vertex to the whole pyramid of which the polygon $EPFQGRHS$ is the base and the point N the vertex

hence also the pyramid of which $ATBUCV DW$ is the base and the point L the vertex has to the pyramid of which the polygon $EPFQGRHS$ is the base and the point N the vertex the ratio triplicate of that which BD has to FH

But by hypothesis the cone of which the circle $ABCD$ is the base and the point L the vertex has also to the solid O the ratio triplicate of that which BD has to FH

therefore as the cone of which the circle $ABCD$ is the base and the point L the vertex is to the solid O so is the pyramid of which the polygon $ATBUCV DW$ is the base and L the vertex to the pyramid of which the polygon $EPFQGRHS$ is the base and the point N the vertex

therefore alternately as the cone of which the circle $ABCD$ is the base and L the vertex is to the pyramid contained in it of which the polygon $ATBUCV DW$ is the base and L the vertex so is the solid O to the pyramid of which the polygon $EPFQGRHS$ is the base and N the vertex [v 16]

But the said cone is greater than the pyramid in it

for it encloses it

Therefore the solid O is also greater than the pyramid of which the polygon $EPFQGRHS$ is the base and N the vertex

But it is also less

which is impossible

Therefore the cone of which the circle $ABCD$ is the base and L the vertex

has not to any solid less than the cone of which the circle $EFGH$ is the base and the point N the vertex the ratio triplicate of that which BD has to FH

Similarly we can prove that neither has the cone $EFGHN$ to any solid less than the cone $ABCDL$ the ratio triplicate of that which FH has to BD

I say next that neither has the cone $ABCDL$ to any solid greater than the cone $EFGHN$ the ratio triplicate of that which BD has to FH

For if possible let it have that ratio to a greater solid O

Therefore, inversely, the solid O has to the cone $ABCDL$ the ratio triplicate of that which FH has to BD

But as the solid O is to the cone $ABCDL$ so is the cone $EFGHN$ to some solid less than the cone $ABCDL$

Therefore the cone $EFGHN$ also has to some solid less than the cone $ABCDL$ the ratio triplicate of that which FH has to BD

which was proved impossible

Therefore the cone $ABCDL$ has not to any solid greater than the cone $EFGHN$ the ratio triplicate of that which BD has to FH

But it was proved that neither has it this ratio to a less solid than the cone $EFGHN$

Therefore the cone $ABCDL$ has to the cone $EFGHN$ the ratio triplicate of that which BD has to FH

But, as the cone is to the cone so is the cylinder to the cylinder, for the cylinder which is on the same base as the cone and of equal height with it is triple of the cone, [XII 10]

therefore the cylinder also has to the cylinder the ratio triplicate of that which BD has to FH

Therefore etc

Q E D


PROPOSITION 13

If a cylinder be cut by a plane which is parallel to its opposite planes then, as the cylinder is to the cylinder so will the axis be to the axis

For let the cylinder AD be cut by the plane GH which is parallel to the opposite planes AB CD

and let the plane GH meet the axis at the point K ,

I say that as the cylinder BG is to the cylinder GD , so is the axis EK to the axis KF



For let the axis EF be produced in both directions to the points L M

and let there be set out any number whatever of axes EN NL equal to the axis EK

and any number whatever FO OM equal to FK

and let the cylinder PW on the axis LM be conceived of which the circles PQ IJ are the bases

Let planes be carried through the points N , O parallel to AB CD and to the bases of the cylinder PW

and let them produce the circles RS TU about the centres N , O

Then since the axes LN NE EK are equal to one another,

therefore the cylinders QR , RB , BG are to one another as their bases [xii 11]

But the bases are equal,

therefore the cylinders QR , RB , BG are also equal to one another

Since then the axes LN , NE , EK are equal to one another

and the cylinders QR , RB , BG are also equal to one another, and the multitude of the former is equal to the multitude of the latter, therefore whatever multiple the axis KL is of the axis EK , the same multiple also will the cylinder QG be of the cylinder BG

For the same reason whatever multiple the axis MK is of the axis KF , the same multiple also is the cylinder MG of the cylinder GD

And if the axis KL is equal to the axis KM , the cylinder QG will also be equal to the cylinder GW ,

if the axis is greater than the axis the cylinder will also be greater than the cylinder

and if less less

Thus there being four magnitudes the axes EK , KF and the cylinders BG , GD

there have been taken equimultiples of the axis EK and of the cylinder BG , namely the axis LA and the cylinder QG ,

and equimultiples of the axis KF and of the cylinder GD , namely the axis LM and the cylinder GW ,

and it has been proved that,

if the axis KL is in excess of the axis KM the cylinder QG is also in excess of the cylinder GW ,

if equal equal

and if less less

Therefore as the axis EK is to the axis KF so is the cylinder BG to the cylinder GD

[v Def 5]

q.e.d.

PROPOSITION 14

Cones and cylinders which are on equal bases are to one another as their heights

For let EB , FD be cylinders on equal bases, the circles AB , CD

I say that as the cylinder EB is to the cylinder FD so is the axis GH to the axis KL

For let the axis KL be produced to the point N

let LN be made equal to the axis GH , and let the cylinder CM be conceived about LN as axis


Since then the cylinders EB , CM are of the same height they are to one another as their bases [xii 11]

But the bases are equal to one another

therefore the cylinders EB , CM are also equal

And since the cylinder FM has been cut by the plane CD which is parallel to its opposite planes

therefore as the cylinder CM is to the cylinder FD so is the axis LN to the axis KL [xii 13]



But the cylinder CM is equal to the cylinder EB ,
 and the axis LN to the axis GH ,
 therefore as the cylinder EB is to the cylinder FD , so is the axis GH to the axis KL

But as the cylinder EB is to the cylinder FD , so is the cone ABG to the cone CDA [xi 10]

Therefore also as the axis GH is to the axis KL so is the cone ABG to the cone CDA and the cylinder EB to the cylinder FD Q E D


PROPOSITION 15

In equal cones and cylinders the bases are reciprocally proportional to the heights and those cones and cylinders in which the bases are reciprocally proportional to the heights are equal

Let there be equal cones and cylinders of which the circles $ABCD$, $EFGH$ are the bases,

let AC , EG be the diameters of the bases
 and AL MN the axes which are also the heights of the cones or cylinders,
 let the cylinders AO , EP be completed

I say that in the cylinders AO , EP the bases are reciprocally proportional to the heights,
 that is as the base $ABCD$ is to the base $EFGH$ so is the height MN to the height AL



For the height LK is either equal to the height MA or not equal

First let it be equal

Now the cylinder AO is also equal to the cylinder EP

But cones and cylinders which are of the same height are to one another as their bases [xii 11]

therefore the base $ABCD$ is also equal to the base $EFGH$

Hence also reciprocally as the base $ABCD$ is to the base $EFGH$, so is the height MN to the height KL

Next, let the height LK not be equal to MN ,
 but let MN be greater,

from the height MN let QN be cut off equal to KL ,
 through the point Q let the cylinder EP be cut by the plane TUS parallel to the planes of the circles $EFGH$ RP
 and let the cylinder ES be conceived erected from the circle $EFGH$ as base and with height NQ

Now since the cylinder AO is equal to the cylinder EP
 therefore as the cylinder AO is to the cylinder ES so is the cylinder EP to the cylinder ES [v 7]

But as the cylinder AO is to the cylinder ES , so is the base $ABCD$ to the base $EFGH$,

for the cylinders AO , ES are of the same height [xi 11]

and, as the cylinder EP is to the cylinder ES so is the height MN to the height QV ,

for the cylinder EP has been cut by a plane which is parallel to its opposite planes [xi 13]

Therefore also as the base $ABCD$ is to the base $EFGH$, so is the height MV to the height QN [v 11]

But the height QV is equal to the height KL ,
therefore as the base $ABCD$ is to the base $EFGH$, so is the height MV to the height KL

Therefore in the cylinders AO , EP the bases are reciprocally proportional to the heights

Next in the cylinders AO , EP let the bases be reciprocally proportional to the heights

that is as the base $ABCD$ is to the base $EFGH$, so let the height MA be to the height KL

I say that the cylinder AO is equal to the cylinder EP

For with the same construction

since as the base $ABCD$ is to the base $EFGH$ so is the height MN to the height KL ,

while the height KL is equal to the height QV

therefore as the base $ABCD$ is to the base $EFGH$, so is the height MV to the height QV

But as the base $ABCD$ is to the base $EFGH$ so is the cylinder AO to the cylinder ES

for they are of the same height [xi 11]

and as the height MA is to QV so is the cylinder EP to the cylinder ES [xi 13]

therefore as the cylinder AO is to the cylinder ES so is the cylinder EP to the cylinder ES [v 11]

Therefore the cylinder AO is equal to the cylinder EP [v 9]

And the same is true for the cones also

Q E D

PROPOSITION 16


Given two circles about the same centre to inscribe in the greater circle an equilateral polygon with an even number of sides which does not touch the lesser circle

Let $ABCD$, $EFGH$ be the two given circles about the same centre K

thus it is required to inscribe in the greater circle $ABCD$ an equilateral polygon with an even number of sides which does not touch the circle $EFGH$

For let the straight line BAD be drawn through the centre K

and from the point G let GA be drawn at right angles to the straight line BD and carried through to C



therefore AC touches the circle $EFGH$ [III 16 Por.]

Then bisecting the circumference BAD bisecting the half of it and doing this continually, we shall leave a circumference less than AD [5.1]

Let such be left and let it be LD ,
from L let LM be drawn perpendicular to BD and carried through to N
and let LD DN be joined
therefore LD is equal to DN [III 3, 1 4]


Now, since LN is parallel to AC ,
and AC touches the circle $EFGH$,
therefore LN does not touch the circle $EFGH$
therefore LD DN are far from touching the circle $EFGH$

If then we fit into the circle $ABCD$ straight lines equal to the straight line LD and placed continuously, there will be inscribed in the circle $ABCD$ an equilateral polygon with an even number of sides which does not touch the lesser circle $EFGH$ Q E F

PROPOSITION 17

Given two spheres about the same centre, to inscribe in the greater sphere a polyhedral solid which does not touch the lesser sphere at its surface

Let two spheres be conceived about the same centre A
thus it is required to inscribe in the greater sphere a polyhedral solid which
does not touch the lesser sphere at its surface



Let the spheres be cut by any plane through the centre
then the sections will be circles

inasmuch as the sphere was produced by the diameter remaining fixed and the semicircle being carried round it, [xi Def 14]

hence in whatever position we conceive the semicircle to be the plane carried through it will produce a circle on the circumference of the sphere

And it is manifest that this circle is the greatest possible inasmuch as the diameter of the sphere which is of course the diameter both of the semicircle and of the circle is greater than all the straight lines drawn across in the circle or the sphere

Let then $BCDE$ be the circle in the greater sphere and FGH the circle in the lesser sphere let two diameters in them BD CE be drawn at right angles to one another, then given the two circles $BCDE$ FGH about the same centre, let there be inscribed in the greater circle $BCDE$ an equilateral polygon with an even number of sides which does not touch the lesser circle FGH

let BA AL LM ME be its sides in the quadrant BE , let AI be joined and carried through to N let AO be set up from the point A at right angles to the plane of the circle $BCDE$ and let it meet the surface of the sphere at O and through IO and each of the straight lines BD AN let planes be carried they will then make greatest circles on the surface of the sphere for the reason stated

Let them make such and in them let BOD AOV be the semicircles on BD AN Now since OI is at right angles to the plane of the circle $BCDE$ therefore all the planes through OI are also at right angles to the plane of the circle $BCDE$ [xi 18] hence the semicircles BOD AOV are also at right angles to the plane of the circle $BCDE$

And since the semicircles BED BOD AON are equal for they are on the equal diameters BD AN therefore the quadrants BL BO AO are also equal to one another Therefore there are as many straight lines in the quadrants BO AO equal to the straight lines BA AL LM ME as there are sides of the polygon in the quadrant BE

Let them be inscribed and let them be BI PQ QR RO and AS ST , TU LO

let SP TQ UR be joined and from P Q let perpendiculars be drawn to the plane of the circle $BCDE$ [xi 11]

these will fall on BD AN the common sections of the planes inasmuch as the planes of BOD AOV are also at right angles to the plane of the circle $BCDE$ [cf xi Def 4]

Let them so fall and let them be PV SB and let BI be joined Now since in the equal semicircles BOD AOV equal straight lines BP , AS have been cut off

and the perpendiculars PV SB have been drawn therefore PI is equal to SB and BI to AB [iii 2^o, 1^o] But the whole BI is also equal to the whole AI

therefore the remainder VA is also equal to the remainder WA ,

therefore as BV is to VA so is KW to WA

therefore WV is parallel to KB [VI 2]

And since each of the straight lines PV , SW is at right angles to the plane of the circle $BCDF$,

therefore PV is parallel to SW [XI 6]

But it was also proved equal to it

therefore WV , SP are also equal and parallel [I 33]

And, since WV is parallel to SP ,

while WV is parallel to KB ,

therefore SP is also parallel to KB [XI 9]

And BP , KS join their extremities,

therefore the quadrilateral $KBPS$ is in one plane

inasmuch as if two straight lines be parallel, and points be taken at random on each of them, the straight line joining the points is in the same plane with the parallels [XI 7]

For the same reason

each of the quadrilaterals $SPQT$, $TQRU$ is also in one plane

But the triangle URO is also in one plane [XI 2]

If then we conceive straight lines joined from the points P , S , Q , T , R , U to A , there will be constructed a certain polyhedral solid figure between the circumferences BO , KO , consisting of pyramids of which the quadrilaterals $KBPS$, $SPQT$, $TQRU$ and the triangle URO are the bases and the point A the vertex

And if we make the same construction in the case of each of the sides AL , LM , ME as in the case of BK and further in the case of the remaining three quadrants

there will be constructed a certain polyhedral figure inscribed in the sphere and contained by pyramids, of which the said quadrilaterals and the triangle URO , and the others corresponding to them are the bases and the point A the vertex

I say that the said polyhedron will not touch the lesser sphere at the surface on which the circle FGH is

Let AX be drawn from the point A perpendicular to the plane of the quadrilateral $KBPS$ and let it meet the plane at the point X [XI 11]

let AB , XA be joined

Then since AX is at right angles to the plane of the quadrilateral $KBPS$, therefore it is also at right angles to all the straight lines which meet it and are in the plane of the quadrilateral [XI Def 3]

Therefore AX is at right angles to each of the straight lines BX , XA

And since AB is equal to AA

the square on AB is also equal to the square on AA

And the squares on AX , XB are equal to the square on AB

for the angle at X is right [I 47]

and the squares on AX , XA are equal to the square on AA [Id.]

Therefore the squares on AX , XB are equal to the squares on AX , XA

Let the square on AX be subtracted from each

therefore the remainder the square on BX is equal to the remainder the square on XA ,

therefore BX is equal to XA

Similarly we can prove that the straight lines joined from Λ to P S are equal to each of the straight lines $B\Lambda$ ΛK

Therefore the circle described with centre Λ and distance one of the straight lines ΛB , ΛA will pass through P S also

and ΛBPS will be a quadrilateral in a circle

Now, since ΛB is greater than $H V$

while $H V$ is equal to SP

therefore ΛB is greater than SP

But ΛB is equal to each of the straight lines ΛS BP

therefore each of the straight lines ΛS BP is greater than SP

And since ΛBPS is a quadrilateral in a circle

and KB BP , ΛS are equal and PS less

and $B\Lambda$ is the radius of the circle

therefore the square on KB is greater than double of the square on $B\Lambda$

Let ΛZ be drawn from Λ perpendicular to BV

Then since BD is less than double of DZ

and as BD is to DZ so is the rectangle DB BZ to the rectangle DZ ZB
if a square be described upon BZ and the parallelogram on ZD be completed
then the rectangle DB BZ is also less than double of the rectangle DZ ZB

And if ΛD be joined

the rectangle DB BZ is equal to the square on BK ,

and the rectangle DZ ZB equal to the square on ΛZ (III 31 vi 8 and Per)

therefore the square on ΛB is less than double of the square on ΛZ

But the square on ΛB is greater than double of the square on $B\Lambda$

therefore the square on ΛZ is greater than the square on $B\Lambda$

And since $B\Lambda$ is equal to ΛI

the square on BA is equal to the square on ΛA

And the squares on $B\Lambda$ ΛA are equal to the square on BA

and the squares on ΛZ ZA equal to the square on ΛA , (I 4)

therefore the squares on $B\Lambda$ ΛA are equal to the squares on ΛZ ZA

and of these the square on ΛZ is greater than the square on $B\Lambda$

therefore the remainder the square on ZA is less than the square on ΛA

Therefore ΛA is greater than ΛZ

therefore $\Lambda \Lambda$ is much greater than AG

And $\Lambda \Lambda$ is the perpendicular on one base of the polyhedron

and AG on the surface of the lesser sphere

hence the polyhedron will not touch the lesser sphere on its surface

Therefore given two spheres about the same centre a polyhedral solid has been inscribed in the greater sphere which does not touch the lesser sphere at its surface

PROPOSITION But if in another sphere also a polyhedral solid be inscribed similar to the solid in the sphere $BCDE$

the polyhedral solid in the sphere $BCDE$ has to the polyhedral solid in the other sphere the ratio triplicate of that which the diameter of the sphere $BCDE$ has to the diameter of the other sphere

For the solids being divided into their pyramids similar in multitude and arrangement the pyramids will be similar

But similar pyramids are to one another in the triplicate ratio of their corresponding sides (XII 8 Per)

therefore the pyramid of which the quadrilateral $KBPS$ is the base and the point A the vertex has to the similarly arranged pyramid in the other sphere the ratio triplicate of that which the corresponding side has to the corresponding side that is of that which the radius AB of the sphere about A as centre has to the radius of the other sphere

Similarly also each pyramid of those in the sphere about A as centre has to each similarly arranged pyramid of those in the other sphere the ratio triplicate of that which AB has to the radius of the other sphere

And as one of the antecedents is to one of the consequents, so are all the antecedents to all the consequents, [v 12]

hence the whole polyhedral solid in the sphere about A as centre has to the whole polyhedral solid in the other sphere the ratio triplicate of that which AB has to the radius of the other sphere that is of that which the diameter BD has to the diameter of the other sphere Q E D

PROPOSITION 18

Spheres are to one another in the triplicate ratio of their respective diameters

Let the spheres ABC , DEF be conceived


and let BC , EF be their diameters,

I say that the sphere ABC has to the sphere DEF the ratio triplicate of that which BC has to EF

For if the sphere ABC has not to the sphere DEF the ratio triplicate of that which BC has to EF

then the sphere ABC will have either to some less sphere than the sphere DEF , or to a greater the ratio triplicate of that which BC has to EF

First let it have that ratio to a less sphere GHA ,



let DEF be conceived about the same centre with GHA ,
let there be inscribed in the greater sphere DEF a polyhedral solid which does not touch the lesser sphere GHA at its surface [xii 17]

and let there also be inscribed in the sphere ABC a polyhedral solid similar to the polyhedral solid in the sphere DEF ,
 therefore the polyhedral solid in ABC has to the polyhedral solid in DEF the ratio triplicate of that which BC has to EF [xii 17, Per]

But the sphere ABC also has to the sphere GHK the ratio triplicate of that which BC has to EF ,
 therefore as the sphere ABC is to the sphere GHK , so is the polyhedral solid in the sphere ABC to the polyhedral solid in the sphere DEF ,
 and alternately as the sphere ABC is to the polyhedron in it so is the sphere GHK to the polyhedral solid in the sphere DEF [v 16]

But the sphere ABC is greater than the polyhedron in it
 therefore the sphere GHA is also greater than the polyhedron in the sphere DEF

But it is also less

for it is enclosed by it

Therefore the sphere ABC has not to a less sphere than the sphere DEF the ratio triplicate of that which the diameter BC has to EF

Similarly we can prove that neither has the sphere DEF to a less sphere than the sphere ABC the ratio triplicate of that which EF has to BC

I say next that neither has the sphere ABC to any greater sphere than the sphere DEF the ratio triplicate of that which BC has to EF

For if possible let it have that ratio to a greater LMN ,
 therefore inversely the sphere LMN has to the sphere ABC the ratio triplicate of that which the diameter EF has to the diameter BC

But inasmuch as LMN is greater than DEF
 therefore as the sphere LMN is to the sphere ABC so is the sphere DEF to some less sphere than the sphere ABC as was before proved [xii 2 Lemma]

Therefore the sphere DEF also has to some less sphere than the sphere ABC the ratio triplicate of that which EF has to BC

which was proved impossible

Therefore the sphere ABC has not to any sphere greater than the sphere DEF the ratio triplicate of that which BC has to EF

But it was proved that neither has it that ratio to a less sphere

Therefore the sphere ABC has to the sphere DEF the ratio triplicate of that which BC has to EF Q E D

BOOK THIRTEEN

PROPOSITIONS

PROPOSITION 1

If a straight line be cut in extreme and mean ratio the square on the greater segment added to the half of the whole is five times the square on the half

For let the straight line AB be cut in extreme and mean ratio at the point C ,
and let AC be the greater segment,

let the straight line AD be produced in a straight line with CA
and let AD be made half of AB ,

I say that the square on CD is five times the square on AD

For let the squares AE , DF be described on AB , DC

and let the figure in DF be drawn,
let FC be carried through to G

Now, since AB has been cut in extreme and mean ratio at C ,
therefore the rectangle AB BC is equal to the square on AC [vi Def 3 vi 17]

And CE is the rectangle AB , BC and FH the square on AC ,

therefore CE is equal to FH

And since BA is double of AD ,
while BA is equal to KA and AD to AH ,
therefore KA is also double of AH

But as KA is to AH so is CK to CH , [vi 1]
therefore CK is double of CH

But LH , HC are also double of CH

Therefore KC is equal to LH HC

But CE was also proved equal to HF

therefore the whole square AE is equal to the gnomon MNO

And since BA is double of AD

the square on BA is quadruple of the square on AD ,

that is, AE is quadruple of DH

But AE is equal to the gnomon MNO

therefore the gnomon MNO is also quadruple of AP


therefore the whole DF is five times AP

And DF is the square on DC , and AP the square on DA ,

therefore the square on CD is five times the square on DA

Therefore etc

Q E D



PROPOSITION 2

If the square on a straight line be five times the square on a segment of it then, when the double of the said segment is cut in extreme and mean ratio the greater segment is the remaining part of the original straight line

For let the square on the straight line AB be five times the square on the segment AC of it

and let CD be double of AC .

I say that when CD is cut in extreme and mean ratio the greater segment is CB

Let the squares AF CG be described on AB CD respectively

let the figure in AF be drawn

and let BE be drawn through

Now since the square on BA is five times the square on AC

AF' is five times AH

Therefore the gnomon MNO is quadruple of AH

And since DC is double of CA
therefore the square on DC is quadruple of the
square on CA that is CG is quadruple of AH

But the gnomon MNO was also proved quadruple of AM

therefore the gnomon MNO is equal to CG

And since DC is double of CA

while DC is equal to CH and AC to CH

therefore AB is also double of BH

But LH HB are also double of HB

therefore AB is equal to $LH + HB$

But the whole gnomon $MANO$ was also proved equal to the whole CG ,

therefore the remainder HF is equal to BG

And BG is the rectangle $\{D, DB$

for CD is equal to DG ,

and HF is the square on CB

therefore the rectangle $CD \cdot DB$ is equal to the square on CB

Therefore as DC is to CB so is C to BD


But DC is greater than CB

therefore $\angle B$ is also greater than $\angle D$

Therefore when the straight line CD is cut in extreme and mean ratio CB is the greater segment

Therefore etc

Q E D



LEMMA

That the double of AC is greater than BC is to be proved thus

If not let BC be if possible double of CA

Therefore the square on BC is quadruple of the square on C'

therefore the squares on BC CA are five times the square on CA

But by hypothesis the square on BA is also five times the square on CA .

therefore the square on BA is equal to the squares on BC , CA
which is impossible

(II 4)

Therefore CB is not double of AC

Similarly we can prove that neither is a straight line less than CB double of CA ,

for the absurdity is much greater

Therefore the double of AC is greater than CB

Q E D

PROPOSITION 3

If a straight line be cut in extreme and mean ratio the square on the lesser segment added to the half of the greater segment is five times the square on the half of the greater segment

For let any straight line AB be cut in extreme and mean ratio at the point C ,

let AC be the greater segment

and let AC be bisected at D ,

I say that the square on BD is five times the square on DC

For let the square AE be described on AB
and let the figure be drawn double

Since AC is double of DC

therefore the square on AC is quadruple of the square on DC ,

that is RS is quadruple of FG

And since the rectangle AB , BC is equal to the square on AC

and CE is the rectangle AB , BC

therefore CE is equal to RS

But RS is quadruple of FG

therefore CE is also quadruple of FG

Again since AD is equal to DC

HK is also equal to KF

Hence the square GF is also equal to the square HL

Therefore GK is equal to KL that is MN to NE ,

hence MF is also equal to FE

But MF is equal to CG ,

therefore CG is also equal to FE

Let CA be added to each

therefore the gnomon OPQ is equal to CE

But CE was proved quadruple of GF ,

therefore the gnomon OPQ is also quadruple of the square FG

Therefore the gnomon OPQ and the square FG are five times FG

But the gnomon OPQ and the square FG are the square DN

And DN is the square on DB and GF the square on DC

Therefore the square on DB is five times the square on DC

Q E D

PROPOSITION 4

If a straight line be cut in extreme and mean ratio the square on the whole and the square on the lesser segment together are triple of the square on the greater segment

Let AB be a straight line,

let it be cut in extreme and mean ratio at C , and let AC be the greater segment,

I say that the squares on AB , BC are triple of the square on CA

For let the square $ADEB$ be described on AB
and let the figure be drawn

Since then, AB has been cut in extreme and mean ratio at C ,

and AC is the greater segment

therefore the rectangle AB , BC is equal to the square on AC [vi Def 3 vi 17]

And AK is the rectangle AB , BC and HG the square on AC ,

therefore AK is equal to HG

And since AF is equal to FE

let CK be added to each,

therefore the whole AK is equal to the whole CE ,

therefore AK , CE are double of AK

But AK , CE are the gnomon LMN and the square CA

therefore the gnomon LMN and the square CK are double of AK

But further AK was also proved equal to HG


therefore the gnomon LMN and the squares CK , HG are triple of the square HG

And the gnomon LMN and the squares CA , HG are the whole square AE and CA , which are the squares on AB , BC

while HG is the square on AC

Therefore the squares on AB , BC are triple of the square on AC

Q E D



PROPOSITION 5

If a straight line be cut in extreme and mean ratio and there be added to it a straight line equal to the greater segment the whole straight line has been cut in extreme and mean ratio and the original straight line is the greater segment

For let the straight line AB be cut in extreme and mean ratio at the point C , let AC be the greater segment and let AD be equal to AC

I say that the straight line DB has been cut in extreme and mean ratio at A and the original straight line AB is the greater segment

For let the square AE be described on AB and let the figure be drawn


Since AB has been cut in extreme and mean ratio at C

therefore the rectangle AB , BC is equal to the square on AC [vi Def 3 vi 17]

And CE is the rectangle AB , BC and CH the square on AC ,

therefore CE is equal to HC

But HE is equal to CE



and DH is equal to HC ,

therefore DH is also equal to HE

Therefore the whole DK is equal to the whole AE

And DK is the rectangle BD, DA ,

for AD is equal to DL ,

and AE is the square on AB ,

therefore the rectangle BD, DA is equal to the square on AB

Therefore as DB is to BA , so is BA to AD [vi 17]

And DB is greater than BA ,

therefore BA is also greater than AD [v 14]

Therefore DB has been cut in extreme and mean ratio at A and AB is the greater segment Q E D

PROPOSITION 6

If a rational straight line be cut in extreme and mean ratio, each of the segments is the irrational straight line called apotome

Let AB be a rational straight line

let it be cut in extreme and mean ratio at C ,

and let AC be the greater segment

I say that each of the straight lines AC, CB is the irrational straight line called apotome

For let BA be produced and let AD be made half of BA

Since then the straight line AB has been cut in extreme and mean ratio

and to the greater segment AC is added AD which is half of AB

therefore the square on CD is five times the square on DA [xiii 1]

Therefore the square on CD has to the square on DA the ratio which a number has to a number

therefore the square on CD is commensurable with the square on DA [x 6]

But the square on DA is rational

for DA is rational, being half of AB which is rational

therefore the square on CD is also rational, [x Def 4]

therefore CD is also rational

And, since the square on CD has not to the square on DA the ratio which a square number has to a square number,

therefore CD is incommensurable in length with DA , [x 9]

therefore CD, DA are rational straight lines commensurable in square only

therefore AC is an apotome [x 73]

Again since AB has been cut in extreme and mean ratio

and AC is the greater segment,

therefore the rectangle AB, BC is equal to the square on AC [vi Def 3 vi 17]

Therefore the square on the apotome AC if applied to the rational straight line AB produces BC as breadth

But the square on an apotome if applied to a rational straight line produces as breadth a first apotome [x 97]

therefore CB is a first apotome

And CA was also proved to be an apotome

Therefore etc

Q E D

PROPOSITION 7

If three angles of an equilateral pentagon, taken either in order or not in order be equal, the pentagon will be equiangular

For in the equilateral pentagon $ABCDE$ let, first three angles taken in order those at A, B, C be equal to one another,

I say that the pentagon $ABCDE$ is equiangular

For let AC, BE, FD be joined

Now since the two sides CB, BA are equal to the two sides BA, AE respectively,

and the angle CBA is equal to the angle BAE

therefore the base AC is equal to the base BE ,

the triangle ABC is equal to the triangle ABE

and the remaining angles will be equal to the remaining angles namely those which the equal sides subtend [1 4]

that is the angle BCA to the angle BEA and the angle ABE to the angle CAB ,

hence the side AF is also equal to the side BF [1 6]

But the whole AC was also proved equal to the whole BE

therefore the remainder FC is also equal to the remainder FE

But CD is also equal to DE

Therefore the two sides FC, CD are equal to the two sides FE, ED ,

and the base FD is common to them

therefore the angle FCD is equal to the angle FED [1 8]

But the angle BCA was also proved equal to the angle AEB ,

therefore the whole angle BCD is also equal to the whole angle AED

But by hypothesis the angle BCD is equal to the angles at A, B

therefore the angle AED is also equal to the angles at A, B

Similarly we can prove that the angle CDE is also equal to the angles at A, B, C

therefore the pentagon $ABCDE$ is equiangular

Next let the given equal angles not be angles taken in order, but let the angles at the points A, C, D be equal

I say that in this case too the pentagon $ABCDE$ is equiangular

For let BD be joined

Then since the two sides BA, AE are equal to the two sides BC, CD

and they contain equal angles

therefore the base BE is equal to the base BD

the triangle ABE is equal to the triangle BCD

and the remaining angles will be equal to the remaining angles, namely those which the equal sides subtend [1 4]

therefore the angle AEB is equal to the angle CDB

But the angle BFD is also equal to the angle BDE [1 5]

since the side BE is also equal to the side BD

Therefore the whole angle AED is equal to the whole angle CDE


But the angle CDE is by hypothesis equal to the angles at A, C ,

therefore the angle AED is also equal to the angles at A, C

For the same reason

the angle ABC is also equal to the angles at A, C, D


Therefore the pentagon $ABCDE$ is equiangular



PROPOSITION 8

If in an equilateral and equiangular pentagon straight lines subtend two angles taken in order, they cut one another in extreme and mean ratio and their greater segments are equal to the side of the pentagon

For in the equilateral and equiangular pentagon $ABCDE$ let the straight lines AC , BE , cutting one another at the point H subtend two angles taken in order, the angles at A B



I say that each of them has been cut in extreme and mean ratio at the point H and their greater segments are equal to the side of the pentagon

For let the circle $ABCDE$ be circumscribed about the pentagon $ABCDE$ [iv 14]

Then since the two straight lines EA , AB are equal to the two AB BC , and they contain equal angles

therefore the base BE is equal to the base AC

the triangle ABE is equal to the triangle ABC

and the remaining angles will be equal to the remaining angles respectively namely those which the equal sides subtend [i 4]

Therefore the angle BAC is equal to the angle ABE ,

therefore the angle AHE is double of the angle BAH [i 32]

But the angle EAC is also double of the angle BAC

inasmuch as the circumference EDC is also double of the circumference CB [iii 28 vi 33]

therefore the angle HAE is equal to the angle AHE

hence the straight line HE is also equal to EA that is to AB [i 6]

And since the straight line BA is equal to AE

the angle ABE is also equal to the angle AEB [i 5]

But the angle ABE was proved equal to the angle BAH

therefore the angle BEH is also equal to the angle BAH

And the angle ABE is common to the two triangles ABE and ABH , therefore the remaining angle BAE is equal to the remaining angle AHB [i 32]

therefore the triangle ABE is equiangular with the triangle ABH

therefore proportionally as EB is to BA so is AB to BH [vi 4]

But BA is equal to EH

therefore as BE is to EH so is EH to HB

And BE is greater than EH

therefore EH is also greater than HB [v 14]

Therefore BE has been cut in extreme and mean ratio at H and the greater segment HE is equal to the side of the pentagon

Similarly we can prove that AC has also been cut in extreme and mean ratio at H and its greater segment CH is equal to the side of the pentagon Q.E.D.

PROPOSITION 9

If the side of the hexagon and that of the decagon inscribed in the same circle be added together the whole straight line has been cut in extreme and mean ratio and its greater segment is the side of the hexagon

squares on the side of the hexagon and on that of the decagon inscribed in the circle $ABCDE$

For let the centre of the circle, the point F , be taken

let AF be joined and carried through to the point G ,

let FB be joined

let FH be drawn from F perpendicular to AB and be carried through to K ,

let AK , KB be joined

let FL be again drawn from F perpendicular to AK and be carried through to M ,

and let AN be joined

Since the circumference $ABCG$ is equal to the circumference $AEDG$

and in them ABC is equal to AED

therefore the remainder the circumference

CG is equal to the remainder GD

But CD belongs to a pentagon,

therefore CG belongs to a decagon

And since FA is equal to FB

and FH is perpendicular,

therefore the angle AFK is also equal to the angle KFB [I 5, I 26]

Hence the circumference AK is also equal to KB [III 26]

therefore the circumference AB is double of the circumference BA ,

therefore the straight line AK is a side of a decagon

For the same reason

AK is also double of KM

Now since the circumference AB is double of the circumference BA ,

while the circumference CD is equal to the circumference AB

therefore the circumference CD is also double of the circumference BK

But the circumference CD is also double of CG

therefore the circumference CG is equal to the circumference BK

But BK is double of KM since KA is also

therefore CG is also double of KM

But, further the circumference CB is also double of the circumference BA ,

for the circumference CB is equal to BA

Therefore the whole circumference GB is also double of BM

hence the angle GFB is also double of the angle BFM [VI 33]

But the angle GFB is also double of the angle FAB

for the angle FAB is equal to the angle ABF

Therefore the angle BFN is also equal to the angle FAB

But the angle ABF is common to the two triangles ABF and BFN ,

therefore the remaining angle AFB is equal to the remaining angle BNF [I 32]


therefore the triangle ABF is equiangular with the triangle BFN

Therefore proportionally as the straight line AB is to BF so is FB to BN

[VI 4]

therefore the rectangle $AB BN$ is equal to the square on BF [VI 17]

Again since AL is equal to LA ,



while LA is common and at right angles

therefore the base KN is equal to the base AN [1 4]

therefore the angle LKN is also equal to the angle LNA

But the angle LNA is equal to the angle LBN ,

therefore the angle LKN is also equal to the angle LBN

And the angle at A is common to the two triangles AKB and ALN

Therefore the remaining angle AKB is equal to the remaining angle ALN [1 32]

therefore the triangle AKB is equiangular with the triangle ALN

Therefore proportionally as the straight line BA is to AK , so is LA to AN [VI 4]

therefore the rectangle BA, AN is equal to the square on LA [VI 17]

But the rectangle AB, BN was also proved equal to the square on BF

therefore the rectangle AB, BN together with the rectangle BA, AN that is the square on BA [II 2] is equal to the square on BF together with the square on LA

And BA is a side of the pentagon BF of the hexagon [IV 15 Por] and LA of the decagon

Therefore etc

Q E D

PROPOSITION 11

If in a circle which has its diameter rational an equilateral pentagon be inscribed the side of the pentagon is the irrational straight line called minor

For in the circle $ABCDE$ which has its diameter rational let the equilateral pentagon $ABCDE$ be inscribed

I say that the side of the pentagon is the irrational straight line called minor

For let the centre of the circle the point F be taken

let AF, FB be joined and carried through to the points G, H

let AC be joined

and let FK be made a fourth part of AF

Now AF is rational

therefore FK is also rational

But BF is also rational

therefore the whole BA is rational

And since the circumference ACG is equal to the circumference ADG

and in them ABC is equal to AED

therefore the remainder CG is equal to the remainder GD


And if we join AD we conclude that the angles at L are right and CD is double of CL

For the same reason

the angles at M are also right

and AC is double of CM

Since then the angle ALC is equal to the angle AMF ,



and the angle LAC is common to the two triangles ACL and AMF
 therefore the remaining angle ACL is equal to the remaining angle MFA , [I 32]

therefore the triangle ACL is equiangular with the triangle AMF ,

therefore, proportionally as LC is to CA , so is MF to FA

And the doubles of the antecedents may be taken,

therefore as the double of LC is to CA , so is the double of MF to FA

But as the double of MF is to FA , so is MF to the half of FA ,

therefore also as the double of LC is to CA , so is MF to the half of FA

And the halves of the consequents may be taken,

therefore as the double of LC is to the half of CA , so is MF to the fourth of FA

And DC is double of LC CM is half of CA , and FK a fourth part of FA ,

therefore as DC is to CM so is MF to FK

Componendo also as the sum of DC CM is to CM , so is MA to KF , [V 18]
 therefore also as the square on the sum of DC , CM is to the square on CM , so is the square on MA to the square on KF

And since when the straight line subtending two sides of the pentagon as AC is cut in extreme and mean ratio the greater segment is equal to the side of the pentagon that is to DC [XIII 8]

while the square on the greater segment added to the half of the whole is five times the square on the half of the whole [XIII 1]

and CM is half of the whole AC ,

therefore the square on DC CM taken as one straight line is five times the square on CM

But it was proved that as the square on DC CM taken as one straight line is to the square on CM so is the square on MA to the square on KF

therefore the square on MA is five times the square on KF

But the square on KF is rational

for the diameter is rational

therefore the square on MA is also rational

therefore MA is rational

And since BF is quadruple of FK

therefore BK is five times KF

therefore the square on BA is twenty five times the square on KF

But the square on MA is five times the square on KF

therefore the square on BA is five times the square on MA

therefore the square on BA has not to the square on MA the ratio which a square number has to a square number

therefore BA is incommensurable in length with MA [X 9]

And each of them is rational

Therefore BK MA are rational straight lines commensurable in square only

But if from a rational straight line there be subtracted a rational straight line which is commensurable with the whole in square only the remainder is irrational namely an apotome

therefore MB is an apotome and MA the annex to it [X 73]

I say next that MB is also a fourth apotome

Let the square on A be equal to that by which the square on BA is greater than the square on MA

therefore the square on BK is greater than the square on MA by the square on N

And, since KF is commensurable with FB ,

componendo also KB is commensurable with FB [x 15]

But BF is commensurable with BH ,

therefore BK is also commensurable with BH [x 12]

And, since the square on BK is five times the square on AM ,
therefore the square on BA has to the square on KM the ratio which 5 has to 1

Therefore, *convertendo* the square on BK has to the square on N the ratio which 5 has to 4 [v 19 Por], and this is not the ratio which a square number has to a square number

therefore BK is incommensurable with N , [x 9]

therefore the square on BA is greater than the square on KM by the square on a straight line incommensurable with BK

Since then the square on the whole BK is greater than the square on the annex KM by the square on a straight line incommensurable with BK ,
and the whole BA is commensurable with the rational straight line, BH , set out

therefore MB is a fourth apotome [x Def III 4]

But the rectangle contained by a rational straight line and a fourth apotome is irrational

and its square root is irrational and is called minor [x 94]

But the square on AB is equal to the rectangle HB BM ,
because when AH is joined the triangle ABH is equiangular with the triangle ABM and as HB is to BA so is AB to BM

Therefore the side AB of the pentagon is the irrational straight line called minor Q E D

PROPOSITION 12

If an equilateral triangle be inscribed in a circle the square on the side of the triangle is triple of the square on the radius of the circle

Let ABC be a circle

and let the equilateral triangle ABC be inscribed in it

I say that the square on one side of the triangle ABC is triple of the square on the radius of the circle

For let the centre D of the circle ABC be taken

let AD be joined and carried through to E


and let BE be joined

Then since the triangle ABC is equilateral
therefore the circumference BEC is a third part of the circumference of the circle ABC

Therefore the circumference BE is a sixth part of the circumference of the circle

therefore the straight line BE belongs to a hexagon

therefore it is equal to the radius DE [iv 15 Por]



And since $1E$ is double of DE
the square on AE is quadruple of the square on ED that is of the square on BE

But the square on AE is equal to the squares on AB BE [iii 31 I 47]
therefore the squares on AB BE are quadruple of the square on BE

Therefore *separando* the square on AB is triple of the square on BE

But BE is equal to DE ,

therefore the square on AB is triple of the square on DE

Therefore the square on the side of the triangle is triple of the square on the radius

Q E D

PROPOSITION 13

To construct a pyramid, to comprehend it in a given sphere and to prove that the square on the diameter of the sphere is one and a half times the square on the side of the pyramid

Let the diameter AB of the given sphere be set out,

and let it be cut at the point C so that AC is double of CB ,

let the semicircle ADB be described on AB ,

let CD be drawn from the point C at right angles to AB ,

and let DA be joined,

let the circle EFG which has its radius equal to DC be set out,

let the equilateral triangle EFG be inscribed in the circle EFG , [IV 2]



let the centre of the circle the point H , be taken [III 1]

let EH HF , HG be joined,

from the point H let HA be set up at right angles to the plane of the circle EFG , [XI 12]

let HK equal to the straight line AC be cut off from HA ,

and let KE , KF , KG be joined



Now since HA is at right angles to the plane of the circle EFG therefore it will also make right angles with all the straight lines which meet it and are in the plane of the circle EFG [XI Def 3]

But each of the straight lines HE HF HG meets it

therefore HK is at right angles to each of the straight lines HE , HF HG

And, since AC is equal to HK , and CD to HE

and they contain right angles

therefore the base DA is equal to the base KE [I 4]

For the same reason

each of the straight lines KF KG is also equal to DA

therefore the three straight lines KE KF , KG are equal to one another

And since AC is double of CB ,

therefore AB is triple of BC

But, as AB is to BC so is the square on AD to the square on DC , as will be proved afterwards

Therefore the square on AD is triple of the square on DC

But the square on FE is also triple of the square on EH , [III 12]
and DC is equal to EH ,

therefore DA is also equal to EF

But DA was proved equal to each of the straight lines KE , KF , KG
therefore each of the straight lines EF , FG , GE is also equal to each of the straight lines KE , KF , KG ,

therefore the four triangles EFG , KEF , KFG , KEG are equilateral

Therefore a pyramid has been constructed out of four equilateral triangles the triangle EFG being its base and the point K its vertex

It is next required to comprehend it in the given sphere and to prove that the square on the diameter of the sphere is one and a half times the square on the side of the pyramid

For let the straight line HL be produced in a straight line with KH ,
and let HL be made equal to CB

Now, since as AC is to CD , so is CD to CB [VI 8 Por]
while AC is equal to KH , CD to HE , and CB to HL ,

therefore, as KH is to HE , so is EH to HL

therefore the rectangle KH , HL is equal to the square on EH [VI 14]

And each of the angles KHE EHL is right,

therefore the semicircle described on KL will pass through F also

[cf VI 8, III 31]

If then KL remaining fixed the semicircle be carried round and restored to the same position from which it began to be moved, it will also pass through the points F G ,

since if FL LG be joined the angles at F G similarly become right angles,
and the pyramid will be comprehended in the given sphere

For KL the diameter of the sphere is equal to the diameter AB of the given sphere, inasmuch as KH was made equal to AC and HL to CB

I say next that the square on the diameter of the sphere is one and a half times the square on the side of the pyramid

For since AC is double of CB ,

therefore AB is triple of BC ,

and *convertendo* BA is one and a half times AC

But as BA is to AC so is the square on BA to the square on AD

Therefore the square on BA is also one and a half times the square on AD

And BA is the diameter of the given sphere and AD is equal to the side of the pyramid

Therefore the square on the diameter of the sphere is one and a half times the square on the side of the pyramid Q E D

LEMMA

It is to be proved that as AB is to BC , so is the square on AD to the square on DC

For let the figure of the semicircle be set out

let DB be joined

let the square EC be described on AC ,

and let the parallelogram FB be completed

Since then because the triangle DAB is equiangular with the triangle DAC ,
as BA is to AD , so is DA to AC [vi 8 vi 4]
therefore the rectangle BA, AC is equal to
the square on AD [vi 17]

And since as AB is to BC , so is EB to BF
[vi 1]

and EB is the rectangle BA, AC , for EA is
equal to AC ,

and BF is the rectangle AC, CB ,
therefore as AB is to BC , so is the rectangle
 BA, AC to the rectangle AC, CB

And the rectangle BA, AC is equal to the
square on AD and the rectangle AC, CB to
the square on DC ,

for the perpendicular DC is a mean propor-
tional between the segments AC, CB of the
[vi 8 Por.]

base because the angle ADB is right

Therefore as AB is to BC , so is the square on AD to the square on DC

Q E D

PROPOSITION 14

To construct an octahedron and comprehend it in a sphere as in the preceding ca e
and to prove that the square on the diameter of the sphere is double of the square on
the side of the octahedron

Let the diameter AB of the given sphere be set out

and let it be bisected at C

let the semicircle ADB be described on AB

let CD be drawn from C at right angles to AB



let DB be joined

let the square $EFGH$, having each of its sides equal to DB be set out

let HF, EG be joined

from the point K let the straight line KL be set up at right angles to the plane
of the square $EFGH$ [xi 12] and let it be carried through to the other side of
the plane as KM

from the straight lines KL, KM let KL, KM be respectively cut off equal to
one of the straight lines EK, FK, GK, HK



and let LE, LF, LG, LH
 ME, MF, MG, MH be
joined

Then, since KE is equal
to KH ,
and the angle EKH is right
therefore the square on HE
is double of the square on
 EK [i 47]

Again, since LK is equal
to KE ,

and the angle LKE is right

therefore the square on EL is double of the square on EK [14]

But the square on HE was also proved double of the square on EK ,
therefore the square on LE is equal to the square on EH ,
therefore LE is equal to EH

For the same reason

LH is also equal to HE ,

therefore the triangle LEH is equilateral

Similarly we can prove that each of the remaining triangles of which the sides of the square $FFGH$ are the bases, and the points L, M the vertices, is equilateral,
therefore an octahedron has been constructed which is contained by eight equilateral triangles

It is next required to comprehend it in the given sphere, and to prove that the square on the diameter of the sphere is double of the square on the side of the octahedron

For since the three straight lines LK, KM, KE are equal to one another,
therefore the semicircle described on LM will also pass through E

And for the same reason,

if LM remaining fixed the semicircle be carried round and restored to the same position from which it began to be moved

it will also pass through the points F, G, H ,

and the octahedron will have been comprehended in a sphere

I say next that it is also comprehended in the given sphere

For since LK is equal to KM ,

while KE is common

and they contain right angles

therefore the base LE is equal to the base EM [14]

And since the angle LFM is right for it is in a semicircle [11 31]

therefore the square on LM is double of the square on LE [14 31]

Again since AC is equal to CB ,

AB is double of BC

But as AB is to BC so is the square on AB to the square on BD ,

therefore the square on AB is double of the square on BD

But the square on LM was also proved double of the square on LE

And the square on DB is equal to the square on LE for EH was made equal to DB

Therefore the square on AB is also equal to the square on LM ,

therefore AB is equal to LM

And AB is the diameter of the given sphere

therefore LM is equal to the diameter of the given sphere

Therefore the octahedron has been comprehended in the given sphere and it has been demonstrated at the same time that the square on the diameter of the sphere is double of the square on the side of the octahedron Q E D

PROPOSITION 15

To construct a cube and comprehend it in a sphere like the pyramid and to prove that the square on the diameter of the sphere is triple of the square on the side of the cube

Let the diameter AB of the given sphere be set out,

and let it be cut at C so that AC is double of CB ,

let the semicircle ADB be described on AB

let CD be drawn from C at right angles to AB ,

and let DB be joined,



let the square $EFGH$ having its side equal to DB be set out

from $E F G H$ let $EK FL GM HN$ be drawn at right angles to the plane of the square $EFGH$

from EK, FL, GM, HN let $EA, FL GM HN$ respectively be cut off equal to one of the straight lines $EF FG, GH, HE$,

and let AL, LM, MN, NK be joined

therefore the cube FN has been constructed which is contained by six equal squares



It is then required to comprehend it in the given sphere and to prove that the square on the diameter of the sphere is triple of the square on the side of the cube

For let $AG EG$ be joined

Then since the angle KEG is right, because KE is also at right angles to the plane EG

[XI Def 3]

and of course to the straight line EG also,

therefore the semicircle described on AG will also pass through the point E

Again since GF is at right angles to each of the straight lines FL, FE

GF is also at right angles to the plane FA

hence also if we join FA GF will be at right angles to FK ,

and for this reason again the semicircle described on GA will also pass through F

Similarly it will also pass through the remaining angular points of the cube

If then AG remaining fixed the semicircle be carried round and restored to the same position from which it began to be moved,

the cube will be comprehended in a sphere

I say next that it is also comprehended in the given sphere

For since GF is equal to FE ,

and the angle at F is right

therefore the square on EG is double of the square on EF

But EF is equal to EA ,

therefore the square on EG is double of the square on EA

hence the squares on $GE EA$ that is the square on GA [I 47] is triple of the square on EA

And since AB is triple of BC

while as AB is to BC so is the square on AB to the square on BD

therefore the square on AB is triple of the square on BD

But the square on GA was also proved triple of the square on AE

And AE was made equal to DB

therefore AG is also equal to AB

And AB is the diameter of the given sphere

therefore AG is also equal to the diameter of the given sphere

Therefore the cube has been comprehended in the given sphere, and it has been demonstrated at the same time that the square on the diameter of the sphere is triple of the square on the side of the cube Q E D

PROPOSITION 16

To construct an icosahedron and comprehend it in a sphere, like the aforesaid figures and to prove that the side of the icosahedron is the irrational straight line called minor


Let the diameter AB of the given sphere be set out

and let it be cut at C so that AC is quadruple of CB ,

let the semicircle ADB be described on AB ,

let the straight line CD be drawn from C at right angles to AB ,

and let DB be joined



let the circle $EFGHA$ be set out and let its radius be equal to DB
let the equilateral and equiangular pentagon $EFGHA$ be inscribed in the circle $EFGHA$

let the circumferences EF , FG , GH , HA , AE be bisected at the points L , M , N , O , P

and let LM , MN , NO , OP , PL , EP be joined

Therefore the pentagon $LMNOP$ is also equilateral

and the straight line EP belongs to a decagon

Now from the points E , F , G , H , A let the straight lines EQ , FR , GS , HT , AL be set up at right angles to the plane of the circle and let them be equal to the radius of the circle $EFGHA$

let QR , RS , ST , TU , UQ , QL , LR , RM , MS , SN , NT , TO , OU , UP , PQ be joined

Now, since each of the straight lines EQ KU is at right angles to the same plane

therefore EQ is parallel to KU [XI 6]

But it is also equal to it,
and the straight lines joining those extremities of equal and parallel straight lines which are in the same direction are equal and parallel [I 33]

Therefore QU is equal and parallel to EK

But FK belongs to an equilateral pentagon,
therefore QU also belongs to the equilateral pentagon inscribed in the circle $EFGHA$

For the same reason

each of the straight lines QR , RS ST TU also belongs to the equilateral pentagon inscribed in the circle $EFGHA$,

therefore the pentagon $QRSTU$ is equilateral

And since QL belongs to a hexagon,

and EP to a decagon

and the angle QEP is right,

therefore QP belongs to a pentagon,

for the square on the side of the pentagon is equal to the square on the side of the hexagon and the square on the side of the decagon inscribed in the same circle [XIII 10]

For the same reason

PU is also a side of a pentagon

But QU also belongs to a pentagon,

therefore the triangle QPU is equilateral

For the same reason

each of the triangles QLR , RMS SNT , TOU is also equilateral

And since each of the straight lines QL QP was proved to belong to a pentagon

and LP also belongs to a pentagon,

therefore the triangle QLP is equilateral

For the same reason

each of the triangles LRM MSN NTO OUP is also equilateral

Let the centre of the circle $EFGHA$ the point V , be taken

from V let VZ be set up at right angles to the plane of the circle

let it be produced in the other direction as VX

let there be cut off VW the side of a hexagon, and each of the straight lines VX WZ being sides of a decagon

and let QZ QW UZ EV LV LX AM be joined

Now since each of the straight lines VW QV is at right angles to the plane of the circle

therefore VW is parallel to QE [XI 6]

But they are also equal

therefore EV QW are also equal and parallel [I 33]

But EV belongs to a hexagon

therefore QW also belongs to a hexagon

And, since QW belongs to a hexagon

and WZ to a decagon

and the angle QWZ is right

therefore QZ belongs to a pentagon

[xiii 10]

For the same reason

UZ also belongs to a pentagon,

inasmuch as, if we join VK , WU , they will be equal and opposite, and VK , being a radius, belongs to a hexagon,

[iv 15 Por]

therefore WU also belongs to a hexagon

But WZ belongs to a decagon

and the angle UWZ is right,

therefore UZ belongs to a pentagon

[xiii 10]

But QU also belongs to a pentagon,

therefore the triangle QUZ is equilateral

For the same reason

each of the remaining triangles of which the straight lines QR , RS , ST , TU are the bases and the point Z the vertex is also equilateral

Again since VL belongs to a hexagon,

and VX to a decagon,

and the angle $L VX$ is right,

therefore LX belongs to a pentagon

[xiii 10]

For the same reason

if we join MV , which belongs to a hexagon,

MX is also inferred to belong to a pentagon

But LM also belongs to a pentagon,

therefore the triangle LMX is equilateral

Similarly it can be proved that each of the remaining triangles of which MN , NO , OP , PL are the bases and the point Y the vertex is also equilateral

Therefore an icosahedron has been constructed which is contained by twenty equilateral triangles

It is next required to comprehend it in the given sphere and to prove that the side of the icosahedron is the irrational straight line called minor

For since VW belongs to a hexagon

and WZ to a decagon

therefore VZ has been cut in extreme and mean ratio at W ,

and VW is its greater segment,

[xiii 9]

therefore as ZV is to VW so is VW to WZ

But VW is equal to VE and WZ to VX

therefore as ZV is to VF so is EV to VX

And the angles ZVE , EVX are right,

therefore if we join the straight line EZ , the angle AEZ will be right because of the similarity of the triangles XEZ , VEZ

For the same reason

since as ZV is to VW so is VW to WZ ,

and ZV is equal to VW and VW to WQ

therefore as VW is to WQ so is QW to WZ

And for this reason again

if we join QV the angle at Q will be right,

[vi 8]

therefore the semicircle described on AZ will also pass through Q

[iii 31]

And if AZ remaining fixed the semicircle be carried round and restored to the same position from which it began to be moved, it will also pass through Q and the remaining angular points of the icosahedron

and the icosahedron will have been comprehended in a sphere

I say next that it is also comprehended in the given sphere

For let VW be bisected at A'

Then, since the straight line VZ has been cut in extreme and mean ratio at W

and ZW is its lesser segment

therefore the square on ZW added to the half of the greater segment that is WA' , is five times the square on the half of the greater segment, [XIII 3]

therefore the square on ZA' is five times the square on $A'W$

And $Z\lambda$ is double of ZA' and VW double of $A'W$,

therefore the square on ZX is five times the square on WV

And, since AC is quadruple of CB ,

therefore AB is five times BC

But, as AB is to BC , so is the square on AB to the square on BD , [VI 8 v Def 9]

therefore the square on AB is five times the square on BD

But the square on ZY was also proved to be five times the square on VW

And DB is equal to VW ,

for each of them is equal to the radius of the circle $EFGHA$,

therefore AB is also equal to λZ

And AB is the diameter of the given sphere,

therefore λZ is also equal to the diameter of the given sphere

Therefore the icosahedron has been comprehended in the given sphere

I say next that the side of the icosahedron is the irrational straight line called minor

For, since the diameter of the sphere is rational

and the square on it is five times the square on the radius of the circle $EFGHA$,

therefore the radius of the circle $EFGHK$ is also rational

hence its diameter is also rational

But if an equilateral pentagon be inscribed in a circle which has its diameter rational the side of the pentagon is the irrational straight line called minor [XIII 11]

And the side of the pentagon $EFGHK$ is the side of the icosahedron

Therefore the side of the icosahedron is the irrational straight line called minor

PORISM From this it is manifest that the square on the diameter of the sphere is five times the square on the radius of the circle from which the icosahedron has been described and that the diameter of the sphere is composed of the side of the hexagon and two of the sides of the decagon inscribed in the same circle Q E D

PROPOSITION 17

To construct a dodecahedron and comprehend it in a sphere, like the aforesaid figures and to prove that the side of the dodecahedron is the irrational straight line called apotome

Let $ABCD$ $CBEF$, two planes of the aforesaid cube at right angles to one another, be set out

let the sides AB BC , CD DA , EF , EB FC be bisected at G H , K , L M N , O respectively

let GA, HL, MH, VO be joined

let the straight lines NP, PO, HQ be cut in extreme and mean ratio at the points R, S, T respectively,

and let RP, PS, TQ be their greater segments,

from the points R, S, T let RU, SI, TW be set up at right angles to the planes of the cube towards the outside of the cube,

let them be made equal to RP, PS, TQ ,

and let UB, BW, WC, CV, VU be joined

I say that the pentagon $UBWCV$ is equilateral, and in one plane, and is further equiangular

For let RB, SB, VB be joined

Then since the straight line VP has been cut in extreme and mean ratio at R

and RP is the greater segment, therefore the squares on PN, NR are triple of the square on RP [XIII. 4]

But PN is equal to NB , and PR to RU ,

therefore the squares on BN, NR are triple of the square on RU

But the square on BR is equal to the squares on BN, NR , [I. 47]

therefore the square on BR is triple of the square on RU ,

hence the squares on BR, RU are quadruple of the square on RU

But the square on BU is equal to the squares on BR, RU ,

therefore the square on BU is quadruple of the square on RU ,

therefore BU is double of RU

But VU is also double of UR

inasmuch as SR is also double of PR , that is, of RU ,

therefore BU is equal to UV

Similarly it can be proved that each of the straight lines BW, WC, CV is also equal to each of the straight lines BU, UV

Therefore the pentagon $BUVCW$ is equilateral

I say next that it is also in one plane

For let $P\Lambda$ be drawn from P parallel to each of the straight lines RU, SI and towards the outside of the cube and let $\Lambda H, HW$ be joined,

I say that ΛHW is a straight line

For since HQ has been cut in extreme and mean ratio at T and QT is its greater segment

therefore as HQ is to QT , so is QT to TH


But HQ is equal to HP and QT to each of the straight lines $TW, P\Lambda$,

therefore as HP is to $P\Lambda$ so is HT to TH

And HP is parallel to TW

for each of them is at right angles to the plane BD ,

and TH is parallel to $P\Lambda$, [XI. 6]



for each of them is at right angles to the plane BF [id]

But if two triangles as $\triangle PH, \triangle TW$, which have two sides proportional to two sides be placed together at one angle so that their corresponding sides are also parallel

the remaining straight lines will be in a straight line, [vi 32]

therefore $\triangle H$ is in a straight line with HW

But every straight line is in one plane, [xi 1]

therefore the pentagon $UBW CV$ is in one plane

I say next that it is also equiangular

For since the straight line NP has been cut in extreme and mean ratio at R and PR is the greater segment

while PR is equal to PS

therefore NS has also been cut in extreme and mean ratio at P

and NP is the greater segment [xiii 5]

therefore the squares on $NS SP$ are triple of the square on NP [xiii 4]

But NP is equal to NB and PS to SV ,

therefore the squares on NS, SV are triple of the square on NB ,
hence the squares on $VS SN NB$ are quadruple of the square on NB

But the square on SB is equal to the squares on SN, NB ,
therefore the squares on $BS SV$, that is the square on BV —for the angle $\angle SB$ is right—is quadruple of the square on NB ,

therefore BV is double of BN

But BC is also double of BN ,

therefore BV is equal to BC

And since the two sides BU, UV are equal to the two sides $BW WC$,

and the base BV is equal to the base BC ,

therefore the angle BUV is equal to the angle BWC [i 8]

Similarly we can prove that the angle UVC is also equal to the angle BWC ,
therefore the three angles BWC, BUV, UVC are equal to one another

But if in an equilateral pentagon three angles are equal to one another the pentagon will be equiangular [xiii 7]

therefore the pentagon $BUVCW$ is equiangular

And it was also proved equilateral

therefore the pentagon $BUVCW$ is equilateral and equiangular and it is on one side BC of the cube

Therefore if we make the same construction in the case of each of the twelve sides of the cube

a solid figure will have been constructed which is contained by twelve equilateral and equiangular pentagons and which is called a dodecahedron

It is then required to comprehend it in the given sphere and to prove that the side of the dodecahedron is the irrational straight line called apotome

For let AP be produced and let the produced straight line be AZ
therefore PZ meets the diameter of the cube and they bisect one another
for this has been proved in the last theorem but one of the eleventh book

[xi 38]

Let them cut at Z

therefore Z is the centre of the sphere which comprehends the cube,
and ZP is half of the side of the cube

Let UZ be joined

Now, since the straight line NS has been cut in extreme and mean ratio at P ,

and NP is its greater segment

therefore the squares on NS , SP are triple of the square on NP [xiii 4]

But NS is equal to λZ

inasmuch as NP is also equal to PZ , and XP to PS

But further, PS is also equal to λU ,

since it is also equal to RP ,

therefore the squares on $Z\lambda$ XU are triple of the square on NP

But the square on UZ is equal to the squares on $Z\lambda$, λU ,

therefore the square on UZ is triple of the square on NP

But the square on the radius of the sphere which comprehends the cube is also triple of the square on the half of the side of the cube for it has previously been shown how to construct a cube and comprehend it in a sphere and to prove that the square on the diameter of the sphere is triple of the square on the side of the cube [xiii 15]

But if whole is so related to whole so is half to half also,

and NP is half of the side of the cube,

therefore UZ is equal to the radius of the sphere which comprehends the cube

And Z is the centre of the sphere which comprehends the cube,

therefore the point U is on the surface of the sphere

Similarly we can prove that each of the remaining angles of the dodecahedron is also on the surface of the sphere,

therefore the dodecahedron has been comprehended in the given sphere

I say next that the side of the dodecahedron is the irrational straight line called apotome

For since when NP has been cut in extreme and mean ratio RP is the greater segment

and when PO has been cut in extreme and mean ratio PS is the greater segment

therefore when the whole NO is cut in extreme and mean ratio RS is the greater segment

[Thus since as NP is to PR so is PR to RN

the same is true of the doubles also

for parts have the same ratio as their equimultiples, [v 15]

therefore as NO is to RS so is RS to the sum of NR SO

But NO is greater than RS

therefore RS is also greater than the sum of NR SO ,

therefore NO has been cut in extreme and mean ratio

and RS is its greater segment]

But RS is equal to UV

therefore when NO is cut in extreme and mean ratio, UV is the greater segment

And since the diameter of the sphere is rational

and the square on it is triple of the square on the side of the cube

therefore NO being a side of the cube is rational

[But if a rational line be cut in extreme and mean ratio each of the segments is an irrational apotome]

Therefore UV being a side of the dodecahedron is an irrational apotome [xiii 6]

PORISM From this it is manifest that when the side of the cube is cut in extreme and mean ratio, the greater segment is the side of the dodecahedron

Q E D

PROPOSITION 18

To set out the sides of the five figures and to compare them with one another

Let AB , the diameter of the given sphere, be set out,
and let it be cut at C so that AC is equal to CB and at D so that AD is double of DB ,
let the semicircle AEB be described on AB ,
from C , D let CE , DF be drawn at right angles to AB


and let AF , FB , EB be joined

Then since AD is double of DB

therefore AB is triple of BD

Converting, therefore BA is one and a half times AD

But as BA is to AD so is the square on BA to the square on AD , [v Def 9 vi 8]
for the triangle AFB is equiangular with the triangle AFD ,



therefore the square on BA is one and a half times the square on AD

But the square on the diameter of the sphere is also one and a half times the square on the side of the pyramid [XIII 13]

And AB is the diameter of the sphere,

therefore AF is equal to the side of the pyramid

Again since AD is double of DB

therefore AB is triple of BD

But as AB is to BD so is the square on AB to the square on BD

[vi 8 v Def 9]

therefore the square on AB is triple of the square on BD

But the square on the diameter of the sphere is also triple of the square on the side of the cube [XIII 15]

And AB is the diameter of the sphere,

therefore BF is the side of the cube

And since AC is equal to CB ,

therefore AB is double of BC

But as AB is to BC so is the square on AB to the square on BC

therefore the square on AB is double of the square on BC

But the square on the diameter of the sphere is also double of the square on the side of the octahedron [XIII 14]

And AB is the diameter of the given sphere

therefore BE is the side of the octahedron

Next let AG be drawn from the point A at right angles to the straight line AB ,

let AG be made equal to AB

let GC be joined

and from H let HK be drawn perpendicular to AB

Then, since GA is double of AC ,

for GA is equal to AB ,
 and, as GA is to AC , so is HK to AC ,
 therefore HK is also double of AC

Therefore the square on HK is quadruple of the square on AC
 therefore the squares on HK , AC , that is the square on HC , is five times the square on AC

But HC is equal to CB ,

therefore the square on BC is five times the square on CA

And, since AB is double of CB ,

and in them AD is double of DB ,

therefore the remainder BD is double of the remainder DC

Therefore BC is triple of CD ,

therefore the square on BC is nine times the square on CD

But the square on BC is five times the square on CA ,

therefore the square on CA is greater than the square on CD ,

therefore CA is greater than CD

Let CL be made equal to CA

from L let LM be drawn at right angles to AB ,

and let MB be joined

Now since the square on BC is five times the square on CA ,

and AB is double of BC , and AL double of CA ,

therefore the square on AB is five times the square on AL

But the square on the diameter of the sphere is also five times the square on the radius of the circle from which the icosahedron has been described

[xiii 16 Por]

And AB is the diameter of the sphere

therefore AL is the radius of the circle from which the icosahedron has been described

therefore AL is a side of the hexagon in the said circle

[iv 15 Por]

And since the diameter of the sphere is made up of the side of the hexagon and two of the sides of the decagon inscribed in the same circle

[xiii 16 Por]

and AB is the diameter of the sphere

while AL is a side of the hexagon

and AA is equal to LB

therefore each of the straight lines AA LB is a side of the decagon inscribed in the circle from which the icosahedron has been described

And since LB belongs to a decagon and MI to a hexagon

for MI is equal to AI since it is also equal to HA being the same distance from the centre and each of the straight lines HA AL is double of AC

therefore MB belongs to a pentagon

[xiii 10]

But the side of the pentagon is the side of the icosahedron,

[xiii 16]

therefore MB belongs to the icosahedron

Now, since FB is a side of the cube

let it be cut in extreme and mean ratio at N ,

and let NB be the greater segment

therefore NB is a side of the dodecahedron

[xiii 17 Por]

And since the square on the diameter of the sphere was proved to be one and a half times the square on the side AF of the pyramid double of the square on

the side BC of the octahedron and triple of the side FB of the cube therefore of parts of which the square on the diameter of the sphere contains six the square on the side of the pyramid contains four, the square on the side of the octahedron three and the square on the side of the cube two

Therefore the square on the side of the pyramid is four thirds of the square on the side of the octahedron and double of the square on the side of the cube and the square on the side of the octahedron is one and a half times the square on the side of the cube

The said sides therefore of the three figures I mean the pyramid, the octahedron and the cube, are to one another in rational ratios

But the remaining two, I mean the side of the icosahedron and the side of the dodecahedron are not in rational ratios either to one another or to the aforesaid sides

for they are irrational, the one being minor [XIII 16] and the other an apotome [XIII 17]

That the side MB of the icosahedron is greater than the side AB of the dodecahedron we can prove thus

For, since the triangle FDB is equiangular with the triangle FAB [VI 8]
proportionally, as DB is to BF , so is BI to BA [VI 4]

And since the three straight lines are proportional,
as the first is to the third so is the square on the first to the square on the second [V Def 9 VI 20, Por]

therefore as DB is to BA so is the square on DB to the square on BF ,
therefore inversely as AB is to BD , so is the square on FB to the square on BD

But AB is triple of BD

therefore the square on FB is triple of the square on BD

But the square on AD is also quadruple of the square on DB

for AD is double of DB ,

therefore the square on AD is greater than the square on FB ,

therefore AD is greater than FB ,

therefore AL is by far greater than FB

And when AL is cut in extreme and mean ratio

KL is the greater segment

inasmuch as LK belongs to a hexagon and KA to a decagon [XIII 9]

and when FB is cut in extreme and mean ratio NB is the greater segment

therefore KL is greater than NB

But KL is equal to LM ,

therefore LM is greater than NB

Therefore MB which is a side of the icosahedron is by far greater than AB
which is a side of the dodecahedron Q E D

I say next that no other figure besides the said five figures can be constructed which is contained by equilateral and equiangular figures equal to one another

For a solid angle cannot be constructed with two triangles or indeed planes

With three triangles the angle of the pyramid is constructed, with four the angle of the octahedron and with five the angle of the icosahedron

but a solid angle cannot be formed by six equilateral and equiangular triangles placed together at one point,
 for the angle of the equilateral triangle being two-thirds of a right angle, the six will be equal to four right angles
 which is impossible, for any solid angle is contained by angles less than four right angles [XI 21]

For the same reason, neither can a solid angle be constructed by more than six plane angles

By three squares the angle of the cube is contained but by four it is impossible for a solid angle to be contained,

for they will again be four right angles

By three equilateral and equiangular pentagons the angle of the dodecahedron is contained,
 but by four such it is impossible for any solid angle to be contained
 for the angle of the equilateral pentagon being a right angle and a fifth the four angles will be greater than four right angles
 which is impossible

Neither again will a solid angle be contained by other polygonal figures by reason of the same absurdity

Therefore etc

Q E D

LEMMA

But that the angle of the equilateral and equiangular pentagon is a right angle and a fifth we must prove thus

Let $ABCDF$ be an equilateral and equiangular pentagon,
 let the circle $ABCDE$ be circumscribed
 about it,

let its centre F be taken

and let FA, FB, FC, FD, FE be joined

Therefore they bisect the angles of the pentagon at A, B, C, D, E

And since the angles at F are equal to four right angles and are equal


therefore one of them as the angle AFB is one right angle less a fifth,

therefore the remaining angles FAB, ABF consist of one right angle and a fifth

But the angle FAB is equal to the angle FBC ,

therefore the whole angle ABC of the pentagon consists of one right angle and a fifth

Q E D



THE WORKS OF ARCHIMEDES
INCLUDING THE METHOD

BIOGRAPHICAL NOTE

ARCHIMEDES, c. 287-212 B C

ARCHIMEDES was a citizen of Syracuse in Sicily, where he was born around the year 287 B C. He was intimate with Hiero, King of Syracuse, and with his son, Gelo. Plutarch says that he was related to them. In his *Sand Reckoner*, which was dedicated to Gelo, Archimedes speaks of his father, Phidias, as an astronomer who investigated the sizes and distances of the sun and moon.

As a young man Archimedes seems to have spent some time in Egypt, where he invented the water-screw as a means of drawing water out of the Nile for irrigating the fields, though it is also said that he invented this machine to drain bilge water from a huge ship built for King Hiero. He may have studied with the pupils of Euclid in Alexandria. It was probably there that he made the friendship of Conon of Samos and Eratosthenes. To Conon he was in the habit of communicating his discoveries before their publication, and it was for Eratosthenes that he wrote the *Method*, and through him that he addressed the famous *Cattle Problem* to the mathematicians of Alexandria—if the tradition is to be credited that associates Archimedes with this problem. After the death of Conon, Archimedes sent his discoveries to Conon's friend and pupil, Dositheus of Pelusium, to whom four of the extant treatises are dedicated.

His mechanical inventions won great fame for Archimedes and figure largely in the traditions about him. After discovering the solution of the problem *To move a given weight by a given force*, he boasted to King Hiero: 'Give me a place to stand on and I can move the earth.' Asked for a practical demonstration, he contrived a machine by which, with the use of only one arm, he drew out of the dock a large ship laden with passengers and goods, which the combined strength of the Syracusans could scarcely move. From that day Hiero ordered that

Archimedes was to be believed in everything he might say. At the King's request Archimedes then made for him catapults, battering rams, cranes, and many other engines of war, which were later used with such success in the defense of Syracuse against the Romans that they were unable to take the city except by treachery. There is also a story in Lucian that Archimedes set fire to the Roman ships by an arrangement of burning glasses.

Although Archimedes acquired by his mechanical inventions 'the renown of more than human sagacity' according to Plutarch, he would not deign to leave behind him any commentary or writing on such subjects, since he considered them "sordid and ignoble." He did, however, write a description now lost of an apparatus composed of concentric glass spheres moved by water power, representing the Eudoxian system of the world. This astronomical machine, which survived to be seen and described by Cicero in his *Republic*, was sufficiently accurate to show the eclipses of the sun and the moon. Except for this lost work *On Sphere making*, Archimedes wrote only on strictly mathematical subjects. He took all the mathematical sciences for his province: arithmetic, geometry, astronomy, mechanics, and hydrostatics. Unlike Euclid and Apol-

lonius he wrote no textbooks. Of his writings, although some have been lost the most important have survived.

The absorption of Archimedes in his mathematical investigations was so great that he forgot his food and neglected his person, and when carried by force to the bath Plutarch records ' he used to trace geometrical figures in the ashes of the fire and diagrams in the oil on his body. Asked by Hiero to discover whether a goldsmith had alloyed with silver the gold of his crown Archimedes found the answer while bathing by considering the water displaced by his body whereupon he is reported to have run home in his excitement with out his clothes, shouting, Eureka (I have found it).

Archimedes' preoccupation with mathematics is even said to have been the cause of his death. In the general massacre which followed the capture of Syracuse by Marcellus in 212 B.C., Archimedes was so intent upon a mathematical diagram that he took no notice, and when ordered by a soldier to attend the victorious general he refused until he should have solved his problem whereupon he was slain by the enraged soldier. No blame attaches to the Roman general, Marcellus, since he had given orders to spare the house and person of the mathematician and in the midst of his triumph he lamented the death of Archimedes, provided him with an honorable burial and befriended his surviving relatives. In accordance with the expressed desire of Archimedes, his family and friends inscribed on his tomb the figure of his favorite theorem on the sphere and the circumscribed cylinder and the ratio of the containing solid to the contained. When Cicero was in Sicily as quaestor in 75 B.C., he discovered the neglected and forgotten tomb of Archimedes near the Agrigentine Gate and piously restored it.

CONTENTS

BIOGRAPHICAL NOTE,	399
ON THE SPHERE AND CYLINDER, BOOK I,	403
ON THE SPHERE AND CYLINDER, BOOK II	434
MEASUREMENT OF A CIRCLE	447
ON CONOIDS AND SPHEROIDS,	452
ON SPIRALS,	482
ON THE EQUILIBRIUM OF PLANES, BOOK I	502
ON THE EQUILIBRIUM OF PLANES BOOK II,	510
THE SAND RECKONER	520
QUADRATURE OF THE PARABOLA,	527
ON FLOATING BODIES, BOOK I,	538
ON FLOATING BODIES, BOOK II	543
BOOK OF LEMMAS,	561
THE METHOD TREATING OF MECHANICAL PROBLEMS	569

ON THE SPHERE AND CYLINDER

BOOK ONE

ARCHIMEDES to DOSITHEUS greeting

"On a former occasion I sent you the investigations which I had up to that time completed including the proofs showing that any segment bounded by a straight line and a section of a right angled cone [a parabola] is four thirds of the triangle which has the same base with the segment and equal height. Since then certain theorems not hitherto demonstrated have occurred to me, and I have worked out the proofs of them. They are these: first that the surface of any sphere is four times its greatest circle, next that the surface of any segment of a sphere is equal to a circle whose radius is equal to the straight line drawn from the vertex of the segment to the circumference of the circle which is the base of the segment, and, further that any cylinder having its base equal to the greatest circle of those in the sphere, and height equal to the diameter of the sphere is itself [i.e. in content] half as large again as the sphere and its surface also [including its bases] is half as large again as the surface of the sphere. Now these properties were all along naturally inherent in the figures referred to but remained unknown to those who were before my time engaged in the study of geometry. Having however, now discovered that the properties are true of these figures I cannot feel any hesitation in setting them side by side both with my former investigations and with those of the theorems of Eudoxus on solids which are held to be most irrefragably established namely, that any pyramid is one third part of the prism which has the same base with the pyramid and equal height, and that any cone is one third part of the cylinder which has the same base with the cone and equal height. For though these properties also were naturally inherent in the figures all along yet they were in fact unknown to all the many able geometers who lived before Eudoxus and had not been observed by any one. Now however it will be open to those who possess the requisite ability to examine these discoveries of mine. They ought to have been published while Conon was still alive for I should conceive that he would best have been able to grasp them and to pronounce upon them the appropriate verdict but as I judge it well to communicate them to those who are conversant with mathematics I send them to you with the proofs written out which it will be open to mathematicians to examine. Farewell.

'I first set out the axioms and the assumptions which I have used for the proofs of my propositions

DEFINITIONS

1 "There are in a plane certain terminated bent lines which either lie wholly on the same side of the straight lines joining their extremities, or have no part of them on the other side

2 "I apply the term *concave in the same direction* to a line such that, if any two points on it are taken either all the straight lines connecting the points fall on the same side of the line, or some fall on one and the same side while others fall on the line itself but none on the other side "

3 "Similarly also there are certain terminated surfaces not themselves being in a plane but having their extremities in a plane and such that they will either be wholly on the same side of the plane containing their extremities or have no part of them on the other side

4 "I apply the term *concave in the same direction* to surfaces such that, if any two points on them are taken the straight lines connecting the points either all fall on the same side of the surface, or some fall on one and the same side of it while some fall upon it but none on the other side

5 "I use the term *olid sector* when a cone cuts a sphere, and has its apex at the centre of the sphere to denote the figure comprehended by the surface of the cone and the surface of the sphere included within the cone "

6 "I apply the term *solid rhombus* when two cones with the same base have their apices on opposite sides of the plane of the base in such a position that their axes lie in a straight line to denote the solid figure made up of both the cones "

ASSUMPTIONS

1 *Of all lines which have the same extremities the straight line is the least "*


2 "Of other lines in a plane and having the same extremities, [any two] such are unequal whenever both are concave in the same direction and one of them is either wholly included between the other and the straight line which has the same extremities with it or is partly included by, and is partly common with, the other and that [line] which is included is the lesser [of the two]

3 *Similarly of surfaces which have the same extremities if those extremities are in a plane the plane is the least [in area]*

4 "Of other surfaces with the same extremities the extremities being in a plane [any two] such are unequal whenever both are concave in the same direction and one surface is either wholly included between the other and the plane which has the same extremities with it or is partly included by and partly common with the other and that [surface] which is included is the lesser [of the two in area]

5 Further of unequal lines unequal surfaces and unequal solids the greater exceed the less by such a magnitude as when added to itself can be made to exceed any assigned magnitude among those which are comparable with it and with one another

"These things being premised if a polygon be inscribed in a circle it is plain that the perimeter of the inscribed polygon is less than the circumference of the circle for each of the sides of the polygon is less than that part of the circumference of the circle which is cut off by it



PROPOSITION 1

If a polygon be circumscribed about a circle the perimeter of the circumscribed polygon is greater than the perimeter of the circle

Let any two adjacent sides meeting in A touch the circle at P, Q respectively

Then [Assumptions 2]

$$PA + AQ > (\text{arc } PQ)$$

A similar inequality holds for each angle of the polygon and by addition, the required result follows

PROPOSITION 2

Given two unequal magnitudes it is possible to find two unequal straight lines such that the greater straight line has to the less a ratio less than the greater magnitude has to the less

Let AB, D represent the two unequal magnitudes AB being the greater

Suppose BC measured along BA equal to D and let GH be any straight line

Then if CA be added to itself a sufficient number of times the sum will exceed D . Let AF be this sum and take E on GH produced such that GH is the same multiple of HE that AF is of AC

$$\text{Thus } EH : HG = AC : AF$$

But since $AF > D$ (or CB)

$$\frac{AC}{AF} < \frac{AC}{CB}$$



Therefore *componendo*

$$\frac{EG}{GH} < \frac{AB}{D}$$

Hence EG, GH are two lines satisfying the given condition

PROPOSITION 3

Given two unequal magnitudes and a circle it is possible to inscribe a polygon in the circle and to describe another about it so that the side of the circumscribed polygon may have to the side of the inscribed polygon a ratio less than that of the greater magnitude to the less



Let A, B represent the given magnitudes A being the greater

Find [Prop 2] two straight lines F, AL of which F is the greater such that

$$F : AL < A : B$$

(1)

Draw LM perpendicular to LA and of such length that $KM = F$

In the given circle let CE DG be two diameters at right angles. Then, bisecting the angle DOC , bisecting the half again and so on we shall arrive ultimately at an angle (as NOC) less than twice the angle LKM

Join NC which (by the construction) will be the side of a regular polygon inscribed in the circle. Let OP be the radius of the circle bisecting the angle NOC (and therefore bisecting NC at right angles in H , say), and let the tangent at P meet OC , ON produced in S T respectively

Now since $\angle COV < 2\angle LKM$,
 $\angle HOC < \angle LKM$,

and the angles at H , L are right

$$\text{therefore } MK/LA > OC/OH \\ > OP/OH$$

$$\text{Hence } ST/CN < MK/LK \\ < F/LA,$$

therefore *a fortiori*, by (1)


$$ST/CN < A/B$$

Thus two polygons are found satisfying the given condition

PROPOSITION 4

Again, given two unequal magnitudes and a sector it is possible to describe a polygon about the sector and to inscribe another in it so that the side of the circumscribed polygon may have to the side of the inscribed polygon a ratio less than the greater magnitude has to the less




[The "inscribed polygon" found in this proposition is one which has for two sides the two radii bounding the sector, while the remaining sides (the number of which is by construction some power of 2) subtend equal parts of the arc of the sector the circumscribed polygon is formed by the tangents parallel to the sides of the inscribed polygon and by the two bounding radii produced]



In this case we make the same construction as in the last proposition except that we bisect the angle COD of the sector, instead of the right angle between two diameters then bisect the half again, and so on. The proof is exactly similar to the preceding one

PROPOSITION 5

Given a circle and two unequal magnitudes, to describe a polygon about the circle and inscribe another in it so that the circumscribed polygon may have to the inscribed a ratio less than the greater magnitude has to the less



Let A be the given circle and B C the given magnitudes B being the greater

Take two unequal straight lines D E of which D is the greater, such that $D : E < B : C$ [Prop 2] and let F be a mean proportional between D E so that D is also greater than F

Describe (in the manner of Prop 3) one polygon about the circle and inscribe another in it, so that the side of the former has to the side

of the latter a ratio less than the ratio $D : F$

Thus the duplicate ratio of the side of the former polygon to the side of the latter is less than the ratio $D^2 : F^2$

But the said duplicate ratio of the sides is equal to the ratio of the areas of the polygons since they are similar,

therefore the area of the circumscribed polygon has to the area of the inscribed polygon a ratio less than the ratio $D^2 : F^2$ or $D : E$, and a fortiori less than the ratio $B : C$

PROPOSITION 6

Similarly we can show that given two unequal magnitudes and a sector it is possible to circumscribe a polygon about the sector and inscribe in it another similar one so that the circumscribed may have to the inscribed a ratio less than the greater magnitude has to the less

And it is likewise clear that if a circle or a sector as well as a certain area be given it is possible by inscribing regular polygons in the circle or sector and by continually inscribing such in the remaining segments to leave segments of the circle or sector which are [together] less than the given area For this is proved in the *Elements* [Eucl xiv 2]

But it is yet to be proved that given a circle or sector and an area it is possible to describe a polygon about the circle or sector such that the area remaining between the circumference and the circumscribed figure is less than the given area

The proof for the circle (which as Archimedes says can be equally applied to a sector) is as follows

Let A be the given circle and B the given area


Now there being two unequal magnitudes $A + B$ and A let a polygon (C) be circumscribed about the circle and a polygon (I) inscribed in it [as in Prop 5]

so that

$$C : I < A + B : A$$

The circumscribed polygon (C) shall be that required

(1)



For the circle (A) is greater than the inscribed polygon (I)

Therefore, from (1), *a fortiori*

$$C - A < A + B - A,$$

whence

$$C < A + B,$$

or

$$C - A < B$$

PROPOSITION 7

If in an isosceles cone [i.e. a right circular cone] a pyramid be inscribed having an equilateral base, the surface of the pyramid excluding the base is equal to a triangle having its base equal to the perimeter of the base of the pyramid and its height equal to the perpendicular drawn from the apex on one side of the base

Since the sides of the base of the pyramid are equal it follows that the perpendiculars from the apex to all the sides of the base are equal, and the proof of the proposition is obvious

PROPOSITION 8

If a pyramid be circumscribed about an isosceles cone, the surface of the pyramid excluding its base is equal to a triangle having its base equal to the perimeter of the base of the pyramid and its height equal to the side [i.e. a generator] of the cone

The base of the pyramid is a polygon circumscribed about the circular base of the cone and the line joining the apex of the cone or pyramid to the point of contact of any side of the polygon is perpendicular to that side. Also all these perpendiculars being generators of the cone, are equal whence the proposition follows immediately

PROPOSITION 9

If in the circular base of an isosceles cone a chord be placed and from its extremities straight lines be drawn to the apex of the cone the triangle so formed will be less than the portion of the surface of the cone intercepted between the lines drawn to the apex

Let ABC be the circular base of the cone, and O its apex

Draw a chord AB in the circle and join O to OB . Bisect the arc ACB in C and join AC , BC , OC

Then

$$\triangle OAC + \triangle OBC > \triangle OAB$$

Let the excess of the sum of the first two triangles over the third be equal to the area D


Then D is either less than the sum of the segments AEC , CFB or not less

I Let D be not less than the sum of the segments referred to

We have now two surfaces

(1) that consisting of the portion OAC of the surface of the cone together with the segment AEC and

(2) the triangle OAC



and, since the two surfaces have the same extremities (the perimeter of the triangle OAC), the former surface is greater than the latter, which is *included* by it [Assumptions 3 or 4]

Hence (surface $OAEC$) + (segment AEC) > $\triangle OAC$

Similarly (surface $OCFB$) + (segment CFB) > $\triangle OBC$

Therefore, since D is not less than the sum of the segments we have, by addition,

$$\begin{aligned} (\text{surface } OAECFB) + D &> \triangle OAC + \triangle OBC \\ &> \triangle OAB + D, \text{ by hypothesis} \end{aligned}$$

Taking away the common part D , we have the required result

II Let D be less than the sum of the segments AEC CFB

If now we bisect the arcs AC , CB , then bisect the halves and so on we shall ultimately leave segments which are together less than D [Prop 6]

Let AGE , EHC , CKF , FLB be those segments, and join OE OF

Then as before,

$$(\text{surface } OAGE) + (\text{segment } AGE) > \triangle OAE$$

$$\text{and } (\text{surface } OEHC) + (\text{segment } EHC) > \triangle OEC$$

$$\text{Therefore } (\text{surface } OAGHC) + (\text{segments } AGE, EHC)$$

$$> \triangle OAE + \triangle OEC$$

$$> \triangle OAC, \text{ a fortiori}$$

Similarly for the part of the surface of the cone bounded by OC OB and the arc CFB

Hence by addition

$$(\text{surface } OAGEHCKFLB) + (\text{segments } AGE, EHC, CKF, FLB)$$

$$> \triangle OAC + \triangle OBC$$

$$> \triangle OAB + D, \text{ by hypothesis}$$

But the sum of the segments is less than D and the required result follows

PROPOSITION 10

If in the plane of the circular base of an isosceles cone two tangents be drawn to the circle meeting in a point and the points of contact and the point of concurrence of the tangents be respectively joined to the apex of the cone the sum of the two triangles formed by the joining lines and the two tangents are together greater than the included portion of the surface of the cone

Let ABC be the circular base of the cone, O its apex AD BD the two tangents to the circle meeting in D Join OA OB , OD

Let ECF be drawn touching the circle at C , the middle point of the arc ACB , and therefore parallel to AB Join OE OF

$$\text{Then } ED + DF > EF,$$

and, adding $AE + FB$ to each side

$$AD + DB > AE + EF + FB$$

Now OA OC , OB being generators of the cone, are equal and they are respectively perpendicular to the tangents at A , C , B

It follows that

$$\triangle OAD + \triangle ODB > \triangle OAE + \triangle OEF + \triangle OFB$$

Let the area G be equal to the excess of the first sum over the second

G is then either less or not less than the sum of the spaces $EAHC$ $FCKB$ remaining between the circle and the tangents which sum we will call L

I Let G be not less than L

We have now two surfaces

(1) that of the pyramid with apex O and base $AEFB$, excluding the face OAB ,

(2) that consisting of the part $OACB$ of the surface of the cone together with the segment ACB

These two surfaces have the same extremities, viz the perimeter of the triangle OAB , and, since the former includes the latter, the former is the greater [Assumptions, 4]

That is the surface of the pyramid exclusive of the face OAB is greater than the sum of the surface $OACB$ and the segment ACB

Taking away the segment from each sum, we have

$$\triangle OAE + \triangle OEF + \triangle OFB + L > \text{the surface } OAHCKB$$

And G is not less than L

It follows that

$$\triangle OAE + \triangle OEF + \triangle OFB + G,$$

which is by hypothesis equal to $\triangle OAD + \triangle ODB$ is greater than the same surface

II Let G be less than L

If we bisect the arcs AC , CB and draw tangents at their middle points, then bisect the halves and draw tangents, and so on, we shall lastly arrive at a polygon such that the sum of the parts remaining between the sides of the polygon and the circumference of the segment is less than G

Let the remainders be those between the segment and the polygon $APQRSB$ and let their sum be M
Join OP OQ etc

Then as before

$$\triangle OAE + \triangle OEF + \triangle OFB > \triangle OAP + \triangle OPQ + \dots + \triangle OSB$$

Also as before

(surface of pyramid $OAPQRSB$ excluding the face OAB) > the part $OACB$ of the surface of the cone together with the segment ACB

Taking away the segment from each sum

$$\triangle OAP + \triangle OPQ + \dots + M > \text{the part } OACB \text{ of the surface of the cone}$$


Hence a fortiori

$$\triangle OAP + \triangle OEF + \triangle OFB + G$$

which is by hypothesis equal to

$$\triangle OAD + \triangle ODB$$

is greater than the part $OACB$ of the surface of the cone



PROPOSITION 11

If a plane parallel to the axis of a right cylinder cut the cylinder the part of the surface of the cylinder cut off by the plane is greater than the area of the parallel plane in which the plane cuts it

PROPOSITION 12

If at the extremities of two generators of any right cylinder tangents be drawn to the circular bases in the planes of those bases respectively and if the pairs of tangents meet, the parallelograms formed by each generator and the two corresponding tangents respectively are together greater than the included portion of the surface of the cylinder between the two generators

[The proofs of these two propositions follow exactly the methods of Props 9 10 respectively, and it is therefore unnecessary to reproduce them]

"From the properties thus proved it is clear (1) that, if a pyramid be inscribed in an isosceles cone the surface of the pyramid excluding the base is less than the surface of the cone [excluding the base], and (2) that, if a pyramid be circumscribed about an isosceles cone, the surface of the pyramid excluding the base is greater than the surface of the cone excluding the base

It is also clear from what has been proved both (1) that if a prism be inscribed in a right cylinder the surface of the prism made up of its parallelograms [i.e. excluding its bases] is less than the surface of the cylinder excluding its bases and (2) that if a prism be circumscribed about a right cylinder the surface of the prism made up of its parallelograms is greater than the surface of the cylinder excluding its bases'

PROPOSITION 13

The surface of any right cylinder excluding the bases is equal to a circle whose radius is a mean proportional between the side [i.e. a generator] of the cylinder and the diameter of its base

Let the base of the cylinder be the circle A and make CD equal to the diameter of this circle and EF equal to the height of the cylinder

Let H be a mean proportional between CD EF , and B a circle with radius equal to H

Then the circle B shall be equal to the surface of the cylinder (excluding the bases), which we will call S

For, if not B must be either greater or less than S


I Suppose $B < S$

Then it is possible to circumscribe a regular polygon about B and to inscribe another in it such that the ratio of the former to the latter is less than the ratio $S : B$

Suppose this done and circumscribe about A a polygon

similar to that described about B then erect on the polygon about A a prism of the same height as the cylinder The prism will therefore be circumscribed to the cylinder

Let AD perpendicular to CD and FL perpendicular to EF be each equal to the perimeter of the polygon about A Bisect CD in M , and join MA



Then $\triangle KDM =$ the polygon about A

Also $\square EL =$ surface of prism (excluding bases)

Produce FE to N so that $FE = EN$, and join NL

Now the polygons about A, B , being similar, are in the duplicate ratio of the radii of A, B

Thus

$$\begin{aligned}\triangle KDM \text{ (polygon about } B) &= MD \cdot H^2 \\ &= MD^2 \cdot CD \cdot EF \\ &= MD \cdot NF \\ &= \triangle KDM \cdot \triangle LFN \quad (\text{since } DK = FL)\end{aligned}$$

Therefore (polygon about B) $= \triangle LFN$

$$= \square EL$$

$=$ (surface of prism about A),

from above

But (polygon about B) (polygon in B) $< S \cdot B$

Therefore

$$(\text{surface of prism about } A) \cdot (\text{polygon in } B) < S \cdot B,$$

and alternately,

$$(\text{surface of prism about } A) \cdot S < (\text{polygon in } B) \cdot B,$$

which is impossible since the surface of the prism is greater than S , while the polygon inscribed in B is less than B

Therefore $B < S$

II Suppose $B > S$

Let a regular polygon be circumscribed about B and another inscribed in it so that

$$(\text{polygon about } B) \cdot (\text{polygon in } B) < B \cdot S$$

Inscribe in A a polygon similar to that inscribed in B and erect a prism on the polygon inscribed in A of the same height as the cylinder

Again let DK, FL drawn as before be each equal to the perimeter of the polygon inscribed in A

Then, in this case

$$\triangle KDM > (\text{polygon inscribed in } A)$$

(since the perpendicular from the centre on a side of the polygon is less than the radius of A)

Also $\triangle LFN = \square EL =$ surface of prism (excluding bases)

Now

$$\begin{aligned}(\text{polygon in } A) \cdot (\text{polygon in } B) &= MD^2 \cdot H^2, \\ &= \triangle KDM \cdot \triangle LFN, \text{ as before}\end{aligned}$$

And

$$\triangle KDM > (\text{polygon in } A)$$

Therefore

$$\triangle LFN \text{ or (surface of prism)} > (\text{polygon in } B)$$

But this is impossible because

$$\begin{aligned}(\text{polygon about } B) \cdot (\text{polygon in } B) &< B \cdot S \\ &< (\text{polygon about } B) \cdot S \text{ a fortiori,}\end{aligned}$$

so that

$$\begin{aligned}(\text{polygon in } B) &> S \\ &> (\text{surface of prism}) \text{ a fortiori}\end{aligned}$$


Hence B is neither greater nor less than S , and therefore

$$B = S$$

PROPOSITION 14

The surface of any isosceles cone excluding the base is equal to a circle whose radius is a mean proportional between the side of the cone [a generator] and the radius of the circle which is the base of the cone

Let the circle A be the base of the cone draw C equal to the radius of the circle, and D equal to the side of the cone and let E be a mean proportional between C, D



Draw a circle B with radius equal to E

Then shall B be equal to the surface of the cone (excluding the base) which we will call S

If not, B must be either greater or less than S

I Suppose $B < S$

Let a regular polygon be described about B and a similar one inscribed in it such that the former has to the latter a ratio less than the ratio $S : B$

Describe about A another similar polygon and on it set up a pyramid with apex the same as that of the cone

Then (polygon about A) (polygon about B)

$$= C^2 : E^2$$

$$= C : D$$

$$= (\text{polygon about } A) : (\text{surface of pyramid excluding base})$$

Therefore

$$(\text{surface of pyramid}) = (\text{polygon about } B)$$

$$\text{Now } (\text{polygon about } B) : (\text{polygon in } B) < S : B$$

Therefore

$$(\text{surface of pyramid}) : (\text{polygon in } B) < S : B$$

which is impossible (because the surface of the pyramid is greater than S , while the polygon in B is less than B)

$$\text{Hence } B < S$$

II Suppose $B > S$

Take regular polygons circumscribed and inscribed to B such that the ratio of the former to the latter is less than the ratio $B : S$

Inscribe in A a similar polygon to that inscribed in B and erect a pyramid on the polygon inscribed in A with apex the same as that of the cone

In this case

$$(\text{polygon in } A) : (\text{polygon in } B) = C^2 : E^2 \\ = C : D$$

$$> (\text{polygon in } A) : (\text{surface of pyramid excluding base})$$

This is clear because the ratio of C to D is greater than the ratio of the perpendicular from the centre of A on a side of the polygon to the perpendicular from the apex of the cone on the same side

Therefore

$$(\text{surface of pyramid}) > (\text{polygon in } B)$$

But

$$(\text{polygon about } B) : (\text{polygon in } B) < B : S$$

Therefore, a *fo hori*,
 (polygon about B) (surface of pyramid) $< B S$,
 which is impossible

Since therefore B is neither greater nor less than S ,
 $B = S$

PROPOSITION 15

The surface of any iso celes cone has the same ratio to its base as the side of the cone has to the radius of the base

By Prop 14, the surface of the cone is equal to a circle whose radius is a mean proportional between the side of the cone and the radius of the base

Hence since circles are to one another as the squares of their radii, the proposition follows

PROPOSITION 16


If an iso celes cone be cut by a plane parallel to the base the portion of the surface of the cone between the parallel planes is equal to a circle whose radius is a mean proportional between (1) the portion of the side of the cone intercepted by the parallel planes and (2) the line which is equal to the sum of the radii of the circles in the parallel planes

Let OAB be a triangle through the axis of a cone, DE its intersection with the plane cutting off the frustum and OFC the axis of the cone

Then the surface of the cone OAB is equal to a circle whose radius is equal to $\sqrt{OA \cdot AC}$ [Prop 14]

Similarly the surface of the cone ODE is equal to a circle whose radius is equal to $\sqrt{OD \cdot DF}$

And the surface of the frustum is equal to the difference between the two circles



Now

$$OA \cdot AC - OD \cdot DF = DA \cdot AC + OD \cdot AC - OD \cdot DF$$

But

$$OD \cdot AC = OD \cdot DF$$

since

$$OA \cdot AC = OD \cdot DF$$

Hence

$$OA \cdot AC - OD \cdot DF = DA \cdot AC + DA \cdot DF \\ = DA \cdot (AC + DF)$$

And since circles are to one another as the squares of their radii it follows that the difference between the circles whose radii are $\sqrt{OA \cdot AC}$, $\sqrt{OD \cdot DF}$ respectively is equal to a circle whose radius is $\sqrt{DA \cdot (AC + DF)}$

Therefore the surface of the frustum is equal to this circle

LEMMAS

"1 Cones having equal height have the same ratio as their bases and those having equal bases have the same ratio as their heights"

2 If a cylinder be cut by a plane parallel to the base then, as the cylinder is to the cylinder so is the axis to the axis"

"Euclid XII 11 Cones and cylinders of equal height are to one another as their bases"

Euclid XII 14 Cones and cylinders on equal bases are to one another as their heights

"Euclid XII 13 If a cylinder be cut by a plane parallel to the opposite planes [the bases], then, as the cylinder is to the cylinder so will the axis be to the axis"

3 The cones which have the same bases as the cylinders [and equal height] are in the same ratio as the cylinders

4 Also the bases of equal cones are reciprocally proportional to their heights and those cones whose bases are reciprocally proportional to their heights are equal¹

5 Also the cones the diameters of whose bases have the same ratio as their axes are to one another in the triplicate ratio of the diameters of the bases²

And all these propositions have been proved by earlier geometers³

PROPOSITION 17

If there be two isosceles cones, and the surface of one cone be equal to the base of the other while the perpendicular from the centre of the base [of the first cone] on the side of that cone is equal to the height [of the second] the cones will be equal

Let OAB , DEF be triangles through the axes of two cones respectively, C G the centres of the respective bases GH the perpendicular from G on FD and suppose that the base of the cone OAB is equal to the surface of the cone DEF , and that $OC = GH$

Then since the base of OAB is equal to the surface of DEF
(base of cone OAB) (base of cone DEF)

$=$ (surface of DEF) (base of DEF)

$= DF \cdot FG$ [Prop 15]

$= DG \cdot GH$ by similar triangles,

$= DG \cdot OC$

Therefore the bases of the cones are reciprocally proportional to their heights whence the cones are equal [Lemma 4]

PROPOSITION 18

Any solid rhombus consisting of isosceles cones is equal to the cone which has its base equal to the surface of one of the cones composing the rhombus and its height equal to the perpendicular drawn from the apex of the second cone to one side of the first cone

Let the rhombus be $OABD$ consisting of two cones with apices O D and with a common base (the circle about AB as diameter)

Let FHK be another cone with base equal to the surface of the cone OAB and height FG equal to DE the perpendicular from D on OB

Then shall the cone FHK be equal to the rhombus

Construct a third cone LMA with base (the circle about MN) equal to the base of OAB and height LP equal to OD

¹Euclid xii 15 The bases of equal cones and cylinders are reciprocally proportional to their heights and those cones and cylinders whose bases are reciprocally proportional to their heights are equal.

²Euclid xii 12 Similar cones and cylinders are to one another in the triplicate ratio of the diameters of their bases



Then, since

$$IP = OD,$$

$$LP \cdot CD = OD \cdot CD$$

But [Lemma 1] $OD \cdot CD = (\text{rhombus } OADB) \text{ (cone } DAB),$
and $LP \cdot CD = (\text{cone } LMN) \text{ (cone } DAB)$

It follows that



$$(\text{rhombus } OADB) = (\text{cone } LMN) \quad (1)$$

Again, since $AB = MN$, and

$$(\text{surface of } OAB) = (\text{base of } FHK),$$

$$\begin{aligned} (\text{base of } FHK) \cdot (\text{base of } LMN) &= (\text{surface of } OAB) \cdot (\text{base of } OAB) \\ &= OB \cdot BC \quad [\text{Prop 15}] \\ &= OD \cdot DE, \text{ by similar triangles,} \\ &= LP \cdot FG, \text{ by hypothesis} \end{aligned}$$

Thus, in the cones FHK , LMN the bases are reciprocally proportional to the heights

Therefore the cones FHK , LMN are equal

and hence, by (1) the cone FHK is equal to the given solid rhombus

PROPOSITION 19

If an isosceles cone be cut by a plane parallel to the base and on the resulting circular section a cone be described having as its apex the centre of the base [of the first cone] and if the rhombus so formed be taken away from the whole cone, the part remaining will be equal to the cone with base equal to the surface of the portion of the first cone between the parallel planes and with height equal to the perpendicular drawn from the centre of the base of the first cone on one side of that cone

Let the cone OAB be cut by a plane parallel to the base in the circle on DF as diameter. Let C be the centre of the base of the cone and with C as apex and the circle about DE as base describe a cone making with the cone ODF the rhombus $ODCE$


Take a cone FGH with base equal to the surface of the frustum $DABF$ and height equal to the perpendicular (CH) from C on AO

Then shall the cone FGH be equal to the difference between the cone OAB and the rhombus $ODCE$

Take (1) a cone LMN with base equal to the surface of the cone OAB and height equal to CH

(2) a cone PQR with base equal to the surface of the cone ODE and height equal to CH

Now, since the surface of the cone OAB is equal to the surface of the cone ODE together with that of the frustum $DABE$, we have, by the construction,



$$(\text{base of } LMN) = (\text{base of } FGH) + (\text{base of } PQR)$$

and since the heights of the three cones are equal


$$(\text{cone } LMN) = (\text{cone } FGH) + (\text{cone } PQR)$$

But the cone LMN is equal to the cone OAB [Prop. 17] and the cone PQR is equal to the rhombus $ODCE$ [Prop. 18]

Therefore $(\text{cone } OAB) = (\text{cone } FGH) + (\text{rhombus } ODCE)$ and the proposition is proved

PROPOSITION 20

If one of the two isosceles cones forming a rhombus be cut by a plane parallel to the base and on the resulting circular section a cone be described having the same apex as the second cone and if the resulting rhombus be taken from the whole rhombus the remainder will be equal to the cone with base equal to the surface of the portion of the cone between the parallel planes and with height equal to the perpendicular drawn from the apex of the second cone to the side of the first cone



Let the rhombus be $OACB$ and let the cone OAB be cut by a plane parallel to its base in the circle about DE as diameter. With this circle as base and C

as apex describe a cone which therefore with ODE forms the rhombus $ODCE$

Take a cone FGH with base equal to the surface of the frustum $DABE$ and height equal to the perpendicular (CK) from C on OA

The cone FGH shall be equal to the difference between the rhombi $OACB$, $ODCE$

For take (1) a cone LMN with base equal to the surface of OAB and height equal to CK ,

(2) a cone PQR , with base equal to the surface of ODE , and height equal to CK

Then, since the surface of OAB is equal to the surface of ODE together with that of the frustum $DABE$ we have, by construction

$$(\text{base of } LMN) = (\text{base of } PQR) + (\text{base of } FGH),$$

and the three cones are of equal height

therefore $(\text{cone } LMN) = (\text{cone } PQR) + (\text{cone } FGH)$

But the cone LMN is equal to the rhombus $OACB$, and the cone PQR is equal to the rhombus $ODCE$ [Prop 18]

Hence the cone FGH is equal to the difference between the two rhombi $OACB$ $ODCE$

PROPOSITION 21

A regular polygon of an even number of sides being inscribed in a circle, as $ABC \dots A' \dots C'BA$ so that AA' is a diameter if two angular points next but one to each other as B, B' be joined, and the other lines parallel to BB' and joining pairs of angular points be drawn, as CC', DD' , then

$(BB' + CC' + \dots) AA' = A'B \cdot BA$

Let $BB' \dots CC' \dots DD'$, meet AA' in $F, G, H \dots$, and let $CB' \dots DC'$, be joined meeting AA' in K, L , respectively

Then clearly CB, DC' are parallel to one another and to AB

Hence by similar triangles,

$$BF : FA = BF : FK$$

$$= CG : GK$$


$$= CG : GL$$

$$= E'I : IA'$$

and summing the antecedents and consequents respectively we have

$$(BB' + CC' + \dots) AA' = BF : FA$$

$$= A'B : BA$$




PROPOSITION 22

If a polygon be inscribed in a segment of a circle LAL' so that all its sides excluding the base are equal and their number even as $LK \dots A \dots K'L'$, A being the middle point of the segment and if the lines $BB' \dots CC'$, parallel to the base LL' and joining pairs of angular points be drawn then

$$(BB' + CC' + \dots + LM) AM = A'B \cdot BA$$

where M is the middle point of LL' and AA is the diameter through M



Joining CB, DC', LK' , as in the last proposition, and supposing that they meet AM in P, Q, R while BB', CC', KH' meet AM in F, G, H we have, by similar triangles

$$\begin{aligned} BF : FA &= BF : FP \\ &= CG : PG \\ &= C'G : GQ \end{aligned}$$


$$= LM : RM,$$

and, summing the antecedents and consequents we obtain

$$\begin{aligned} (BB' + CC' + LM) : AM &= BF : FA \\ &= A'B : BA \end{aligned}$$

PROPOSITION 23

Take a great circle ABC of a sphere and inscribe in it a regular polygon whose sides are a multiple of four in number. Let AA', MM' be diameters at right angles and joining opposite angular points of the polygon



Then, if the polygon and great circle revolve together about the diameter AA' , the angular points of the polygon, except A, A' will describe circles on the surface of the sphere at right angles to the diameter AA' . Also the sides of the polygon will describe portions of conical surfaces e.g. BC will describe a surface forming part of a cone whose base is a circle about CC' as diameter and whose apex is the point in which CB, CB' produced meet each other and the diameter AA' .

Comparing the hemisphere MAM' and that half of the figure described by the revolution of the polygon

which is included in the hemisphere we see that the surface of the hemisphere and the surface of the inscribed figure have the same boundaries in one plane (viz the circle on MM' as diameter) the former surface entirely includes the latter and they are both concave in the same direction

Therefore [Assumptions 4] the surface of the hemisphere is greater than that of the inscribed figure and the same is true of the other halves of the figures

Hence the surface of the sphere is greater than the surface described by the revolution of the polygon inscribed in the great circle about the diameter of the great circle

PROPOSITION 24

If a regular polygon $AB A' B'A$, the number of whose sides is a multiple of four, be inscribed in a great circle of a sphere, and if BB' subtending two sides be joined, and all the other lines parallel to BB' and joining pairs of angular points be drawn, then the surface of the figure inscribed in the sphere by the revolution of the polygon about the diameter AA' is equal to a circle the square of whose radius is equal to the rectangle

$$BA(BB + CC' + \dots)$$


The surface of the figure is made up of the surfaces of parts of different cones

Now the surface of the cone ABB' is equal to a circle whose radius is $\sqrt{BA \cdot \frac{1}{2}BB}$ [Prop 14]

The surface of the frustum $BB' C' C$ is equal to a circle of radius $\sqrt{BC \cdot \frac{1}{2}(BB + CC')}$, [Prop 16] and so on

It follows since $BA = BC = \dots$, that the whole surface is equal to a circle whose radius is equal to

$$\sqrt{BA(BB + CC + \dots + MM + \dots + YY')}$$



PROPOSITION 25

The surface of the figure inscribed in a sphere as in the last propositions, consisting of portions of conical surfaces, is less than four times the greatest circle in the sphere

Let $AB A' B'A$ be a regular polygon inscribed in a great circle, the number of its sides being a multiple of four

As before let BB' be drawn subtending two sides, and CC' , YY' parallel to BB'

Let R be a circle such that the square of its radius is equal to

$$1B(BB + CC + \dots + YY'),$$

so that the surface of the figure inscribed in the sphere is equal to R

[Prop 24]


Now

$$(BB + CC + \dots + YY') \cdot AA' = AB \cdot AB \quad [\text{Prop 21}]$$

$$\text{whence } AB(BB + CC + \dots + YY') = AA' \cdot AB$$

$$\text{Hence } (\text{radius of } R)^2 = AB \cdot AB < AA' \cdot AB$$


Therefore the surface of the inscribed figure or the circle R is less than four times the circle $1MA M'$



PROPOSITION 26

The figure inscribed as above in a sphere is equal [in volume] to a cone whose base is a circle equal to the surface of the figure inscribed in the sphere and whose height is equal to the perpendicular drawn from the centre of the sphere to one side of the polygon

Suppose, as before that $AB A' B'A$ is the regular polygon inscribed in a great circle, and let $BB', CC',$ be joined



With apex O construct cones whose bases are the circles on $BB', CC',$ as diameters in planes perpendicular to AA'

Then $OBAB$ is a solid rhombus, and its volume is equal to a cone whose base is equal to the surface of the cone ABB' and whose height is equal to the perpendicular from O on AB [Prop 18] Let the length of the perpendicular be p

Again, if $CB C'B$ produced meet in T , the portion of the solid figure which is described by the revolution of the triangle

angle BOC about AA' is equal to the difference between the rhombi $OCTC$ and $OBTB$ i.e. to a cone whose base is equal to the surface of the frustum $BB'CC'$ and whose height is p [Prop 20]

Proceeding in this manner, and adding we prove that since cones of equal height are to one another as their bases the volume of the solid of revolution is equal to a cone with height p and base equal to the sum of the surfaces of the cone BAB' , the frustum $BB'CC'$ etc i.e. a cone with height p and base equal to the surface of the solid

PROPOSITION 27

The figure inscribed in the sphere as before is less than four times the cone whose base is equal to a great circle of the sphere and whose height is equal to the radius of the sphere

By Prop 26 the volume of the solid figure is equal to a cone whose base is equal to the surface of the solid and whose height is p the perpendicular from O on any side of the polygon Let R be such a cone

Take also a cone S with base equal to the great circle and height equal to the radius of the sphere

Now, since the surface of the inscribed solid is less than four times the great circle [Prop 25] the base of the cone R is less than four times the base of the cone S


Also the height (p) of R is less than the height of S

Therefore the volume of R is less than four times that of S and the proposition is proved

PROPOSITION 28

Let a regular polygon, whose sides are a multiple of four in number, be circumscribed about a great circle of a given sphere, as $AB A' B'A$, and about the polygon describe another circle, which will therefore have the same centre as the great circle of the sphere. Let AA' bisect the polygon and cut the sphere in a, a'

If the great circle and the circumscribed polygon revolve together about AA' the great circle will describe the surface of a sphere the angular points of the polygon except A, A' will move round the surface of a larger sphere, the points of contact of the sides of the polygon with the great circle of the inner sphere will describe circles on that sphere in planes perpendicular to AA' , and the sides of the polygon themselves will describe portions of conical surfaces. *The circumscribed figure will thus be greater than the sphere itself*



Let any side as BM touch the inner circle in K , and let A' be the point of contact of the circle with BW'

Then the circle described by the revolution of KA' about AA' is the boundary in one plane of two surfaces

- (1) the surface formed by the revolution of the circular segment KA' , and
- (2) the surface formed by the revolution of the part $KB A B'A'$ of the polygon

Now the second surface entirely includes the first, and they are both concave in the same direction

therefore [Assumptions 4] the second surface is greater than the first

The same is true of the portion of the surface on the opposite side of the circle on AA' as diameter

Hence adding we see that *the surface of the figure circumscribed to the given sphere is greater than that of the sphere itself*

PROPOSITION 29

In a figure circumscribed to a sphere in the manner shown in the previous proposition the surface is equal to a circle the square on whose radius is equal to

$$AB(BB + CC + \dots)$$

For the figure circumscribed to the sphere is inscribed in a larger sphere, and the proof of Prop 24 applies

PROPOSITION 30

The surface of a figure circumscribed as before about a sphere is greater than four times the great circle of the sphere

Let $AB A' B'A$ be the regular polygon of $4n$ sides which by its revolu

tion about AA' describes the figure circumscribing the sphere of which $ama m'$ is a great circle. Suppose aa' , AA' to be in one straight line.

Let R be a circle equal to the surface of the circumscribed solid.

Now

$$(BB' + CC' + \dots) AA' = A'B \cdot BA \quad [\text{as in Prop 21}]$$

so that

$$\begin{aligned} AB(BB' + CC' + \dots) &= AA' \cdot AB \\ \text{Hence (radius of } R) &= \sqrt{AA' \cdot AB} \quad [\text{Prop 29}] \\ &> AB \end{aligned}$$

But $AB = 2OP$ where P is the point in which AB touches the circle $ama m'$.

Therefore (radius of R) > (diameter of circle $ama m'$)

whence R , and therefore the surface of the circumscribed solid is greater than four times the great circle of the given sphere.

PROPOSITION 31

The solid of revolution circumscribed as before about a sphere is equal to a cone whose base is equal to the surface of the solid and whose height is equal to the radius of the sphere.

The solid is as before a solid inscribed in a larger sphere and, since the perpendicular on any side of the revolving polygon is equal to the radius of the inner sphere, the proposition is identical with Prop 26.

Con. *The solid circumscribed about the smaller sphere is greater than four times the cone whose base is a great circle of the sphere and whose height is equal to the radius of the sphere.*

For since the surface of the solid is greater than four times the great circle of the inner sphere [Prop 30] the cone whose base is equal to the surface of the solid and whose height is the radius of the sphere is greater than four times the cone of the same height which has the great circle for base [Lemma 1].

Hence by the proposition the volume of the solid is greater than four times the latter cone.

PROPOSITION 32

If a regular polygon with $4n$ sides be inscribed in a great circle of a sphere as $ab a' b a'$ and a similar polygon $AB A' B A'$ be described about the great circle and if the polygons revolve with the great circle about the diameters aa' , AA' respectively so that they describe the surfaces of solid figures inscribed in and circumscribed to the sphere respectively then

(1) *the surfaces of the circumscribed and inscribed figures are to one another in the duplicate ratio of their sides and*

(2) *the figures themselves [i.e. their volumes] are in the triplicate ratio of their sides*

(1) Let AA' , aa' be in the same straight line, and let $MmOm'M'$ be a diameter at right angles to them

Join BB' , CC' , and bb' , cc' , which will all be parallel to one another and MM'

Suppose R , S to be circles such that

R = (surface of circumscribed solid),

S = (surface of inscribed solid)

Then (radius of R)² = $AB(BB' + CC' + \dots)$ [Prop 29]

(radius of S)² = $ab(bb' + cc' + \dots)$ [Prop 24]

And, since the polygons are similar the rectangles in these two equations are similar and are therefore in the ratio of

$$AB^2 : ab^2$$

Hence

(surface of circumscribed solid) : (surface of inscribed solid) = $AB^2 : ab^2$

(2) Take a cone V whose base is the circle R and whose height is equal to Oa and a cone W whose base is the circle S and whose height is equal to the perpendicular from O on ab which we will call p

Then V , W are respectively equal to the volumes of the circumscribed and inscribed figures


Now since the polygons are similar

$$AB : ab = Oa : p$$

= (height of cone V) : (height of cone W),

and, as shown above the bases of the cones (the circles R , S) are in the ratio of AB^2 to ab^2

Therefore $V : W = AB^2 : ab^2$



[Props 31, 26]

PROPOSITION 33

The surface of any sphere is equal to four times the greatest circle in it

Let C be a circle equal to four times the great circle

Then if C is not equal to the surface of the sphere it must either be less or greater

1 Suppose C less than the surface of the sphere


It is then possible to find two lines β , γ of which β is the greater, such that $\beta : \gamma < (\text{surface of sphere}) : C$ [Prop 2]

Take such lines and let δ be a mean proportional between them

Suppose similar regular polygons with $4n$ sides circumscribed about and inscribed in a great circle such that the ratio of their sides is less than the ratio $\beta : \delta$ [Prop 3]

Let the polygons with the circle revolve together about a diameter common to all, describing solids of revolution as before

Then (surface of outer solid) (surface of inner solid)
 = (side of outer)² (side of inner)² [Prop 32]
 $< \beta^2 \delta^2$, or $\beta \gamma$
 $< (\text{surface of sphere}) \ C$, *a fortiori*



But this is impossible since the surface of the circumscribed solid is greater than that of the sphere [Prop 28], while the surface of the inscribed solid is less than C [Prop 25]

Therefore C is not less than the surface of the sphere

II Suppose C greater than the surface of the sphere

Take lines β, γ , of which β is the greater, such that

$$\beta \gamma < C \text{ (surface of sphere)}$$

Circumscribe and inscribe to the great circle similar regular polygons, as before such that their sides are in a ratio less than that of β to δ , and suppose solids of revolution generated in the usual manner

Then, in this case,

$$\begin{aligned} &(\text{surface of circumscribed solid}) \quad (\text{surface of inscribed solid}) \\ &\qquad\qquad\qquad < C \quad (\text{surface of sphere}) \end{aligned}$$

But this is impossible, because the surface of the circumscribed solid is greater than C [Prop 30] while the surface of the inscribed solid is less than that of the sphere [Prop 23]

Thus C is not greater than the surface of the sphere

Therefore since it is neither greater nor less C is equal to the surface of the sphere

PROPOSITION 34

Any sphere is equal to four times the cone which has its base equal to the greatest circle in the sphere and its height equal to the radius of the sphere

Let the sphere be that of which $ama m$ is a great circle

If now the sphere is not equal to four times the cone described it is either greater or less

I If possible let the sphere be greater than four times the cone

Suppose V to be a cone whose base is equal to four times the great circle and whose height is equal to the radius of the sphere

Then by hypothesis the sphere is greater than V , and two lines β, γ can be found (of which β is the greater) such that


$$\beta \cdot \gamma < (\text{volume of sphere}) \cdot V$$

Between β and γ place two arithmetic means δ, ϵ

As before let similar regular polygons with sides $4n$ in number be circumscribed about and inscribed in the great circle, such that their sides are in a ratio less than $\beta : \delta$

Imagine the diameter aa' of the circle to be in the same straight line with a diameter of both polygons and imagine the latter to revolve with the circle about aa' describing the surfaces of two solids of revolution. The volumes of these solids are therefore in the triplicate ratio of their sides [Prop 32]

Thus $(\text{vol of outer solid}) : (\text{vol of inscribed solid})$
 $< \beta^3 : \delta^3$, by hypothesis,
 $< \beta : \gamma$, *a fortiori* (since $\beta : \gamma > \beta^3 : \delta^3$),
 $< (\text{volume of sphere}) : V$, *a fortiori*



But this is impossible since the volume of the circumscribed solid is greater than that of the sphere [Prop 28] while the volume of the inscribed solid is less than V [Prop 27]

Hence the sphere is not greater than V or four times the cone described in the enunciation

II If possible let the sphere be less than V

In this case we take $\beta : \gamma$ (β being the greater) such that

$$\beta \cdot \gamma < V \quad (\text{volume of sphere})$$

The rest of the construction and proof proceeding as before we have finally
 $(\text{volume of outer solid}) : (\text{volume of inscribed solid})$
 $< V : (\text{volume of sphere})$

But this is impossible because the volume of the outer solid is greater than V [Prop 31 Cor.] and the volume of the inscribed solid is less than the volume of the sphere

Hence the sphere is not less than V

Since then the sphere is neither less nor greater than V , it is equal to V , or to four times the cone described in the enunciation

COR From what has been proved it follows that every cylinder whose base is the greatest circle in a sphere and whose height is equal to the diameter of the sphere is $\frac{2}{3}$ of the sphere, and its surface together with its bases is $\frac{2}{3}$ of the surface of the sphere

For the cylinder is three times the cone with the same base and height [Eucl XII 10] i.e. six times the cone with the same base and with height equal to the radius of the sphere

But the sphere is four times the latter cone [Prop 34] Therefore the cylinder is $\frac{2}{3}$ of the sphere

Again the surface of a cylinder (excluding the bases) is equal to a circle whose radius is a mean proportion between the height of the cylinder and the diameter of its base [Prop 13]

In this case the height is equal to the diameter of the base and therefore the circle is that whose radius is the diameter of the sphere or a circle equal to four times the great circle of the sphere

Therefore the surface of the cylinder with the bases is equal to six times the great circle

And the surface of the sphere is four times the great circle [Prop 33] whence
(surface of cylinder with bases) = $\frac{3}{2}$ (surface of sphere)

PROPOSITION 35

If in a segment of a circle LAL (where A is the middle point of the arc) a polygon $LHAK'L'$ be inscribed of which LL' is one side while the other sides are $2n$ in number and all equal, and if the polygon revolve with the segment about the diameter AM generating a solid figure inscribed in a segment of a sphere then the surface of the inscribed solid is equal to a circle the square on whose radius is equal to the rectangle

$$AB \left(BB' + CC' + \dots + KK' + \frac{LL'}{2} \right)$$

The surface of the inscribed figure is made up of portions of surfaces of cones

If we take these successively the surface of the cone BAB' is equal to a circle whose radius is

$$\sqrt{AB \cdot \frac{1}{2}BB'} \quad [\text{Prop 14}]$$

The surface of the frustum of a cone $BCC'B'$ is equal to a circle whose radius is


$$\sqrt{AB \cdot \frac{BB' + CC'}{2}}, \quad [\text{Prop 16}]$$

and so on

Proceeding in this way and adding we find since circles are to one another as the squares of their radii that the

surface of the inscribed figure is equal to a circle whose radius is

$$\sqrt{AB \left(BB' + CC' + \dots + KK' + \frac{LL'}{2} \right)}$$



PROPOSITION 36

The surface of the figure inscribed as before in the segment of a sphere is less than that of the segment of the sphere

This is clear because the circular base of the segment is a common boundary of each of two surfaces, of which one, the segment, includes the other, the solid, while both are concave in the same direction [Assumptions, 4]

PROPOSITION 37

The surface of the solid figure inscribed in the segment of the sphere by the revolution of Lh A $K'L'$ about AM is less than a circle with radius equal to AL

Let the diameter AM meet the circle of which LAL' is a segment again in A . Join $A'B$.

As in Prop 35, the surface of the inscribed solid is equal to a circle the square on whose radius is

$$AB(BB' + CC' + \dots + KK' + LM)$$


But this rectangle

$$= A'B \cdot AM \quad [\text{Prop 22}]$$

$$< A'A \cdot AM$$

$$< AL^2$$

Hence the surface of the inscribed solid is less than the circle whose radius is AL .



PROPOSITION 38


The solid figure described as before in a segment of a sphere less than a hemisphere, together with the cone whose base is the base of the segment and whose apex is the centre of the sphere is equal to a cone whose base is equal to the surface of the inscribed solid and whose height is equal to the perpendicular from the centre of the sphere on any side of the polygon

Let O be the centre of the sphere, and p the length of the perpendicular from O on AB .

Suppose cones described with O as apex and with the circles on BB' CC' as diameters as bases.

Then the rhombus $OBAB$ is equal to a cone whose base is equal to the surface of the cone $B'AB$ and whose height is p . [Prop 18]

Again if CB $C'B$ meet in T the solid described by the triangle BOC as the polygon revolves about AO is the difference between the rhombi $OCTC'$ and $OBTB$ and is therefore equal to a cone whose base is equal to the surface of



the frustum $BCCB$ and whose height is p

[Prop 20]

Similarly for the part of the solid described by the triangle COD as the poly-
gon revolves and so on

Hence, by addition the solid figure inscribed in the segment together with
the cone OLL' is equal to a cone whose base is the surface of the inscribed solid
and whose height is p

COR The cone whose base is a circle with radius equal to AL and whose height
is equal to the radius of the sphere is greater than the sum of the inscribed solid and
the cone OLL'

For by the proposition, the inscribed solid together with the cone OLL' is
equal to a cone with base equal to the surface of the solid and with height p

This latter cone is less than a cone with height equal to OA and with base
equal to the circle whose radius is AL because the height p is less than OA
while the surface of the solid is less than a circle with radius AL [Prop 37]


PROPOSITION 39

Let lal be a segment of a great circle of a sphere being less than a semicircle
Let O be the centre of the sphere and join Ol , Ol Suppose a polygon circum-
scribed about the sector $Olal$ such that its sides excluding the two radii are $2n$
in number and all equal as LK , BA , AB , KL and let OA be that
radius of the great circle which bisects the segment lal

The circle circumscribing the polygon will then have the same centre O as
the given great circle

Now suppose the polygon and the two circles to revolve together about OA
The two circles will describe spheres the angular points except A will describe

circles on the outer sphere with
diameters BB etc the points
of contact of the sides with the
inner segment will describe
circles on the inner sphere the
sides themselves will describe
the surfaces of cones or frusta
of cones and the whole figure
circumscribed to the segment of
the inner sphere by the revolu-
tion of the equal sides of the
polygon will have for its base
the circle on LL as diameter



The surface of the solid figure so circumscribed about the sector of the sphere
[excluding its base] will be greater than that of the segment of the sphere whose base
is the circle on LL as diameter

I or draw the tangents IT lT to the inner segment at l l These with the
sides of the polygon will describe by their revolution a solid whose surface is
greater than that of the segment [Assumptions 4]

But the surface described by the revolution of IT is less than that described
by the revolution of LT since the angle TLL is a right angle and therefore
 $LT > IT$

Hence a fortiori the surface described by LK A K L is greater than
that of the segment

CON *The surface of the figure so described about the sector of the sphere is equal to a circle the square on whose radius is equal to the rectangle*

$$AB (BB' + CC' + \dots + KK' + \frac{1}{2}LL')$$

For the circumscribed figure is inscribed in the outer sphere, and the proof of Prop 35 therefore applies

PROPOSITION 40

The surface of the figure circumscribed to the sector as before is greater than a circle whose radius is equal to al

Let the diameter AaO meet the great circle and the circle circumscribing the revolving polygon again in a' , A' . Join $A'B$, and let ON be drawn to N , the point of contact of AB with the inner circle

Now by Prop 39 Cor, the surface of the solid figure circumscribed to the sector $OlAl$ is equal to a circle the square on whose radius is equal to the rectangle

$$AB \left(BB' + CC' + \dots + KK' + \frac{LL'}{2} \right)$$

But this rectangle is equal to $A'B \ AM$ [as in Prop 22]

Next since $AL' \ al$ are parallel the triangles AML' and aml are similar. And $AL' > al'$ therefore $AM > am$

$$\text{Also } A'B = 2ON = aa'$$

$$\text{Therefore } A'B \ AM > am \ aa' > al^2$$

Hence the surface of the solid figure circumscribed to the sector is greater than a circle whose radius is equal to al or al


COR 1 *The volume of the figure circumscribed about the sector together with the cone whose apex is O and base the circle on LL' as diameter is equal to the volume of a cone whose base is equal to the surface of the circumscribed figure and whose height is ON*

For the figure is inscribed in the outer sphere which has the same centre as the inner. Hence the proof of Prop 38 applies

COR 2 *The volume of the circumscribed figure with the cone OLL' is greater than the cone whose base is a circle with radius equal to al and whose height is equal to the radius (Oa) of the inner sphere*

For the volume of the figure with the cone OLL' is equal to a cone whose base is equal to the surface of the figure and whose height is equal to OV

And the surface of the figure is greater than a circle with radius equal to al [Prop 40] while the heights $Oa \ OV$ are equal



PROPOSITION 41

Let lal be a segment of a great circle of a sphere which is less than a semicircle

Suppose a polygon inscribed in the sector $Olal$ such that the sides la , ab , ba , al are $2n$ in number and all equal. Let a similar polygon be circumscribed about the sector so that its sides are parallel to those of the first polygon and draw the circle circumscribing the outer polygon.

Now let the polygons and circles revolve together about OaA the radius bisecting the segment lal .

Then (1) the surfaces of the outer and inner solids of revolution so described are in the ratio of AB^2 to ab^2 , and (2) their volumes together with the corresponding cones with the same base and with apex O in each case are as AB^3 to ab^3 .

(1) For the surfaces are equal to circles the squares on whose radii are equal respectively to

$$AB \left(BB + CC' + \dots + AA' + \frac{LL'}{2} \right), \quad [\text{Prop 39 Cor}]$$

$$\text{and} \quad ab \left(bb + cc + \dots + kk + \frac{ll}{2} \right) \quad [\text{Prop 35}]$$

But these rectangles are in the ratio of AB^2 to ab^2 . Therefore so are the surfaces.

(2) Let OnN be drawn perpendicular to ab and AB , and suppose the circles which are equal to the surfaces of the outer and inner solids of revolution to be denoted by S , s respectively.

Now the volume of the circumscribed solid together with the cone OLL is equal to a

cone whose base is S and whose height is OV [Prop 40 Cor 1]

And the volume of the inscribed figure with the cone Oll is equal to a cone with base s and height On [Prop 38]

$$\text{But} \quad S/s = AB/ab^2$$


$$\text{and} \quad ON/On = AB/ab$$

Therefore the volume of the circumscribed solid together with the cone OLL is to the volume of the inscribed solid together with the cone Oll as AB^3 is to ab^3 [Lemma 5]

PROPOSITION 42

If lal be a segment of a sphere less than a hemisphere and Oa the radius perpendicular to the base of the segment the surface of the segment is equal to a circle whose radius is equal to al .

Let R be a circle whose radius is equal to al . Then the surface of the segment which we will call S must if it be not equal to R be either greater or less than R .



I Suppose, if possible, $S > R$

Let lal' be a segment of a great circle which is less than a semicircle Join Ol , Ol' , and let similar polygons with $2n$ equal sides be circumscribed and inscribed to the sector, as in the previous propositions, but such that (circumscribed polygon) (inscribed polygon) $< S \ R$

[Prop 6]

Let the polygons now revolve with the segment about OaA , generating solids of revolution circumscribed and inscribed to the segment of the sphere

Then

$$\begin{aligned} & \text{(surface of outer solid)} \quad \text{(surface of inner solid)} \\ & = AB^2 \quad ab^2 \\ & = \text{(circumscribed polygon)} \quad \text{(inscribed polygon)} \\ & < S \ R \text{ by hypothesis} \end{aligned} \quad [\text{Prop 41}]$$

But the surface of the outer solid is greater than S [Prop 39]

Therefore the surface of the inner solid is greater than R , which is impossible, by Prop 37

II Suppose if possible $S < R$

In this case we circumscribe and inscribe polygons such that their ratio is less than $R \ S$ and we arrive at the result that

$$\begin{aligned} & \text{(surface of outer solid)} \quad \text{(surface of inner solid)} \\ & < R \ S \end{aligned}$$

But the surface of the outer solid is greater than R [Prop 40] Therefore the surface of the inner solid is greater than S which is impossible [Prop 36]

Hence since S is neither greater nor less than R ,

$$S = R$$

PROPOSITION 43

Even if the segment of the sphere is greater than a hemisphere, its surface is still equal to a circle whose radius is equal to al



For let lal' be a great circle of the sphere aa' being the diameter perpendicular to ll' and let $la'l'$ be a segment less than a semicircle

Then by Prop 42 the surface of the segment $la'l'$ of the sphere is equal to a circle with radius equal to al

Also the surface of the whole sphere is equal to a circle with radius equal to aa' [Prop 33]

But $aa'^2 - a'l'^2 = al^2$ and circles are to one another as the squares on their radii


Therefore the surface of the segment $la'l'$ being the difference between the surfaces of the sphere and of $la'l$ is equal to a circle with radius equal to al



PROPOSITION 44

The volume of any sector of a sphere is equal to a cone whose base is equal to the surface of the segment of the sphere included in the sector, and whose height is equal to the radius of the sphere

Let R be a cone whose base is equal to the surface of the segment lal of a sphere and whose height is equal to the radius of the sphere, and let S be the volume of the sector Oal



β —————
 δ —————
 ϵ —————
 γ —————

Then if S is not equal to R it must be either greater or less

I Suppose, if possible, that $S > R$

Find two straight lines β , γ , of which β is the greater, such that

$$\beta \cdot \gamma < S \cdot R$$

and let δ, ϵ be two arithmetic means between β, γ

Let lal be a segment of a great circle of the sphere Join Ol Ol' and let similar polygons with $2n$ equal sides be circumscribed and inscribed to

the sector of the circle as before but such that their sides are in a ratio less than $\beta : \delta$ [Prop 4]

Then let the two polygons revolve with the segment about OaA generating two solids of revolution

Denoting the volumes of these solids by V & respectively we have

$$\begin{aligned} (V + \text{cone } OLL') - (v + \text{cone } Oll') &= AB^3 - ab^3 & [\text{Prop 41}] \\ &< \beta^3 - \delta^3 \\ &< \beta \cdot \gamma, \text{ a fortiori,} \\ &< S \cdot R \text{ by hypothesis} \end{aligned}$$

Now

$$(V + \text{cone } OLL') > S$$

Therefore also

$$(v + \text{cone } Oll') > R$$

But this is impossible by Prop 38 Cor combined with Props 42 43

Hence

$$S > R$$

II Suppose if possible that $S < R$

In this case we take β, γ such that

$$\beta \cdot \gamma < R \cdot S$$

and the rest of the construction proceeds as before

We thus obtain the relation

$$(V + \text{cone } OLL') - (v + \text{cone } Oll') < R \cdot S$$

Now

$$(v + \text{cone } Oll') < S$$

Therefore

$$(V + \text{cone } OLL') < R$$

which is impossible by Prop 40 Cor 2 combined with Props 42 43

Since then S is neither greater nor less than R

$$S = R$$

ON THE SPHERE AND CYLINDER

BOOK TWO

ARCHIMEDES to Dositheus greeting

'On a former occasion you asked me to write out the proofs of the problems the enunciations of which I had myself sent to Conon. In point of fact they depend for the most part on the theorems of which I have already sent you the demonstrations, namely (1) that the surface of any sphere is four times the greatest circle in the sphere (2) that the surface of any segment of a sphere is equal to a circle whose radius is equal to the straight line drawn from the vertex of the segment to the circumference of its base (3) that the cylinder whose base is the greatest circle in any sphere and whose height is equal to the diameter of the sphere is itself in magnitude half as large again as the sphere while its surface [including the two bases] is half as large again as the surface of the sphere, and (4) that any solid sector is equal to a cone whose base is the circle which is equal to the surface of the segment of the sphere included in the sector and whose height is equal to the radius of the sphere. Such then of the theorems and problems as depend on these theorems I have written out in the book which I send herewith, those which are discovered by means of a different sort of investigation, those namely which relate to spirals and the conoids. I will endeavour to send you soon

The first of the problems was as follows *Given a sphere to find a plane area equal to the surface of the sphere*

The solution of this is obvious from the theorems aforesaid. For four times the greatest circle in the sphere is both a plane area and equal to the surface of the sphere.

The second problem was the following

PROPOSITION 1 (PROBLEM)

Given a cone or a cylinder to find a sphere equal to the cone or to the cylinder


If Γ be the given cone or cylinder we can make a cylinder equal to Γ . Let this cylinder be the cylinder whose base is the circle on AB as diameter and whose height is OD .

Now if we could make another cylinder equal to the cylinder (OD) but such that its height is equal to the diameter of its base the problem would be solved because this latter cylinder would be equal to Γ and the sphere whose diameter is equal to the height (or to the diameter of the base) of the same cylinder would then be the sphere required [I. 31 Cor.]

Suppose the problem solved and let the cylinder (CG) be equal to the cylinder (OD) , while EF the diameter of the base is equal to the height CG .

Then since in equal cylinders the heights and bases are reciprocally proportional,

$$\frac{AB^3}{EF^3} = \frac{CG}{OD} \cdot \frac{OD}{EF} \quad (1)$$



Suppose MN to be such a line that

$$\frac{EF}{AB} = \frac{AB}{MN} \quad (2)$$

Hence

$$\frac{AB}{EF} = \frac{EF}{MN},$$

and combining (1) and (2), we have

$$\frac{AB}{MN} = \frac{EF}{OD}$$

or

$$\frac{AB}{EF} = \frac{MN}{OD}$$

Therefore

$$\frac{AB}{EF} = \frac{EF}{MN} = \frac{MN}{OD}$$

and EF , MN are two mean proportionals between AB , OD

The synopsis of the problem is therefore as follows. Take two mean proportionals EF , MN between AB and OD and describe a cylinder whose base is a circle on EF as diameter and whose height CG is equal to EF

Then since

$$\frac{AB}{EF} = \frac{EF}{MN} = \frac{MN}{OD}$$

$$EF^2 = AB \cdot MN$$

and therefore

$$\frac{AB^3}{EF^3} = \frac{AB}{MN}$$

$$= \frac{EF}{OD}$$

$$= \frac{CG}{OD}$$

whence the bases of the two cylinders (OD) (CG) are reciprocally proportional to their heights

Therefore the cylinders are equal and it follows that

$$\text{cylinder } (CG) = \frac{2}{3}V$$

The sphere on EF as diameter is therefore the sphere required being equal to $\frac{2}{3}V$

PROPOSITION 2


If BAB be a segment of a sphere BB a diameter of the base of the segment and O the centre of the sphere and if AA be the diameter of the sphere bisecting BB in M then the volume of the segment is equal to that of a cone whose base is the same as that of the segment and whose height is h where

$$h \cdot AM = OA + AM \cdot A'M$$

Measure MH along MA equal to h and MH along MA equal to h where

$$h \cdot AM = OA + AM \cdot A'M$$

Suppose the three cones constructed which have O, H, H' for their apices and the base (BB') of the segment for their common base. Join $AB, A'B$



Let C be a cone whose base is equal to the surface of the segment BAB' of the sphere i.e. to a circle with radius equal to $4B$ [I 42] and whose height is equal to OA

Then the cone C is equal to the solid sector $OBAB'$ [I 44]

Now since $HM \cdot MA = OA' + 4M \cdot A'M$,
dividendo $HA \cdot AM = OA \cdot A'M$,
 and, alternately $HA \cdot AO = AM \cdot MA'$,
 so that

$$\begin{aligned} HO \cdot OA &= AA' \cdot A'M \\ &= AB^2 \cdot BM^2 \\ &= (\text{base of cone } C) \cdot (\text{circle on } BB \text{ as diameter}) \end{aligned}$$

But OA is equal to the height of the cone C , therefore since cones are equal if their bases and heights are reciprocally proportional it follows that the cone C (or the solid sector $OBAB'$) is equal to a cone whose base is the circle on BB' as diameter and whose height is equal to OH

And this latter cone is equal to the sum of two others having the same base and with heights OM, MH i.e. to the solid rhombus $OBHB'$

Hence the sector $OBAB'$ is equal to the rhombus $OBHB'$

Taking away the common part, the cone OBB' ,
 the segment BAB = the cone HBB'

Similarly by the same method we can prove that
 the segment $BA'B'$ = the cone $H'BB'$

Alternative proof of the latter property

Suppose D to be a cone whose base is equal to the surface of the whole sphere and whose height is equal to OA

Thus D is equal to the volume of the sphere

[I 33 31]

Now since $OA + AM \cdot 4M = HM \cdot MA$
dividendo and alternando, as before,

$$OA \cdot AH = A'M \cdot MA$$

Again since $HM \cdot MA' = OA + AM \cdot AM$,

$$H \cdot A' \cdot OA = A \cdot M \cdot MA$$

$$= OA \cdot AH, \text{ from above}$$

Componendo, $HO \cdot OA = OH \cdot H' A$

Alternately $H'O \cdot OH = OA \cdot AH$

(1)

(2)

and *componendo*,

$$\begin{aligned} HH' \quad HO &= OH \quad HA, \\ &= H'O \quad OA \text{ from (1),} \end{aligned}$$

whence

$$HH \quad OA = HO \quad OH \quad (3)$$

Next since

$$\begin{aligned} HO \quad OH &= OA \quad AH \text{ by (2)} \\ &= A'M \quad MA \end{aligned}$$

$$(H'O + OH)^2 \quad HO \quad OH = (A'M + MA)^2 \quad A'M \quad MA,$$

whence, by means of (3),

$$HH \quad HH \quad OA = AA' \quad A'M \quad MA$$

or

$$HH \quad OA = AA'^2 \quad BM^2$$

Now the cone D which is equal to the sphere has for its base a circle whose radius is equal to AA' , and for its height a line equal to OA

Hence this cone D is equal to a cone whose base is the circle on BB as diameter and whose height is equal to HH

therefore

the cone D = the rhombus $HBH B$,

or

the rhombus $HBH B$ = the sphere

But

the segment BAB = the cone HBB


therefore the remaining segment $BA B$ = the cone HBB

COR The segment BAB is to a cone with the same base and equal height in the ratio of $OA + A'M$ to $A'M$

PROPOSITION 3 (PROBLEM)

To cut a given sphere by a plane so that the surfaces of the segments may have to one another a given ratio

Suppose the problem solved Let AA' be a diameter of a great circle of the sphere and suppose that a plane perpendicular to AA cuts the plane of the great circle in the straight line BB and AA in M , and that it divides the sphere so that the surface of the segment BAB has to the surface of the segment $BA B$ the given ratio



Now these surfaces are respectively equal to circles with radii equal to AB , $A'B$ [I 42 43]

Hence the ratio $AB^2 : A'B^2$ is equal to the given ratio i.e. AM is to MA in the given ratio

Accordingly the synthesis proceeds as follows

If $H : A$ be the given ratio divide $A 1$ in M so that

$$AM : MA = H : A$$

Then $AM : MA = AB^2 : A'B^2$

$$= (\text{circle with radius } AB) : (\text{circle with radius } A'B)$$

$$= (\text{surface of segment } BAB) : (\text{surface of segment } BAB)$$


Thus the ratio of the surfaces of the segments is equal to the ratio $H : K$

PROPOSITION 4 (PROBLEM)

To cut a given sphere by a plane so that the volumes of the segments are to one another in a given ratio

Suppose the problem solved and let the required plane cut the great circle

ABA' at right angles in the line BB' let AA' be that diameter of the great circle which bisects BB' at right angles (in M), and let O be the centre of the sphere



Take H on OA produced, and H' on OA' produced such that

$$OA + A'M : A'M = HM : MA, \quad (1)$$

and
$$OA + AM : AM = H'M : MA' \quad (2)$$

Join $BH, B'H, BH', B'H'$

Then the cones $HBB' : H'B'B'$ are respectively equal to the segments $BAB', BA'B'$ of the sphere [Prop 2]

Hence the ratio of the cones and therefore of their altitudes, is given, i.e.

$$HM : H'M = \text{the given ratio} \quad (3)$$

We have now three equations (1) (2) (3) in which there appear three as yet undetermined points M, H, H' , and it is first necessary to find, by means of them, another equation in which only one of these points (M) appears, i.e. we have so to speak to eliminate H, H'

Now, from (3), it is clear that $HH' : H'M$ is also a given ratio, and Archimedes' method of elimination is first, to find values for each of the ratios $A'H' : H'M$ and $HH' : H'A'$ which are alike independent of H, H' , and then secondly, to equate the ratio compounded of these two ratios to the known value of the ratio $HH' : H'M$

(a) To find such a value for $A'H' : H'M$

It is at once clear from equation (2) above that

$$A'H' : H'M = OA : OA' + AM \quad (4)$$

(b) To find such a value for $HH' : H'A'$

From (1) we derive

$$\begin{aligned} AM : MA &= OA' + A'M : HM \\ &= OA' : AH' \end{aligned} \quad (5)$$

and from (2) $A'M : MA = H'M : OA' + AM$

$$= H'M : OA' \quad (6)$$

Thus $HA : AO = OA' : AH'$,

whence $OH : OA' = OH' : A'H'$,

or $OH : OH' = OA' : A'H'$

It follows that

$$HH' : OH' = OH' : A'H',$$

or $HH' : H'A' = OH'^2$

Therefore $HH' : H'A' = OH'^2 : H'A'^2$
 $= AA'^2 : A'M^2$ by means of (6)

(c) To express the ratios $A'H' : H'M$ and $HH' : H'M$ more simply we make

the following construction Produce OA to D so that $OA = AD$ (D will lie beyond H , for $A'H > MA$ and therefore by (5) $OA > AH$)

$$\begin{aligned} \text{Then} \quad AH \cdot H'M &= OA \cdot OA + AM \\ &= AD \cdot DM \end{aligned} \quad (7)$$

Now divide AD at E so that

$$HH \cdot H'M = AD \cdot DE \quad (8)$$

Thus using equations (8) (7) and the value of $HH \cdot H'A'$ above found we have

$$\begin{aligned} AD \cdot DE &= HH \cdot H'M \\ &= (HH \cdot HA') (AH \cdot HM) \\ &= (AA^2 - AM^2) (AD \cdot DM) \end{aligned}$$

$$\text{But} \quad AD \cdot DE = (DM \cdot DE) (AD \cdot DM) \quad (9)$$

Therefore $MD \cdot DE = AA^2 - A'M^2$
And D is given since $AD = OA$. Also $AD \cdot DE$ (being equal to $HH \cdot H'M$) is a given ratio. Therefore DE is given.

Hence the problem reduces itself to the problem of dividing $A'D$ into two parts at M so that

$$MD \text{ (a given length)} = (\text{a given area}) \cdot A'M$$

Archimedes adds 'If the problem is propounded in this general form it requires a *διορισμός* [i.e. it is necessary to investigate the limits of possibility] but if there be added the conditions subsisting in the present case it does not require a *διορισμός*'

In the present case the problem is

Given a straight line $A'A$ produced to D so that $AA = 2AD$ and given a point E on AD to cut AA in a point M so that

$$AA^2 - A'M^2 = MD \cdot DE$$

And the analysis and synthesis of both problems will be given at the end.¹

The synthesis of the main problem will be as follows. Let $R \cdot S$ be the given ratio R being less than S , AA being a diameter of a great circle and O the centre produce OA to D so that $OA = AD$ and divide AD in E so that

$$AE \cdot ED = R \cdot S$$

Then cut AA in M so that

$$MD \cdot DE = AA \cdot A'M^2$$

Through M erect a plane perpendicular to AA this plane will then divide the sphere into segments which will be to one another as R to S .

Take H on AA produced and H on AA produced so that

$$OA + AM \cdot AM = HM \cdot MA \quad (1)$$

$$OA + HM \cdot AM = HM \cdot MA \quad (2)$$

We have then to show that

$$HM \cdot MH = R \cdot S \text{ or } AE \cdot ED$$

(a) We first find the value of $HH \cdot HA$ as follows

As was shown in the analysis (b)

$$HH \cdot HA = OH^2$$

$$\text{or} \quad HH \cdot HA = OH^2 - H'A'^2$$

$$= AA^2 - A'M^2$$

$$= MD \cdot DE \text{ by construction}$$

¹As Archimedes' commentator Eutocius notes we do not find the promise kept in any of the copies. Sir Thomas Heath's translation of Eutocius' note on the matter along with the solutions of Dionysodorus and Dioctes is omitted from this edition.—Ed

(β) Next we have

$$\begin{aligned} H'A' \cdot H'M &= OA \cdot OA + AM \\ &= AD \cdot DM \end{aligned}$$

$$\begin{aligned} \text{Therefore } HH' \cdot HM &= (HH' \cdot H'A') (H'A' \cdot H'M) \\ &= (MD \cdot DE) (AD \cdot DM) \\ &= AD \cdot DE \end{aligned}$$

$$\begin{aligned} \text{whence } HM \cdot MH' &= AE \cdot ED \\ &= R \cdot S \end{aligned}$$

Q.E.D.


PROPOSITION 5 (PROBLEM)

To construct a segment of a sphere similar to one segment and equal in volume to another

Let ABP' be one segment whose vertex is A and whose base is the circle on BB' as diameter and let DEF be another segment whose vertex is D and whose base is the circle on FF' as diameter. Let AA' , DD' be diameters of the great circles passing through BB' , FF' respectively, and let O , C be the respective centres of the spheres.

Suppose it required to draw a segment similar to DEF and equal in volume to ABP' .

Analysis. Suppose the problem solved, and let def be the required segment, d being the vertex and ef the diameter of the base. Let dd' be the diameter of the sphere which bisects ef at right angles, c the centre of the sphere.



Let M , C , g be the points where BB' , FF' , ef are bisected at right angles by AA' , DD' , dd' respectively and produce OA , CD , cd respectively to H , K , I so that

$$\left. \begin{aligned} OA + A'M \cdot A'M &= HM \cdot MA \\ CD + DG \cdot DG &= KG \cdot GD \\ cd + dg \cdot dg &= Ig \cdot gd \end{aligned} \right\},$$

and suppose cones formed with vertices H , K , I and with the same bases as the respective segments. The cones will then be equal to the segments respectively [Prop 2]

Therefore by hypothesis,

the cone HBB = the cone λef

Hence

(circle on diameter BB) (circle on diameter ef) = λg HM ,

so that $BB^2 \cdot ef^2 = \lambda g \cdot HM$ (1)

But since the segments DEF def are similar so are the cones λEF λef

Therefore $\lambda C \cdot EF = \lambda g \cdot ef$

And the ratio $\lambda G \cdot EF$ is given Therefore the ratio $\lambda g \cdot ef$ is given

Suppose a length R taken such that

$$\lambda g \cdot ef = HM \cdot R \quad (2)$$

Thus R is given

Again since $\lambda g \cdot HM = BB'^2 \cdot ef^2 = ef^2 \cdot R$, by (1) and (2), suppose a length S taken such that

$$ef^2 = BB'^2 \cdot S$$

$$\text{or } BB'^2 \cdot ef^2 = BB^2 \cdot S$$

$$\text{Thus } BB' \cdot ef = ef \cdot S = S \cdot R,$$

and ef , S are two mean proportionals in continued proportion between BB , R

Synthesis Let ABB DEF be great circles AA' , DD' the diameters bisecting BB' , EF at right angles in M , G respectively, and O , C the centres

Take H , λ in the same way as before and construct the cones HBB λEF , which are therefore equal to the respective segments ABB' , DEI

Let R be a straight line such that

$$\lambda G \cdot EF = HM \cdot R$$

and between BB' R take two mean proportionals ef , S

On ef as base describe a segment of a circle with vertex d and similar to the segment of a circle DEF Complete the circle, and let dd' be the diameter through d and c the centre Conceive a sphere constructed of which def is a great circle and through ef draw a plane at right angles to dd'

Then shall def be the required segment of a sphere

For the segments DEF def of the spheres are similar, like the circular segments DEF def

Produce cd to λ so that

$$cd + d'g \cdot dg = \lambda g \cdot gd$$

The cones λEF , λef are then similar

Therefore $\lambda g \cdot ef = \lambda G \cdot EF = HM \cdot R$,

whence $\lambda g \cdot HM = ef \cdot R$

But since BB ef S λ are in continued proportion

$$BB \cdot ef = BB^2 \cdot S$$

$$= ef \cdot R$$

$$= \lambda g \cdot HM$$

Thus the bases of the cones HBB , λef are reciprocally proportional to their heights The cones are therefore equal and def is the segment required being equal in volume to the cone λef [Prop 2]

PROPOSITION 6 (PROBLEM)

Given two segments of spheres to find a third segment of a sphere similar to one of the given segments and having its surface equal to that of the other

Let ABB be the segment to whose surface the surface of the required segment is to be equal AB B the great circle whose plane cuts the plane of the

base of the segment ABB' at right angles in BB' . Let AA' be the diameter which bisects BB' at right angles.

Let DEF be the segment to which the required segment is to be similar, $DED'F$ the great circle cutting the base of the segment at right angles in EF . Let DD' be the diameter bisecting EF at right angles in G .

Suppose the problem solved def being a segment similar to DEF and having its surface equal to that of ABB' and complete the figure for def as for DEF , corresponding points being denoted by small and capital letters respectively.

Join AB, DF, df .

Now, since the surfaces of the segments def, ABB' are equal, so are the circles on df, AB as diameters.

[I 42 43]

that is $df = AB$.

From the similarity of the segments DEF, def we obtain

$$d d dg = D D DG,$$

and $dg df = DG DF$,

whence $d'd df = D'D DF$,

or $d'd AB = D'D DF$.

But $AB, D'D, DF$ are all given.

therefore $d'd$ is given.

Accordingly the synthesis is as follows.

Take $d d$ such that

$$d d AB = D D DF \quad (1)$$

Describe a circle on $d d$ as diameter, and conceive a sphere constructed of which this circle is a great circle.

Divide $d d$ at g so that

$$d'g gd = DG GD$$

and draw through g a plane perpendicular to $d d$ cutting off the segment def of the sphere and intersecting the plane of the great circle in ef . The segments def, DEF are thus similar and

$$dg df = DG DF$$

But from above *componendo*

$$d'd dg = D'D DG$$

Therefore *ex aequali*

$$d'd df = D D DF$$

whence by (1) $df = AB$.

Therefore the segment def has its surface equal to the surface of the segment ABB' [I 42 43], while it is also similar to the segment DEF .

PROPOSITION 7 (PROBLEM)

From a given sphere to cut off a segment by a plane so that the segment may have a given ratio to the cone which has the same base as the segment and equal height.

Let AA' be the diameter of a great circle of the sphere. It is required to draw a plane at right angles to AA' cutting off a segment ABB' , such that the segment ABB' has to the cone ABB' a given ratio.

Analysis

Suppose the problem solved and let the plane of section cut the plane of the great circle in BB' and the diameter AA' in M . Let O be the centre of the sphere.

Produce OA' to H so that

$$OA' + A'M : A'M = HM : MA \quad (1)$$

Thus the cone HBB' is equal to the segment ABB' .

[Prop 2]

Therefore the given ratio must be equal to the ratio of the cone HBB' to the cone ABB' , i.e. to the ratio $HM : MA$.

Hence the ratio $OA' + A'M : A'M$ is given and therefore $A'M$ is given.

διορισμός

Now
so that

$$\begin{aligned} OA' : A'M &> OA' : A'A \\ OA' + A'M : A'M &> OA' + A'A : A'A \\ &> 3 : 2 \end{aligned}$$

Thus in order that a solution may be possible it is a necessary condition that the given ratio must be greater than $3 : 2$.

The synthesis proceeds thus.

Let AA' be a diameter of a great circle of the sphere O the centre.

Take a line DE and a point F on it, such that $DE : EF$ is equal to the given ratio being greater than $3 : 2$.

Now since

$$\begin{aligned} OA' + A'A : A'A &= 3 : 2 \\ DE : EF &> OA' + A'A : A'A \\ DF : FE &> OA' : A'A \end{aligned}$$

so that

Hence a point M can be found on AA' such that

$$DF : FE = OA' : A'M \quad (2)$$

Through M draw a plane at right angles to AA' intersecting the plane of the great circle in BB' and cutting off from the sphere the segment ABB' .

As before take H on OA' produced such that

$$OA' + AM : AM = HM : MA$$

Therefore $HM : MA = DE : EF$ by means of (2).

It follows that the cone HBB' or the segment ABB' is to the cone ABB' in the given ratio $DE : EF$.

PROPOSITION 8

If a sphere be cut by a plane not passing through the centre into two segments ABB' $A'B'B$, of which $A'B'B$ is the greater, then the ratio (segment $A'B'B$) (segment ABB')

$$\begin{aligned} &< (\text{surface of } A'B'B)^2 : (\text{surface of } ABB') \\ &\text{but} > (\text{surface of } A'B'B)^3 : (\text{surface of } ABB')^3 \end{aligned}$$

Let the plane of section cut a great circle $ABAB'$ at right angles in BB' , and let AA' be the diameter bisecting BB' at right angles in M .

Let O be the centre of the sphere.

R will fall between O and M .

Also, since $AB^2 = DE$, $AR = CD$

Produce OA' to K so that $OA' = A'K$, and produce $A'A$ to H so that

$$A'K : A'M = HA : AM,$$

or *componendo* $A'K + A'M : A'M = HA + MA : MA$ (1)

Thus the cone HBB' is equal to the segment ABB' [Prop 2]

Again, produce CD to F so that $CD = DF$, and the cone FEE' will be equal to the hemisphere DEE' [Prop 2]

Now $AR : RA' > AM : MA'$,

and $AR^2 = \frac{1}{2}AB^2 = \frac{1}{2}AM \cdot AA' = AM \cdot A'K$

Hence

$$AR : RA' + RA^2 > AM : MA' + AM \cdot A'K,$$

or $AA' : AR > AM : MK$

$$> HM : A'M, \text{ by (1)}$$

Therefore $AA' : A'M > HM : AR$,

or $AB^2 : BM^2 > HM : AR$,

i.e. $AR^2 : BM^2 > HM : 2AR$ since $AB^2 = 2AR$,

$$> HM : CF$$

Thus since $AR = CD$, or CE ,

$$(\text{circle on diam } EE') : (\text{circle on diam } BB') > HM : CF$$

It follows that



$$(\text{the cone } FEE') > (\text{the cone } HBB'),$$

and therefore the hemisphere DEE' is greater in volume than the segment ABB'

MEASUREMENT OF A CIRCLE

PROPOSITION 1

The area of any circle is equal to a right-angled triangle in which one of the sides about the right angle is equal to the radius, and the other to the circumference, of the circle



Let $ABCD$ be the given circle K the triangle described

Then if the circle is not equal to K it must be either greater or less

I If possible let the circle be greater than K

Inscribe a square $ABCD$ bisect the arcs AB BC CD DA then bisect (if necessary) the halves and so on until the sides of the inscribed polygon whose angular points are the points of division subtend segments whose sum is less than the excess of the area of the circle over K

Thus the area of the polygon is greater than K

Let AE be any side of it and ON the perpendicular on AE from the centre O

Then ON is less than the radius of the circle and therefore less than one of the sides about the right angle in K . Also the perimeter of the polygon is less than the circumference of the circle i.e. less than the other side about the right angle in K

Therefore the area of the polygon is less than K which is inconsistent with the hypothesis

Thus the area of the circle is not greater than K

II If possible let the circle be less than K

Circumscribe a square and let two adjacent sides touching the circle in E H meet in T . Bisect the arcs between adjacent points of contact and draw the

tangents at the points of bisection. Let A be the middle point of the arc EH , and FAG the tangent at A .

Then the angle TAG is a right angle.

Therefore $TG > GA$
 $> GH$

It follows that the triangle FTG is greater than half the area $TEAH$.

Similarly, if the arc AH be bisected and the tangent at the point of bisection be drawn, it will cut off from the area GAH more than one-half.

Thus, by continuing the process we shall ultimately arrive at a circumscribed polygon such that the spaces intercepted between it and the circle are together less than the excess of A over the area of the circle.

Thus the area of the polygon will be less than A .

Now, since the perpendicular from O on any side of the polygon is equal to the radius of the circle while the perimeter of the polygon is greater than the circumference of the circle it follows that the area of the polygon is greater than the triangle A which is impossible.

Therefore the area of the circle is not less than A .

Since then the area of the circle is neither greater nor less than A , it is equal to it.

PROPOSITION 2

The area of a circle is to the square on its diameter as 11 to 14¹

PROPOSITION 3

The ratio of the circumference of any circle to its diameter is less than 3 $\frac{1}{4}$ but greater than 3 $\frac{1}{2}$.

I let AB be the diameter of any circle O its centre AC the tangent at A and let the angle AOC be one-third of a right angle.

Then $OA : AC [= \sqrt{3} : 1] > 265 : 153$ (1)

and $OC : CA [= 2 : 1] = 306 : 163$ (2)

First draw OD bisecting the angle COA and meeting AC in D .

Now $CO : OA = CD : DA$ [Eucl VI 3]

so that $[CO + OA : OA = CD + DA : DA]$ or

$$CO + OA : CA = OA : AD$$

Therefore [by (1) and (2)]

$$OA : AD > 571 : 153 \quad (3)$$

The text of this proposition is not satisfactory and Archimedes cannot have placed it before Proposition 3 as the approximation depends upon the result of that proposition.


In view of the interesting questions arising out of the arithmetical content of this proposition of Archimedes it is necessary in reproducing it to distinguish carefully the actual steps set out in the text as we have it from the intermediate steps (mostly supplied by Eutocius) which it is convenient to put in for the purpose of making the proof easier to follow. Accordingly all the steps not actually appearing in the text have been enclosed in square brackets in order that it may be clearly seen how far Archimedes omits actual calculations and only gives results. It will be observed that he gives two fractional approximations to $\sqrt{3}$ (one being less and the other greater than the real value) without any explanation as to how he arrived at them and in like manner approximations to the square roots of several large numbers which are not complete squares are merely stated.

'Hence

$$\begin{aligned} OD^2 AD &= (OA + AD) AD^2 \\ &> (571 + 153^2) 153 \\ &> 349450 \quad 23409, \end{aligned}$$

so that

$$OD DA > 591\frac{1}{8} 153 \quad (4)$$



Secondly let OE bisect the angle AOD , meeting AD in E

[Then $DO OA = DE EA$]

so that $DO + OA DA = OA AE$

Therefore $OA AE [> (591\frac{1}{8} + 571) 153, \text{ by (3) and (4)}]$
 $> 1162\frac{1}{8} 153$

(5)

[It follows that

$$\begin{aligned} OE^2 EA^2 &> \{ (1162\frac{1}{8})^2 + 153^2 \} 153^2 \\ &> (1350534\frac{1}{4} + 23409) 23409 \\ &> 1373943\frac{3}{4} 23409 \end{aligned}$$

Thus $OE EA > 1172\frac{1}{8} 153$

(6)

Thirdly let OF bisect the angle AOE and meet AE in F

We thus obtain the result [corresponding to (3) and (5) above] that

$$\begin{aligned} OA AF &[> (1162\frac{1}{8} + 1172\frac{1}{8}) 153] \\ &> 2334\frac{1}{4} 153 \end{aligned}$$

(7)

[Therefore $OF^2 FA^2 > \{ (2334\frac{1}{4})^2 + 153^2 \} 153^2$
 $> 5472132\frac{1}{4} 23409$]

Thus $OF FA > 2339\frac{1}{4} 153$

(8)

Fourthly let OG bisect the angle AOF meeting AF in G

We have then

$$\begin{aligned} OA AG &[> (2334\frac{1}{4} + 2339\frac{1}{4}) 153 \text{ by means of (7) and (8)}] \\ &> 4673\frac{1}{2} 153 \end{aligned}$$

Now the angle AOC which is one third of a right angle has been bisected four times and it follows that

$$\angle AOG = \frac{1}{16} \text{ (a right angle)}$$

Make the angle AOH on the other side of O equal to the angle $10C$ and let GA produced meet OH in H

Then $\angle GOH = \frac{1}{16} \text{ (a right angle)}$

Thus GH is one side of a regular polygon of 96 sides circumscribed to the given circle

And, since $OA \ AG > 4673\frac{1}{2} \ 153$,
while $AB = 2OA, GH = 2AG$,¹¹

it follows that

$$AB \text{ (perimeter of polygon of 96 sides)} \left[\begin{array}{l} > 4673\frac{1}{2} \ 153 \times 96 \\ > 4673\frac{1}{2} \ 14688 \end{array} \right]$$

$$\begin{aligned} \text{But} \quad \frac{14688}{4673\frac{1}{2}} &= 3 + \frac{667\frac{1}{2}}{4673\frac{1}{2}} \\ &\left[< 3 + \frac{667\frac{1}{2}}{4672\frac{1}{2}} \right] \\ &< 3\frac{1}{2} \end{aligned}$$

Therefore the circumference of the circle (being less than the perimeter of the polygon) is *a fortiori* less than $3\frac{1}{2}$ times the diameter AB

II Next let AB be the diameter of a circle, and let AC , meeting the circle in C make the angle CAB equal to one-third of a right angle Join BC

Then $AC \ CB [= \sqrt{3} \ 1] < 1351 \ 780$

First let AD bisect the angle BAC and meet BC in d and the circle in D Join BD

$$\begin{aligned} \text{Then} \quad \angle BAD &= \angle DAC \\ &= \angle CBD \end{aligned}$$

and the angles at D, C are both right angles

It follows that the triangles $ADB \ [ACd] \ BDb$ are similar

$$\begin{aligned} \text{Therefore} \quad AD \ DB &= BD \ Dd \\ &[= AC \ Cd] \\ &= AB \ Bd \\ &= AB + AC \ Bd + Cd \\ &= AB + AC \ BC \end{aligned} \quad [\text{Eucl vi } 3]$$

$$\begin{aligned} \text{or} \quad BA + AC \ BC &= AD \ DB \\ \text{[But} \quad AC \ CB &< 1351 \ 780, \text{ from above,} \\ \text{while} \quad B \ 1 \ BC &= 2 \ 1 \\ &= 1560 \ 780 \end{aligned}$$

$$\text{Therefore} \quad AD \ DB < 2911 \ 780 \quad (1)$$

$$\begin{aligned} \text{[Hence} \quad AB^2 \ BD^2 &< (2911^2 + 780^2) \ 780 \\ &< 9082321 \ 608100 \end{aligned}$$

$$\text{Thus} \quad AB \ BD < 3013\frac{1}{2} \ 780 \quad (2)$$

Secondly let $1E$ bisect the angle BAD , meeting the circle in E and let BE be joined

Then we prove in the same way as before that

$$\begin{aligned} 1E \ EB [= BA + AD \ BD] &< (3013\frac{1}{2} + 2911) \ 780 \text{ by (1) and (2)} \\ &< 5924\frac{1}{2} \ 780 \\ &< 5924\frac{1}{2} \times \frac{1}{2} \ 780 \times \frac{1}{2} \\ &< 1823 \ 210 \end{aligned} \quad (3)$$


$$\begin{aligned} \text{[Hence} \quad AB^2 \ BE^2 &< (1823 + 210) \ 210^2 \\ &< 3380929 \ 57600 \end{aligned} \quad (4)$$

$$\text{Therefore} \quad 1B \ BE < 1838\frac{1}{2} \ 210$$

Thirdly, let AF bisect the angle BAE meeting the circle in F

Thus

$$\begin{aligned}
 AF \cdot FB & [= BA + AE \cdot BE] \\
 & < 3661 \frac{1}{11} \text{ or } 240 \text{ by (3) and (4)} \\
 & < 3661 \frac{1}{11} \times \frac{1}{11} \cdot 240 \times \frac{1}{11} \\
 & < 1007 \cdot 66
 \end{aligned}
 \tag{5}$$



[It follows that

$$\begin{aligned}
 AB^2 - BF^2 & < (1007^2 + 66^2) \cdot 66^2 \\
 & < 1018405 \cdot 4356]
 \end{aligned}
 \tag{6}$$

Therefore

$$AB \cdot BF < 1009 \frac{1}{2} \cdot 66$$

Fourthly, let the angle BAF be bisected by AG meeting the circle in G

Then

$$\begin{aligned}
 AG \cdot GB & [= BA + AF \cdot BF] \\
 & < 2016 \frac{1}{2} \cdot 66, \text{ by (5) and (6)}
 \end{aligned}$$

[And

$$\begin{aligned}
 AB^2 - BG^2 & < \{(2016 \frac{1}{2})^2 + 66^2\} \cdot 66^2 \\
 & < 4069284 \frac{1}{4} \cdot 4356]
 \end{aligned}$$

Therefore

$$AB \cdot BG < 2017 \frac{1}{2} \cdot 66$$

whence

$$BG \cdot AB > 66 \cdot 2017 \frac{1}{2}
 \tag{7}$$

[Now the angle BAG which is the result of the fourth bisection of the angle BAC or of one-third of a right angle is equal to one-fortyeighth of a right angle

Thus the angle subtended by BG at the centre is

$$\frac{1}{16} \text{ (a right angle)}$$

Therefore BG is a side of a regular inscribed polygon of 96 sides

It follows from (7) that

$$\begin{aligned}
 (\text{perimeter of polygon}) \cdot AB & [> 96 \times 66 \cdot 2017 \frac{1}{2}] \\
 & > 6336 \cdot 2017 \frac{1}{2}
 \end{aligned}$$

And

$$\frac{6336}{2017 \frac{1}{2}} > 3 \frac{1}{4}$$

Much more then is the circumference of the circle greater than $3 \frac{1}{4}$ times the diameter

Thus the ratio of the circumference to the diameter

$$< 3 \frac{1}{2} \text{ but } > 3 \frac{1}{4}$$

ON CONOIDS AND SPHEROIDS

INTRODUCTION¹

"ARCHIMEDES to Dositheus greeting

"In this book I have set forth and send you the proofs of the remaining theorems not included in what I sent you before and also of some others discovered later which, though I had often tried to investigate them previously, I had failed to arrive at because I found their discovery attended with some difficulty. And this is why even the propositions themselves were not published with the rest. But afterwards, when I had studied them with greater care, I discovered what I had failed in before.

"Now the remainder of the earlier theorems were propositions concerning the right angled conoid [paraboloid of revolution] but the discoveries which I have now added relate to an obtuse-angled conoid [hyperboloid of revolution] and to spheroidal figures some of which I call *oblong* and others *flat*."

I Concerning the *right-angled conoid* it was laid down that, if a section of a right-angled cone [a parabola] be made to revolve about the diameter [axis] which remains fixed and return to the position from which it started, the figure comprehended by the section of the right angled cone is called a *right-angled conoid* and the diameter which has remained fixed is called its *axis*, while its *vertex* is the point in which the axis meets the surface of the conoid. And if a plane touch the right-angled conoid, and another plane drawn parallel to the tangent plane cut off a segment of the conoid the *base* of the segment cut off is defined as the portion intercepted by the section of the conoid on the cutting plane the *vertex* [of the segment] as the point in which the first plane touches the conoid and the *axis* [of the segment] as the portion cut off within the segment from the line drawn through the vertex of the segment parallel to the axis of the conoid.

'The questions propounded for consideration were'

(1) why if a segment of the right angled conoid be cut off by a plane at right angles to the axis will the segment so cut off be half as large again as the cone which has the same base as the segment and the same axis, and

(2) 'why if two segments be cut off from the right angled conoid by planes drawn in any manner will the segments so cut off have to one another the duplicate ratio of their axes

II "Respecting the *obtuse-angled conoid* we lay down the following premises. If there be in a plane a section of an obtuse-angled cone [a hyperbola], its

¹The whole of this introductory matter including the definitions is translated literally from the Greek text in order that the terminology of Archimedes may be faithfully represented. When this has once been set out nothing will be lost by returning to modern phraseology and notation. These will accordingly be employed as usual, when we come to the actual propositions of the treatise.

diameter [axis], and the nearest lines to the section of the obtuse-angled cone [i.e. the asymptotes of the hyperbola], and if, the diameter [axis] remaining fixed, the plane containing the aforesaid lines be made to revolve about it and return to the position from which it started the nearest lines to the section of the obtuse-angled cone [the asymptotes] will clearly comprehend an isosceles cone whose vertex will be the point of concurrence of the nearest lines and whose axis will be the diameter [axis] which has remained fixed. The figure comprehended by the section of the obtuse-angled cone is called an *obtuse-angled conoid* [hyperboloid of revolution] its axis is the diameter which has remained fixed, and its *vertex* the point in which the axis meets the surface of the conoid. The cone comprehended by the nearest lines to the section of the obtuse-angled cone is called [the cone] *enveloping the conoid* and the straight line between the vertex of the conoid and the vertex of the cone enveloping the conoid is called [the line] *adjacent to the axis*. And if a plane touch the obtuse-angled conoid and another plane drawn parallel to the tangent plane cut off a segment of the conoid, the base of the segment so cut off is defined as the portion intercepted by the section of the conoid on the cutting plane the *vertex* [of the segment] as the point of contact of the plane which touches the conoid the axis [of the segment] as the portion cut off within the segment from the line drawn through the vertex of the segment and the vertex of the cone enveloping the conoid, and the straight line between the said vertices is called *adjacent to the axis*.

"Right angled conoids are all similar, but of obtuse-angled conoids let those be called similar in which the cones enveloping the conoids are similar

The following questions are propounded for consideration"

(1) 'why, if a segment be cut off from the obtuse-angled conoid by a plane at right angles to the axis, the segment so cut off has to the cone which has the same base as the segment and the same axis the ratio which the line equal to the sum of the axis of the segment and three times the line adjacent to the axis bears to the line equal to the sum of the axis of the segment and twice the line adjacent to the axis and'

(2) 'why, if a segment of the obtuse-angled conoid be cut off by a plane not at right angles to the axis, the segment so cut off will bear to the figure which has the same base as the segment and the same axis, being a segment of a cone, the ratio which the line equal to the sum of the axis of the segment and three times the line adjacent to the axis bears to the line equal to the sum of the axis of the segment and twice the line adjacent to the axis

III Concerning spheroidal figures we lay down the following premisses. If a section of an acute-angled cone [ellipse] be made to revolve about the greater diameter [major axis] which remains fixed and return to the position from which it started, the figure comprehended by the section of the acute-angled cone is called an *oblong spheroid*. But if the section of the acute-angled cone revolve about the lesser diameter [minor axis] which remains fixed and return to the position from which it started the figure comprehended by the section of the acute-angled cone is called a *flat spheroid*. In either of the spheroids the axis is defined as the diameter [axis] which has remained fixed the *vertex* as the point in which the axis meets the surface of the spheroid the *centre* as the middle point of the axis and the *diameter* as the line drawn through the centre at right angles to the axis. And if parallel planes touch without cutting, either

of the spheroidal figures, and if another plane be drawn parallel to the tangent planes and cutting the spheroid, the base of the resulting segments is defined as the portion intercepted by the section of the spheroid on the cutting plane, their *vertices* as the points in which the parallel planes touch the spheroid and their *axes* as the portions cut off within the segments from the straight line joining their vertices. And that the planes touching the spheroid meet its surface at one point only and that the straight line joining the points of contact passes through the centre of the spheroid we shall prove. Those spheroidal figures are called *similar* in which the axes have the same ratio to the 'diameters'. And let segments of spheroidal figures and conoids be called *similar* if they are cut off from similar figures and have their bases similar, while their axes, being either at right angles to the planes of the bases or making equal angles with the corresponding diameters [axes] of the bases have the same ratio to one another as the corresponding diameters [axes] of the bases.

"The following questions about spheroids are propounded for consideration

(1) 'why if one of the spheroidal figures be cut by a plane through the centre at right angles to the axis each of the resulting segments will be double of the cone having the same base as the segment and the same axis, while if the plane of section be at right angles to the axis without passing through the centre (a) the greater of the resulting segments will bear to the cone which has the same base as the segment and the same axis the ratio which the line equal to the sum of half the straight line which is the axis of the spheroid and the axis of the lesser segment bears to the axis of the lesser segment and (b) the lesser segment bears to the cone which has the same base as the segment and the same axis the ratio which the line equal to the sum of half the straight line which is the axis of the spheroid and the axis of the greater segment bears to the axis of the greater segment

(2) 'why if one of the spheroids be cut by a plane passing through the centre but not at right angles to the axis each of the resulting segments will be double of the figure having the same base as the segment and the same axis and consisting of a segment of a cone

(3) 'But if the plane cutting the spheroid be neither through the centre nor at right angles to the axis (a) the greater of the resulting segments will have to the figure which has the same base as the segment and the same axis the ratio which the line equal to the sum of half the line joining the vertices of the segments and the axis of the lesser segment bears to the axis of the lesser segment and (b) the lesser segment will have to the figure with the same base as the segment and the same axis the ratio which the line equal to the sum of half the line joining the vertices of the segments and the axis of the greater segment bears to the axis of the greater segment. And the figure referred to is in these cases also a segment of a cone

'When the aforesaid theorems are proved there are discovered by means of them many theorems and problems

'Such for example are the theorems

(1) 'that similar spheroids and similar segments both of spheroidal figures and conoids have to one another the triplicate ratio of their axes and

(2) 'that in equal spheroidal figures the squares on the diameters' are reciprocally proportional to the axes and if in spheroidal figures the squares on

the 'diameters' are reciprocally proportional to the axes, the spheroids' are equal

Such also is the problem, From a given spheroidal figure or conoid to cut off a segment by a plane drawn parallel to a given plane so that the segment cut off is equal to a given cone or cylinder or to a given sphere

"After prefixing therefore the theorems and directions which are necessary for the proof of them, I will then proceed to expound the propositions themselves to you Farewell"

DEFINITIONS

'If a cone be cut by a plane meeting all the sides [generators] of the cone the section will be either a circle or a section of an acute-angled cone [an ellipse] If then the section be a circle, it is clear that the segment cut off from the cone towards the same parts as the vertex of the cone will be a cone But if the section be a section of an acute-angled cone [an ellipse] let the figure cut off from the cone towards the same parts as the vertex of the cone be called a *segment of a cone* Let the *base* of the segment be defined as the plane comprehended by the section of the acute-angled cone its *vertex* as the point which is also the vertex of the cone, and its *axis* as the straight line joining the vertex of the cone to the centre of the section of the acute-angled cone

'And if a cylinder be cut by two parallel planes meeting all the sides [generators] of the cylinder the sections will be either circles or sections of acute-angled cones [ellipses] equal and similar to one another If then the sections be circles it is clear that the figure cut off from the cylinder between the parallel planes will be a cylinder But, if the sections be sections of acute-angled cones [ellipses] let the figure cut off from the cylinder between the parallel planes be called a *frustum of a cylinder* And let the *bases* of the frustum be defined as the planes comprehended by the sections of the acute-angled cones [ellipses] and the *axis* as the straight line joining the centres of the sections of the acute-angled cones so that the axis will be in the same straight line with the axis of the cylinder'

LEMMA

If in an ascending arithmetical progression consisting of the magnitudes $A_1, A_2 \dots A_n$ the common difference be equal to the least term A_1 , then

$$n A_n < 2(A_1 + A_2 + \dots + A_n),$$

$$\text{and} \quad > 2(A_1 + A_2 + \dots + A_{n-1})$$

[The proof of this is given incidentally in the treatise *On Spirals* Prop 11 By placing lines side by side to represent the terms of the progression and then producing each so as to make it equal to the greatest term Archimedes gives the equivalent of the following proof

$$\begin{aligned} \text{If} \quad S_n &= A_1 + A_2 + \dots + A_{n-1} + A_n \\ \text{we have also} \quad S &= A_n + A_{n-1} + A_{n-2} + \dots + A_1 \\ \text{And} \quad A_1 + A_{n-1} &= A_2 + A_{n-2} = \dots = A_n \end{aligned}$$

$$\begin{aligned} \text{Therefore} \quad 2S &= (n+1)A_n, \\ \text{whence} \quad n A_n &< 2S \\ \text{and} \quad n A_1 &> 2S_{n-1} \end{aligned}$$

Thus if the progression is a $2a, na$

$$S = \frac{n(n+1)}{2}a,$$

and
but

$$n^2a < 2S \\ > 2S_{n-1}]$$

PROPOSITION 1

If $A_1, B_1, C_1, \dots, K_1$ and $A_2, B_2, C_2, \dots, K_2$ be two series of magnitudes such that

$$\left. \begin{aligned} A_1, B_1 &= A, B_2 \\ B_1, C_1 &= B, C_2 \text{ and so on} \end{aligned} \right\} \quad (\alpha)$$

and if $A_3, B_3, C_3, \dots, K_3$ and $A_4, B_4, C_4, \dots, K_4$ be two other series such that

$$\left. \begin{aligned} A_3, A_4 &= A_2, A_4 \\ B_3, B_4 &= B_2, B_4 \text{ and so on} \end{aligned} \right\} \quad (\beta)$$

then

$$\begin{aligned} (A_1 + B_1 + C_1 + \dots + K_1) (A_2 + B_2 + C_2 + \dots + K_2) \\ = (A_3 + B_3 + C_3 + \dots + K_3) (A_4 + B_4 + C_4 + \dots + K_4) \end{aligned}$$

The proof is as follows

Since $A_1, A_2 = A_4, A_2$
and $A_1, B_1 = A_2, B_2$
while $B_1, B_2 = B_2, B_4$
we have *ex aequali* $A_2, B_1 = A_4, B_4$
Similarly $B_3, C_3 = B_4, C_4$ and so on } (γ)

Again it follows from equations (α) that

$$A_1, A_2 = B_1, B_2 = C_1, C_2 =$$

Therefore

$A_1, A_4 = (A_1 + B_1 + C_1 + \dots + K_1) (A_2 + B_2 + C_2 + \dots + K_2),$
or $(A_1 + B_1 + C_1 + \dots + K_1) A_1 = (A_2 + B_2 + C_2 + \dots + K_2) A_4$
and $A_1, A_4 = A_2, A_4$

while from equations (γ) it follows in like manner that

$$A_2, (A_2 + B_2 + C_2 + \dots + K_2) = A_4, (A_4 + B_4 + C_4 + \dots + K_4)$$

By the last three equations *ex aequali*

$$\begin{aligned} (A_1 + B_1 + C_1 + \dots + K_1) (A_2 + B_2 + C_2 + \dots + K_2) \\ = (A_1 + B_1 + C_1 + \dots + K_1) (A_4 + B_4 + C_4 + \dots + K_4) \end{aligned}$$

Con. If any terms in the third and fourth series corresponding to terms in the first and second be left out the result is the same. For example if the last terms A_3, A_4 are absent

$$\begin{aligned} (A_1 + B_1 + C_1 + \dots + K_1) (A_2 + B_2 + C_2 + \dots + I_2) \\ = (A_1 + B_1 + C_1 + \dots + K_1) (A_4 + B_4 + C_4 + \dots + I_4) \end{aligned}$$

where I immediately precedes A in each series

LEMMA TO PROPOSITION 2

[On Spirals Prop 10]

If $1, 4, 9, \dots, 1$ be n lines forming an ascending arithmetical progression in which the common difference is equal to the least term A_1 , then

$$(n+1)1^2 + 1_1(1_1 + A_1 + 1_2 + \dots + A_n) = 3(A_1^2 + A_2^2 + A_3^2 + \dots + A_n^2)$$

Let the lines $1, 1_{n-1}, 1_{n-2}, \dots, A_1$ be placed in a row from left to right. Produce $1_{n-1}, 1_{n-2}, \dots, 1$ until they are each equal to A so that the parts produced are respectively equal to $A_1, 1_2, \dots, A_{n-1}$.

Taking each line successively we have

$$21^2 = 2A_1^2$$

$$(1_1 + 1_{n-1})^2 = 1_1^2 + A_1^2_{n-1} + 21_1 A_{n-1}$$

$$(1 + 1_{n-2})^2 = 1^2 + A_1^2_{n-2} + 2A_2 1_{n-2}$$

$$(A_{-1} + A_1)^2 = A^2_{-1} + A_1^2 + 2A_{-1}A_1$$

And, by addition

$$(n+1)A^2 = 2(A_1^2 + A_2^2 + \dots + A_n^2) + 2A_1A_{-1} + 2A_2A_{-2} + \dots + 2A_{n-1}A_1 + 2(A_1A_{-1} + A_2A_{-2} + \dots + A_{n-1}A_1) + A_1(A_1 + A_2 + A_3 + \dots + A_n) = A_1^2 + A_2^2 + \dots + A_n^2 \quad (\alpha)$$

Now

$$2A_1A_{-1} = A_1^2 + A_{-1}^2 \text{ because } A_{-1} = 2A_1 \\ 2A_2A_{-2} = A_2^2 + A_{-2}^2 \text{ because } A_{-2} = 3A_2$$

$$2A_{-1}A_1 = A_1^2 + 2(n-1)A_1^2$$

It follows that

$$2(A_1A_{-1} + A_2A_{-2} + \dots + A_{n-1}A_1) + A_1(A_1 + A_2 + \dots + A_n) = A_1\{1 + 3A_{-1} + 5A_{-2} + \dots + (2n-1)A_1\}$$

And this last expression can be proved to be equal to

$$A_1^2 + A_2^2 + \dots + A_n^2$$

For

$$A = A_1(n+1) \\ = A_1\{1 + (n-1)A\} \\ = A_1\{1 + 2(A_{-1} + A_{-2} + \dots + A_1)\} \\ \text{because } (n-1)A = A_{-1} + A_1 \\ + A_{-2} + A_2 \\ + \dots + A_{-n} + A_n$$

$$\text{Similarly } A^2_{-1} = A_1\{A_{-1} + 2(A_{-2} + A_{-3} + \dots + A_1)\}$$

$$A_2^2 = A_2(A_2 + 2A_1)$$

$$A_1^2 = A_1A_1$$

whence by addition

$$A_1^2 + A_2^2 + A_3^2 + \dots + A_n^2 = A_1\{A + 3A_{-1} + 5A_{-2} + \dots + (2n-1)A_1\}$$

Thus the equation marked (α) above is true and it follows that

$$(n+1)A^2 + A_1(A_1 + 1 + A_3 + \dots + A_n) = 3(A_1^2 + A_2^2 + \dots + A_n^2)$$

Cor 1 From this it is evident that

$$n+1 < 3(1^2 + 1^2 + \dots + A^2) \quad (1)$$

Also $A = A_1\{1 + 2(A_{-1} + A_{-2} + \dots + A_1)\}$ as above

so that $A > A_1(A + A_{-1} + \dots + A_1)$

and therefore

$$A + A_1(A_1 + A_2 + \dots + A_n) < 2A^2$$

It follows from the proposition that

$$n+1 > 3(1^2 + 1^2 + \dots + A^2_{-1}) \quad (2)$$

Cor 2 All these results will hold if we substitute similar figures for squares on all the lines for similar figures are in the duplicate ratio of their sides

PROPOSITION 2

If A_1, A_2, \dots, A_n be any number of areas such that

$$A_1 = ax + x$$

$$A_2 = a \cdot 2x + (2x)^2,$$

$$A_3 = a \cdot 3x + (3x)^2$$

$$A_n = a \cdot nx + (nx)^2$$

then $n A_1 (A_1 + A_2 + \dots + A_n) < (a + nx) \left(\frac{a}{2} + \frac{nx}{3} \right),$

and $n A_1 (A_1 + A_2 + \dots + A_{n-1}) > (a + nx) \left(\frac{a}{2} + \frac{nx}{3} \right)$

For by the Lemma immediately preceding Prop 1,

$$n \cdot anx < (ax + a \cdot 2x + \dots + a \cdot nx)$$

and $> 2(ax + a \cdot 2x + \dots + a \cdot (n-1)x)$

Also by the Lemma preceding this proposition

$$n \cdot (nx)^2 < 3\{x^2 + (2x)^2 + (3x)^2 + \dots + (nx)^2\}$$

and $> 3\{x + (2x)^2 + \dots + (n-1)x^2\}$

Hence

$$\frac{an^2x}{2} + \frac{n(nx)}{3} < [(ax + x^2) + \{a \cdot 2x + (2x)^2\} + \dots + \{a \cdot nx + (nx)^2\}],$$

and

$$> [(ax + x) + \{a \cdot 2x + (2x)^2\} + \dots + \{a \cdot (n-1)x + (n-1)x^2\}],$$

or $\frac{an^2x}{2} + \frac{n(nx)}{3} < A_1 + A_2 + \dots + A_n$

and $> A_1 + A_2 + \dots + A_{n-1}$

It follows that

$$n-1 \cdot (A_1 + A_2 + \dots + A_n) < n\{a \cdot nx + (nx)^2\} \left\{ \frac{an \cdot x}{2} + \frac{n(nx)^2}{3} \right\}$$

or $n-1 \cdot (A_1 + A_2 + \dots + A_n) < (a + nx) \left(\frac{a}{2} + \frac{nx}{3} \right)$

also $n-1 \cdot (A_1 + A_2 + \dots + A_{n-1}) > (a + nx) \left(\frac{a}{2} + \frac{nx}{3} \right)$

PROPOSITION 3

(1) If TP, TP be two tangents to any conic meeting in T and if Qq, Qq be any two chords parallel respectively to TP, TP and meeting in O then

$$QO \cdot Oq \cdot QO \cdot Oq = TP \cdot TP^2$$

And this is proved in the elements of conics ¹

(2) If QQ be a chord of a parabola bisected in I by the diameter PI , and if PI be of constant length then the areas of the triangle PQQ and of the segment PQQ are both constant whatever be the direction of QQ

Let IBB be the particular segment of the parabola whose vertex is I so that BB is bisected perpendicularly by the axis at the point H where $4HI = PI$

Draw QD perpendicular to I

¹In the treatises on conics by Aristaeus and Euclid

Let p be the parameter of the principal ordinates and let p be another line of such length that

$$QV \cdot QD = p \cdot p,$$

it will then follow that p is equal to the parameter of the ordinates to the diameter PV i.e. those which are parallel to QV .

"For this is proved in the conics."

$$\text{Thus } QV^2 = p \cdot PV$$

$$\text{And } BH^2 = p \cdot AH \text{ while } AH = PV$$

$$\text{Therefore } QV^2 \cdot BH = p \cdot p$$

$$\text{But } QV^2 \cdot QD = p \cdot p,$$

$$\text{hence } BH = QD$$

$$\text{Thus } BH \cdot AH = QD \cdot PV$$

$$\text{and therefore } \triangle ABB = \triangle PQQ,$$

that is the area of the triangle PQQ is constant so long as PV is of constant length

Hence also the area of the segment PQQ is constant under the same conditions for the segment is equal to $\frac{1}{2} \triangle PQQ$ [*Quadrature of the Parabola*, Prop 17 or 24]

PROPOSITION 4


The area of any ellipse is to that of the auxiliary circle as the minor axis to the major

Let AA' be the major and BB' the minor axis of the ellipse and let BB' meet the auxiliary circle in b, b'

Suppose O to be such a circle that

$$(\text{circle } AbA'b') \quad O = CA \cdot CB$$

Then shall O be equal to the area of the ellipse



For if not O must be either greater or less than the ellipse

I. If possible let O be greater than the ellipse

We can then inscribe in the circle O an equilateral polygon of $4n$ sides such that its area is greater than that of the ellipse [cf. *On the Sphere and Cylinder* I 6]

The theorem which is here assumed by Archimedes as known is easily deduced from Apollonius I 49

Let this be done and inscribe in the auxiliary circle of the ellipse the polygon $AefbghA'$ similar to that inscribed in O . Let the perpendiculars eM, fN , on AA' meet the ellipse in E, F , respectively. Join AF, EF, FB ,

Suppose that P denotes the area of the polygon inscribed in the auxiliary circle and P' that of the polygon inscribed in the ellipse.

Then, since all the lines eM, fN , are cut in the same proportions at E, F ,

$$eM : EM = fN : FN = bC : BC$$

the pairs of triangles, as eAM, EAM and the pairs of trapeziums, as $eMNF, EMNI$, are all in the same ratio to one another as bC to BC , or as CA to CB .

Therefore, by addition

$$P' : P = CA : CB$$

Now P (polygon inscribed in O)

$$= (\text{circle } AbA'b') : O$$

$$= CA : CB, \text{ by hypothesis}$$

Therefore P is equal to the polygon inscribed in O .

But this is impossible because the latter polygon is by hypothesis greater than the ellipse, and *a fortiori* greater than P .

Hence O is not greater than the ellipse.

II. If possible, let O be less than the ellipse.

In this case we inscribe in the ellipse a polygon P with $4n$ equal sides such that $P > O$.

Let the perpendiculars from the angular points on the axis AA' be produced to meet the auxiliary circle and let the corresponding polygon (P') in the circle be formed.

Inscribe in O a polygon similar to P .

Then $P' : P = CA : CB$

$$= (\text{circle } AbA'b') : O, \text{ by hypothesis,}$$

$$= P' : (\text{polygon inscribed in } O)$$

Therefore the polygon inscribed in O is equal to the polygon P , which is impossible because $P > O$.

Hence O being neither greater nor less than the ellipse is equal to it, and the required result follows.

PROPOSITION 5

If AA', BB' be the major and minor axis of an ellipse respectively, and if d be the diameter of any circle then

$$(\text{area of ellipse}) : (\text{area of circle}) = AA' : BB' : d^2$$

For

$$(\text{area of ellipse}) : (\text{area of auxiliary circle}) = BB' : AA' \quad [\text{Prop. 4}]$$

$$= AA' : BB' : AA'^2$$

And

$$(\text{area of aux. circle}) : (\text{area of circle with diam. } d) = AA'^2 : d^2$$

Therefore the required result follows *ex aequali*.

PROPOSITION 6

The areas of ellipses are as the rectangles under their axes.

Thus follows at once from Prop. 4 & 5.

Cor. The areas of similar ellipses are as the squares of corresponding axes.

PROPOSITION 7

Given an ellipse with centre C , and a line CO drawn perpendicular to its plane it is possible to find a circular cone with vertex O and such that the given ellipse is a section of it [or, in other words to find the circular sections of the cone with vertex O passing through the circumference of the ellipse]

Conceive an ellipse with BB' as its minor axis and lying in a plane perpendicular to that of the paper. Let CO be drawn perpendicular to the plane of the ellipse and let O be the vertex of the required cone. Produce OB OC OB' and in the same plane with them draw BED meeting OC OB produced in E , D respectively and in such a direction that

$$BE \cdot ED \cdot EO^2 = CA^2 \cdot CO,$$

where CA is half the major axis of the ellipse

And this is possible, since


$$BE \cdot ED \cdot EO > BC \cdot CB \cdot CO^2$$

[Both the construction and this proposition are assumed as known]

Now conceive a circle with BD as diameter lying in a plane at right angles to that of the paper, and describe a cone with this circle for its base and with vertex O

We have therefore to prove that the given ellipse is a section of the cone or if P be any point on the ellipse, that P lies on the surface of the cone

Draw PN perpendicular to BB' . Join ON and produce it to meet BD in V and let MQ be drawn in the plane of the circle on BD as diameter perpendicular to BD and meeting the circle in Q . Also let FG , HK be drawn through E , M respectively parallel to BB'



We have then

$$\begin{aligned} QM^2 \cdot HM \cdot MK &= BV \cdot MD \cdot HM \cdot MK \\ &= BE \cdot ED \cdot FE \cdot EG \\ &= (BE \cdot ED \cdot EO) \cdot (EO^2 \cdot FE \cdot EG) \\ &= (CA^2 \cdot CO) \cdot (CO^2 \cdot BC \cdot CB) \\ &= CA^2 \cdot CB^2 \\ &= PN^2 \cdot BN \cdot NB \end{aligned}$$

$$\text{Therefore } QM^2 \cdot PN^2 = HM \cdot MK \cdot BN \cdot NB \\ = OM^2 \cdot ON^2$$

whence since PN QM are parallel OPQ is a straight line

But Q is on the circumference of the circle on BD as diameter, therefore OQ is a generator of the cone and hence P lies on the cone

Thus the cone passes through all points on the ellipse

PROPOSITION 8

Given an ellipse, a plane through one of its axes AA' and perpendicular to the plane of the ellipse and a line CO drawn from C the centre in the given plane through A' but not perpendicular to AA' it is possible to find a cone with vertex

O such that the given ellipse is a section of it [or, in other words to find the circular sections of the cone with vertex O whose surface passes through the circumference of the ellipse]

By hypothesis OA OA are unequal Produce $O1'$ to D so that $O1 = OD$ Join $1D$ and draw FG through C parallel to it

The given ellipse is to be supposed to lie in a plane perpendicular to the plane of the paper Let BB be the other axis of the ellipse

Conceive a plane through $1D$ perpendicular to the plane of the paper, and in it describe either (a), if $CB^2 = FC \cdot CG$ a circle with diameter AD , or (b), if not an ellipse on AD as axis such that if d be the other axis

$$d^2 \cdot AD = CB^2 \cdot FC \cdot CG$$

Take a cone with vertex O whose surface passes through the circle or ellipse just drawn This is possible even when the curve is an ellipse because the line from O to the middle point of AD is perpendicular to the plane of the ellipse and the construction is effected by means of Prop 7

Let P be any point on the given ellipse and we have only to prove that P lies on the surface of the cone so described

Draw PN perpendicular to AA Join ON and produce it to meet AD in M Through M draw HA parallel to AA

Lastly draw MQ perpendicular to the plane of the paper (and therefore perpendicular to both HA and AD) meeting the ellipse or circle about AD (and therefore the surface of the cone) in Q

Then

$$\begin{aligned} QM^2 \cdot HM \cdot MK &= (QM^2 \cdot DM \cdot MA) \cdot (DM \cdot MA \cdot HM \cdot MA) \\ &= (d^2 \cdot AD^2) \cdot (FC \cdot CG \cdot AC \cdot CA) \\ &= (CB^2 \cdot FC \cdot CG) \cdot (FC \cdot CG \cdot AC \cdot CA) \\ &= CB^2 \cdot C1^2 \\ &= P1^2 \cdot AN \cdot NA \end{aligned}$$

Therefore alternately


$$\begin{aligned} QM^2 \cdot P1^2 &= HM \cdot MK \cdot AN \cdot NA \\ &= OM^2 \cdot ON^2 \end{aligned}$$

Thus since $P1$ QM are parallel OPQ is a straight line and Q being on the surface of the cone it follows that P is also on the surface of the cone

Similarly all points on the ellipse are also on the cone and the ellipse is therefore a section of the cone

PROPOSITION 9

Given an ellipse a plane through one of its axes and perpendicular to that of the ellipse and a straight line CO drawn from the centre C of the ellipse in the given plane through the axis but not perpendicular to that axis it is possible to find a



cylinder with axis OC such that the ellipse is a section of it [or in other words, to find the circular sections of the cylinder with axis OC whose surface passes through the circumference of the given ellipse]

Let AA' be an axis of the ellipse and suppose the plane of the ellipse to be perpendicular to that of the paper, so that OC lies in the plane of the paper

Draw AD AE parallel to CO , and let DE be the line through O perpendicular to both AD and $A'E$

We have now three different cases according as the other axis BB' of the ellipse is (1) equal to, (2) greater than or (3) less than, DE

(1) Suppose $BB' = DE$

Draw a plane through DE at right angles to OC and in this plane describe a circle on DE as diameter Through

this circle describe a cylinder with axis OC

This cylinder shall be the cylinder required or its surface shall pass through every point P of the ellipse

For if P be any point on the ellipse draw PN perpendicular to AA' , through N draw NM parallel to CO meeting DE in M and through M , in the plane of the circle on DE as diameter draw MQ perpendicular to DE meeting the circle in Q

Then since

$$DE = BB'$$

And

$$PN^2 = AN \cdot NA = DO^2 = AC \cdot CA'$$

since DN CM CO $A'E$ are parallel

Therefore

$$PN^2 = DM \cdot ME \\ = QM^2$$

by the property of the circle

Hence since PN QM are equal as well as parallel PQ is parallel to MN and therefore to CO It follows that PQ is a generator of the cylinder, whose surface accordingly passes through P

(2) If $BB' > DE$ we take E on AE such that $DE = BB'$ and describe a circle on DE as diameter in a plane perpendicular to that of the paper, and the rest of the construction and proof is exactly similar to those given for case (1)

(3) Suppose $BB' < DE$

Take a point K on CO produced such that

$$DO^2 - CB^2 = OK^2$$



From K draw KR perpendicular to the plane of the paper and equal to CB

Thus $OR^2 = OK^2 + CB^2 = OD^2$

In the plane containing DE OR describe a circle on DE as diameter Through this circle (which must pass through R) draw a cylinder with axis OC

We have then to prove that if P be

any point on the given ellipse P lies on the cylinder so described



Draw PN perpendicular to AA' , and through N draw NM parallel to CO meeting DE in M . In the plane of the circle on DE as diameter draw MQ perpendicular to DE and meeting the circle in Q .

Lastly, draw QH perpendicular to NM produced. QH will then be perpendicular to the plane containing AC , DE , i.e. the plane of the paper.

Now $QH^2 = QM^2 = KR \cdot OR$ by similar triangles

And $QM \cdot AN = DM \cdot ME \cdot AN \cdot NA'$
 $= OD^2 \cdot CA^2$

Hence *ex aequali* since $OR = OD$

$$\begin{aligned} QH \cdot 1N \cdot N1' &= KR^2 \cdot CA^2 \\ &= CB^2 \cdot CA^2 \\ &= PN^2 \cdot AA' \cdot NA' \end{aligned}$$

Thus $QH = PN$. And QH , PN are also parallel. Accordingly PQ is parallel to MN and therefore to CO so that PQ is a generator, and the cylinder passes through P .

PROPOSITION 10

It was proved by the earlier geometers that any two cones have to one another the ratio compounded of the ratios of their bases and of their heights.¹ The same method of proof will show that any segments of cones have to one another the ratio compounded of the ratios of their bases and of their heights.

The proposition that any frustum of a cylinder is triple of the conical segment which has the same base as the frustum and equal height is also proved in the same manner as the proposition that the cylinder is triple of the cone which has the same base as the cylinder and equal height.²

PROPOSITION 11

(1) If a paraboloid of revolution be cut by a plane through or parallel to the axis the section will be a parabola equal to the original parabola which by its revolution generates the paraboloid. And the axis of the section will be the intersection between the cutting plane and the plane through the axis of the paraboloid at right angles to the cutting plane.

If the paraboloid be cut by a plane at right angles to its axis the section will be a circle whose centre is on the axis.

(2) If a hyperboloid of revolution be cut by a plane through the axis parallel to the axis or through the centre the section will be a hyperbola (a) if the section be through the axis equal (b) if parallel to the axis similar (c) if through the centre not similar to the original hyperbola which by its revolution generates the hyperboloid. And the axis of the section will be the intersection of the cutting plane and the plane through the axis of the hyperboloid at right angles to the cutting plane.

Any section of the hyperboloid by a plane at right angles to the axis will be a circle whose centre is on the axis.

(3) If any of the spheroidal figures be cut by a plane through the axis or parallel to the axis the section will be an ellipse (a) if the section be through the axis, equal (b) if parallel to the axis similar to the ellipse which by its revolution gen

¹This follows from Eucl vii 11 and 11 taken together. Cf. *On the Sphere and Cylinder* 1, Lemma 1.

²This proposition is proved by Eucl viii as stated in the preface to *On the Sphere and Cylinder*. Cf. Eucl vii 10.

erates the figure. And the axis of the section will be the intersection of the cutting plane and the plane through the axis of the spheroid at right angles to the cutting plane.

If the section be by a plane at right angles to the axis of the spheroid, it will be a circle whose centre is on the axis.

(4) If any of the said figures be cut by a plane through the axis, and if a perpendicular be drawn to the plane of section from any point on the surface of the figure but not on the section that perpendicular will fall within the section.

And the proofs of all these propositions are evident.

PROPOSITION 12

If a paraboloid of revolution be cut by a plane neither parallel nor perpendicular to the axis and if the plane through the axis perpendicular to the cutting plane intersect it in a straight line of which the portion intercepted within the paraboloid is RR the section of the paraboloid will be an ellipse whose major axis is RR and whose minor axis is equal to the perpendicular distance between the lines through R R parallel to the axis of the paraboloid.

Suppose the cutting plane to be perpendicular to the plane of the paper and let the latter be the plane through the axis ANF of the paraboloid which intersects the cutting plane at right angles in RR . Let RH be parallel to the axis of the paraboloid, and $R'H$ perpendicular to RH .

Let Q be any point on the section made by the cutting plane and from Q draw QM perpendicular to RR . QM will therefore be perpendicular to the plane of the paper.

Through M draw $DMTE$ perpendicular to the axis ANF meeting the parabolic section made by the plane of the paper in D E . Then QM is perpendicular to DE and if a plane be drawn through DE QM , it will be perpendicular to the axis and will cut the paraboloid in a circular section.

Since Q is on this circle

$$QM^2 = DM \cdot ME$$

Again if PT be that tangent to the parabolic section in the plane of the paper which is parallel to RR and if the tangent at A meet PT in O then from the property of the parabola

$$DM \cdot ME \cdot RM \cdot MR = AO \cdot OP^2 \quad [\text{Prop 3 (1)}]$$

$$= AO \cdot OT^2 \text{ since } AN = 1T$$

$$\text{Therefore } QM \cdot RM \cdot MR = AO^2 \cdot OT^2$$

$$= RH^2 \cdot RR^2$$

by similar triangles


Hence Q lies on an ellipse whose major axis is RR and whose minor axis is equal to RH .

PROPOSITIONS 13 14

If a hyperboloid of revolution be cut by a plane meeting all the generators of the enveloping cone or if an oblong spheroid be cut by a plane not perpendicular to

the axis¹ and if a plane through the axis intersect the cutting plane at right angles in a straight line on which the hyperboloid or spheroid intercepts a length RR , then the section by the cutting plane will be an ellipse whose major axis is RR .

Suppose the cutting plane to be at right angles to the plane of the paper, and suppose the latter plane to be that through the axis ANF which intersects the



cutting plane at right angles in RR . The section of the hyperboloid or spheroid by the plane of the paper is thus a hyperbola or ellipse having ANF for its transverse or major axis.

Take any point on the section made by the cutting plane as Q and draw QM perpendicular to RR . QM will then be perpendicular to the plane of the paper.

Through M draw DPE at right angles to the axis ANF meeting the hyperbola or ellipse in D , E and through QM , DE let a plane be described. This plane will accordingly be perpendicular to the axis and will cut the hyperboloid or spheroid in a circular section.

Thus $QM = DM = ME$

Let PT be that tangent to the hyperbola or ellipse which is parallel to RR , and let the tangent at A meet PT in O .

Then by the property of the hyperbola or ellipse

$$DM \cdot ME \cdot RM \cdot MR = OA^2 \cdot OP^2$$

or $QM^2 \cdot RM \cdot MR = OA^2 \cdot OP^2$

Now (1) in the hyperbola $OA < OP$ because $AT < AV$ and accordingly $OT < OP$ while $OA < OT$

(2) in the ellipse if AA' be the diameter parallel to RR , and BB' the minor axis

$$BC \cdot CB \cdot KC \cdot CK = OA^2 \cdot OP^2$$

and $BC \cdot CB < KC \cdot CK$ so that $OA < OP$

Hence in both cases the locus of Q is an ellipse whose major axis is RR .

CON 1 If the spheroid be a flat spheroid the section will be an ellipse, and everything will proceed as before except that LR will in this case be the minor axis.

¹Archimedes begins Fr. p. 14 for the spheroid with the remark that when the cutting plane passes through or is parallel to the axis the case is clear Cf Prop. 11 (3)

COR 2 In all conoids or spheroids parallel sections will be similar, since the ratio $OA^2 : OP^2$ is the same for all the parallel sections

PROPOSITION 15

(1) *If from any point on the surface of a conoid a line be drawn, in the case of the paraboloid, parallel to the axis and, in the case of the hyperboloid parallel to any line passing through the vertex of the enveloping cone the part of the straight line which is in the same direction as the convexity of the surface will fall without it, and the part which is in the other direction within it*

For, if a plane be drawn, in the case of the paraboloid, through the axis and the point and in the case of the hyperboloid through the given point and through the given straight line drawn through the vertex of the enveloping cone, the section by the plane will be (a) in the paraboloid a parabola whose axis is the axis of the paraboloid, (b) in the hyperboloid a hyperbola in which the given line through the vertex of the enveloping cone is a diameter¹

[Prop 11]


Hence the property follows from the plane properties of the conics

(2) *If a plane touch a conoid without cutting it, it will touch it at one point only, and the plane drawn through the point of contact and the axis of the conoid will be at right angles to the plane which touches it*

For if possible let the plane touch at two points. Draw through each point a parallel to the axis. The plane passing through both parallels will therefore either pass through or be parallel to the axis. Hence the section of the conoid made by this plane will be a conic [Prop 11 (1), (2)], the two points will lie on this conic and the line joining them will lie within the conic and therefore within the conoid. But this line will be in the tangent plane since the two points are in it. Therefore some portion of the tangent plane will be within the conoid which is impossible since the plane does not cut it.

Therefore the tangent plane touches in one point only.

That the plane through the point of contact and the axis is perpendicular to the tangent plane is evident in the particular case where the point of contact is the vertex of the conoid. For if two planes through the axis cut it in two comes the tangents at the vertex in both conics will be perpendicular to the axis of the conoid. And all such tangents will be in the tangent plane which must therefore be perpendicular to the axis and to any plane through the axis.



If the point of contact P is not the vertex draw the plane passing through the axis AN and the point P . It will cut the conoid in a conic whose axis is AN and the tangent plane in a line DPE touching the conic at P . Draw PNP perpendicular to the axis and draw a plane through it also perpendicular to the axis. This plane will make a circular section and meet the tangent plane in a tangent to the circle which will therefore be at right angles to PN . Hence the tangent to the circle will be at right angles to the plane containing PN, AN , and it follows that this last plane is perpendicular to the tangent plane.

¹There seems to be some error in the text here which says that the diameter (i.e. axis) of the hyperbola is the straight line drawn in the conoid from the vertex of the cone. But this straight line is not in general the axis of the section.

PROPOSITION 16

(1) *If a plane touch any of the spheroidal figures without cutting it, it will touch at one point only, and the plane through the point of contact and the axis will be at right angles to the tangent plane*

This is proved by the same method as the last proposition

(2) *If any conoid or spheroid be cut by a plane through the axis and if through any tangent to the resulting conic a plane be erected at right angles to the plane of section the plane so erected will touch the conoid or spheroid in the same point as that in which the line touches the conic*

For it cannot meet the surface at any other point. If it did the perpendicular from the second point on the cutting plane would be perpendicular also to the tangent to the conic and would therefore fall outside the surface. But it must fall within it [Prop 11 (4)]

(3) *If two parallel planes touch any of the spheroidal figures the line joining the points of contact will pass through the centre of the spheroid*

If the planes are at right angles to the axis the proposition is obvious. If not, the plane through the axis and one point of contact is at right angles to the tangent plane at that point. It is therefore at right angles to the parallel tangent plane and therefore passes through the second point of contact. Hence both points of contact lie on one plane through the axis and the proposition is reduced to a plane one

PROPOSITION 17

If two parallel planes touch any of the spheroidal figures and another plane be drawn parallel to the tangent planes and passing through the centre the line drawn through any point of the circumference of the resulting section parallel to the chord of contact of the tangent planes will fall outside the spheroid

This is proved at once by reduction to a plane proposition

Archimedes adds that it is evident that if the plane parallel to the tangent planes does not pass through the centre a straight line drawn in the manner described will fall without the spheroid in the direction of the smaller segment but within it in the other direction

PROPOSITION 18

Any spheroidal figure which is cut by a plane through the centre is divided both as regards its surface and its volume into two equal parts by that plane

To prove this Archimedes takes another equal and similar spheroid divides it similarly by a plane through the centre and then uses the method of application

PROPOSITIONS 19 20

Given a segment cut off by a plane from a paraboloid or hyperboloid of revolution or a segment of a spheroid less than half the spheroid also cut off by a plane it is possible to inscribe in the segment one solid figure and to circumscribe about it another solid figure each made up of cylinders or frusta of cylinders of equal height, and such that the circumscribed figure exceeds the inscribed figure by a volume less than that of any given solid

Let the plane base of the segment be perpendicular to the plane of the paper

and let the plane of the paper be the plane through the axis of the conoid or spheroid which cuts the base of the segment at right angles in BC . The section in the plane of the paper is then a conic BAC [Prop 11]

Let EAF be that tangent to the conic which is parallel to BC , and let I be the point of contact. Through EAF draw a plane parallel to the plane through BC bounding the segment. The plane so drawn will then touch the conoid or spheroid at A [Prop 16]

(1) If the base of the segment is at right angles to the axis of the conoid or spheroid A will be the vertex of the conoid or spheroid and its axis AD will bisect BC at right angles

(2) If the base of the segment is not at right angles to the axis of the conoid or spheroid, we draw AD

(a) in the paraboloid, parallel to the axis

(b) in the hyperboloid, through the centre (or the vertex of the enveloping cone),

(c) in the spheroid, through the centre

and in all the cases it will follow that AD bisects BC in D


Then A will be the vertex of the segment, and AD will be its axis

Further, the base of the segment will be a circle or an ellipse with BC as diameter or as an axis respectively and with centre D . We can therefore describe through this circle or ellipse a cylinder or a 'frustum' of a cylinder whose axis is AD [Prop 9]

Dividing this cylinder or frustum continually into equal parts by planes parallel to the base we shall at length arrive at a cylinder or frustum less in volume than any given solid

Let this cylinder or frustum be that whose axis is OD , and let AD be divided into parts equal to OD at L, M

Through L, M draw lines parallel to



BC meeting the conic in P, Q and through these lines draw planes parallel to the base of the segment. These will cut the conoid or spheroid in circles or similar ellipses. On each of these circles or ellipses describe two cylinders or frusta of cylinders each with axis equal to OD one of them lying in the direction of A and the other in the direction of D as shown in the figure

Then the cylinders or frusta of cylinders drawn in the direction of A make up a circumscribed figure and those in the direction of D an inscribed figure in relation to the segment

Also the cylinder or frustum PG in the circumscribed figure is equal to the cylinder or frustum PH in the inscribed figure. QI in the circumscribed figure is equal to QA in the inscribed figure, and so on

Therefore by addition

(circumscribed fig) = (inscr fig) + (cylinder or frustum whose axis is OD)

But the cylinder or frustum whose axis is OD is less than the given solid figure whence the proposition follows

'Having set out these preliminary propositions let us proceed to demonstrate the theorems propounded with reference to the figures'

it follows that (inscribed figure) $> \lambda$ (a)

Next comparing successively the cylinders or frusta with heights equal to OD and respectively forming parts of the complete cylinder or frustum EC and of the inscribed figure, we have

(first cylinder or frustum in EC) (first in inscr fig)

$$= BD \quad RO$$

$$= 4D \quad 4O$$

$$= BD \quad TO \text{ where } AB \text{ meets } OR \text{ in } T$$

And (second cylinder or frustum in EC) (second in inscr fig)

$$= HO \quad SN \text{ in like manner,}$$

and so on

Hence [Prop 1] (cylinder or frustum EC) (inscribed figure)

$$= (BD + HO + \dots) (TO + SN + \dots),$$

where $BD \quad HO$, are all equal, and $BD, TO \quad SN$, diminish in arithmetical progression

But [Lemma preceding Prop 1]

$$BD + HO + \dots > 2(TO + SN + \dots)$$

Therefore (cylinder or frustum EC) > 2 (inscribed fig),

or $\lambda > \text{(inscribed fig)}$,

which is impossible, by (a) above

II If possible, let the segment be less than λ

In this case we inscribe and circumscribe figures as before but such that

$$(\text{circumscriber fig}) - (\text{inscr fig}) < \lambda - (\text{segment}),$$

whence it follows that

$$(\text{circumscribed figure}) < \lambda \quad (\beta)$$

And, comparing the cylinders or frusta making up the complete cylinder or frustum CE and the circumscribed figure respectively we have

(first cylinder or frustum in CE) (first in circumscriber fig)

$$= BD \quad BD^2$$

$$= BD \quad BD$$

(second in CF) (second in circumscriber fig)

$$= HO^2 \quad RO^2$$

$$= AD \quad AO$$

$$= HO \quad TO,$$

and so on

Hence [Prop 1]

(cylinder or frustum CE) (circumscribed fig)

$$= (BD + HO + \dots) (BD + TO + \dots)$$

$$< 2 \quad 1$$

[Lemma preceding Prop 1]

and it follows that

$$\lambda < (\text{circumscribed fig})$$


which is impossible, by (b)

Thus the segment being neither greater nor less than λ is equal to it, and therefore to $\frac{2}{3}$ (cone or segment of cone ABC)

PROPOSITION 23

If from a paraboloid of revolution two segments be cut off one by a plane perpendicular to the axis the other by a plane not perpendicular to the axis and if the axes of the segments are equal the segments will be equal in volume

Let the two planes be supposed perpendicular to the plane of the paper, and let the latter plane be the plane through the axis of the paraboloid cutting the other two planes at right angles in BB' , QQ' respectively and the paraboloid itself in the parabola $QPQ B$



Let $1A$ PI be the equal axes of the segments and $1, P$ their respective vertices

Draw QI parallel to AN or PV and QL perpendicular to QL

Now, since the segments of the parabolic section cut off by BB' , QQ have equal axes, the triangles ABB , PQQ are equal [Prop 3]

Also if QD be perpendicular to PV , $QD = BV$ (as in the same Prop 3)

Conceive two cones drawn with the same bases as the segments and with A, P as vertices respectively. The height of the cone PQQ is then PA where PK is perpendicular to QQ

Now the cones are in the ratio compounded of the ratios of their bases and of their heights i.e. the ratio compounded of (1) the ratio of the circle about BB' to the ellipse about QQ , and (2) the ratio of $1V$ to PA

That is to say, we have by means of Props 5, 12,

$$(\text{cone } 1BB) (\text{cone } PQQ) = (BB' : QQ) (QL) (1A : PA)$$

$$\text{And } BB' = 2BA = 2QD = QL \text{ while } QQ = 2QI$$

Therefore

$$\begin{aligned} (\text{cone } 1BB) (\text{cone } PQQ) &= (QD : QV) (1A : PA) \\ &= (PA : PI) (1A : PA) \\ &= 1A : PI \end{aligned}$$

Since $1N = PI$, the ratio of the cones is a ratio of equality, and it follows that the segments being each half as large again as the respective cones [Prop 22], are equal


PROPOSITION 24

If from a paraboloid of revolution two segments be cut off by planes drawn in any manner, the segments will be to one another as the squares on their axes

For let the paraboloid be cut by a plane through the axis in the parabolic section $P P A p p'$, and let the axis of the parabola and paraboloid be ANN'

Measure along ANN' the lengths AN , AN' equal to the respective axes of the given segments and through N N' draw planes perpendicular to the axis making circular sections on Pp , $P p'$ as diameters respectively. With these circles as bases and with the common vertex A let two cones be described

Now the segments of the paraboloid whose bases are the circles about Pp , $P p'$ are equal to the given segments respectively, since their respective axes are



equal [Prop 23] and, since the segments $4Pp$ $4P'p$ are half as large again as the cones $4Pp$ $4P'p$ respectively we have only to show that the cones are in the ratio of $4V$ to $4V'$.

But

$$\begin{aligned} (\text{cone } 4Pp) (\text{cone } 4P'p) &= (PV'' P'V'') (4V \quad 4V') \\ &= (4V \quad 4V') (4V \quad 4V') \\ &= 4V'' \quad 4V'' \end{aligned}$$

thus the proposition is proved.

PROPOSITION 23 2b

In any hyperboloid of revolution, if 4 be the vertex and $4D$ the axis of any segment cut off by a plane and if $C4$ be the semidiameter of the hyperboloid through 4 ($C4$ being of course in the same straight line with $4D$) then

$$\frac{(\text{segment})}{(\text{cone with same base and axis})} = (4D+3C4) \quad (4D+2C4)$$

Let the plane cutting off the segment be perpendicular to the plane of the paper and let the latter plane be the plane through the axis of the hyperboloid which intersects the cutting plane at right angles in BB' , and makes the hyperbolic segment $B4B'$. Let C be the centre of the hyperboloid (or the vertex of the enveloping cone)

Let EF be that tangent to the hyperbolic section which is parallel to BB' . Let EF touch at A and join $C4$. Then $C4$ produced will bisect BB' at D . $C4$ will be a semi-diameter of the hyperboloid. A will be the vertex of the segment and AD its axis. Produce AC to A' and H so that $AC = C4 = A'H$.

Through EF draw a plane parallel to the base of the segment. This plane will touch the hyperboloid at A .

Then (1) if the base of the segment is at right angles to the axis of the hyperboloid 4 will be the vertex and $4D$ the axis of the hyperboloid as well as of the segment, and the base of the segment will be a circle on BB' as diameter.

(2) If the base of the segment is not perpendicular to the axis of the hyperboloid the base will be an ellipse on BB' as major axis. [Prop 13]

Then we can draw a cylinder or a frustum of a cylinder $FBB'I$ passing through the circle or ellipse about BB' and having $4D$ for its axis and so we can describe a cone or a segment of a cone through the circle or ellipse and having 4 for its vertex.

We have to prove that

$$(\text{segment } ABB') (\text{cone or segment of cone } ABB') = HD \quad 4'D$$

Let V be a cone such that

$$V (\text{cone or segment of cone } ABB') = HD \quad 4'D \quad (\alpha)$$

and we have to prove that V is equal to the segment

Now

$$(\text{cylinder or frustum } EB) (\text{cone or segment of cone } ABB') = 3 \quad 1$$

$$\text{Therefore, by means of } (\alpha) \quad (\text{cylinder or frustum } EB) \quad V = 4D \quad \frac{HD}{3} \quad (\beta)$$

If the segment is not equal to V it must either be greater or less.

I. If possible let the segment be greater than V .



Inscribe and circumscribe to the segment figures made up of cylinders or frusta of cylinders with axes along AD and all equal to one another such that

(circumscribed fig) - (inscr fig) < (segmt) - I,

whence

(inscribed figure) > V (7)

Produce all the planes forming the bases of the cylinders or frusta of cylinders to meet the surface of the complete cylinder or frustum EB'



Then if AD be the axis of the greatest cylinder or frustum in the circumscribed figure the complete cylinder will be divided into cylinders or frusta each equal to this greatest cylinder or frustum

Let there be a number of straight lines a equal to AA' and as many in number as the parts into which AD is divided by the bases of the cylinders or frusta. To each line a apply a rectangle which shall overlap it by a square and let the greatest of the rectangles be equal to the rectangle $AD A'D$ and the least equal to the rectangle $1L A'L$ also let the sides of the overlapping squares b, p, q, l be in descending arithmetical progression. Thus b, p, q, l will be respectively equal to AD, AN, AM, AL and the rectangles $(ab + b^2)$, $(ap + p^2)$, $(al + l^2)$ will be respectively equal to $AD A'D, AN A'N, AL A'L$.

Suppose further that we have a series of spaces S each equal to the largest rectangle $AD A'D$ and as many in number as the diminishing rectangles.

Comparing now the successive cylinders or frusta (1) in the complete cylin

der or frustum EB' and (2) in the inscribed figure, beginning from the base of the segment, we have

$$\begin{aligned} & \text{(first cylinder or frustum in } FB) \quad \text{(first in inscr figure)} \\ &= BD^2 \quad PA^2 \\ &= AD \cdot A'D \quad AN \cdot A'N, \text{ from the hyperbola} \\ &= S \quad (ap + p^2) \end{aligned}$$

Again

$$\begin{aligned} & \text{(second cylinder or frustum in } EB) \quad \text{(second in inscr fig)} \\ &= BD^2 \quad QM^2 \\ &= AD \cdot A'D \quad 1M \cdot 4'M \\ &= S \quad (aq + q^2) \end{aligned}$$

and so on

The last cylinder or frustum in the complete cylinder or frustum EB' has no cylinder or frustum corresponding to it in the inscribed figure

Combining the proportions, we have [Prop 1]

$$\begin{aligned} & \text{(cylinder or frustum } EB') \quad \text{(inscribed figure)} \\ &= (\text{sum of all the spaces } S) \quad (ap + p^2) + (aq + q^2) + \\ &> (a+b) \quad \left(\frac{a}{2} + \frac{b}{3}\right) \quad \text{[Prop 2]} \end{aligned}$$

$$> A'D \quad \frac{HD}{3} \quad \text{since } a = AA' \quad b = AD,$$

$$> (EB) \quad V \quad \text{by } (\beta) \text{ above}$$

Hence

(inscribed figure) $< V$

But this is impossible because by (γ) above the inscribed figure is greater than V

II Next suppose if possible, that the segment is less than V

In this case we circumscribe and inscribe figures such that

$$(\text{circumscribed fig}) - (\text{inscribed fig}) < V - (\text{segment}),$$

whence we derive

$$V > (\text{circumscribed figure}) \quad (\delta)$$

We now compare successive cylinders or frusta in the complete cylinder or frustum and in the circumscribed figure and we have

$$\begin{aligned} & \text{(first cylinder or frustum in } EB') \quad \text{(first in circumscribed fig)} \\ &= S \quad S \\ &= S \quad (ab + b^2) \\ & \text{(second in } EB) \quad \text{(second in circumscribed fig)} \\ &= S \quad (ap + p^2) \end{aligned}$$

and so on

Hence [Prop 1]

$$\begin{aligned} & \text{(cylinder or frustum } EB) \quad \text{(circumscribed fig)} \\ &= (\text{sum of all the spaces } S) \quad (ab + b^2) + (ap + p^2) + \end{aligned}$$

$$< (a+b) \quad \left(\frac{a}{2} + \frac{b}{3}\right) \quad \text{[Prop 2]}$$

$$< A'D \quad \frac{HD}{3}$$

$$< (EB) \quad V \quad \text{by } (\beta) \text{ above}$$

Hence the circumscribed figure is greater than I which is impossible, by (8) above.

Thus the segment is neither greater nor less than I, and is therefore equal to it.

Therefore by (α),

$$\begin{aligned} & (\text{segment } ABB) \quad (\text{cone or segment of cone } ABB') \\ & = (AD + 3CI) \quad (AD + 2C4) \end{aligned}$$

PROPOSITIONS 27, 28, 29, 30

(1) In any spheroid whose centre is C, if a plane meeting the axis cut off a segment not greater than half the spheroid and having A for its vertex and AD for its axis, and if A'D be the axis of the remaining segment of the spheroid then

$$\begin{aligned} & (\text{first segmt}) \quad (\text{cone or segmt of cone with same base and axis}) \\ & = CI + AD \quad A'D \\ & [= 3CI - ID \quad 2CI - AD] \end{aligned}$$



(2) As a particular case, if the plane passes through the centre so that the segment is half the spheroid, half the spheroid is double of the cone or segment of a cone which has the same vertex and axis.

Let the plane cutting off the segment be at right angles to the plane of the paper and let the latter plane be the plane through the axis of the spheroid which intersects the cutting plane in BB' and makes the elliptic section BB'B.

Let EF EF' be the two tangents to the ellipse which are parallel to BB' let them touch it in A and through the tangents draw planes parallel to the base of the segment. These planes will touch the spheroid at I I' which will be the vertices of the two segments into which it is divided. Also AA' will pass through the centre C and bisect BB' in D.

Then (1) if the base of the segments be perpendicular to the axis of the spheroid AA' will be the vertices of the spheroid as well as of the segments. AA' will be the axis of the spheroid and the base of the segments will be a circle on BB' as diameter.

(2) if the base of the segments be not perpendicular to the axis of the spheroid the base of the segments will be an ellipse of which BB' is one axis and ID I'D will be the axes of the segments respectively.



We can now draw a cylinder or a frustum of a cylinder $EBB'F$ through the circle or ellipse about BB' and having AD for its axis, and we can also draw a cone or a segment of a cone passing through the circle or ellipse about BB' and having A for its vertex.

We have then to show that if CA' be produced to H so that $CA' = A'H$,
(segment ABB) (cone or segment of cone ABB') = $HD \cdot A'D$.

Let V be such a cone that

$$V \text{ (cone or segment of cone } ABB') = HD \cdot A'D \quad (\alpha)$$

and we have to show that the segment ABB is equal to V

But, since

$$\text{(cylinder or frustum } EB) \text{ (cone or segment of cone } ABB') = 3 \cdot I,$$

we have, by the aid of (α)

$$\text{(cylinder or frustum } EB) \cdot V = A'D \cdot \frac{HD}{3} \quad (\beta)$$

Now, if the segment ABB is not equal to V , it must be either greater or less

I Suppose if possible, that the segment is greater than V

Let figures be inscribed and circumscribed to the segment consisting of cylinders or frusta of cylinders, with axes along AD and all equal to one another, such that

$$\text{(circumscribed fig)} - \text{(inscribed fig)} < (\text{segment}) - V$$

whence it follows that

$$\text{(inscribed fig)} > V \quad (\gamma)$$

Produce all the planes forming the bases of the cylinders or frusta to meet the surface of the complete cylinder or frustum EB' . Thus if ND be the axis of the greatest cylinder or frustum of a cylinder in the circumscribed figure the complete cylinder or frustum EB' will be divided into cylinders or frusta of cylinders each equal to the greatest of those in the circumscribed figure.

Take straight lines da' each equal to $A'D$ and as many in number as the parts into which AD is divided by the bases of the cylinders or frusta, and measure da along da' equal to AD . It follows that $aa' = 2CD$.

Apply to each of the lines $a'd$ rectangles with height equal to ad and draw the squares on each of the lines ad as in the figure. Let S denote the area of each complete rectangle.

From the first rectangle take away a gnomon with breadth equal to AN (i.e. with each end of a length equal to AN) take away from the second rectangle a gnomon with breadth equal to AM and so on the last rectangle having no gnomon taken from it.

Then

$$\begin{aligned} \text{the first gnomon} &= AD \cdot ID - ND \cdot (AD - AN) \\ &= A'D \cdot AN + ND \cdot AN \\ &= AN \cdot ID \end{aligned}$$

Similarly

$$\text{the second gnomon} = IM \cdot ID$$

and so on

And the last gnomon (that in the last rectangle but one) is equal to $AL \cdot IL$.

Also after the gnomons are taken away from the successive rectangles the remainders (which we will call R_1, R_2, \dots, R_n where n is the number of rectangles and accordingly $R_n = S$) are rectangles applied to straight lines each of

length aa' and "exceeding by squares" whose sides are respectively equal to DN, DM, DA

For brevity let DV be denoted by x , and aa' or $2CD$ by c , so that

$$R_1 = cx + x^2 \quad R_2 = c \cdot 2x + (2x)^2,$$

Then, comparing successively the cylinders or frusta of cylinders (1) in the complete cylinder or frustum EB and (2) in the inscribed figure, we have

(first cylinder or frustum in EB) (first in inscribed fig)

$$= BD^2 - PV^2$$

$$= AD \cdot A'D - AN \cdot A'N$$

$$= S \quad (\text{first gnomon}),$$

(second cylinder or frustum in EB') (second in inscribed fig)

$$= S \quad (\text{second gnomon})$$

and so on

The last of the cylinders or frusta in the cylinder or frustum EB has none corresponding to it in the inscribed figure, and there is no corresponding gnomon

Combining the proportions, we have [by Prop. 1]

(cylinder or frustum EB) (inscribed fig)

$$= (\text{sum of all spaces } S) \quad (\text{sum of gnomons})$$

Now the differences between S and the successive gnomons are R_1, R_2, \dots, R_n , while

$$R_1 = cx + x^2,$$

$$R_2 = c \cdot 2x + (2x)^2,$$

$$R_n = cb + b^2 = S,$$

where $b = nx = AD$

Hence [Prop. 2]

$$(\text{sum of all spaces } S) \quad (R_1 + R_2 + \dots + R_n) < (c + b) \quad \left(\frac{c}{2} + \frac{b}{3}\right)$$

It follows that

$$\begin{aligned} (\text{sum of all spaces } S) \quad (\text{sum of gnomons}) &> (c + b) \quad \left(\frac{c}{2} + \frac{2b}{3}\right) \\ &> 1'D \quad \frac{HD}{3} \end{aligned}$$

Thus (cylinder or frustum EB) (inscribed fig)

$$> AD \quad \frac{HD}{3}$$

$$> (\text{cylinder or frustum } EB) \quad V,$$

from (β) above

Therefore (inscribed fig) $< V$,

which is impossible by (γ) above

Hence the segment ABB' is not greater than V

II If possible let the segment ABB' be less than V

We then inscribe and circumscribe figures such that

$$(\text{circumscribed fig}) - (\text{inscribed fig}) < 1' - (\text{segment}),$$

whence

$$1' > (\text{circumscribed fig})$$

(δ)

In this case we compare the cylinders or frusta in (EB) with those in the circumscribed figure

Thus
 (first cylinder or frustum in EB') (first in circumscribed fig)
 $= S$
 (second in EB') (second in circumscribed fig)
 $= S$ (first gnomon),

and so on

Lastly (last in EB') (last in circumscribed fig)
 $= S$ (last gnomon)

Now

$$\{S + (\text{all the gnomons})\} = nS - (R_1 + R_2 + \dots + R_{n-1})$$

And $nS - R_1 - R_2 - \dots - R_{n-1} > (c+b) \left(\frac{c}{2} + \frac{b}{3}\right)$ [Prop 2]

so that

$$nS - \{S + (\text{all the gnomons})\} < (c+b) \left(\frac{c}{2} + \frac{2b}{3}\right)$$

It follows that, if we combine the above proportions as in Prop 1, we obtain
 (cylinder or frustum EB') (circumscribed fig)

$$< (c+b) \left(\frac{c}{2} + \frac{2b}{3}\right)$$

$$< A'D \frac{HD}{3}$$

$$< (EB) V, \text{ by } (\beta) \text{ above}$$

Hence the circumscribed figure is greater than V , which is impossible by (δ) above

Thus since the segment ABB is neither greater nor less than V , it is equal to it and the proposition is proved

(2) The particular case [Props 27, 28] where the segment is half the spheroid differs from the above in that the distance CD or $c/2$ vanishes, and the rectangles $cb+b$ are simply squares (b) so that the gnomons are simply the differences between b^2 and x^2 , b^2 and $(2x)^2$, and so on

Instead therefore of Prop 2 we use the Lemma to Prop 2, Cor 1 given above [On Spirals, Prop 10] and instead of the ratio $(c+b) \left(\frac{c}{2} + \frac{2b}{3}\right)$ we obtain the ratio $3/2$ whence (segment ABB) (cone or segment of cone ABB) = $2/1$

PROPOSITIONS 31, 32


If a plane divide a spheroid into two unequal segments and if AN $A'N$ be the axes of the lesser and greater segments respectively while C is the centre of the spheroid then

(greater segmt) (cone or segmt of cone with same base and axis)
 $= CA + AN$ AN

Let the plane dividing the spheroid be that through PP' perpendicular to the plane of the paper and let the latter plane be that through the axis of the spheroid which intersects the cutting plane in PP and makes the elliptic section $PAP A$

Draw the tangents to the ellipse which are parallel to PP let them touch the ellipse at A A' , and through the tangents draw planes parallel to the base of the segments These planes will touch the spheroid at A , A' , the line AA'

will pass through the centre C and bisect PP' in N , while AN , $A'N$ will be the axes of the segments



Then (1) if the cutting plane be perpendicular to the axis of the spheroid, AA' will be that axis and A, A' will be the vertices of the spheroid as well as of the segments. Also the sections of the spheroid by the cutting plane and all planes parallel to it will be circles.

(2) If the cutting plane be not perpendicular to the axis, the base of the segments will be an ellipse of which PP' is an axis and the sections of the spheroid by all planes parallel to the cutting plane will be similar ellipses.

Draw a plane through C parallel to the base of the segments and meeting the plane of the paper in BB' .

Construct three cones or segments of cones, two having A for their common vertex and the plane sections through PP' , BB' for their respective bases, and a third having the plane section through PP' for its base and A' for its vertex.

Produce CA to H and CA' to H' so that

$$AH = A'H' = CA$$

We have then to prove that

$$\begin{aligned} (\text{segment } APP') &= (\text{cone or segment of cone } A'PP') \\ &= CA + AN \quad AN \\ &= NH \quad HN \end{aligned}$$

Now half the spheroid is double of the cone or segment of a cone ABB' [Props 27-28]. Therefore

$$(\text{the spheroid}) = 4(\text{cone or segment of cone } ABB')$$

But

$$\begin{aligned} (\text{cone or segmt of cone } ABB') &= (\text{cone or segmt of cone } APP') \\ &= (CA \quad AN) (BC^2 \quad PN^2) \\ &= (CA \quad AN) (CA \quad CA' \quad AN \quad A'N) \end{aligned} \quad (\alpha)$$

If we measure AK along AA' so that

$$\frac{AK}{AN} = \frac{AC}{AN} = \frac{AC}{AN} = \frac{CA}{AN},$$

and the compound ratio in (α) becomes

$$\frac{(AK \quad AN \quad CA \quad A'N) (CA \quad CA' \quad AN \quad A'N)}{AK \quad CA' \quad AN \quad AN}$$

Thus

$$\begin{aligned} & (\text{cone or segmt of cone } ABB') \quad (\text{cone or segmt of cone } APP) \\ & \quad = AK \quad CA' \quad AN \quad A'N \end{aligned}$$

$$\begin{aligned} \text{But} \quad & (\text{cone or segment of cone } APP) \quad (\text{segment } APP) \\ & \quad = A'N \quad NH' \\ & \quad = AN \quad A'N \quad AN \quad NH' \end{aligned} \quad [\text{Props } 29, 30]$$

Therefore *ex aequali*

$$\begin{aligned} & (\text{cone or segment of cone } ABB) \quad (\text{segment } APP') \\ & \quad = AK \quad CA' \quad AN \quad NH', \end{aligned}$$

$$\begin{aligned} \text{so that} \quad & (\text{spheroid}) \quad (\text{segment } APP) \\ & \quad = HH' \quad AK \quad AN \quad NH', \end{aligned}$$

$$\text{since} \quad HH' = 4CA'$$

$$\begin{aligned} \text{Hence} \quad & (\text{segment } A'PP) \quad (\text{segment } APP') \\ & = (HH \quad AK - AN \quad NH) \quad AN \quad NH' \\ & = (AK \quad NH + NH' \quad NK) \quad AN \quad NH' \end{aligned}$$

Further,

$$\begin{aligned} & (\text{segment } APP) \quad (\text{cone or segment of cone } APP') \\ & \quad = NH \quad A'N \\ & \quad = AN \quad NH' \quad AN \quad A'N, \end{aligned}$$

and

$$\begin{aligned} & (\text{cone or segmt of cone } APP') \quad (\text{cone or segmt of cone } A'PP') \\ & \quad = AN \quad A'N \\ & \quad = AN \quad A'N \quad A'N^2 \end{aligned}$$

From the last three proportions we obtain *ex aequali*

$$\begin{aligned} & (\text{segment } A'PP') \quad (\text{cone or segment of cone } A'PP') \\ & \quad = (AK \quad NH + NH' \quad NK) \quad A'N^2 \\ & \quad = (AK \quad NH + NH' \quad NK) \quad (CA^2 + NH' \quad CN) \\ & \quad = (AK \quad NH + NH' \quad NK) \quad (AK \quad AN + NH \quad CN) \end{aligned} \quad (\beta)$$

But

$$\begin{aligned} AK \quad NH \quad AK \quad AN &= NH \quad AN \\ &= CA + AN \quad AN \\ &= AK + CA \quad CA \quad (\text{since } AK \quad AC = AC \quad AN) \\ &= HK \quad CA \\ &= HK - NH \quad CA - AN \\ &= NK \quad CN \\ &= NH \quad NK \quad NH \quad CN \end{aligned}$$

Hence the ratio in (β) is equal to the ratio

$$AK \quad NH \quad AK \quad AN, \text{ or } NH \quad AN$$

Therefore

$$\begin{aligned} & (\text{segment } A'PP) \quad (\text{cone or segment of cone } A'PP') \\ & \quad = NH \quad AN \\ & \quad = CA + AN \quad AN \end{aligned}$$

ON SPIRALS

"ARCHIMEDES to Dositheus greeting

Of most of the theorems which I sent to Conon, and of which you ask me from time to time to send you the proofs the demonstrations are already before you in the books brought to you by Heracleides and some more are also contained in that which I now send you. Do not be surprised at my taking a considerable time before publishing these proofs. This has been owing to my desire to communicate them first to persons engaged in mathematical studies and anxious to investigate them. In fact, how many theorems in geometry which have seemed at first impracticable are in time successfully worked out! Now Conon died before he had sufficient time to investigate the theorems referred to otherwise he would have discovered and made manifest all these things and would have enriched geometry by many other discoveries besides. For I know well that it was no common ability that he brought to bear on mathematics and that his industry was extraordinary. But though many years have elapsed since Conon's death I do not find that any one of the problems has been stirred by a single person. I wish now to put them in review one by one, particularly as it happens that there are two included among them which are impossible of realisation (and which may serve as a warning) how those who claim to discover everything but produce no proofs of the same may be confuted as having actually pretended to discover the impossible.

What are the problems I mean, and what are those of which you have already received the proofs, and those of which the proofs are contained in this book respectively, I think it proper to specify. The first of the problems was Given a sphere, to find a plane area equal to the surface of the sphere and this was first made manifest on the publication of the book concerning the sphere, for when it is once proved that the surface of any sphere is four times the greatest circle in the sphere, it is clear that it is possible to find a plane area equal to the surface of the sphere. The second was Given a cone or a cylinder to find a sphere equal to the cone or cylinder, the third To cut a given sphere by a plane so that the segments of it have to one another an assigned ratio the fourth To cut a given sphere by a plane so that the segments of the surface have to one another an assigned ratio the fifth To make a given segment of a sphere similar to a given segment of a sphere,¹ the sixth Given two segments of either the same or different spheres to find a segment of a sphere which shall be similar to one of the segments and have its surface equal to the surface of the other segment. The seventh was From a given sphere to cut off a segment by a plane so that the segment bears to the cone which has the same base

¹Cf. *On the Sphere and Cylinder* II. 5

as the segment and equal height an assigned ratio greater than that of three to two. Of all the propositions just enumerated Heracleides brought you the proofs. The proposition stated next after these was wrong viz that, if a sphere be cut by a plane into unequal parts the greater segment will have to the less the duplicate ratio of that which the greater surface has to the less. That this is wrong is obvious by what I sent you before, for it included this proposition. If a sphere be cut into unequal parts by a plane at right angles to any diameter in the sphere the greater segment of the surface will have to the less the same ratio as the greater segment of the diameter has to the less while the greater segment of the sphere has to the less a ratio less than the duplicate ratio of that which the greater surface has to the less but greater than the sesqui alterate¹ of that ratio. The last of the problems was also wrong viz that, if the diameter of any sphere be cut so that the square on the greater segment is triple of the square on the lesser segment, and if through the point thus arrived at a plane be drawn at right angles to the diameter and cutting the sphere, the figure in such a form as is the greater segment of the sphere is the greatest of all the segments which have an equal surface. That this is wrong is also clear from the theorems which I before sent you. For it was there proved that the hemisphere is the greatest of all the segments of a sphere bounded by an equal surface.

After these theorems the following were propounded concerning the cone: If a section of a right-angled cone [a parabola] in which the diameter [axis] remains fixed be made to revolve so that the diameter [axis] is the axis [of revolution] let the figure described by the section of the right angled cone be called a *conoid*. And if a plane touch the conoidal figure and another plane drawn parallel to the tangent plane cut off a segment of the conoid, let the base of the segment cut off be defined as the cutting plane and the vertex as the point in which the other plane touches the conoid. Now, if the said figure be cut by a plane at right angles to the axis it is clear that the section will be a circle, but it needs to be proved that the segment cut off will be half as large again as the cone which has the same base as the segment and equal height. And if two segments be cut off from the conoid by planes drawn in any manner, it is clear that the sections will be sections of acute-angled cones [ellipses] if the cutting planes be not at right angles to the axis but it needs to be proved that the segments will bear to one another the ratio of the squares on the lines drawn from their vertices parallel to the axis to meet the cutting planes. The proofs of these propositions are not yet sent to you.

After these came the following propositions about the *spiral* which are as it were another sort of problem having nothing in common with the foregoing, and I have written out the proofs of them for you in this book. They are as follows. If a straight line of which one extremity remains fixed be made to revolve at a uniform rate in a plane until it returns to the position from which it started and if at the same time as the straight line revolves a point move at a uniform rate along the straight line starting from the fixed extremity the point will describe a *spiral* in the plane. I say then that the area bounded by the spiral and the straight line which has returned to the position from which it started is a third part of the circle described with the fixed point as centre and with radius the length traversed by the point along the straight line during

¹See On the Sphere and Cylinder II 8

²This should be presumably the *conoid* not the cone

the one revolution. And, if a straight line touch the spiral at the extreme end of the spiral and another straight line be drawn at right angles to the line which has revolved and resumed its position from the fixed extremity of it, so as to meet the tangent. I say that the straight line so drawn to meet it is equal to the circumference of the circle. Again, if the revolving line and the point moving along it make several revolutions and return to the position from which the straight line started. I say that the area added by the spiral in the third revolution will be double of that added in the second that in the fourth three times, that in the fifth four times and generally the areas added in the later revolutions will be multiples of that added in the second revolution according to the successive numbers while the area bounded by the spiral in the first revolution is a sixth part of that added in the second revolution. Also if on the spiral described in one revolution two points be taken and straight lines be drawn joining them to the fixed extremity of the revolving line and if two circles be drawn with the fixed point as centre and radii the lines drawn to the fixed extremity of the straight line and the shorter of the two lines be produced. I say that (1) the area bounded by the circumference of the greater circle in the direction of (the part of) the spiral included between the straight lines the spiral (itself) and the produced straight line will be to (2) the area bounded by the circumference of the lesser circle the same (part of the) spiral and the straight line joining their extremities the ratio which (3) the radius of the lesser circle together with two thirds of the excess of the radius of the greater circle over the radius of the lesser bears to (4) the radius of the lesser circle together with one third of the said excess.

The proofs then of these theorems and others relating to the spiral are given in the present book. Prefixed to them, after the manner usual in other geometrical works are the propositions necessary to the proofs of them. And here too as in the books previously published I assume the following lemma, that if there be (two) unequal lines or (two) unequal areas the excess by which the greater exceeds the less can by being [continually] added to itself be made to exceed any given magnitude among those which are comparable with [it and with] one another.

PROPOSITION 1

If a point move at a uniform rate along any line, and two lengths be taken on it, they will be proportional to the times of describing them.

Two unequal lengths are taken on a straight line and two lengths on another straight line representing the times and they are proved to be proportional by taking equimultiples of each length and the corresponding time after the manner of Eucl. V Def. 5.

PROPOSITION 2

If each of two points on different lines respectively move along them each at a uniform rate and if lengths be taken one on each line forming pairs, such that each pair are described in equal times the lengths will be proportionals.

This is proved at once by equating the ratio of the lengths taken on one line to that of the times of description which must also be equal to the ratio of the lengths taken on the other line.

PROPOSITION 3

Given any number of circles it is possible to find a straight line greater than the sum of all their circumferences

For we have only to describe polygons about each and then take a straight line equal to the sum of the perimeters of the polygons

PROPOSITION 4

Given two unequal lines, viz a straight line and the circumference of a circle, it is possible to find a straight line less than the greater of the two lines and greater than the less

For by the Lemma, the excess can, by being added a sufficient number of times to itself, be made to exceed the lesser line

Thus e.g., if $c > l$ (where c is the circumference of the circle and l the length of the straight line) we can find a number n such that

$$n(c-l) > l$$

Therefore
$$c-l > \frac{l}{n}$$

and
$$c > l + \frac{l}{n} > l$$

Hence we have only to divide l into n equal parts and add one of them to l . The resulting line will satisfy the condition

PROPOSITION 5

Given a circle with centre O and the tangent to it at a point A it is possible to draw from O a straight line OPF , meeting the circle in P and the tangent in F , such that, if c be the circumference of any given circle whatever

$$FP \cdot OP < (\text{arc } AP) \cdot c$$

Take a straight line as D greater than the circumference c [Prop 3]

Through O draw OH parallel to the given tangent and draw through A a line APH meeting the circle in P and OH in H such that the portion PH intercepted between the circle and the line OH may be equal to D . Join OP and produce it to meet the tangent in F .

$$\begin{aligned} \text{Then } FP \cdot OP &= AP \cdot PH \text{ by parallels} \\ &= AP \cdot D \\ &< (\text{arc } AP) \cdot c \end{aligned}$$


PROPOSITION 6

Given a circle with centre O , a chord AB less than the diameter and OM the perpendicular on AB from O it is possible to draw a straight line OFP meeting the chord AB in F and the circle in P such that

$$FP \cdot PB = D \cdot E$$

where $D \cdot E$ is any given ratio less than $BM \cdot MO$

Draw OH parallel to AB and BT perpendicular to BO meeting OH in T



Then the triangles BMO , OBT are similar, and therefore

$$BM : MO = OB : BT,$$

whence

$$D : E < OB : BT$$

Suppose that a line PH (greater than BT) is taken such that



$$D : E = OB : PH$$

and let PH be so placed that it passes through B and P lies on the circumference of the circle, while H is on the line OH (PH will fall outside BT , because $PH > BT$)

Join OP meeting AB in F

We now have

$$\begin{aligned} FI : PB &= OP : PH \\ &= OB : PH \\ &= D : E \end{aligned}$$



PROPOSITION 7

Given a circle with centre O a chord AB less than the diameter, and OM the perpendicular on it from O it is possible to draw from O a straight line OPF , meeting the circle in P and AB produced in F such that

$$FP : PB = D : E$$

where $D : E$ is any given ratio greater than $BM : MO$

Draw OT parallel to AB and BT perpendicular to BO meeting OT in T

In this case

$$D : E > BM : MO$$

$$> OB : BT, \text{ by similar triangles}$$



Take a line PH (less than BT) such that

$$D : E = OB : PH,$$

and place PH so that P , H are on the circle and on OT respectively while HP produced passes through B

Then

$$\begin{aligned} FP : PB &= OP : PH \\ &= D : E \end{aligned}$$



PROPOSITION 8

Given a circle with centre O , a chord AB less than the diameter the tangent at B , and the perpendicular OM from O on AB it is possible to draw from O a straight line OFP , meeting the chord AB in F the circle in P and the tangent in G , such that

$$FP : BC = D : E$$

where $D : E$ is any given ratio less than $BM : MO$

If OT be drawn parallel to AB meeting the tangent at B in T ,

$$BM : MO = OB : BT,$$

so that


$$D : E < OB : BT$$

Take a point C on TB produced such that

$$D : E = OB : BC,$$

whence

$$BC > BT$$



or

$$\begin{array}{rcl}
 CG & GT & OF \\
 CG & OF & GT = OG \\
 & & = BG \\
 & & = BC \\
 & & = BC \\
 OP & OF & = BC \\
 PF & OP & = BG \\
 PF & BG & = OP \\
 & & = OB \\
 & & = D
 \end{array}
 \begin{array}{rcl}
 GT = OG & GQ & OG \\
 GT = OG & GT & BT \\
 BT & BT & by construction, \\
 OB & OB & \\
 OP & OP & \\
 CG & CG & \\
 BC & BC & \\
 BC & BC & \\
 BC & BC & \\
 E
 \end{array}$$


Hence
and therefore
or

PROPOSITION 9

Given a circle with centre O a chord AB less than the diameter the tangent at B , and the perpendicular OM from O on AB it is possible to draw from O a straight line $OPGF$, meeting the circle in P , the tangent in G , and AB produced in F , such that

$$FP : BG = D : E,$$

where $D : E$ is any given ratio greater than $BM : MO$



in Q such that $GQ = BK$. Let OQ meet the original circle in P and AB produced in F

Through the points O, T, C describe a circle and let OB be produced to meet this circle in K

Then, since $BC > BT$, and OB is perpendicular to CT it is possible to draw from O a straight line OGQ meeting CT in G and the circle about OTC in Q , such that $GQ = BK$

Let OGQ meet AB in F and the original circle in P

Now $CG : GT = OG : GQ$,
and $OF : OG = BT : GT$,
so that $OF : GT = OG : BT$

It follows that

$$\begin{array}{rcl}
 CG & GT & OF \\
 CG & OF & GT = OG \\
 & & = BG \\
 & & = BC \\
 & & = BC \\
 OP & OF & = BC \\
 PF & OP & = BG \\
 PF & BG & = OP \\
 & & = OB \\
 & & = D
 \end{array}
 \begin{array}{rcl}
 GT = OG & GQ & OG \\
 GT = OG & GT & BT \\
 BT & BT & by construction, \\
 OB & OB & \\
 OP & OP & \\
 CG & CG & \\
 BC & BC & \\
 BC & BC & \\
 BC & BC & \\
 E
 \end{array}$$

Let OT be drawn parallel to AB meeting the tangent at B in T

Then

$$D : E > BM : MO$$

$> OB : BT$ by similar triangles

Produce TB to C so that

$$D : E = OB : BC$$

whence $BC < BT$

Describe a circle through the points O, T, C and produce OB to meet this circle in K

Then since $TB > BC$, and OB is perpendicular to CT it is possible to draw from O a line OGQ meeting CT in G and the circle about OTC

We now prove, exactly as in the last proposition, that

$$\begin{array}{l} CG \quad OF = BA \quad BT \\ \quad \quad \quad = BC \quad OP \end{array}$$

Thus as before,

$$OP \quad OF = BC \quad CG,$$

and
whence

$$\begin{array}{l} OP \quad PF = BC \quad BG, \\ PF \quad BC = OP \quad HC \\ \quad \quad \quad = OB \quad BC \\ \quad \quad \quad = D \quad F \end{array}$$

PROPOSITION 10

If $A_1, A_2, A_3 \dots A_n$ be n lines forming an ascending arithmetical progression in which the common difference is equal to A_1 , the least term, then

$$(n+1)A_1^2 + A_1(A_1 + A_2 + \dots + A_n) = 3(1^2 + 2^2 + \dots + n^2)$$

[Archimedes' proof of this proposition is given above, pp 456-7, and it is there pointed out that the result is equivalent to

$$1^2 + 2^2 + 3^2 + \dots + n^2 = \frac{n(n+1)(2n+1)}{6}]$$

CON 1 It follows from this proposition that

$$n A_1^2 < 3(1^2 + 2^2 + \dots + n^2),$$

and also that

$$n A_n^2 > 3(A_1^2 + 1^2 + \dots + A_n^2)$$

[For the proof of the latter inequality see p 457 above]

CON 2 All the results will equally hold if similar figures are substituted for squares

PROPOSITION 11

If $A_1, A_2, A_3 \dots A_n$ be n lines forming an ascending arithmetical progression [in which the common difference is equal to the least term A_1], then


$$(n-1)A_1^2 \{A_1 + 1^2 A_{n-1}^2 + \dots + A_n^2\} < A_n^2 \{A_1 + 1^2(A_1 - A_1)^2\}$$

but

$$(n-1)A_n \{A_{n-1} + A_{n-2} + \dots + A_1\} > 1 \{A_1 + 1^2(1 - A_1)^2\}$$

[Archimedes sets out the terms side by side in the manner shown in the figure where $BC = A$, $DE = A_{n-1}$, $RS = A_1$ and produces DE , FG , RS until they are respectively equal to BC or A so that EH , GI , SU in the figure are respectively equal to A_1 , A_2 , A_{n-1} . He further measures lengths BA , DL , FM , PL along BC , DE , FG , PQ respectively each equal to RS

The figure makes the relations between the terms easier to see with the eye but the use of so large a number of letters makes the proof somewhat difficult to follow, and it may be more clearly represented as follows]



It is evident that $(A_n - A_1) = A_{n-1}$

The following proportion is therefore obviously true viz

$$(n-1)A_n^2 \{ (n-1)(A_1 + \frac{1}{2}A_{n-1}) \} = A_n^2 \{ 1 \cdot A_1 + \frac{1}{2}(A_n - A_1)^2 \}$$

In order therefore to prove the desired result we have only to show that

$$(n-1) \frac{1}{2} A_1 + \frac{1}{2}(n-1)A_{n-1} < (A_1^2 + A_{n-1}^2 + A_1^2) \\ \text{but} > (A_{n-1}^2 + A_{n-2}^2 + A_1^2)$$

I To prove the first inequality we have

$$(n-1)A_1 + \frac{1}{2}(n-1)A_{n-1} \\ = (n-1) \frac{1}{2} A_1^2 + (n-1)A_1 \frac{1}{2} A_{n-1} + \frac{1}{2}(n-1)A_{n-1}^2 \quad (1)$$

And

$$A_n^2 + A_{n-1}^2 + A_1^2 = (A_{n-1} + A_1)^2 + (A_{n-2} + A_1)^2 + (A_1 + A_1)^2 \\ = (A_{n-1}^2 + A_{n-2}^2 + A_1^2) \\ + (n-1)A_1^2 \\ + 2A_1(A_{n-1} + A_{n-2} + A_1) \\ = (A_{n-1}^2 + A_{n-2}^2 + A_1^2) \\ + (n-1)A_1^2 \\ + A_1\{A_{n-1} + A_{n-2} + A_{n-3} + \dots + A_1 \\ + A_1 + A_1 + \dots + A_{n-2} + A_{n-1}\} \\ = (A_{n-1}^2 + A_{n-2}^2 + A_1^2) \\ + (n-1)A_1^2 \\ + nA_1 A_{n-1} \quad (2)$$

Comparing the right hand sides of (1) and (2), we see that $(n-1)A_1$ is common to both sides, and

$$(n-1)A_1 A_{n-1} < nA_1 A_{n-1}$$

while by Prop 10 Cor 1

$$\frac{1}{2}(n-1)A_{n-1}^2 < A_{n-1}^2 + A_{n-2}^2 + A_1^2$$

It follows therefore that

$$(n-1)A_1 + \frac{1}{2}(n-1)A_{n-1}^2 < (A_n^2 + A_{n-1}^2 + A_1^2)$$

and hence the first part of the proposition is proved

II We have now in order to prove the second result, to show that

$$(n-1)A_n + \frac{1}{2}(n-1)A_{n-1}^2 > (A_{n-1}^2 + A_{n-2}^2 + A_1^2)$$

The right hand side is equal to

$$(A_{n-2} + A_1)^2 + (A_{n-3} + A_1)^2 + \dots + (A_1 + A_1)^2 + A_1 \\ = A_{n-2}^2 + A_{n-3}^2 + \dots + A_1^2 \\ + (n-1)A_1 \\ + 2A_1(A_{n-2} + A_{n-3} + \dots + A_1) \\ = (A_{n-2}^2 + A_{n-3}^2 + \dots + A_1^2) \\ + (n-1)A_1^2 \\ + A_1\{A_{n-2} + A_{n-3} + \dots + A_1 \\ + A_1 + A_2 + \dots + A_{n-2}\} \\ = (A_{n-2}^2 + A_{n-3}^2 + \dots + A_1^2) \\ + (n-1)A_1^2 \\ + (n-2)A_1 A_{n-1} \quad (3)$$

Comparing this expression with the right hand side of (1) above, we see that $(n-1)A_1$ is common to both sides and

$$(n-1)A_1 A_{n-1} > (n-2)A_1 A_{n-1}$$

while by Prop 10 Cor 1,

$$\frac{1}{2}(n-1)A_{n-1}^2 > (A_{n-2}^2 + A_{n-3}^2 + \dots + A_1^2)$$

Hence $(n-1)A_n + \frac{1}{2}(n-1)A_{n-1}^2 > (A_{n-1}^2 + A_{n-2}^2 + A_1^2)$,

and the second required result follows

COR The results in the above proposition are equally true if similar figures be substituted for squares on the several lines

DEFINITIONS

1 If a straight line drawn in a plane revolve at a uniform rate about one extremity which remains fixed and return to the position from which it started, and if, at the same time as the line revolves a point move at a uniform rate along the straight line beginning from the extremity which remains fixed the point will describe a *spiral* (ὄλιξ) in the plane

2 Let the extremity of the straight line which remains fixed while the straight line revolves be called the *origin* of the spiral

3 And let the position of the line from which the straight line began to revolve be called the *initial line* in the revolution

4 Let the length which the point that moves along the straight line describes in one revolution be called the *first distance* that which the same point describes in the second revolution the *second distance*, and similarly let the distances described in further revolutions be called after the number of the particular revolution

5 Let the area bounded by the spiral described in the first revolution and the *first distance* be called the *first area* that bounded by the spiral described in the second revolution and the *second distance* the *second area*, and similarly for the rest in order

6 If from the origin of the spiral any straight line be drawn let that side of it which is in the same direction as that of the revolution be called *forward* (πρὸσφυόμενα), and that which is in the other direction *backward* (εὐρόμενα)

7 Let the circle drawn with the *origin* as centre and the *first distance* as radius be called the *first circle* that drawn with the same centre and twice the radius the *second circle*, and similarly for the succeeding circles

PROPOSITION 12

If any number of straight lines drawn from the origin to meet the spiral make equal angles with one another the lines will be in arithmetical progression

[The proof is obvious]

PROPOSITION 13

If a straight line touch the spiral it will touch it in one point only

Let O be the origin of the spiral and BC a tangent to it

If possible let BC touch the spiral in two points P Q Join OP OQ and bisect the angle POQ by the straight line OR meeting the spiral in R


Then [Prop 12] OR is an arithmetic mean between OP and OQ or

$$OP + OQ = 2OR$$

But in any triangle POQ if the bisector of the angle POQ meets PQ in K


$$OP + OQ > 2OK$$

Therefore $OK < OR$ and it follows that some point on BC between P and Q lies within the spiral Hence BC cuts the spiral, which is contrary to the hypothesis



PROPOSITION 14

If O be the origin, and P, Q two points on the first turn of the spiral and if OP, OQ produced meet the 'first circle' $AKPQ$ in P', Q' respectively, OA being the initial line, then



$$OP : OQ = (\text{arc } AKP) : (\text{arc } AKQ)$$

For while the revolving line OA moves about O the point A on it moves uniformly along the circumference of the circle $AKP'Q'$ and at the same time the point describing the spiral moves uniformly along OA .

Thus while A describes the arc AKP' , the moving point on OA describes the length OP , and while A describes the arc AKQ' the moving point on OA describes the distance OQ .

Hence

$$OP : OQ = (\text{arc } AKP) : (\text{arc } AKQ)$$

[Prop 2]

PROPOSITION 15

If P, Q be points on the second turn of the spiral, and OP, OQ meet the 'first circle' $AKPQ$ in P', Q' , as in the last proposition and if c be the circumference of the 'first circle' then

$$OP : OQ = c + (\text{arc } AKP') : c + (\text{arc } AKQ')$$


For, while the moving point on OA describes the distance OP the point A describes the whole of the circumference of the first circle together with the arc AKP' and while the moving point on OA describes the distance OQ the point A describes the whole circumference of the first circle together with the arc AKQ' .

Cor. Similarly, if P, Q are on the n th turn of the spiral

$$OP : OQ = (n-1)c + (\text{arc } AKP') : (n-1)c + (\text{arc } AKQ')$$

PROPOSITIONS 16 17

If BC be the tangent at P , any point on the spiral PC being the forward part of BC , and if OP be joined, the angle OPC is obtuse while the angle OPB is acute.



I Suppose P to be on the first turn of the spiral.

Let OA be the initial line AKP the first circle. Draw the circle DLP with centre O and radius OP meeting OA in D . This circle must then in the forward direction from P fall within the spiral and in the backward direction outside it since the radii vectores of the spiral are on the forward side greater and on the 'backward' side less than OP . Hence the angle OPC cannot be acute since it cannot be less than the angle between OP and

the tangent to the circle at P which is a right angle

It only remains therefore to prove that OPC is not a right angle

If possible let it be a right angle BC will then touch the circle at P

Therefore [Prop 5] it is possible to draw a line OQC meeting the circle through P in Q and BC in C such that

$$CQ \cdot OQ < (\text{arc } PQ) \cdot (\text{arc } DLP) \quad (1)$$

Suppose that OC meets the spiral in R and the first circle in R and produce OP to meet the "first circle" in P'

From (1) it follows *componendo* that


$$\begin{aligned} CO \cdot OQ &< (\text{arc } DLQ) \cdot (\text{arc } DLP) \\ &< (\text{arc } AAK') \cdot (\text{arc } IAK') \\ &< OR \cdot OP \quad [\text{Prop 14}] \end{aligned}$$

But this is impossible because $OQ = OP$, and $OR < OC$

Hence the angle OPC is not a right angle. It was also proved not to be acute

Therefore the angle OPC is obtuse and the angle OPB consequently acute

II If P is on the second or the n th turn the proof is the same except that in the proportion (1) above we have to substitute for the arc DLP an arc equal to $(p + \text{arc } DLP)$ or $(n-1)p + \text{arc } DLP$ where p is the perimeter of the circle DLP through P . Similarly in the later steps p or $(n-1)p$ will be added to each of the arcs DLQ and DLP and c or $(n-1)c$ to each of the arcs AAK' , IAK' where c is the circumference of the first circle IAK'



PROPOSITIONS 18 19

I If $O 1$ be the initial line A the end of the first turn of the spiral, and if the tangent to the spiral at A be drawn the straight line OB drawn from O perpendicular to $O 1$ will meet the said tangent in some point B and OB will be equal to the circumference of the first circle

II If A be the end of the second turn the perpendicular OB will meet the tangent at A in some point B and OB will be equal to 2 (circumference of second circle)

III Generally if A be the end of the n th turn and OB meet the tangent at A in B then $OB = nc$ where c is the circumference of the n th circle

I Let IAK be the first circle. Then meet the backward angle between OA and the tangent at A is acute [Prop 16], the tangent will meet the first circle in a second point C . And the angles $C 1 O$, BOA are together less than two right angles therefore OB will meet $1C$ produced in some point B

Then if c be the circumference of the first circle we have to prove that

$$OB = c$$


If not OB must be either greater or less than c

(1) If possible suppose $OB > c$

Measure along OB a length OD less than OB but greater than c

We have then a circle IAK a chord AC in it less than the diameter and a ratio $AO : OD$ which is greater than the ratio $1O : OR$ or (what is by similar

triangles equal to it) the ratio of $\frac{1}{2}AC$ to the perpendicular from O on AC . Therefore [Prop 7] we can draw a straight line OPF , meeting the circle in P and CA produced in F , such that



$FP \quad PA=AO \quad OD$

Thus alternately, since $AO=PO$,

$$FP \quad PO=PA \quad OD$$

$\angle(\arg PA) \in$

since $(\text{arc } PA) > PA$ and $OD > c$

Componendo

$$FO \quad PO < (c + \text{arc } PA) \quad c$$

$\leq 00 \quad 0_A$

where OF meets the spiral in Q

[Prop 15]

Therefore since $OA = OP$, $FO < OQ$ which is impossible

Hence $OB \geq c$

(2) If possible suppose $OB < c$

Measure OE along OB so that OE is greater than OB but less than c

In this case since the ratio $AO : OE$ is less than the ratio $AO : OB$ (or the ratio of $\frac{1}{2}AC$ to the perpendicular from O on AC), we can [Prop 8] draw a line OPG meeting AC in F' the circle in P' and the tangent at A to the circle in G , such that

$$\Gamma P \quad AG=AO \quad OE$$

Let $OP'G$ cut the spiral in Q

Then we have alternately

$$FP \quad PO = AG \quad OE$$

$$> (\text{arc } AI) \quad c$$

because $AG > (\text{arc } AP)$ and $OE < c$

Therefore

$$\Gamma O \perp O \angle (\text{arc } AKP) \in$$

$\leq 00 \quad 0A$

But this is impossible since $O I = OP$ and $OQ < OF$

Hence $OB \nless c$

Since therefore OB is neither greater nor less than c


$$OB = c$$

II Let AAC be the second circle AC being the tangent to the spiral

II Generally, if P be a point on the n th turn, and the notation be as before, while p represents the circumference of the circle with radius OP ,

$$OT = (n-1)p + \text{arc } KP \text{ (measured 'forward')}$$

I Let P be a point on the first turn of the spiral OA the initial line, PR the tangent at P taken in the 'backward' direction



Then [Prop 16] the angle OPR is acute. Therefore PR meets the circle through P in some point R , and also OT will meet PR produced in some point F .

If now OT is not equal to the arc KRP , it must be either greater or less.

(1) If possible let OT be greater than the arc KRP .

Measure OU along OT less than OT but greater than the arc KRP .

Then, since the ratio $PO : OU$ is greater than the ratio $PO : OT$, or (what is by similar triangles,

equal to it) the ratio of $\frac{1}{2}PR$ to the perpendicular from O on PR , we can draw a line OQF meeting the circle in Q and RP produced in F such that

$$FQ : PQ = PO : OU \quad [\text{Prop 7}]$$

Let OF meet the spiral in Q .

We have then

$$FQ : QO = PQ : OU \\ < (\text{arc } PQ) : (\text{arc } KRP), \text{ by hypothesis}$$

Componendo

$$FO : QO < (\text{arc } KRQ) : (\text{arc } KRP) \\ < OQ : OP \quad [\text{Prop 14}] \\ QO = OP$$

But

Therefore $FO < OQ$ which is impossible.

Hence

$$OT = (\text{arc } KRP)$$

(2) The proof that $OT < (\text{arc } KRP)$ follows the method of Prop 18 I (2), exactly as the above follows that of Prop 18 I (1).

Since then OT is neither greater nor less than the arc KRP it is equal to it.

II If P be on the second turn the same method shows that

$$OT = p + (\text{arc } KRP)$$

and, similarly we have for a point P on the n th turn

$$OT = (n-1)p + (\text{arc } KRP)$$

PROPOSITIONS 21, 22, 23


Given an area bounded by any arc of a spiral and the lines joining the extremities of the arc to the origin it is possible to circumscribe about the area one figure, and to inscribe in it another figure each consisting of similar sectors of circles and such that the circumscribed figure exceeds the inscribed by less than any assigned area.

For let BC be any arc of the spiral O the origin. Draw the circle with centre

O and radius OC , where C is the "forward" end of the arc

Then, by bisecting the angle BOC bisecting the resulting angles and so on continually we shall ultimately arrive at an angle COr cutting off a sector of the circle less than any assigned area. Let COr be this sector

Let the other lines dividing the angle BOC into equal parts meet the spiral in P, Q and let Or meet it in R . With O as centre and radii OB, OP, OQ, OR respectively describe arcs of circles Bp, bBq, pQr, qRc each meeting the adjacent radii as shown in the figure. In each case the arc in the 'forward' direction from each point will fall within and the arc in the backward direction outside the spiral.



We have now a circumscribed figure and an inscribed figure each consisting of similar sectors of circles. To compare their areas we take the successive sectors of each beginning from OC and compare them.

The sector OCr in the circumscribed figure stands alone
And

$$(\text{sector } ORq) = (\text{sector } ORc),$$

$$(\text{sector } OQp) = (\text{sector } OQr)$$

$$(\text{sector } OPb) = (\text{sector } OPq)$$

while the sector OBp in the inscribed figure stands alone

Hence if the equal sectors be taken away the difference between the circumscribed and inscribed figures is equal to the difference between the sectors OCr and OBp and this difference is less than the sector OCr which is itself less than any assigned area.


The proof is exactly the same whatever be the number of angles into which the angle BOC is divided the only difference being that when the arc begins from the origin the smallest sectors OPb, OPq in each figure are equal and there is therefore no inscribed sector standing by itself so that the difference between the circumscribed and inscribed figures is equal to the sector OCr itself.

Thus the proposition is universally true.

CON. Since the area bounded by the spiral is intermediate in magnitude between the circumscribed and inscribed figures it follows that

(1) a figure can be circumscribed to the area such that it exceeds the area by less than any assigned space.

(2) a figure can be inscribed such that the area exceeds it by less than any assigned space.



PROPOSITION 24

The area bounded by the first turn of the spiral and the initial line is equal to one third of the first circle [$= \frac{1}{3}\pi(2\pi a)^2$ where the spiral is $r = a\theta$]

[The same proof shows equally that, if OP be any radius vector in the first turn of the spiral, the area of the portion of the spiral bounded thereby is equal to one third of that sector of the circle drawn with radius OP which is bounded by the initial line and OP , measured in the "forward" direction from the initial line]

Let O be the origin, OA the initial line, A the extremity of the first turn

Draw the "first circle," i.e. the circle with O as centre and OA as radius

Then if C_1 be the area of the first circle R_1 that of the first turn of the spiral bounded by OA , we have to prove that

$$R_1 = \frac{1}{3}C_1$$


For, if not, R_1 must be either greater or less than C_1

I If possible suppose $R_1 < \frac{1}{3}C_1$

We can then circumscribe a figure about R_1 made up of similar sectors of circles such that, if F be the area of this figure

$$F - R_1 < \frac{1}{3}C_1 - R_1$$

whence $F < \frac{1}{3}C_1$



Let OP OQ , be the radii of the circular sectors beginning from the smallest. The radius of the largest is of course OA

The radii then form an ascending arithmetical progression in which the common difference is equal to the least term OP . If n be the number of the sectors, we have [by Prop 10 Cor 1]

$n \cdot OA^2 < 3(OP^2 + OQ^2 + \dots + OA^2)$,
and since the similar sectors are proportional to the squares on their radii it follows that

$$C_1 < 3F,$$

$$F > \frac{1}{3}C_1$$

or

But this is impossible since F was less than $\frac{1}{3}C_1$

Therefore $R_1 < \frac{1}{3}C_1$

II If possible, suppose $R_1 > \frac{1}{3}C_1$

We can then inscribe a figure made up of similar sectors of circles such that, if f be its area

$$R_1 - f < R_1 - \frac{1}{3}C_1$$

whence $f > \frac{1}{3}C_1$

If there are $(n-1)$ sectors their radii as OP OQ , , form an ascending arithmetical progression in which the least term is equal to the common difference and the greatest term as OY , is equal to $(n-1)OP$

Thus [Prop 10 Cor 1]

$$n \cdot OA^2 > 3(OP^2 + OQ^2 + \dots + OY^2)$$

whence

$$C_1 > 3f$$

or

$$f < \frac{1}{3}C_1$$

which is impossible since

$$f > \frac{1}{3}C_1$$

Therefore

$$R_1 = \frac{1}{3}C_1$$

Since then R_1 is neither greater nor less than $\frac{1}{3}C_1$,

$$R_1 = \frac{1}{3}C_1$$

PROPOSITIONS 25, 26, 27

[Prop 25] If A_2 be the end of the second turn of the spiral the area bounded by the second turn and OA_2 is to the area of the "second circle" in the ratio of 7 to 12 being the ratio of $\{r_2 r_1 + \frac{1}{3}(r_2 - r_1)^2\}$ to r_2^2 , where r_1, r_2 are the radii of the "first" and "second" circles respectively

[Prop 26] If BC be any arc measured in the 'forward' direction on any turn of a spiral not being greater than the complete turn, and if a circle be drawn with O as centre and OC as radius meeting OB in B' then

$$\begin{aligned} & (\text{area of spiral between } OB \text{ } OC) : (\text{sector } OB'C) \\ &= \{OC \cdot OB + \frac{1}{3}(OC - OB)^2\} : OC^2 \end{aligned}$$

[Prop 27] If R_1 be the area of the first turn of the spiral bounded by the initial line R_2 the area of the ring added by the second complete turn R_3 that of the ring added by the third turn, and so on then


$$R_3 = 2R_2, R_4 = 3R_2, R_5 = 4R_2, \dots, R_n = (n-1)R_2$$

Also

$$R_2 = 6R_1$$

[Archimedes' proof of Prop 25 is *mutatis mutandis* the same as his proof of the more general Prop 26. The latter will accordingly be given here and applied to Prop 25 as a particular case.]

Let BC be an arc measured in the 'forward' direction on any turn of the spiral CKB the circle drawn with O as centre and OC as radius



Take a circle such that the square of its radius is equal to

$$OC \cdot OB + \frac{1}{3}(OC - OB)^2,$$

and let σ be a sector in it whose central angle is equal to the angle BOC

Thus σ (sector $OB'C$) = $\{OC \cdot OB + \frac{1}{3}(OC - OB)^2\} : OC^2$

and we have therefore to prove that

$$(\text{area of spiral } OBC) = \sigma$$

For if not the area of the spiral OBC (which we will call S) must be either greater or less than σ

I Suppose if possible $S < \sigma$

Circumscribe to the area S a figure made up of similar sectors of circles such that if F be the area of the figure

$$F - S < \sigma - S$$

$$F < \sigma$$

whence

Let the radii of the successive sectors starting from OB be OP, OQ, OC
Produce OP, OQ to meet the circle CKB' ,

If then the lines OB, OP, OQ, OC be n in number, the number of sectors in the circumscribed figure will be $(n-1)$, and the sector $OB'C$ will also be divided into $(n-1)$ equal sectors. Also OB, OP, OQ, OC will form an ascending arithmetical progression of n terms.

Therefore [see Prop 11 and Cor.]

$$(n-1)OC^2 (OP^2 + OQ^2 + \dots + OC^2) < OC \{OC \cdot OB + \frac{1}{2}(OC - OB)^2\} \\ < (\text{sector } OB'C) \cdot \sigma \text{ by hypothesis}$$

Hence, since similar sectors are as the squares of their radii

$$(\text{sector } OB'C) \cdot F < (\text{sector } OB'C) \cdot \sigma$$

so that

$$F > \sigma$$

But this is impossible because

$$F < \sigma$$

Therefore

$$S < \sigma$$

II Suppose if possible $S > \sigma$

Inscribe in the area S a figure made up of similar sectors of circles such that if f be its area,

$$S - f < S - \sigma,$$

whence

$$f > \sigma$$

Suppose OB, OP, OY to be the radii of the successive sectors making up the figure f being $(n-1)$ in number.

We shall have in this case [see Prop 11 and Cor.]

$$(n-1)OC^2 (OB^2 + OP^2 + \dots + OY^2) > OC \{OC \cdot OB + \frac{1}{2}(OC - OB)^2\},$$

whence

$$(\text{sector } OB'C) \cdot f > (\text{sector } OB'C) \cdot \sigma$$

so that

$$f < \sigma$$

But this is impossible because

$$f > \sigma$$

Therefore

$$S > \sigma$$

Since then S is neither greater nor less than σ , it follows that

$$S = \sigma$$

In the particular case where B coincides with A_1 the end of the first turn of the spiral, and C with A_2 the end of the second turn the sector $OB'C$ becomes the complete 'second circle' that, namely with OA_2 (or r_2) as radius

Thus (area of spiral bounded by $O A_1$) ('second circle')

$$= \{r_2 r_1 + \frac{1}{2}(r_2 - r_1)^2\} \cdot r \\ = (2 + \frac{1}{2}) \cdot 4 \quad (\text{since } r_2 = 2r_1) \\ = 7 \cdot 12$$

Again the area of the spiral bounded by OA_2 is equal to $R_1 + R_2$ (i.e. the area bounded by the first turn and $O A_1$ together with the ring added by the second turn). Also the 'second circle' is four times the 'first circle' and therefore equal to $12 R_1$.

Hence

$$(R_1 + R_2) \cdot 12 R_1 = 7 \cdot 12$$

or

$$R_1 + R_2 = 7 R_1$$

Thus

$$R_2 = 6 R_1$$

(1)

Next, for the third turn we have

$$(R_1 + R_2 + R_3) \cdot (\text{third circle}) = \{r_3 r_2 + \frac{1}{2}(r_3 - r_2)^2\} \cdot r_3^2 \\ = (3 \cdot 2 + \frac{1}{2}) \cdot 3^2 \\ = 19 \cdot 27,$$

and

$$(\text{'third circle'}) = 9(\text{'first circle'}) \\ = 27 R_1$$

therefore

$$R_1 + R_2 + R_3 = 19R_1,$$

and, by (1) above it follows that

$$\begin{aligned} R_3 &= 12R_1 \\ &= 2R_2 \end{aligned} \quad (2)$$

and so on

Generally, we have

$$\begin{aligned} (R_1 + R_2 + \dots + R_n) \text{ (nth circle)} &= \{r_n r_{n-1} + \frac{1}{2}(r_n - r_{n-1})^2\} r_n^2, \\ (R_1 + R_2 + \dots + R_{n-1}) \text{ (n-1th circle)} &= \{r_{n-1} r_{n-2} + \frac{1}{2}(r_{n-1} - r_{n-2})^2\} r_{n-1}^2, \\ \text{and} \quad \frac{(R_1 + R_2 + \dots + R_n) \text{ (nth circle)}}{(R_1 + R_2 + \dots + R_{n-1}) \text{ (n-1th circle)}} &= r_n^2 : r_{n-1}^2 \end{aligned}$$

Therefore

$$\begin{aligned} \frac{(R_1 + R_2 + \dots + R_n) \text{ (nth circle)}}{(R_1 + R_2 + \dots + R_{n-1}) \text{ (n-1th circle)}} &= \frac{\{n(n-1) + \frac{1}{2}\} \{(n-1)(n-2) + \frac{1}{2}\}}{\{3n(n-1) + 1\} \{3(n-1)(n-2) + 1\}} \end{aligned}$$

Diminendo,

$$R \text{ (nth circle)} : (R_1 + R_2 + \dots + R_{n-1}) = 6(n-1) : \{3(n-1)(n-2) + 1\} \quad (\alpha)$$

Similarly

$$R_{n-1} \text{ (n-1th circle)} : (R_1 + R_2 + \dots + R_{n-2}) = 6(n-2) : \{3(n-2)(n-3) + 1\},$$

from which we derive

$$\begin{aligned} \frac{R_{n-1} \text{ (n-1th circle)}}{(R_1 + R_2 + \dots + R_{n-2}) \text{ (n-2th circle)}} &= \frac{6(n-2) \{6(n-2) + 3(n-2)(n-3) + 1\}}{6(n-2) \{3(n-1)(n-2) + 1\}} \\ &= \frac{6(n-2) \{6(n-2) + 3(n-2)(n-3) + 1\}}{6(n-2) \{3(n-1)(n-2) + 1\}} \end{aligned} \quad (\beta)$$

Combining (α) and (β) we obtain

$$R : R_{n-1} = (n-1) : (n-2)$$

Thus

$$R_1 : R_2 : R_3 : \dots : R \text{ are in the ratio of the successive numbers } 1 : 2 : 3 : \dots : (n-1)$$

PROPOSITION 28

If O be the origin and BC any arc measured in the 'forward' direction on any turn of the spiral let two circles be drawn (1) with centre O and radius OB meeting OC in C and (2) with centre O and radius OC , meeting OB produced in B . Then if E denote the area bounded by the larger circular arc BC , the line BB and the spiral BC while F denotes the area bounded by the smaller arc BC , the line CC and the spiral BC

$$E : F = \{OB + \frac{1}{2}(OC - OB)\} : \{OB + \frac{1}{2}(OC - OB)\}$$

Let σ denote the area of the lesser sector OBC then the larger sector $OB'C$ is equal to $\sigma + F + E$

Thus [Prop 26]

$$\begin{aligned} (\sigma + F) : (\sigma + F + E) &= \\ &= \{OC : OB + \frac{1}{2}(OC - OB)\}^2 : OC^2 \end{aligned} \quad (1)$$

whence


$$\begin{aligned} E : (\sigma + F) &= \{OC(OC - OB) - \frac{1}{2}(OC - OB)^2\} : \\ &= \{OC : OB + \frac{1}{2}(OC - OB)\}^2 : \\ &= \{OB(OC - OB) + \frac{1}{2}(OC - OB)^2\} : \\ &= \{OC : OB + \frac{1}{2}(OC - OB)\}^2 \end{aligned} \quad (2)$$

Again

$$(\sigma + F + E) : \sigma = OC^2 : OB^2$$

Therefore by the first proportion above *ex aequali*

$$(\sigma + F) : \sigma = \{OC : OB + \frac{1}{2}(OC - OB)\}^2 : OB^2,$$



whence

$$\begin{aligned} (\sigma + F) \quad F = & \{OC \quad OB + \frac{1}{2}(OC - OB)^2\} \\ & \{OB(OC - OB) + \frac{1}{2}(OC - OB)^2\} \end{aligned}$$

Combining this with (2) above we obtain

$$\begin{aligned} E \quad F = & \{OB(OC - OB) + \frac{1}{2}(OC - OB)^2\} \quad \{OB(OC - OB) + \frac{1}{2}(OC - OB)^2\} \\ = & \{OB + \frac{1}{2}(OC - OB)\} \quad \{OB + \frac{1}{2}(OC - OB)\} \end{aligned}$$

ON THE EQUILIBRIUM OF PLANES OR THE CENTRES OF GRAVITY OF PLANES

BOOK ONE

"I POSTULATE the following

1 Equal weights at equal distances are in equilibrium, and equal weights at unequal distances are not in equilibrium but incline towards the weight which is at the greater distance '

2 If, when weights at certain distances are in equilibrium, something be added to one of the weights they are not in equilibrium but incline towards that weight to which the addition was made "

3 ' Similarly if anything be taken away from one of the weights they are not in equilibrium but incline towards the weight from which nothing was taken '

4 "When equal and similar plane figures coincide if applied to one another, their centres of gravity similarly coincide '

5 ' In figures which are unequal but similar, the centres of gravity will be similarly situated By points similarly situated in relation to similar figures I mean points such that, if straight lines be drawn from them to the equal angles they make equal angles with the corresponding sides

6 "If magnitudes at certain distances be in equilibrium, (other) magnitudes equal to them will also be in equilibrium at the same distances

7 "In any figure whose perimeter is concave in (one and) the same direction the centre of gravity must be within the figure

PROPOSITION 1

Weights which balance at equal distances are equal

For, if they are unequal take away from the greater the difference between the two The remainders will then not balance [Post 3], which is absurd

Therefore the weights cannot be unequal

PROPOSITION 2


Unequal weights at equal distances will not balance but will incline towards the greater weight

For take away from the greater the difference between the two The equal remainders will therefore balance [Post 1] Hence if we add the difference again the weights will not balance but incline towards the greater [Post 2]

PROPOSITION 3

Unequal weights will balance at unequal distances, the greater weight being at the less distance

Let A B be two unequal weights (of which A is the greater) balancing about C at distances AC BC respectively



Then shall AC be less than BC
 For, if not, take away from A the weight $(A-B)$ The remainders will then incline towards B [Post 3] But this is impossible for (1) if $AC=CB$ the equal remainders will balance or (2) if $AC>CB$, they will incline towards A at the greater distance [Post 1]

Hence $AC < CB$

Conversely, if the weights balance, and $AC < CB$, then $A > B$

PROPOSITION 4

If two equal weights have not the same centre of gravity, the centre of gravity of both taken together is at the middle point of the line joining their centres of gravity
 [Proved from Prop 3 by *reductio ad absurdum*]

PROPOSITION 5

If three equal magnitudes have their centres of gravity on a straight line at equal distances, the centre of gravity of the system will coincide with that of the middle magnitude

[This follows immediately from Prop 4]

COR 1 The same is true of any odd number of magnitudes if those which are at equal distances from the middle one are equal while the distances between their centres of gravity are equal


COR 2 If there be an even number of magnitudes with their centres of gravity situated at equal distances on one straight line and if the two middle ones be equal while those which are equidistant from them (on each side) are equal respectively, the centre of gravity of the system is the middle point of the line joining the centres of gravity of the two middle ones

PROPOSITIONS 6, 7

Two magnitudes whether commensurable [Prop 6] or incommensurable [Prop 7] balance at distances reciprocally proportional to the magnitudes

I Suppose the magnitudes A, B to be commensurable, and the points A, B to be their centres of gravity Let DE be a straight line so divided at C that

$$A \cdot B = DC \cdot CE$$



We have then to prove that if A be placed at E and B at D C is the centre of gravity of the two taken together

Since A, B are commensurable so are DC, CE Let N be a common measure of DC, CE Make DH, DA each equal to CE and EL (on CE produced) equal to CD Then $EH = CD$, since $DH = CE$ Therefore LH

is bisected at E as HK is bisected at D

Thus LH, HK must each contain N an even number of times

Take a magnitude O such that O is contained as many times in A as Λ is contained in LH , whence

$$\begin{array}{l} \text{But} \quad A \quad O = LH \quad N \\ \quad \quad B \quad A = CE \quad DC \\ \quad \quad \quad = HK \quad LH \end{array}$$

Hence *ex aequali* $B \quad O = HK \quad N$, or O is contained in B as many times as Λ is contained in HK

Thus O is a common measure of $A \quad B$

Divide LH , HK into parts each equal to Λ and $A \quad B$ into parts each equal to O . The parts of A will therefore be equal in number to those of LH , and the parts of B equal in number to those of HK . Place one of the parts of A at the middle point of each of the parts N of LH and one of the parts of B at the middle point of each of the parts N of HK .

Then the centre of gravity of the parts of A placed at equal distances on LH will be at E the middle point of LH [Prop 5 Cor 2] and the centre of gravity of the parts of B placed at equal distances along HK will be at D the middle point of HK .

Thus we may suppose A itself applied at E and B itself applied at D

But the system formed by the parts O of A and B together is a system of equal magnitudes even in number and placed at equal distances along LK . And since $LE = CD$ and $EC = DK$ $LC = CK$ so that C is the middle point of LK . Therefore C is the centre of gravity of the system ranged along LK .

Therefore A acting at E and B acting at D balance about the point C

II Suppose the magnitudes to be incommensurable, and let them be $(1+a)$ and B respectively. Let DE be a line divided at C so that

$$(1+a) \quad B = DC \quad CE$$

Then if $(1+a)$ placed at E and B placed at D do not balance about C $(1+a)$ is either too great to balance B or not great enough.

Suppose if possible that $(1+a)$ is too great to balance B . Take from $(1+a)$ a magnitude a smaller than the deduction which would make the remainder balance B but such that the remainder 1 and the magnitude B are commensurable.

Then since $A \quad B$ are commensurable and

$$A \quad B < DC \quad CE$$

A and B will not balance [Prop 6] but D will be depressed.


But this is impossible since the deduction a is an insufficient deduction from $(1+a)$ to produce equilibrium so that E was still depressed.


Therefore $(1+a)$ is not too great to balance B and similarly it may be proved that B is not too great to balance $(1+a)$.

Hence $(1+a) \quad B$ taken together have their centre of gravity at C .

PROPOSITION 8

If AB be a magnitude whose centre of gravity is C and AD a part of it whose centre of gravity is F then the centre of gravity of the remaining part will be a point G on FC produced such that





$$GC = CF = (AD) = (DE)$$


For, if the centre of gravity of the remainder (DE) be not G , let it be a point H . Then an absurdity follows at once from Props 6, 7

PROPOSITION 9

The centre of gravity of any parallelogram lies on the straight line joining the middle points of opposite sides

Let $ABCD$ be a parallelogram, and let EF join the middle points of the opposite sides AD, BC

If the centre of gravity does not lie on EF suppose it to be H , and draw HA parallel to AD or BC meeting EF in K



Then it is possible by bisecting ED then bisecting the halves and so on continually to arrive at a length EL less than KH . Divide both AE and ED into parts each equal to EL and through the points of division draw parallels to AB or CD

We have then a number of equal and similar parallelograms, and if any one be applied

to any other their centres of gravity coincide [Post 4]. Thus we have an even number of equal magnitudes whose centres of gravity lie at equal distances along a straight line. Hence the centre of gravity of the whole parallelogram will lie on the line joining the centres of gravity of the two middle parallelograms [Prop 5, Cor 2]

But this is impossible for H is outside the middle parallelograms

Therefore the centre of gravity cannot but lie on EF


PROPOSITION 10

The centre of gravity of a parallelogram is the point of intersection of its diagonals

For by the last proposition the centre of gravity lies on each of the lines which bisect opposite sides. Therefore it is at the point of their intersection, and this is also the point of intersection of the diagonals

Alternative proof

Let $ABCD$ be the given parallelogram and BD a diagonal. Then the triangles ABD, CDB are equal and similar, so that [Post 4] if one be applied to the other their centres of gravity will fall one upon the other



Suppose F to be the centre of gravity of the triangle ABD . Let G be the middle point of BD . Join FG and produce it to H , so that $FG = GH$

If we then apply the triangle ABD to the triangle CDB so that AD falls on CB and AB on CD the point F will fall on H

But [by Post 4] F will fall on the centre of gravity of CDB . Therefore H is the centre of gravity of CDB

Hence since F, H are the centres of gravity of the two equal triangles the centre of gravity of the whole parallelogram is at the middle point of FH i.e. at the middle point of BD which is the intersection of the two diagonals

PROPOSITION 11


If abc ABC be two similar triangles and g G two points in them similarly situated with respect to them respectively, then if g be the centre of gravity of the triangle abc G must be the centre of gravity of the triangle ABC

Suppose

$$ab : bc : ca = AB : BC : CA$$

The proposition is proved by an obvious *reductio ad absurdum*. For if G be not the centre of gravity of the triangle ABC suppose H to be its centre of gravity.

Post 5 requires that g H shall be similarly situated with respect to the triangles respectively and this leads at once to the absurdity that the angles HAB , GAB are equal



PROPOSITION 12

Given two similar triangles abc ABC and d D the middle points of bc BC respectively then, if the centre of gravity of abc lie on ad , that of ABC will lie on AD

Let g be the point on ad which is the centre of gravity of abc

Take G on AD such that

$$ad : ag = AD : AG$$

and join gb , gc GB GC

Then since the triangles are similar and bd BD are the halves of bc BC respectively

$$ab : bd = AB : BD$$

and the angles abd ABD are equal

Therefore the triangles abd ABD are similar and

$$\angle bad = \angle BAD$$

Also

$$ba : ad = BA : AD$$

while from above

$$ad : ag = AD : AG$$

Therefore $ba : ag = BA : AG$ while the angles bag BAG are equal

Hence the triangles bag BAG are similar and

$$\angle abg = \angle ABG$$

And since the angles abd ABD are equal it follows that

$$\angle gbd = \angle GBD$$


In exactly the same manner we prove that

$$\angle gac = \angle GAC$$

$$\angle acg = \angle ACG$$

$$\angle gcd = \angle GCD$$

Therefore g G are similarly situated with respect to the triangles respectively whence [Prop 11] G is the centre of gravity of ABC




PROPOSITION 13

In any triangle the centre of gravity lies on the straight line joining any angle to the middle point of the opposite side

Let ABC be a triangle and D the middle point of BC Join AD Then shall the centre of gravity lie on AD

For if possible let this not be the case and let H be the centre of gravity Draw HI parallel to CB meeting AD in I



Then if we bisect DC then bisect the halves, and so on, we shall at length arrive at a length as DE less than HI Divide both BD and DC into lengths each equal to DE and through the points of division draw lines each parallel to DA meeting BA and AC in points as K, L, M and N, P, Q respectively

Join MN, LP, KQ which lines will then be each parallel to BC

We have now a series of parallelograms as FQ, TP, SN and AD bisects opposite sides in each Thus the centre of gravity of each parallelogram lies on AD [Prop 9] and therefore the centre of gravity of the figure made up of them all lies on AD

Let the centre of gravity of all the parallelograms taken together be O Join OH and produce it, also draw CV parallel to DA meeting OH produced in V

Now if n be the number of parts into which AC is divided

$$\begin{aligned}\triangle ADC \text{ (sum of triangles on } AN, NP, \dots) &= AC^2 (AN^2 + NP^2 + \dots) \\ &= n^2 \cdot n \\ &= n \cdot 1 \\ &= AC \cdot AN\end{aligned}$$

Similarly

$$\triangle ABD \text{ (sum of triangles on } AM, ML, \dots) = AB \cdot AM$$

And

$$AC \cdot AN = AB \cdot AM$$

It follows that

$$\begin{aligned}\triangle ABC \text{ (sum of all the small } \triangle s) &= CA \cdot AN \\ &> VO \cdot OH \text{ by parallels}\end{aligned}$$

Suppose OV produced to λ so that

$$\triangle ABC \text{ (sum of small } \triangle s) = \lambda O \cdot OH$$

whence *dividendo*

$$\text{(sum of parallelograms) (sum of small } \triangle s) = \lambda H \cdot HO$$

Since then the centre of gravity of the triangle ABC is at H and the centre of gravity of the part of it made up of the parallelograms is at O it follows from Prop 8 that the centre of gravity of the remaining portion consisting of all the small triangles taken together is at λ

But this is impossible since all the triangles are on one side of the line through λ parallel to AD

Therefore the centre of gravity of the triangle cannot but lie on AD

Alternative proof

Suppose if possible that H not lying on AD , is the centre of gravity of the triangle ABC . Join AH , BH , CH . Let E , F be the middle points of CA , AB respectively and join DE , CF , FD . Let EF meet AD in M .


Draw FA , EL parallel to AH meeting BH , CH in K , L respectively. Join KD , HD , LD , KL . Let KL meet DH in N and join MN .

Since DE is parallel to AB the triangles ABC , EDC are similar.

And, since $CE=EA$ and EL is parallel to AH it follows that $CL=LH$. And $CD=DB$. Therefore BH is parallel to DL .

Thus in the similar and similarly situated triangles ABC , EDC the straight lines AH , BH are respectively parallel to EL , DL and it follows that H , L are similarly situated with respect to the triangles respectively.

But H is by hypothesis the centre of gravity of ABC . Therefore L is the centre of gravity of EDC . [Prop 11]



Similarly the point K is the centre of gravity of the triangle FBD .

And the triangles FBD , EDC are equal so that the centre of gravity of both together is at the middle point of KL i.e. at the point N .

The remainder of the triangle ABC after the triangles FBD , EDC are deducted is the parallelogram $AFDE$ and the centre of gravity of this parallelogram is at M the intersection of its diagonals.

It follows that the centre of gravity of the whole triangle ABC must lie on MN that is MN must pass through H which is impossible (since MN is parallel to AH).

Therefore the centre of gravity of the triangle ABC cannot but lie on AD .

PROPOSITION 14

It follows at once from the last proposition that the centre of gravity of any triangle is at the intersection of the lines drawn from any two angles to the middle points of the opposite sides respectively.

PROPOSITION 15

If AD , BC be the two parallel sides of a trapezium $ABCD$, AD being the smaller, and if AD , BC be bisected at E , F respectively then the centre of gravity of the trapezium is at a point G on EF such that

$$GE:GF = (2BC + AD) : (2AD + BC)$$

Produce BA , CD to meet at O . Then FE produced will also pass through O , since $AE=ED$ and $BF=FC$.

Now the centre of gravity of the triangle OAD will lie on OE and that of the triangle OBC will lie on OF . [Prop 13]

It follows that the centre of gravity of the remainder the trapezium $ABCD$ will also lie on OF . [Prop 8]

Join BD and divide it at L , M into three equal parts. Through L , M draw PQ , RS parallel to BC meeting BA in P , R , FE in W , V and CD in Q , S respectively.

Join DF , BE meeting PQ in H and RS in K respectively

Now, since

$$BL = \frac{1}{3}BD,$$

$$FH = \frac{1}{3}FD$$


Therefore H is the centre of gravity of the triangle DBC

Similarly since $EK = \frac{1}{3}BE$, it follows that K is the centre of gravity of the triangle ADB

Therefore the centre of gravity of the triangles DBC , ADB together, i.e. of the trapezium, lies on the line HK

But it also lies on OF

Therefore if OF HK meet in G G is the centre of gravity of the trapezium



Hence [Props 6, 7]

$$\triangle DBC \quad \triangle ABD = HG \quad GH$$

$$= VG \quad GW$$

$$\triangle DBC \quad \triangle ABD = BC \quad AD$$

$$BC \quad AD = VG \quad GW$$

But

Therefore

It follows that

$$(2BC + AD) \quad (2AD + BC) = (2VG + GW) \quad (2GW + VG) \\ = EG \quad GF$$

QED

ON THE EQUILIBRIUM OF PLANES


BOOK TWO

PROPOSITION I

If P, P' be two parabolic segments and D, E their centres of gravity respectively the centre of gravity of the two segments taken together will be at a point C on DE determined by the relation

$$P \cdot P' = CE \cdot CD$$

In the same straight line with DE measure EH, EL each equal to DC , and DK equal to DH whence it follows at once that $DK = CE$, and also that $AC = CL$



Apply a rectangle MN equal in area to the parabolic segment P to a base equal to KH , and place the rectangle so that KH bisects it, and is parallel to its base

Then D is the centre of gravity of MN since $KD = DH$

Produce the sides of the rectangle which are parallel to KH , and complete the rectangle NO whose base is equal to HL . Then E is the centre of gravity of the rectangle NO

Now

$$\begin{aligned} (MN) \cdot (NO) &= KH \cdot HL \\ &= DH \cdot EH \\ &= CE \cdot CD \\ &= P \cdot P' \end{aligned}$$

But

$$\begin{aligned} (MN) &= P \\ (NO) &= P' \end{aligned}$$

Therefore

Also since C is the middle point of KL C is the centre of gravity of the whole parallelogram made up of the two parallelograms $(MN) (NO)$, which are equal to, and have the same centres of gravity as P, P' respectively

Hence C is the centre of gravity of P, P' taken together

DEFINITION AND LEMMAS PRELIMINARY TO PROPOSITION 2

If in a segment bounded by a straight line and a section of a right angled cone [a parabola] a triangle be inscribed having the same base as the segment and equal height, if again triangles be inscribed in the remaining segments having the same bases as the segments and equal height and if in the remaining segments triangles be inscribed in the same manner let the resulting figure be said to be *inscribed in the recognised manner* in the segment

"And it is plain

(1) that the lines joining the two angles of the figure so inscribed which are nearest to the vertex of the segment and the next pairs of angles in order will be parallel to the base of the segment"

(2) "that the said lines will be bisected by the diameter of the segment, and

(3) ' that they will cut the diameter in the proportions of the successive odd numbers, the number one having reference to [the length adjacent to] the vertex of the segment

"And these properties will have to be proved in their proper places'

PROPOSITION 2

If a figure be "inscribed in the recognised manner' in a parabolic segment, the centre of gravity of the figure so inscribed will lie on the diameter of the segment

For, in the figure of the foregoing lemmas, the centre of gravity of the trapezium $BRRb$ must lie on AO , that of the trapezium $RQqr$ on WX , and so on while the centre of gravity of the triangle PAP lies on AV

Hence the centre of gravity of the whole figure lies on AO

PROPOSITION 3

If BAB , bab' be two similar parabolic segments whose diameters are AO ao respectively and if a figure be inscribed in each segment in the recognised manner' the number of sides in each figure being equal, the centres of gravity of the inscribed figures will divide AO ao in the same ratio'

Suppose $BRQPAP$ $Q R' B$ $brqpap$ $q r' b'$ to be the two figures inscribed "in the recognised manner Join PP' , QQ' RR' meeting AO in L M , N , and pp' qq' rr' meeting ao in l m n

Then [Lemma (3)]

$$\begin{aligned} AL \quad LM \quad MN \quad NO &= 1 \quad 3 \quad 5 \quad 7 \\ &= al \quad lm \quad mn \quad no \end{aligned}$$

so that AO ao are divided in the same proportion

Also, by reversing the proof of Lemma (3) we see that


$$PP' \quad pp' = QQ' \quad qq' = RR' \quad rr' = BB' \quad bb'$$

Since then $RR' \quad BB' = rr' \quad bb'$, and these ratios respectively determine the proportion in which AO , ao are divided by the centres of gravity of the trapezia $BRRB$ $brrb$ [1 15] it follows that the centres of gravity of the trapezia divide NO no in the same ratio

Similarly the centres of gravity of the trapezia $RQQR'$ $rqqr'$ divide MN mn in the same ratio respectively, and so on

¹Archimedes enunciates this proposition as true of *similar* segments but it is equally true of segments which are not similar as the course of the proof will show

Lastly, the centres of gravity of the triangles PAP , pap' divide AL , al respectively in the same ratio



Moreover the corresponding trapezia and triangles are each to each in the same proportion (since their sides and heights are respectively proportional), while AO ao are divided in the same proportion

Therefore the centres of gravity of the complete inscribed figures divide AO ao in the same proportion

PROPOSITION 4

The centre of gravity of any parabolic segment cut off by a straight line lies on the diameter of the segment


Let BAB be a parabolic segment A its vertex and AO its diameter

Then if the centre of gravity of the segment does not lie on AO suppose it to be if possible the point F Draw FE parallel to AO meeting BB in E

Inscribe in the segment the triangle ABB having the same vertex and height as the segment and take an area S such that

$$\triangle ABB \quad S \approx BE \cdot EO$$

We can then inscribe in the segment ' in the recognised manner ' a figure such that the segments of the parabola left over are together less than S '



'For Prop. 20 of the Quadrature of the Parabola proves that if in any segment the triangle with the same base and height be inscribed the triangle is greater than half the segment whence it appears that each time that we increase the number of the sides of the figure inscribed in the recognised manner we take away more than half of the remaining segments

Let the inscribed figure be drawn accordingly, its centre of gravity then lies on AO [Prop 2] Let it be the point H

Join HF and produce it to meet in K the line through B parallel to AO

Then we have

$$\begin{aligned} (\text{inscribed figure}) (\text{remainder of segmt}) &> \triangle ABB \quad S \\ &> BE \quad EO \\ &> HF \quad FH \end{aligned}$$

Suppose L taken on HK produced so that the former ratio is equal to the ratio $LF \quad FH$

Then since H is the centre of gravity of the inscribed figure, and F that of the segment, L must be the centre of gravity of all the segments taken together which form the remainder of the original segment [I 8]

But this is impossible since all these segments lie on one side of the line drawn through L parallel to AO (Cl Post 7)

Hence the centre of gravity of the segment cannot but lie on AO

PROPOSITION 5

If in a parabolic segment a figure be inscribed¹ in the recognised manner, the centre of gravity of the segment is nearer to the vertex of the segment than the centre of gravity of the inscribed figure is

Let BAB be the given segment and AO its diameter First let ABB be the triangle inscribed² in the recognised manner

Divide AO in F so that $AF = 2FO$ F is then the centre of gravity of the triangle ABB

Bisect AB in D AB' in D' respectively and join DD' meeting AO in E Draw $DQ \quad D'Q$ parallel to OA to meet the curve $QD \quad Q'D$ will then be the diameters of the segments whose bases are $AB \quad AB'$, and the centres of gravity of those segments will lie respectively on $QD \quad Q'D$ [Prop 4] Let them be H, H' and join HH' meeting AO in A

Now $QD \quad Q'D$ are equal³ and therefore the segments of which they are the diameters are equal [On Conoids and Spheroids Prop 3]

Also since $QD \quad Q'D$ are parallel and $DE = ED'$ A is the middle point of HH'

Hence the centre of gravity of the equal segments $AQB \quad AQB'$ taken together is K where A lies between E and A And the centre of gravity of the triangle ABB is F

It follows that the centre of gravity of the whole segment BAB lies between A and F and is therefore nearer to the vertex A than F is

Secondly take the five sided figure $BQAQ'B$ inscribed in the recognised manner $QD \quad Q'D$ being as before the diameters of the segments $AQB \quad AQB'$

Then by the first part of this proposition the centre of gravity of the segment AQB (lying of course on QD) is nearer to Q than the centre of gravity of

¹This may either be inferred from Lemma (1) above (since $QQ' \quad DD'$ are both parallel to BB') or from Prop 19 of the *Quadrature of the Parabola* which applies equally to Q or Q'

the triangle AQB is I . Let the centre of gravity of the segment be H and that of the triangle I .

Similarly let H' be the centre of gravity of the segment AQB , and I' that of the triangle AQB' .

It follows that the centre of gravity of the two segments AQB , AQB' taken together is K , the middle point of HH' and that of the two triangles AQB , AQB' is L , the middle point of II' .

If now the centre of gravity of the triangle ABB be F , the centre of gravity of the whole segment BAB (i.e. that of the triangle ABB and the two segments AQB , AQB' taken together) is a point G on KF determined by the proportion

$$(\text{sum of segments } AQB, AQB') : \triangle ABB = KF : KG \quad [I\ 6\ 7]$$

And the centre of gravity of the inscribed figure $BQAQB$ is a point F on LI determined by the proportion

$$(\triangle AQB + \triangle AQB') : \triangle ABB = LF : FL \quad [I\ 6\ 7]$$

$$[\text{Hence} \quad FG : GK > LF : FL]$$

$$\text{or} \quad GK : FG < LF : FL$$

$$\text{and componendo} \quad FK : FG < FL : FL \quad \text{while } FK > FL]$$

Therefore $FG > FL$ or G lies nearer than F to the vertex A .

Using this last result and proceeding in the same way we can prove the proposition for any figure inscribed in the recognised manner.

PROPOSITION 6

Given a segment of a parabola cut off by a straight line it is possible to inscribe in it in the recognised manner a figure such that the distance between the centres of gravity of the segment and of the inscribed figure is less than any assigned length.

Let BAB be the segment AO its diameter C its centre of gravity and ABB the triangle inscribed in the recognised manner.

Let D be the assigned length and S an area such that

$$AG : D = \triangle ABB : S$$



In the segment inscribe in the recognised manner a figure such that the sum of the segments left over is less than S . Let F be the centre of gravity of the inscribed figure.

We shall prove that $FG < D$.

For if not FG must be either equal to or greater than D .

And clearly

$$\begin{aligned} & (\text{inscribed fig}) : (\text{sum of remaining segments}) \\ & > \triangle ABB : S \\ & > AG : D \\ & > AG : FG \text{ by hypothesis (since } FG < D) \end{aligned}$$



Let the first ratio be equal to the ratio $AG : FG$ (where A lies on GA produced), and it follows that K is the centre of gravity of the small segments taken together [I 8]

But this is impossible, since the segments are all on the same side of a line drawn through A parallel to BB'

Hence FG cannot but be less than D

PROPOSITION 7

If there be two similar parabolic segments their centres of gravity divide their diameters in the same ratio

Let $BAB' : bab'$ be the two similar segments AO, ao their diameters and G, g their centres of gravity respectively

Then if G, g do not divide AO, ao respectively in the same ratio, suppose H to be such a point on AO that

$$AH : HO = ag : go$$

and inscribe in the segment BAB' "in the recognised manner" a figure such that if F be its centre of gravity,

$$GF < GH \quad [\text{Prop 6}]$$

Inscribe in the segment bab' "in the recognised manner" a similar




figure then if f be the centre of gravity of this figure

$$ag < af$$

[Prop 5]

And by Prop 3 $af : fo = AF : FO$

But $AF : FO < AH : HO$

$$< ag : go \text{ by hypothesis}$$

Therefore $af : fo < ag : go$ which is impossible

It follows that G, g cannot but divide AO, ao in the same ratio

PROPOSITION 8

If AO be the diameter of a parabolic segment, and G its centre of gravity then

$$AG = \frac{3}{2}GO$$

Let the segment be BAB' . Inscribe the triangle ABB' in the recognised manner and let F be its centre of gravity

Bisect AB, AB' in D, D' , and draw $DQ, D'Q$ parallel to OA to meet the curve so that $QD, Q'D'$ are the diameters of the segments $AQB, A'Q'B'$ respectively

Let H, H' be the centres of gravity of the segments $AQB, A'Q'B'$ respectively. Join QQ', HH' meeting AO in V, A' respectively

A is then the centre of gravity of the two segments $AQB, A'Q'B'$ taken together

Now $AG : GO = QH : HD$ [Prop 7]

whence $AO : OG = QD : HD$

But $AO = 4QD$ [as is easily proved by means of Lemma (3) p. 511]

Therefore $OG = 4HD$

and by subtraction $AG = 4QH$

Also by Lemma (2) QQ is parallel to BB' and therefore to DD' . It follows from Prop 7 that HH' is also parallel to QQ or DD' ,

and hence $QH = VK$

Therefore $AG = 4VK$

and $AV + VK = 3VK$

Measuring VK along VK so that $VL = \frac{1}{3}AV$, we have

$$KG = 3LK \quad (1)$$

Again $AO = 4AV$ [Lemma (3)]

$$= 34L, \text{ since } AV = 3VL$$

whence $AL = \frac{1}{3}AO = OF$ (2)

Now, by I 6, 7

$\triangle ABB'$ (sum of segmts $AQB, AQ'B$) = $KG + GF$,

and $\triangle ABB = 3(\text{sum of segments } AQB, AQ'B')$

[since the segment ABB' is equal to $\frac{1}{3}\triangle ABB'$ (Quadrature of the Parabola Props 17, 24)]

Hence $KG = 3GF$

But $KG = 3LK$ from (1) above

Therefore $LF = LK + KG + GF$
 $= 5GF$

And, from (2),

$$LF = (AO - AL - OF) = \frac{1}{3}AO = OF$$

Therefore $OF = 5GF$,

and $OG = 6GF$

But $AO = 3OF = 15GF$

Therefore, by subtraction

$$AG = 9GF \\ = \frac{3}{2}GO$$

PROPOSITION 9 (LEMMA)

If a, b, c, d be four lines in continued proportion and in descending order of magnitude and if

$$d(a-d) = x \cdot \frac{1}{2}(a-c),$$

and $(2a+4b+6c+3d)(5a+10b+10c+5d) = y(a-c)$,

it is required to prove that



$$x+y = \frac{1}{2}a$$

[The following is the proof given by Archimedes with the only difference that it is set out in algebraical instead of geometrical notation. This is done in the particular case simply in order to make the proof easier to follow. Archimedes exhibits his lines in the figure reproduced in the margin but now that it is possible to use algebraical notation there is no advantage in using the figure and the more cumbersome notation which only obscures the course of the proof. The relation between Archimedes' figure and the letters used below is as follows

$$AB = a, \Gamma B = b, \Delta B = c, EB = d, 2H = x, H\Theta = y, \Delta O = z]$$

We have

$$\frac{a}{b} = \frac{b}{c} = \frac{c}{d} \quad (1)$$



whence

$$\frac{a-b}{b} = \frac{b-c}{c} = \frac{c-d}{d}$$

and therefore

$$\frac{a-b}{b-c} = \frac{b-c}{c-d} = \frac{a}{b} = \frac{b}{c} = \frac{c}{d} \quad (2)$$

Now

$$\frac{2(a+b)}{2c} = \frac{a+b}{c} = \frac{a+b}{b} \cdot \frac{b}{c} = \frac{a-c}{b-c} \cdot \frac{b-c}{c-d} = \frac{a-c}{c-d}$$

And in like manner,

$$\frac{b+c}{d} = \frac{b+c}{c} \cdot \frac{c}{d} = \frac{a-c}{c-d}$$

It follows from the last two relations that

$$\frac{a-c}{c-d} = \frac{2a+3b+c}{2c+d} \quad (3)$$

Suppose z to be so taken that

$$\frac{2a+4b+4c+2d}{2c+d} = \frac{a-c}{z} \quad (4)$$

so that $z < (c-d)$

Therefore

$$\frac{a-c+z}{a-c} = \frac{2a+4b+6c+3d}{2(a+d)+4(b+c)}$$

And by hypothesis

$$\frac{a-c}{y} = \frac{5(a+d)+10(b+c)}{2a+4b+6c+3d},$$

so that

$$\frac{a-c+z}{y} = \frac{5(a+d)+10(b+c)}{2(a+d)+4(b+c)} = \frac{5}{2} \quad (5)$$

Again dividing (3) by (4) crosswise we obtain

$$\frac{z}{c-d} = \frac{2a+3b+c}{2(a+d)+4(b+c)},$$

whence

$$\frac{c-d-z}{c-d} = \frac{b+3c+2d}{2(a+d)+4(b+c)} \quad (6)$$

But by (2)

$$\frac{c-d}{d} = \frac{a-b}{b} = \frac{3(b-c)}{3c} = \frac{2(c-d)}{2d},$$

so that

$$\frac{c-d}{d} = \frac{(a-b)+3(b-c)+2(c-d)}{b+3c+2d} \quad (7)$$

Combining (6) and (7) we have

$$\frac{c-d-z}{d} = \frac{(a-b)+3(b-c)+2(c-d)}{2(a+d)+4(b+c)}$$

whence

$$\frac{c-z}{d} = \frac{3a+5b+3c}{2(a+d)+4(b+c)} \quad (8)$$

And since [by (1)]

$$\frac{c-d}{c+d} = \frac{b-c}{b+c} = \frac{a-b}{a+b}$$

we have

$$\frac{c-d}{a-c} = \frac{c+d}{b+c+a+b}$$

whence

$$\frac{a-d}{a-c} = \frac{a+2b+2c+d}{a+2b+c} = \frac{2(a+d)+4(b+c)}{2(a+c)+4b} \quad (9)$$

Thus

$$\frac{a-d}{\frac{2}{3}(a-c)} = \frac{2(a+d)+4(b+c)}{\frac{2}{3}[2(a+c)+4b]},$$

and therefore, by hypothesis

$$\frac{d}{x} = \frac{2(a+d)+4(b+c)}{\frac{2}{3}[2(a+c)+4b]}$$

But by (8),

$$\frac{c-z}{d} = \frac{3a+6b+3c}{2(a+d)+4(b+c)}$$

and it follows *ex aequali*, that

$$\frac{c-z}{x} = \frac{3(a+c)+6b}{\frac{2}{3}[2(a+c)+4b]} = \frac{5}{3} \cdot \frac{3}{2} = \frac{5}{2}$$

And, by (5)

$$\frac{a-c+z}{y} = \frac{5}{2}$$

Therefore

$$\frac{5}{2} = \frac{a}{x+y}$$

or

$$x+y = \frac{2}{5}a$$

PROPOSITION 10

If $PPBB$ be the portion of a parabola intercepted between two parallel chords PP BB' bisected respectively in N O by the diameter ANO (N being nearer than O to A , the vertex of the segments), and if NO be divided into five equal parts of which LM is the middle one (L being nearer than M to N), then, if G be a point on LM such that

$$LG \cdot GM = BO^2 (2PN + BO) \cdot PN^2 (2BO + PN),$$

G will be the centre of gravity of the area $PPBB$

Take a line ao equal to AO and an on it equal to AN . Let p q be points on the line ao such that

$$ao \cdot aq = aq \cdot an \quad (1)$$

$$ao \cdot an = aq \cdot ap \quad (2)$$

[whence $ao \cdot aq = aq \cdot an = an \cdot ap$ or $ao \cdot aq \cdot an \cdot ap$ are lines in continued proportion and in descending order of magnitude]

Measure along GA a length GI' such that

$$op \cdot ap = OL \cdot GF \quad (3)$$

Then since PN BO are ordinates to ANO

$$BO \cdot PN^2 = AO \cdot AN$$

$$= ao \cdot an$$

$$= ao^2 \cdot aq^2 \text{ by (1),}$$

so that

$$BO \cdot PN = ao \cdot aq \quad (4)$$

and

$$BO^3 \cdot PN^3 = ao^3 \cdot aq^3$$

$$= (ao \cdot aq) (aq \cdot an) (an \cdot ap) \quad (5)$$

$$= ao \cdot ap$$

Thus

$$(\text{segment } BAB) \cdot (\text{segment } PAP')$$



$$= \triangle BAB \cdot \triangle PAP$$

$$= BO^3 \cdot PN^3$$

$$= ao \cdot ap$$

whence

$$(\text{area } PPBB) \cdot (\text{segment } PAP) = op \cdot ap$$



$$=OL \quad GF, \text{ by (3)}$$

$$=\frac{1}{2}OL \quad GF \quad (6)$$

Now

$$BO^2 (2PV + BO) = BO^3$$

$$= (2PN + BO) BO$$

$$= (2aq + ao) ao \text{ by (4)}$$

$$BO^3 = PN^3$$

$$= ao \quad ap, \text{ by (5),}$$

and

$$PV^3 = PV (2BO + PN)$$

$$= PV (2BO + PN)$$

$$= aq (2ao + aq), \text{ by (4)}$$

$$= ap (2an + ap) \text{ by (2)}$$

Hence ex aequali

$$BO^3 (2PV + BO) = PN^3$$

$$(2BO + PN) = (2aq + ao)$$

$$(2an + ap),$$

so that by hypothesis

$$LG \quad GM = (2aq + ao) (2an + ap)$$

Componendo and multiplying the antecedents by 5

$$OL \quad GM = \{5(ao + ap) + 10(aq + an)\} (2an + ap)$$

But

$$OL \quad OM = 5 \quad 2 = \{5(ao + ap) + 10(aq + an)\} \{2(ao + ap) + 4(aq + an)\}$$

It follows that

$$OL \quad OG = \{5(ao + ap) + 10(aq + an)\} (2ao + 4aq + 6an + 3ap)$$

Therefore

$$(2ao + 4aq + 6an + 3ap) \{5(ao + ap) + 10(aq + an)\} = OL \quad ON$$

$$= OG \quad on$$

And

$$ap (ao - ap) = ap \quad op$$

$$= GF \quad OL \text{ by hypothesis}$$

$$= GF \quad \frac{1}{2}on$$

while $ao \quad aq \quad an \quad ap$ are in continued proportion

Therefore by Prop 9

$$GF + OG = OF = \frac{1}{2}ao = \frac{1}{2}OA$$

Thus F is the centre of gravity of the segment BAB

[Prop 8]

Let H be the centre of gravity of the segment PAP so that $AH = \frac{1}{2}AP$

And, since $AF = \frac{1}{2}AO$

we have by subtraction

$$HF = \frac{1}{2}OV$$

But by (6) above

$$(\text{area } PPBB) (\text{segment } PAP) = \frac{1}{2}ON \quad GF$$

$$= HF \quad FG$$

Thus since $F \quad H$ are the centres of gravity of the segments $BAB \quad PAP$ respectively it follows [by 1.6.7] that G is the centre of gravity of the area $PBBB$

THE SAND-RECKONER

THERE are some King Gelon who think that the number of the sand is infinite in multitude and I mean by the sand not only that which exists about Syracuse and the rest of Sicily but also that which is found in every region whether inhabited or uninhabited. Again there are some who without regarding it as infinite yet think that no number has been named which is great enough to exceed its multitude. And it is clear that they who hold this view if they imagined a mass made up of sand in other respects as large as the mass of the earth including in it all the seas and the hollows of the earth filled up to a height equal to that of the highest of the mountains would be many times further still from recognising that any number could be expressed which exceeded the multitude of the sand so taken. But I will try to show you by means of geometrical proofs which you will be able to follow that of the numbers named by me and given in the work which I sent to Zeuxippus some exceed not only the number of the mass of sand equal in magnitude to the earth filled up in the way described but also that of a mass equal in magnitude to the universe. Now you are aware that universe is the name given by most astronomers to the sphere whose centre is the centre of the earth and whose radius is equal to the straight line between the centre of the sun and the centre of the earth. Thus is the common account (*τα γράφοντα*) as you have heard from astronomers. But Aristarchus of Samos brought out a book consisting of some hypotheses in which the premisses lead to the result that the universe is many times greater than that now so called. His hypotheses are that the fixed stars and the sun remain unmoved that the earth revolves about the sun in the circumference of a circle the sun lying in the middle of the orbit, and that the sphere of the fixed stars situated about the same centre as the sun is so great that the circle in which he supposes the earth to revolve bears such a proportion to the distance of the fixed stars as the centre of the sphere bears to its surface. Now it is easy to see that this is impossible for since the centre of the sphere has no magnitude we cannot conceive it to bear any ratio whatever to the surface of the sphere. We must however take Aristarchus to mean this since we conceive the earth to be as it were the centre of the universe the ratio which the earth bears to what we describe as the universe is the same as the ratio which the sphere containing the circle in which he supposes the earth to revolve bears to the sphere of the fixed stars. For he adapts the proofs of his results to a hypothesis of this kind and in particular he appears to suppose the magnitude of the sphere in which he represents the earth as moving to be equal to what we call the universe.

I say then that even if a sphere were made up of the sand as great as Aristarchus supposes the sphere of the fixed stars to be I shall still prove that,

of the numbers named in the *Principles*,¹ some exceed in multitude the number of the sand which is equal in magnitude to the sphere referred to, provided that the following assumptions be made "

1 ' *The perimeter of the earth is about 3 000,000 stadia and not greater*

' It is true that some have tried as you are of course aware to prove that the said perimeter is about 300 000 stadia. But I go further and, putting the magnitude of the earth at ten times the size that my predecessors thought it, I suppose its perimeter to be about 3,000 000 stadia and not greater '

2 ' *The diameter of the earth is greater than the diameter of the moon, and the diameter of the sun is greater than the diameter of the earth*

"In this assumption I follow most of the earlier astronomers "

3 ' *The diameter of the sun is about 30 times the diameter of the moon and not greater*

' It is true that, of the earlier astronomers, Eudoxus declared it to be about nine times as great and Phedias my father twelve times while Aristarchus tried to prove that the diameter of the sun is greater than 18 times but less than 20 times the diameter of the moon. But I go even further than Aristarchus, in order that the truth of my proposition may be established beyond dispute, and I suppose the diameter of the sun to be about 30 times that of the moon and not greater '

4 ' *The diameter of the sun is greater than the side of the chiliagon inscribed in the greatest circle in the (sphere of the) universe*

"I make this assumption because Aristarchus discovered that the sun appeared to be about $\frac{1}{180}$ th part of the circle of the zodiac and I myself tried by a method which I will now describe to find experimentally (*ὁργανικῶς*) the angle subtended by the sun and having its vertex at the eye

[Up to this point the treatise has been literally translated because of the historical interest attaching to the *ipsissima verba* of Archimedes on such a subject. The rest of the work can now be more freely reproduced and, before proceeding to the mathematical contents of it it is only necessary to remark that Archimedes next describes how he arrived at a higher and a lower limit for the angle subtended by the sun. This he did by taking a long rod or ruler fastening on the end of it a small cylinder or disc pointing the rod in the direction of the sun just after its rising (so that it was possible to look directly at it), then putting the cylinder at such a distance that it just concealed and just failed to conceal, the sun and lastly measuring the angles subtended by the cylinder. He explains also the correction which he thought it necessary to make because the eye does not see from one point but from a certain area.]

The result of the experiment was to show that the angle subtended by the diameter of the sun was less than $\frac{1}{180}$ th part and greater than $\frac{1}{200}$ th part of a right angle

To prove that (on this assumption) the diameter of the sun is greater than the side of a chiliagon, or figure with 1000 equal sides inscribed in a great circle of the universe

Suppose the plane of the paper to be the plane passing through the centre of the sun the centre of the earth and the eye at the time when the sun has

¹A lost work of Archimedes

just risen above the horizon. Let the plane cut the earth in the circle EHL and the sun in the circle FAG , the centres of the earth and sun being C & O respectively, and E being the position of the eye.

Further let the plane cut the sphere of the 'universe' (i.e. the sphere whose centre is C and radius CO) in the great circle AOB .

Draw from E two tangents to the circle FAG touching it at P, Q and from C draw two other tangents to the same circle touching it in F, G respectively.


Let CO meet the sections of the earth and sun in H, K respectively and let CF, CG produced meet the great circle AOB in A, B .

Join EO, OF, OG, OP, OQ, AB and let AB meet CO in M .

Now $CO > EO$ since the sun is just above the horizon.

Therefore $\angle PEQ > \angle FCG$

And $\angle PEQ > \frac{1}{16}R$
but $< \frac{1}{16}R$ where R represents a right angle



Thus

$\angle FCG < \frac{1}{16}R$ a fortiori

and the chord AB subtends an arc of the great circle which is less than $\frac{1}{16}$ th of the circumference of that circle i.e.

$AB < (\text{side of 656-sided polygon inscribed in the circle})$

Now the perimeter of any polygon inscribed in the great circle is less than $\frac{1}{4}CO$ [Cf. Measurement of a circle Prop 3]

Therefore $AB, CO < 11 \frac{1}{16} 1148$

and a fortiori $AB < \frac{1}{16}CO$ (a)

Again since $CH = CO$ and AM is perpendicular to CO while OF is perpendicular to CF

$AM = OF$

Therefore $AB = 2AM = (\text{diameter of sun})$

Thus $(\text{diameter of sun}) < \frac{1}{16}CO$ by (a),

and a fortiori $(\text{diameter of earth}) < \frac{1}{16}CO$

[Assumption 2]

Hence $CH + OK < \frac{1}{16}CO$

so that $HK > \frac{1}{16}CO$

or $CO \quad HA < 100 \quad 99$
 And $CO > CF$
 while $HA < EQ$
 Therefore $CF \quad EQ < 100 \quad 99$ (β)
 Now in the right-angled triangles $CFO \quad EQO$ of the sides about the right angles $OF = OQ$, but $EQ < CF$ (since $EO < CO$)
 Therefore $\angle OEQ \quad \angle OCF > CO \quad EO$
 but $< CF \quad EQ$
 Doubling the angles

$$\angle PEQ \quad \angle ACB < CF \quad EQ$$

But $< 100 \quad 99$, by (β) above
 Therefore $\angle PEQ > \frac{1}{100} R$, by hypothesis
 $\angle ACB > \frac{1}{100000} R$
 $> \frac{1}{100} R$

It follows that the arc AB is greater than $\frac{1}{100} \pi$ th of the circumference of the great circle AOB

Hence, *a fortiori*

$AB > (\text{side of chiliagon inscribed in great circle})$

and AB is equal to the diameter of the sun as proved above

The following results can now be proved

(diameter of 'universe') $< 10 \, 000$ (diameter of earth)

and (diameter of "universe") $< 10 \, 000 \, 000 \, 000$ stadia

(1) Suppose, for brevity that d represents the diameter of the 'universe' d that of the sun d that of the earth and d_m that of the moon

By hypothesis $d > 30d_m$ [Assumption 3]

and $d > d_m$, [Assumption 2]

therefore $d < 30d$

Now by the last proposition,

$d > (\text{side of chiliagon inscribed in great circle}),$

so that (perimeter of chiliagon) $< 1000d$
 $< 30 \, 000d$

But the perimeter of any regular polygon with more sides than 6 inscribed in a circle is greater than that of the inscribed regular hexagon and therefore greater than three times the diameter Hence

(perimeter of chiliagon) $> 3d$

It follows that $d < 10,000d$

(2) (Perimeter of earth) $> 3 \, 000 \, 000$ stadia [Assumption 1]

and (perimeter of earth) $> 3d$

Therefore $d < 1 \, 000 \, 000$ stadia

whence $d < 10 \, 000 \, 000 \, 000$ stadia

Assumption 5

Suppose a quantity of sand taken not greater than a poppy seed and suppose that it contains not more than 10 000 grains

The proposition here assumed is of course equivalent to the trigonometrical formula which states that, if $\alpha \quad \beta$ are the circular measures of two angles each less than a right angle of which α is the greater then

$$\frac{\tan \alpha}{\tan \beta} > \frac{\alpha}{\beta} > \frac{\sin \alpha}{\sin \beta}$$

Next suppose the diameter of the poppy seed to be not less than $\frac{1}{10}$ th of a finger breadth

ORDERS AND PERIODS OF NUMBERS

I We have traditional names for numbers up to a myriad (10 000), we can therefore express numbers up to a myriad myriads (100 000,000) Let these numbers be called numbers of the *first order*

Suppose the 100 000 000 to be the unit of the *second order* and let the *second order* consist of the numbers from that unit up to (100 000 000)²

Let this again be the unit of the *third order* of numbers ending with (100 000 000)² and so on until we reach the 100 000 000th *order* of numbers ending with (100 000 000)^{100 000 000} which we will call *P*

II Suppose the numbers from 1 to *P* just described to form the *first period*

Let *P* be the unit of the *first order* of the *second period* and let this consist of the numbers from *P* up to 100 000 000*P*

Let the last number be the unit of the *second order* of the *second period* and let this end with (100 000 000)²*P*

We can go on in this way till we reach the 100 000 000th *order* of the *second period* ending with (100 000 000)^{100 000 000} *P* or *P*²

III Taking *P*² as the unit of the *first order* of the *third period* we proceed in the same way till we reach the 100 000 000th *order* of the *third period* ending with *P*³

IV Taking *P*³ as the unit of the *first order* of the *fourth period*, we continue the same process until we arrive at the 100 000 000th *order* of the 100 000 000th *period* ending with *P*^{100 000 000} This last number is expressed by Archimedes as

a myriad myriad units of the myriad myriad th order of the myriad myriad th period (αἱ μυριάκις μυριάδες περιέχον μυριάκις μυριάδων ἀριθμῶν μυριάς μυριάδες) which is easily seen to be 100 000 000 times the product of (100 000 000)^{99 999 999} and *P*^{99 999 999} i.e. *P*^{100 000 000}

OCTADS

Consider the series of terms in continued proportion of which the first is 1 and the second 10 [i.e. the geometrical progression 1 10¹ 10² 10³ ...] The *first octad* of these terms [i.e. 1 10¹ 10² ... 10⁷] fall accordingly under the *first order* of the *first period* above described the *second octad* [i.e. 10⁸ 10⁹ ... 10¹⁵] under the *second order* of the *first period* the first term of the octad being the unit of the corresponding order in each case Similarly for the *third octad*, and so on We can in the same way place any number of octads

THEOREM

If there be any number of terms of a series in continued proportion say *A*₁ *A*₂ *A*₃ ... *A*_{*m*} *A* *A*_{*m*+1} ... of which *A*₁ = 1 *A*₂ = 10 [so that the series forms the geometrical progression 1 10¹ 10² ... 10^{*m*-1} 10⁻¹ 10^{*m*+1} ...] and if any two terms as *A*_{*m*} *A* be taken and multiplied the product *A*_{*m*} *A* will be a term in the same series and will be as many terms distant from *A* as *A*_{*m*} is distant from *A*₁ also it will be distant from *A*₁ by a number of terms less by one than the sum of the numbers of terms by which *A*_{*m*} and *A* respectively are distant from *A*₁

Take the term which is distant from *A*₁ by the same number of terms as *A*_{*m*}

is distant from A_1 . This number of terms is m (the first and last being both counted). Thus the term to be taken is m terms distant from A_1 , and is therefore the term A_{m+1} .

We have therefore to prove that

$$A_m A_1 = A_{m+1}$$

Now terms equally distant from other terms in the continued proportion are proportional

$$\text{Thus} \quad \frac{A_m}{A_1} = \frac{A_{m+1}}{A_1}$$

$$\text{But} \quad A_m = A_m A_1 \quad \text{since } A_1 = 1$$

$$\text{Therefore} \quad A_{m+1} = A_m A_1 \quad (1)$$

The second result is now obvious since A_m is m terms distant from A_1 , A_1 is n terms distant from A_1 and A_{m+1} is $(m+n-1)$ terms distant from A_1 .

APPLICATION TO THE NUMBER OF THE SAND

By Assumption 5 [p. 523],

(diam. of poppy seed) $< \frac{1}{10}$ (finger breadth),

and since spheres are to one another in the triplicate ratio of their diameters it follows that

(sphere of diam. 1 finger breadth) > 64 000 poppy seeds

> 64 000 $\times 10$ 000

> 640 000 000

> 6 units of second order

order + 40,000,000 of

units of first order sand

(a fortiori) < 10 units of second order of numbers

We now gradually increase the diameter of the supposed sphere multiplying it by 100 each time. Thus remembering that the sphere is thereby multiplied by 100^3 or 1 000 000 the number of grains of sand which would be contained in a sphere with each successive diameter may be arrived at as follows

Diameter of sphere	Corresponding number of grains of sand
(1) 100 finger breadths	< 1 000 000 $\times 10$ units of second order $< (7\text{th term of series}) \times (10\text{th term of series})$ $< 16\text{th term of series}$ [i.e. 10^{16}] $< [10^9 \text{ or } 10$ 000 000 units of the second order
(2) 10 000 finger breadths	< 1 000 000 \times (last number) $< (7\text{th term of series}) \times (16\text{th term})$ $< 22\text{nd term of series}$ [i.e. 10^{22}] $< [10^4 \text{ or } 100$ 000 units of third order
(3) 1 stadium (< 10 000 finger breadths)	< 100 000 units of third order
(4) 100 stadia	< 1 000 000 \times (last number) $< (7\text{th term of series}) \times (22\text{nd term})$ $< 28\text{th term of series}$ [10 ²⁸] $< [10^9 \text{ or } 1$ 000 units of fourth order
(5) 10 000 stadia	< 1 000 000 \times (last number) $< (7\text{th term of series}) \times (28\text{th term})$ $< 34\text{th term of series}$ [10 ³⁴] < 10 units of fifth order

(6) 1 000 000 stadia	<(7th term of series) × (34th term)	[10 ³¹]
	<40th term	
	<[10 ⁷ or] 10 000 000 units of <i>fifth order</i>	
(7) 100 000 000 stadia	<(7th term of series) × (40th term)	[10 ³⁴]
	<46th term	
	<[10 ⁸ or] 100 000 units of <i>sixth order</i>	
(8) 10 000 000 000 stadia	<(7th term of series) × (46th term)	[10 ³⁷]
	<52nd term of series	
	<[10 ⁹ or] 1 000 units of <i>seventh order</i>	

But, by the proposition above [p 523]

(diameter of 'universe') < 10,000 000 000 stadia

Hence *the number of grains of sand which could be contained in a sphere of the size of our "universe" is less than 1,000 units of the seventh order of numbers [or 10⁶¹]*

From this we can prove further that *a sphere of the size attributed by Aristarchus to the sphere of the fixed stars would contain a number of grains of sand less than 10 000 000 units of the eighth order of numbers [or 10⁶⁴⁺⁷ = 10⁶³]*

For by hypothesis

(earth) ('universe') = ("universe") (sphere of fixed stars)

And [p 523]

(diameter of "universe") < 10 000 (diam of earth),

whence

(diam of sphere of fixed stars) < 10 000 (diam of 'universe')

Therefore

(sphere of fixed stars) < (10 000)³ ("universe")

It follows that the number of grains of sand which would be contained in a sphere equal to the sphere of the fixed stars

< (10,000)³ × 1 000 units of *seventh order*

< (13th term of series) × (52nd term of series)

< 64th term of series

< [10⁷ or] 10,000,000 units of *eighth order* of numbers

[i.e. 10⁶³]

CONCLUSION

I conceive that these things, King Gelon will appear incredible to the great majority of people who have not studied mathematics but that to those who are conversant therewith and have given thought to the question of the distances and sizes of the earth, the sun and moon and the whole universe the proof will carry conviction And it was for this reason that I thought the subject would be not inappropriate for your consideration

QUADRATURE OF THE PARABOLA

ARCHIMEDES to DOSITHEUS greeting


' When I heard that Conon, who was my friend in his lifetime was dead but that you were acquainted with Conon and withal versed in geometry while I grieved for the loss not only of a friend but of an admirable mathematician I set myself the task of communicating to you, as I had intended to send to Conon a certain geometrical theorem which had not been investigated before but has now been investigated by me and which I first discovered by means of mechanics and then exhibited by means of geometry Now some of the earlier geometers tried to prove it possible to find a rectilineal area equal to a given circle and a given segment of a circle, and after that they endeavoured to square the area bounded by the section of the whole cone and a straight line assuming lemmas not easily conceded so that it was recognised by most people that the problem was not solved But I am not aware that any one of my predecessors has attempted to square the segment bounded by a straight line and a section of a right-angled cone [a parabola], of which problem I have now discovered the solution For it is here shown that every segment bounded by a straight line and a section of a right angled cone {a parabola} is four thirds of the triangle which has the same base and equal height with the segment and for the demonstration of this property the following lemma is assumed that the excess by which the greater of (two) unequal areas exceeds the less can by being added to itself be made to exceed any given finite area The earlier geometers have also used this lemma for it is by the use of this same lemma that they have shown that circles are to one another in the duplicate ratio of their diameters and that spheres are to one another in the triplicate ratio of their diameters and further that every pyramid is one third part of the prism which has the same base with the pyramid and equal height also, that every cone is one third part of the cylinder having the same base as the cone and equal height they proved by assuming a certain lemma similar to that aforesaid And in the result each of the aforesaid theorems has been accepted no less than those proved without the lemma As therefore my work now published has satisfied the same test as the propositions referred to I have written out the proof and send it to you, first as investigated by means of mechanics and afterwards too as demonstrated by geometry Prefixed are also the elementary propositions in conics which are of service in the proof Farewell

PROPOSITION 1

If from a point on a parabola a straight line be drawn which is either itself the axis or parallel to the axis, as PV , and if QQ' be a chord parallel to the tangent to the parabola at P and meeting PV in V then

$$QV = VQ'$$


Conversely, if $QV = VQ'$, the chord QQ' will be parallel to the tangent at P



PROPOSITION 2

If in a parabola QQ' be a chord parallel to the tangent at P , and if a straight line be drawn through P which is either itself the axis or parallel to the axis, and which meets QQ' in V and the tangent at Q to the parabola in T , then

$$PV = PT$$



PROPOSITION 3

If from a point on a parabola a straight line be drawn which is either itself the axis or parallel to the axis as PV and if from two other points Q, Q' on the parabola straight lines be drawn parallel to the tangent at P and meeting PV in V, V' respectively, then

$$PV \cdot PV' = QV^2 \cdot Q'V'^2$$

'And these propositions are proved in the elements of conics''

PROPOSITION 4

If Qq be the base of any segment of a parabola and P the vertex of the segment, and if the diameter through any other point R meet Qq in O and QP (produced if necessary) in F then

$$QV \cdot VO = QF \cdot FR$$


Draw the ordinate RW to PV meeting QP in K

Then $PV \cdot PW = QV^2 \cdot RW^2$,

whence, by parallels $PQ \cdot PK = PQ^2 \cdot PF^2$

¹i.e. in the treatises on conics by Euclid and Aristaeus

In other words, PQ, PF, PK are in continued proportion therefore



$$\begin{aligned} PQ \cdot PF &= PF \cdot PK \\ &= PQ \pm PF \cdot PF \pm PK \\ &= QF \cdot AF \end{aligned}$$

Hence by parallels, $QV \cdot VO = OF \cdot FR$

PROPOSITION 5

If Qq be the base of any segment of a parabola P the vertex of the segment, and PV its diameter and if the diameter of the parabola through any other point R meet Qq in O and the tangent at Q in E , then

$$QO \cdot Oq = ER \cdot RO$$

Let the diameter through R meet QP in F

Then, by Prop 4,

$$QV \cdot VO = OF \cdot FR$$

Since $QV = Vq$ it follows that

$$QV \cdot qO = OF \cdot OR \quad (1)$$


Also if VP meet the tangent in T ,

$$PT = PV, \text{ and therefore } EF = OF$$

Accordingly doubling the antecedents in (1), we have

$$Qq \cdot qO = OE \cdot OR$$

whence $QO \cdot Oq = ER \cdot RO$



PROPOSITIONS 6 7¹

Suppose a lever AOB placed horizontally and supported at its middle point O . Let a triangle BCD in which the angle C is right or obtuse be suspended from B and O , so that C is attached to O and CD is in the same vertical line with O . Then if P be such an area as when suspended from A will keep the system in equilibrium

$$P = \frac{1}{2} \Delta BCD$$

¹In Prop 6 Archimedes takes the separate case in which the angle BCD of the triangle is a right angle so that C coincides with O in the figure and F with E . He then proves in Prop 7 the same property for the triangle in which BCD is an obtuse angle by treating the triangle as the difference between two right-angled triangles BOD BOC and using the result of Prop 6. I have combined the two propositions in one proof for the sake of brevity. The same remark applies to the propositions following Props 6 7.

Take a point E on OB such that $BE = 2OE$, and draw EFH parallel to CD meeting BC , BD in F , H respectively
 Let G be the middle point of FH

Then G is the centre of gravity of the triangle BCD

Hence, if the angular points B , C be set free and the triangle be suspended by attaching F to E , the triangle will hang in the same position as before, because EFG is a vertical straight line "For this is proved"¹

Therefore as before, there will be equilibrium

$$\begin{aligned} \text{Thus } P \triangle BCD &= OE \cdot AO \\ &= 1 \cdot 3 \\ \text{or } P &= \frac{1}{3} \triangle BCD \end{aligned}$$

PROPOSITIONS 8 9

Suppose a lever AOB placed horizontally and supported at its middle point O . Let a triangle BCD right angled or obtuse-angled at C , be suspended from the points B , E on OB the angular point C being so attached to E that the side CD is in the same vertical line with E . Let Q be an area such that

$$AO \cdot OE = \triangle BCD \cdot Q$$

Then if an area P suspended from A keep the system in equilibrium,
 $P < \triangle BCD$ but $> Q$

Take G the centre of gravity of the triangle BCD and draw GH parallel to DC i.e. vertically meeting BO in H

We may now suppose the triangle BCD suspended from H and since there is equilibrium

$$\triangle BCD \cdot P = AO \cdot OH,$$

$$P < \triangle BCD$$

$$\triangle BCD \cdot Q = AO \cdot OE$$

$$\triangle BCD \cdot Q > \triangle BCD \cdot P,$$

$$P > Q$$

(1)

PROPOSITIONS 10 11

Suppose a lever AOB placed horizontally and supported at O its middle point. Let $CDEF$ be a trapezium which can be so placed that its parallel sides CD , FE are vertical while C is vertically below O , and the other sides CF , DE meet in B . Let EF meet BO in H and let the trapezium be suspended by attaching F to H and C to O . Further suppose Q to be an area such that

$$AO \cdot OH = (\text{trapezium } CDEF) \cdot Q$$

Then if P be the area which when suspended from A , keeps the system in equilibrium,


$$P < Q$$

¹Doubtless in the lost book π σι γγω

The same is true in the particular case where the angles at C, F are right and consequently C, F coincide with O, H respectively

Divide OH in K so that

$$(2CD + FE) (2FE + CD) = HK \cdot KO$$



Draw KG parallel to OD and let G be the middle point of the portion of KG intercepted with in the trapezium. Then G is the centre of gravity of the trapezium [On the equilibrium of planes I 15]

Thus we may suppose the trapezium suspended from A and

the equilibrium will remain undisturbed

Therefore $AO \cdot OK = (\text{trapezium } CDEF) \cdot P$

and, by hypothesis $AO \cdot OH = (\text{trapezium } CDEF) \cdot Q$

Since $OK < OH$, it follows that

$$P < Q$$

PROPOSITIONS 12 13

If the trapezium $CDEF$ be placed as in the last propositions except that CD is vertically below a point L on OB instead of being below O and the trapezium is suspended from L, H suppose that Q, R are areas such that

$$AO \cdot OH = (\text{trapezium } CDEF) \cdot Q$$


$$\text{and } AO \cdot OL = (\text{trapezium } CDEF) \cdot R$$

If then an area P suspended from A keep the system in equilibrium

$$P > R \text{ but } < Q$$

Take the centre of gravity G of the trapezium as in the last propositions and let the line through G parallel to DC meet OB in K

Then we may suppose the trapezium suspended from A and there will still be equilibrium



Therefore $(\text{trapezium } CDEF) \cdot P = AO \cdot OK$

Hence $(\text{trapezium } CDEF) \cdot P > (\text{trapezium } CDEF) \cdot Q$

but $< (\text{trapezium } CDEF) \cdot R$

It follows that $P < Q$ but $> R$

PROPOSITIONS 14 15

Let Qq be the base of any segment of a parabola. Then if two lines be drawn from Q, q each parallel to the axis of the parabola and on the same side of Qq as the segment is either (1) the angles so formed at Q, q are both right angles or (2) one is acute and the other obtuse. In the latter case let the angle at q be the obtuse angle


Divide Qq into any number of equal parts at the points O_1, O_2, \dots, O . Draw

through q, O_1, O_2, \dots, O_n diameters of the parabola meeting the tangent at Q in E, E_1, E_2, \dots, E_n and the parabola itself in q, R_1, R_2, \dots, R_n . Join QR_1, QR_2, \dots, QR_n meeting $qE, O_1E_1, O_2E_2, \dots, O_nE_n$ in F, F_1, F_2, \dots, F_n .

Let the diameters $Eq, E_1O_1, E_2O_2, \dots, E_nO_n$ meet a straight line QOA drawn through Q perpendicular to the diameters in the points O, H_1, H_2, \dots, H_n respectively. (In the particular case where Qq is itself perpendicular to the diameters q will coincide with O, O_1 with H_1 and so on.)

It is required to prove that

- (1) $\triangle E q Q < 3(\text{sum of trapezia } FO_1, F_1O_2, \dots, F_{n-1}O \text{ and } \triangle E O Q),$
- (2) $\triangle E q Q > 3(\text{sum of trapezia } R_1O_2, R_2O_3, \dots, R_{n-1}O \text{ and } \triangle R O Q)$



Suppose AO made equal to OQ and conceive QOA as a lever placed horizontally and supported at O . Suppose the triangle $E q Q$ suspended from OQ in the position drawn and suppose that the trapezium EO_1 in the position drawn is balanced by an area P_1 suspended from A the trapezium E_1O_2 in the position drawn is balanced by the area P_2 suspended from A and so on the triangle $E O Q$ being in like manner balanced by P_{n+1} .

Then $P_1 + P_2 + \dots + P_{n+1}$ will balance the whole triangle $E q Q$ as drawn and therefore $P_1 + P_2 + \dots + P_{n+1} = \frac{1}{3} \triangle E q Q$ [Props 6 7]

Again $AO \cdot OH_1 = QO \cdot OH_1$

$$= Qq \cdot qO_1$$

$$= E_1O_1 \cdot O_1R_1 \text{ [by means of Prop 5]}$$

$$= (\text{trapezium } EO_1) \cdot (\text{trapezium } FO_1),$$

whence [Props 10 11]

$$(FO_1) > P_1$$

Next $AO \ OH_1 = E_1O_1 \ O_1R_1$
 $= (E_1O_2) \ (R_1O_2),$ (α)

while $AO \ OH_2 = E_2O_2 \ O_2R_2$
 $= (E_1O_2) \ (F_1O_2)$ (β)

and, since (α) and (β) are simultaneously true we have by Props 12, 13,
 $(F_1O_2) > P_2 > (R_1O_2)$

Similarly it may be proved that
 $(F_2O_3) > P_3 > (R_2O_3),$

and so on

Lastly [Props 8, 9] $\triangle E \ O_n Q > P_{n+1} > \triangle R_n O_n Q$

By addition, we obtain

$$(1) \quad (FO_1) + (F_1O_2) + \dots + (F_{n-1}O) + \triangle E \ O \ Q > P_1 + P_2 + \dots + P_{n+1}$$

$$\text{or} \quad \triangle E q Q < 3(FO_1 + F_1O_2 + \dots + F_{n-1}O + \triangle E \ O \ Q)$$

$$(2) \quad (R_1O_2) + (R_2O_3) + \dots + (R_{n-1}O) + \triangle R_n O_n Q < P_1 + P_2 + \dots + P_{n+1}$$

$$\text{or} \quad \triangle E q Q > 3(R_1O_2 + R_2O_3 + \dots + R_{n-1}O + \triangle R \ O \ Q)$$

PROPOSITION 16

Suppose Qq to be the base of a parabolic segment, q being not more distant than Q from the vertex of the parabola. Draw through q the straight line qE parallel to the axis of the parabola to meet the tangent at Q in E . It is required to prove that

$$(\text{area of segment}) = \frac{1}{3} \triangle E q Q$$

For, if not, the area of the segment must be either greater or less than $\frac{1}{3} \triangle E q Q$

I Suppose the area of the segment greater than $\frac{1}{3} \triangle E q Q$. Then the excess can if continually added to itself be made to exceed $\triangle E q Q$. And it is possible to find a submultiple of the triangle $E q Q$ less than the said excess of the segment over $\frac{1}{3} \triangle E q Q$.

Let the triangle $F q Q$ be such a submultiple of the triangle $E q Q$. Divide $E q$ into equal parts each equal to qF , and let all the points of division including F be joined to Q meeting the parabola in $R_1 \ R_2 \ \dots \ R_n$ respectively. Through $R_1 \ R_2 \ \dots \ R_n$ draw diameters of the parabola meeting qQ in $O_1 \ O_2 \ \dots \ O$ respectively.

Let O_1R_1 meet QR_2 in F_1 .

Let O_2R_2 meet QR_3 in D_1 and QR_4 in F_2 .


Let O_3R_3 meet QR_5 in D_2 and QR_6 in F_3 , and so on.

We have, by hypothesis,

$$\triangle F q Q < (\text{area of segment}) - \frac{1}{3} \triangle E q Q$$

$$\text{or} \quad (\text{area of segment}) - \triangle F q Q > \frac{1}{3} \triangle E q Q \quad (\alpha)$$

Now, since all the parts of qE as qF and the rest are equal $O_1R_1 = R_1F_1$, $O_2D_1 = D_1R_2 = R_2F_2$ and so on therefore



$$\begin{aligned}\Delta FqQ &= (FO_1 + R_1O_2 + D_1O_3 + \dots) \\ &= (FO_1 + F_1D_1 + F_2D_2 + \dots + F_{n-1}D_{n-1} + \Delta E R Q) \quad (\beta)\end{aligned}$$

But (area of segment) $< (FO_1 + F_1O_2 + \dots + F_{n-1}O + \Delta E O Q)$

Subtracting we have

$$(\text{area of segment}) - \Delta FqQ < (R_1O_2 + R_2O_3 + \dots + R_{n-1}O + \Delta R O Q),$$

whence *a fortiori*, by (α)

$$\frac{1}{2}\Delta EqQ < (R_1O_2 + R_2O_3 + \dots + R_{n-1}O + \Delta R O Q)$$

But this is impossible since [Props 14, 15]

$$\frac{1}{2}\Delta EqQ > (R_1O + R_2O_3 + \dots + R_{n-1}O + \Delta R O Q)$$

Therefore (area of segment) $> \frac{1}{2}\Delta EqQ$

II If possible suppose the area of the segment less than $\frac{1}{2}\Delta EqQ$

Take a submultiple of the triangle EqQ as the triangle FqQ less than the excess of $\frac{1}{2}\Delta EqQ$ over the area of the segment, and make the same construction as before

Since $\Delta FqQ < \frac{1}{2}\Delta EqQ - (\text{area of segment})$,
it follows that

$$\begin{aligned}\Delta FqQ + (\text{area of segment}) &< \frac{1}{2}\Delta EqQ \\ &< (FO_1 + F_1O_2 + \dots + F_{n-1}O + \Delta E O Q) \quad [\text{Props 14, 15}]\end{aligned}$$

Subtracting from each side the area of the segment, we have

$$\begin{aligned}\Delta FqQ &< (\text{sum of spaces } qFR_1, R_1F_1R_2, \dots, E R Q) \\ &< (FO_1 + F_1D_1 + \dots + F_{n-1}D_{n-1} + \Delta E R Q), \text{ a fortiori}\end{aligned}$$

which is impossible because, by (β) above

$$\Delta FqQ = FO_1 + F_1D_1 + \dots + F_{n-1}D_{n-1} + \Delta E R Q$$

Hence (area of segment) $< \frac{1}{2}\Delta EqQ$

Since then the area of the segment is neither less nor greater than $\frac{1}{2}\Delta EqQ$ it is equal to it

PROPOSITION 17

It is now manifest that the area of any segment of a parabola is four thirds of the triangle which has the same base as the segment and equal height

Let Qq be the base of the segment P its vertex
Then PQq is the inscribed triangle with the same base as the segment and equal height

Since P is the vertex of the segment the diameter through P bisects Qq . Let V be the point of bisection

Let VP and qE drawn parallel to it, meet the tangent at Q in T E respectively

Then by parallels

$$qE = 2VT,$$


and $PV = PT$ [Prop 2]

so that $VT = 2PV$

Hence $\Delta EqQ = 4\Delta PQq$

But, by Prop 16 the area of the segment is equal to $\frac{1}{2}\Delta EqQ$

Therefore (area of segment) $= \frac{1}{2}\Delta PQq$



DEF In segments bounded by a straight line and any curve I call the

straight line the *base* and the *height* the greatest perpendicular drawn from the curve to the base of the segment, and the *vertex* the point from which the greatest perpendicular is drawn'

PROPOSITION 18

If Qq be the base of a segment of a parabola and V the middle point of Qq , and if the diameter through V meet the curve in P , then P is the vertex of the segment

For Qq is parallel to the tangent at P [Prop 1] Therefore, of all the perpendiculars which can be drawn from points on the segment to the base Qq , that from P is the greatest Hence, by the definition P is the vertex of the segment

PROPOSITION 19

If Qq be a chord of a parabola bisected in V by the diameter PV and if RM be a diameter bisecting QV in M , and RW be the ordinate from R to PV then

$$PV = \frac{4}{3} RM$$

For, by the property of the parabola,

$$\begin{aligned} PV \cdot PIV &= QV^2 \\ &= 4RW^2 = RW^2, \end{aligned}$$

so that
whence

$$\begin{aligned} PV &= 4PW \\ PV &= \frac{4}{3} RM \end{aligned}$$

PROPOSITION 20

If Qq be the base, and P the vertex, of a parabolic segment, then the triangle PQq is greater than half the segment PQq

For the chord Qq is parallel to the tangent at P , and the triangle PQq is half the parallelogram formed by Qq the tangent at P and the diameters through Q q

Therefore the triangle PQq is greater than half the segment

CON It follows that it is possible to inscribe in the segment a polygon such that the segments left over are together less than any assigned area

PROPOSITION 21

If Qq be the base and P the vertex of any parabolic segment and if R be the vertex of the segment cut off by PQ , then

$$\Delta PQq = 8 \Delta PRQ$$

The diameter through R will bisect the chord PQ , and therefore also QV where PV is the diameter bisecting Qq Let the diameter through R bisect PQ in Y and QV in M Join PM

By Prop 19





$$PV = \frac{4}{3} RM$$

Also

$$PV = 2YM$$

Therefore

$$YM = 2R\frac{1}{3}$$



and
Hence
and

$$\begin{aligned}\Delta PQW &= 2\Delta PRQ \\ \Delta PQV &= 4\Delta PRQ, \\ \Delta PQq &= 8\Delta PRQ\end{aligned}$$

Also, if RW , the ordinate from R to PV , be produced to meet the curve again in r ,
and the same proof shows that

$$\Delta PQq = 8\Delta Prq$$

PROPOSITION 22

If there be a series of areas A, B, C, D , each of which is four times the next in order, and if the largest, A be equal to the triangle PQq inscribed in a parabolic segment PQq and having the same base with it and equal height, then

$$(A+B+C+D+\dots) < (\text{area of segment } PQq)$$


For, since $\Delta PQq = 8\Delta PRQ = 8\Delta Prq$ where R, r are the vertices of the segments cut off by PQ, Pq as in the last proposition,

$$\Delta PQq = 4(\Delta PQR + \Delta Pqr)$$

$$\begin{aligned}\text{Therefore since } \Delta PQq &= A, \\ \Delta PQR + \Delta Pqr &= B\end{aligned}$$

In like manner we prove that the triangles similarly inscribed in the remaining segments are together equal to the area C , and so on

Therefore $A+B+C+D+\dots$ is equal to the area of a certain inscribed polygon, and is therefore less than the area of the segment



PROPOSITION 23

Given a series of areas A, B, C, D, \dots, Z of which A is the greatest and each is equal to four times the next in order, then

$$A+B+C+\dots+Z+\frac{1}{3}Z=\frac{1}{3}A$$

Take areas b, c, d such that

$$b=\frac{1}{3}B,$$

$$c=\frac{1}{3}C,$$

$$d=\frac{1}{3}D \text{ and so on}$$

$$\text{Then since } b=\frac{1}{3}B$$

$$\text{and } B=\frac{1}{3}A$$

$$B+b=\frac{1}{3}A$$

$$\text{Similarly } C+c=\frac{1}{3}B$$

Therefore


$$B+C+D+\dots+Z+b+c+d+\dots+z=\frac{1}{3}(A+B+C+\dots+Y)$$

But

$$b+c+d+\dots+y=\frac{1}{3}(B+C+D+\dots+Y)$$

Therefore, by subtraction

$$\begin{aligned}B+C+D+\dots+Z+z &= \frac{1}{3}A \\ \text{or } A+B+C+\dots+Z+\frac{1}{3}Z &= \frac{1}{3}A\end{aligned}$$



PROPOSITION 24

Every segment bounded by a parabola and a chord PQ is equal to four thirds of the triangle which has the same base as the segment and equal height

Suppose $K = \frac{4}{3} \Delta PQq$

where P is the vertex of the segment, and we have then to prove that the area of the segment is equal to K

For if the segment be not equal to K , it must either be greater or less

I Suppose the area of the segment greater than K

If then we inscribe in the segments cut off by PQ Pq triangles which have the same base and equal height, i.e. triangles with the same vertices R r as those of the segments and if in the remaining segments we inscribe triangles in the same manner, and so on, we shall finally have segments remaining whose sum is less than the area by which the segment PQq exceeds K

Therefore the polygon so formed must be greater than the area K , which is impossible since [Prop 23]

$$A + B + C + \dots + Z < \frac{4}{3}A,$$

where

$$A = \Delta PQq$$

Thus the area of the segment cannot be greater than K

II Suppose if possible that the area of the segment is less than K

If then $\Delta PQq = A$, $B = \frac{1}{4}A$, $C = \frac{1}{4}B$, and so on until we arrive at an area X such that X is less than the difference between K and the segment we have

$$A + B + C + \dots + X + \frac{1}{4}X = \frac{4}{3}A \quad [\text{Prop 23}]$$

$$= K$$

Now since K exceeds $A + B + C + \dots + X$ by an area less than X and the area of the segment by an area greater than X it follows that


$$A + B + C + \dots + X > (\text{the segment})$$

which is impossible by Prop 22 above

Hence the segment is not less than K

Thus, since the segment is neither greater nor less than K ,

$$(\text{area of segment } PQq) = K = \frac{4}{3} \Delta PQq$$



ON FLOATING BODIES

BOOK ONE

POSTULATE 1

"Let it be supposed that a fluid is of such a character that, its parts lying evenly and being continuous, that part which is thrust the less is driven along by that which is thrust the more, and that each of its parts is thrust by the fluid which is above it in a perpendicular direction if the fluid be sunk in any thing and compressed by anything else'

PROPOSITION 1

If a surface be cut by a plane always passing through a certain point, and if the section be always a circumference [of a circle] whose centre is the aforesaid point, the surface is that of a sphere

For if not there will be some two lines drawn from the point to the surface which are not equal

Suppose O to be the fixed point, and A, B to be two points on the surface such that OA, OB are unequal. Let the surface be cut by a plane passing through OA, OB . Then the section is by hypothesis a circle whose centre is O .

Thus $OA = OB$, which is contrary to the assumption. Therefore the surface cannot but be a sphere.

PROPOSITION 2

The surface of any fluid at rest is the surface of a sphere whose centre is the same as that of the earth.


Suppose the surface of the fluid cut by a plane through O , the centre of the earth, in the curve $ABCD$.

$ABCD$ shall be the circumference of a circle.

For if not, some of the lines drawn from O to the curve will be unequal. Take one of them OB such that OB is greater than some of the lines from O to the curve and less than others. Draw a circle with OB as radius. Let it be EBF which will therefore fall partly within and partly without the surface of the fluid.

Draw OGH making with OB an angle equal to the angle EOB and meeting the surface in H and the circle in G . Draw also in the plane an arc of a circle PQR with centre O and within the fluid.

Then the parts of the fluid along PQR are uniform and continuous and the part PQ is compressed by the part between it and AB , while the part QR is compressed by the part between QR and BH .



Therefore the parts along PQ , QR will be unequally compressed, and the part which is compressed the less will be set in motion by that which is compressed the more

Therefore there will not be rest, which is contrary to the hypothesis

Hence the section of the surface will be the circumference of a circle whose centre is O and so will all other sections by planes through O

Therefore the surface is that of a sphere with centre O


PROPOSITION 3

Of solids those which are of equal size, are of equal weight with a fluid will if let down into the fluid be immersed so that they do not project above the surface but do not sink lower

If possible, let a certain solid $ETHG$ of equal weight volume for volume with the fluid remain immersed in it so that part of it $EBCF$, projects above the surface

Draw through O the centre of the earth, and through the solid a plane cutting the surface of the fluid in the circle $ABCD$

Conceive a pyramid with vertex O and base a parallelogram at the surface of the fluid such that it includes the immersed portion of the solid. Let this pyramid be cut by the plane of $ABCD$ in OL , OM . Also let a sphere within the fluid and below GH be described with centre O and let the plane of $ABCD$ cut this sphere in PQR



Conceive also another pyramid in the fluid with vertex O , continuous with the former pyramid and equal and similar to it. Let the pyramid so described be cut in OM , ON by the plane of $ABCD$

Lastly let $STUV$ be a part of the fluid within the second pyramid equal and similar to the part $BGHC$ of the solid, and let SV be at the surface of the fluid

Then the pressures on PQ , QR are unequal that on PQ being the greater. Hence the part at QR will be set in motion by that at PQ , and the fluid will not be at rest, which is contrary to the hypothesis

Therefore the solid will not stand out above the surface

Nor will it sink further, because all the parts of the fluid will be under the same pressure

PROPOSITION 4


A solid lighter than a fluid will if immersed in it not be completely submerged but part of it will project above the surface

In this case after the manner of the previous proposition we assume the solid if possible to be completely submerged and the fluid to be at rest in that position and we conceive (1) a pyramid with its vertex at O the centre of the earth, including the solid (2) another pyramid continuous with the former and equal and similar to it with the same vertex O (3) a portion of the fluid within this latter pyramid equal to the immersed solid in the other pyramid (4) a sphere with centre O whose surface is below the immersed solid and the part of the fluid in the second pyramid corresponding thereto. We suppose a plane to be drawn through the centre O cutting the surface of the fluid in the circle

ABC , the solid in S , the first pyramid in OA , OB , the second pyramid in OB , OC , the portion of the fluid in the second pyramid in K , and the inner sphere in PQR

Then the pressures on the parts of the fluid at PQ , QR are unequal, since S is lighter than K . Hence there will not be rest, which is contrary to the hypothesis


Therefore the solid S cannot, in a condition of rest, be completely submerged



PROPOSITION 5

Any solid lighter than a fluid will, if placed in the fluid be so far immersed that the weight of the solid will be equal to the weight of the fluid displaced

For let the solid be $EGHF$, and let $BGHC$ be the portion of it immersed when the fluid is at rest. As in Prop 3, conceive a pyramid with vertex O including the solid, and another pyramid with the same vertex continuous with the former and equal and similar to it. Suppose a portion of the fluid $STUV$ at the base of the second pyramid to be equal and similar to the immersed portion of the solid and let the construction be the same as in Prop 3



Then since the pressure on the parts of the fluid at PQ , QR must be equal in order that the fluid may be at rest, it follows that the weight of the portion $STUV$ of the fluid must be equal to the weight of the solid $EGHF$. And the former is equal to the weight of the fluid displaced by the immersed portion of the solid $BGHC$


PROPOSITION 6

If a solid lighter than a fluid be forcibly immersed in it the solid will be driven upwards by a force equal to the difference between its weight and the weight of the fluid displaced

For let A be completely immersed in the fluid, and let G represent the weight of A and $(G+H)$ the weight of an equal volume of the fluid. Take a solid D , whose weight is H and add it to A . Then the weight of $(A+D)$ is less than that of an equal volume of the fluid, and if $(A+D)$ is immersed in the fluid, it will project so that its weight will be equal to the weight of the fluid displaced. But its weight is $(G+H)$

Therefore the weight of the fluid displaced is $(G+H)$ and hence the volume of the fluid displaced is the volume of the solid A . There will accordingly be rest with A immersed and D projecting

Thus the weight of D balances the upward force exerted by the fluid on A and therefore the latter force is equal to H which is the difference between the weight of A and the weight of the fluid which A displaces




PROPOSITION 7

A solid heavier than a fluid will, if placed in it, descend to the bottom of the fluid, and the solid will, when weighed in the fluid, be lighter than its true weight by the weight of the fluid displaced

(1) The first part of the proposition is obvious, since the part of the fluid under the solid will be under greater pressure, and therefore the other parts will give way until the solid reaches the bottom

(2) Let A be a solid heavier than the same volume of the fluid and let $(G+H)$ represent its weight, while G represents the weight of the same volume of the fluid



Take a solid B lighter than the same volume of the fluid and such that the weight of B is G , while the weight of the same volume of the fluid is $(G+H)$

Let A and B be now combined into one solid and immersed. Then since $(A+B)$ will be of the same weight as the same volume of fluid both weights being equal to $(G+H)+G$, it follows that $(A+B)$ will remain stationary in the fluid

Therefore the force which causes A by itself to sink must be equal to the upward force exerted by the fluid on B by itself. This latter is equal to the difference between $(G+H)$ and G [Prop 6]. Hence A is depressed by a force equal to H , i.e. its weight in the fluid is H or the difference between $(G+H)$ and G

POSTULATE 2

"Let it be granted that bodies which are forced upwards in a fluid are forced upwards along the perpendicular [to the surface] which passes through their centre of gravity

PROPOSITION 8

If a solid in the form of a segment of a sphere and of a substance lighter than a fluid, be immersed in it so that its base does not touch the surface the solid will rest in such a position that its axis is perpendicular to the surface and if the solid be forced into such a position that its base touches the fluid on one side and be then set free it will not remain in that position but will return to the symmetrical position

PROPOSITION 9

If a solid in the form of a segment of a sphere and of a substance lighter than a fluid be immersed in it so that its base is completely below the surface, the solid will rest in such a position that its axis is perpendicular to the surface

[The proof of this proposition has only survived in a mutilated form. It deals moreover with only one case out of three which are distinguished at the beginning viz that in which the segment is greater than a hemisphere]

Suppose first that the segment is greater than a hemisphere. Let it be cut by a plane through its axis and the centre of the earth and if possible let it be at rest in the position shown in the figure where AB is the intersection of

the plane with the base of the segment, DE its axis, C the centre of the sphere of which the segment is a part, O the centre of the earth

The centre of gravity of the portion of the segment outside the fluid as F , lies on OC produced its axis passing through C


Let G be the centre of gravity of the segment Join FG , and produce it to H so that

$FG : GH = (\text{volume of immersed portion}) : (\text{rest of solid})$

Join OH

Then the weight of the portion of the solid outside the fluid acts along FO and the pressure of the fluid on the immersed portion along OH , while the weight of the immersed portion acts along HO and is by hypothesis less than the pressure of the fluid acting along OH

Hence there will not be equilibrium but the part of the segment towards A will ascend and the part towards B descend, until DE assumes a position perpendicular to the surface of the fluid




ON FLOATING BODIES

BOOK TWO

PROPOSITION 1

If a solid lighter than a fluid be at rest in it the weight of the solid will be to that of the same volume of the fluid as the immersed portion of the solid is to the whole

Let $(A+B)$ be the solid, B the portion immersed in the fluid



Let $(C+D)$ be an equal volume of the fluid, C being equal in volume to A and B to D

Further suppose the line E to represent the weight of the solid $(A+B)$ $(F+G)$ to represent the weight of $(C+D)$, and G that of D

Then

$$\frac{\text{weight of } (A+B)}{E} = \frac{\text{weight of } (C+D)}{(F+G)} \quad (1)$$

And the weight of $(A+B)$ is equal to the weight of a volume B of the fluid [I 5], i.e. to the weight of D

That is to say $E=G$

Hence by (1)

$$\begin{aligned} \frac{\text{weight of } (A+B)}{\text{weight of } (C+D)} &= \frac{G}{F+G} \\ &= \frac{D}{C+D} \\ &= \frac{B}{A+B} \end{aligned}$$

PROPOSITION 2

If a right segment of a paraboloid of revolution whose axis is not greater than $\frac{3}{4}p$ (where p is the principal parameter of the generating parabola) and whose specific gravity is less than that of a fluid be placed in the fluid with its axis inclined to the vertical at any angle but so that the base of the segment does not touch the surface of the fluid the segment of the paraboloid will not remain in that position but will return to the position in which its axis is vertical

Let the axis of the segment of the paraboloid be AN and through AN draw a plane perpendicular to the surface of the fluid. Let the plane intersect the paraboloid in the parabola BAB the base of the segment of the paraboloid in BB and the plane of the surface of the fluid in the chord QQ of the parabola

Then since the axis AN is placed in a position not perpendicular to QQ , BB will not be parallel to QQ

Draw the tangent PT to the parabola which is parallel to QQ and let P be the point of contact

¹The rest of the proof is given in brackets as supplied by Commandinus

[From P draw PV parallel to AN meeting QQ' in V . Then PV will be a diameter of the parabola, and also the axis of the portion of the paraboloid immersed in the fluid]

Let C be the centre of gravity of the paraboloid BAB' , and F that of the portion immersed in the fluid. Join FC and produce it to H so that H is the centre of gravity of the remaining portion of the paraboloid above the surface.


Then, since $AN = \frac{3}{2}AC$,
and $AN > \frac{1}{2}p$,

it follows that $AC > \frac{p}{2}$

Therefore if CP be joined the angle CPT is acute. Hence if CA be drawn perpendicular to PT , A will fall between P and T . And if FL , HM be drawn parallel to CA to meet PT , they will each be perpendicular to the surface of the fluid.

Now the force acting on the immersed portion of the segment of the paraboloid will act upwards along LF , while the weight of the portion outside the fluid will act downwards along HM .

Therefore there will not be equilibrium but the segment will turn so that B will rise and B' will fall, until AN takes the vertical position.]



PROPOSITION 3


If a right segment of a paraboloid of revolution whose axis is not greater than $\frac{1}{2}p$ (where p is the parameter) and whose specific gravity is less than that of a fluid, be placed in the fluid with its axis inclined at any angle to the vertical, but so that its base is entirely submerged, the solid will not remain in that position but will return to the position in which the axis is vertical.

Let the axis of the paraboloid be AN , and through AN draw a plane perpendicular to the surface of the fluid intersecting the paraboloid in the parabola BAB' the base of the segment in BNB' and the plane of the surface of the fluid in the chord QQ' of the parabola.

Then since AN as placed is not perpendicular to the surface of the fluid QQ and BB' will not be parallel.

Draw PT parallel to QQ and touching the parabola at P . Let PT meet NA produced in T . Draw the diameter PV bisecting QQ in V . PV is then the axis of the portion of the paraboloid above the surface of the fluid.

Let C be the centre of gravity of the whole segment of the paraboloid, F that of the portion above the surface. Join FC and produce it to H so that H is the centre of gravity of the immersed portion.



Then since $AC > \frac{p}{2}$ the angle CPT is an acute angle, as in the last proposition.

Hence, if CK be drawn perpendicular to PT , K will fall between P and T . Also, if HM , FL be drawn parallel to CK , they will be perpendicular to the surface of the fluid.


And the force acting on the submerged portion will act upwards along HM , while the weight of the rest will act downwards along LF produced.

Thus the paraboloid will turn until it takes the position in which AN is vertical.

PROPOSITION 4

Given a right segment of a paraboloid of revolution whose axis AN is greater than $\frac{2}{3}p$ (where p is the parameter), and whose specific gravity is less than that of a fluid but bears to it a ratio not less than $(AN - \frac{2}{3}p)^2 : AN^2$ if the segment of the paraboloid be placed in the fluid with its axis at any inclination to the vertical but so that its base does not touch the surface of the fluid, it will not remain in that position but will return to the position in which its axis is vertical.

Let the axis of the segment of the paraboloid be AN , and let a plane be drawn through AN perpendicular to the surface of the fluid and intersecting the segment in the parabola BAB' , the base of the segment in BB' and the surface of the fluid in the chord QQ' of the parabola.



Then AN as placed will not be perpendicular to QQ' .

Draw PT parallel to QQ' and touching the parabola at P . Draw the diameter PV bisecting QQ' in V . Thus PV will be the axis of the submerged portion of the solid.

Let C be the centre of gravity of the whole solid, F that of the immersed portion. Join FC and produce it to H so that H is the centre of gravity of the remaining portion.

$$AN = \frac{2}{3}AC$$

$$AN > \frac{2}{3}p$$

$$AC > \frac{p}{2}$$

Now, since
and

it follows that

Measure CO along CA equal to $\frac{p}{2}$ and OR along OC equal to $\frac{1}{2}AO$.

Then since
and
we have by subtraction

$$AN = \frac{2}{3}AC,$$

$$AR = \frac{1}{3}AO$$

$$NR = \frac{2}{3}OC$$

$$AN - AR = \frac{2}{3}OC$$

$$= \frac{1}{3}p$$

$$AR = (AN - \frac{1}{3}p)$$

$$(AN - \frac{1}{3}p)^2 : AN^2 = AR^2 : AN^2$$

and therefore the ratio of the specific gravity of the solid to that of the fluid is, by the enunciation, not less than the ratio $AR^2 : AN^2$.

But, by Prop. 1 the former ratio is equal to the ratio of the immersed portion to the whole solid i.e. to the ratio $PV^2 : AN^2$ [On Conoids and Spheroids Prop. 24]

Hence

$$PV^2 : AN^2 < AR^2 : AN^2$$

or

It follows that

$$\begin{aligned} PV &< AR \\ PF (= \frac{2}{3}PV) &< \frac{2}{3}AR \\ &< AO \end{aligned}$$

If, therefore OA be drawn from O perpendicular to OP , it will meet PF between P and F

Also if CA be joined the triangle ACO is equal and similar to the triangle formed by the normal the subnormal and the ordinate at P (since $CO = \frac{1}{2}p$ or the subnormal and AO is equal to the ordinate)

Therefore CA is parallel to the normal at P , and therefore perpendicular to the tangent at P and to the surface of the fluid

Hence if parallels to CA be drawn through F H they will be perpendicular to the surface of the fluid and the force acting on the submerged portion of the solid will act upwards along the former while the weight of the other portion will act downwards along the latter

Therefore the solid will not remain in its position but will turn until AN assumes a vertical position


PROPOSITION 5

Given a right segment of a paraboloid of revolution such that its axis AN is greater than $\frac{1}{2}p$ (where p is the parameter), and its specific gravity is less than that of a fluid but in a ratio to it not greater than the ratio $\{1N^2 - (AN - \frac{1}{2}p)^2\} : AN^2$, if the segment be placed in the fluid with its axis inclined at any angle to the vertical, but so that its base is completely submerged, it will not remain in that position but will return to the position in which AN is vertical

Let a plane be drawn through AN as placed perpendicular to the surface of the fluid and cutting the segment of the paraboloid in the parabola BAB' , the base of the segment in BB' and the plane of the surface of the fluid in the chord QQ of the parabola

Draw the tangent PT parallel to QQ and the diameter PI bisecting QQ will accordingly be the axis of the portion of the paraboloid above the surface of the fluid

Let F be the centre of gravity of the portion above the surface C that of the whole solid and produce FC to H the centre of gravity of the immersed portion



As in the last proposition $AC > \frac{p}{2}$ and we measure CO along CA equal to $\frac{p}{2}$ and OR along OC equal to $\frac{1}{2}AO$

Then $1N = \frac{2}{3}AC$ and $AR = \frac{2}{3}AO$
and we derive as before $AR = (AN - \frac{1}{2}p)$

Now by hypothesis

$$\begin{aligned} &(\text{spec gravity of solid}) < (\text{spec gravity of fluid}) \\ &> \{1N^2 - (AN - \frac{1}{2}p)^2\} : AN^2 \\ &> (AN^2 - 4R^2) : AN^2 \end{aligned}$$

Therefore

$$\begin{aligned} &(\text{portion submerged}) < (\text{whole solid}) \\ &> (AN^2 - 4R^2) : AN^2 \end{aligned}$$

and (whole solid) (portion above surface)

Thus
whence
and

$$\begin{aligned} &> AN^2 \quad AR^2 \\ AN^2 \quad PV &> 4N^2 \quad AR^2, \\ PV &< AR \\ PF &< \frac{3}{2}AR \\ &< AO \end{aligned}$$

Therefore if a perpendicular to AC be drawn from O it will meet PF in some point K between P and F


And, since $CO = \frac{1}{2}p$ CK will be perpendicular to PT as in the last proposition

Now the force acting on the submerged portion of the solid will act upwards through H and the weight of the other portion downwards through F in directions parallel in both cases to CA whence the proposition follows

PROPOSITION 6

If a right segment of a paraboloid lighter than a fluid be such that its axis AM is greater than $\frac{3}{2}p$ but $AM - \frac{1}{2}p < 15 \cdot 4$ and if the segment be placed in the fluid with its axis so inclined to the vertical that its base touches the fluid it will never remain in such a position that the base touches the surface in one point only

Suppose the segment of the paraboloid to be placed in the position described and let the plane through the axis AM perpendicular to the surface of the fluid intersect the segment of the paraboloid in the parabolic segment BAB and the plane of the surface of the fluid in BQ



Take C on AM such that $AC = 2CM$ (or so that C is the centre of gravity of the segment of the paraboloid) and measure CK along CA such that

$$AM \cdot CK = 15 \cdot 4$$

Thus $AM \cdot CK > AM \cdot \frac{1}{2}p$ by hypothesis therefore $CK < \frac{1}{2}p$

Measure CO along CA equal to $\frac{1}{2}p$. Also draw KR perpendicular to AC meeting the parabola in R

Draw the tangent PT parallel to BQ and through P draw the diameter PV bisecting BQ in V and meeting KR in I

Then

$$PV \cdot PI_{\text{or}} > AM \cdot AK,$$

for this is proved

And

$$CK = \frac{1}{2}AM = \frac{1}{2}AC$$

whence

$$AK = AC - CK = \frac{1}{2}AC = \frac{1}{2}AM$$

Thus

$$AM = \frac{2}{3}AK$$

Therefore

$$AM = \frac{2}{3}AK$$

It follows that

$$PV_{\text{or}} > \frac{2}{3}PI$$

so that



$$PI_{\text{or}} < 2IV$$

Let F be the centre of gravity of the immersed portion of the paraboloid so that $PF = 2FV$. Produce FC to H the centre of gravity of the portion above the surface

Draw OL perpendicular to PV

PROPOSITION 8

Given a solid in the form of a right segment of a paraboloid of revolution whose axis AM is greater than $\frac{2}{3}p$ but such that $AM - \frac{2}{3}p < 15/4$, and whose specific gravity bears to that of a fluid a ratio less than $(AM - \frac{2}{3}p)^2 / AM^2$ then, if the solid be placed in the fluid so that its base does not touch the fluid and its axis is inclined at an angle to the vertical the solid will not return to the position in which its axis is vertical and will not remain in any position except that in which its axis makes with the surface of the fluid a certain angle to be described



Let am be taken equal to the axis AM , and let c be a point on am such that $ac = 2cm$. Measure co along ca equal to $\frac{1}{2}p$ and or along oc equal to $\frac{1}{2}ao$

Let $X+Y$ be a straight line such that

(spec gr of solid) (spec gr of fluid) $= (Y+Y)^2 am^2$, (α)
and suppose $X=2Y$

$$\begin{aligned} \text{Now } ar &= \frac{2}{3}ao = \frac{2}{3}(\frac{2}{3}am - \frac{1}{2}p) \\ &= am - \frac{1}{2}p \\ &= AM - \frac{1}{2}p \end{aligned}$$

Therefore by hypothesis

$(Y+Y)^2 am^2 < ar^2 am^2$,
whence $(Y+Y) < ar$, and therefore
 $X < ao$

Measure ob along oa equal to X , and draw bd perpendicular to ab and of such length that

$$bd^2 = \frac{1}{2}co \cdot ab \quad (\beta)$$

Join ad

Now let the solid be placed in the fluid with its axis AM inclined at an angle to the vertical. Through AM draw a plane perpendicular to the surface of the fluid and let this plane cut the paraboloid in the parabola BAB' and the plane of the surface of the fluid in the chord QQ' of the parabola.

Draw the tangent PT parallel to QQ' touching at P , and let PV be the diameter bisecting QQ' in V (or the axis of the immersed portion of the solid), and PN the ordinate from P .

Measure AO along AM equal to ao and OC along OM equal to oc and draw OL perpendicular to PV .

I Suppose the angle OTP greater than the angle dab

Thus

$$PN^2 \cdot NT > db^2 \cdot ba^2$$

But

$$PN^2 \cdot NT^2 = p \cdot 4AN$$

$$= co \cdot NT$$

and

$$db^2 \cdot ba^2 = \frac{1}{2}co \cdot ab \text{ by } (\beta)$$

Therefore

$$NT < 2ab$$

or

$$AN < ab$$

whence

$$NO > bo \text{ (since } ao = AO)$$

$$> X$$

Now $(Y+Y)^2 am^2 = (\text{spec gr of solid}) (\text{spec gr of fluid})$
 $= (\text{portion immersed}) (\text{rest of solid})$

so that $\lambda + \gamma = PV$
 But $PL (= NO) > \lambda$
 $> \frac{2}{3}(\lambda + \gamma)$, since $\gamma = 2\gamma$,
 $> \frac{2}{3}PV$,
 or $PV < \frac{3}{2}PL$
 and therefore $PL > 2LV$

Take a point F on PV so that $PF = 2FV$, i.e. so that F is the centre of gravity of the immersed portion of the solid

Also $AC = ac = \frac{2}{3}am = \frac{2}{3}AM$, and therefore C is the centre of gravity of the whole solid

Join FC and produce it to H the centre of gravity of the portion of the solid above the surface

Now since $CO = \frac{1}{3}p$ CL is perpendicular to the surface of the fluid therefore so are the parallels to CL through F and H . But the force on the immersed portion acts upwards through F and that on the rest of the solid downwards through H

Therefore the solid will not rest but turn in the direction of diminishing the angle MTP

II Suppose the angle OTP less than the angle dab . In this case we shall have instead of the above results, the following,

$$AN > ab$$


$$NO < \lambda$$

Also $PV > \frac{3}{2}PL$
 and therefore $PL < 2LV$

Make PF equal to $2FV$ so that F is the centre of gravity of the immersed portion

And, proceeding as before we prove in this case that the solid will turn in the direction of increasing the angle MTP

III When the angle MTP is equal to the angle dab , equalities replace inequalities in the results obtained, and L is itself the centre of gravity of the immersed portion. Thus all the forces act in one straight line the perpendicular CL therefore there is equilibrium and the solid will rest in the position described





PROPOSITION 9

Given a solid in the form of a right segment of a paraboloid of revolution whose axis AM is greater than $\frac{1}{2}p$, but such that $1M - \frac{1}{2}p < 15/4$ and whose specific gravity bears to that of a fluid a ratio greater than $\{AM^2 - (AM - \frac{1}{2}p)^2\} / AM^2$, then, if the solid be placed in the fluid with its axis inclined at an angle to the vertical but so that its base is entirely below the surface the solid will not return to the position in which its axis is vertical and will not remain in any position except that in which its axis makes with the surface of the fluid an angle equal to that described in the last proposition

Take am equal to AM and take c on am such that $ac = 2cm$. Measure co along ca equal to $\frac{1}{2}p$ and ar along ac such that $ar = \frac{2}{3}ao$

Let $\lambda + \gamma$ be such a line that

(spec gr of solid) (spec gr of fluid) = $\{am^2 - (\lambda + Y)^2\} am^2$,
and suppose $\lambda = 2Y$



Let PT be the tangent parallel to QQ (PV the diameter bisecting QQ or the axis of the portion of the paraboloid above the surface), PN the ordinate from P

I Suppose the angle MTP greater than the angle dab . Let AM be cut as before in C and O so that $AC = 2CM$, $OC = \frac{1}{2}p$ and accordingly AM , am are equally divided. Draw OL perpendicular to PV

Then we have, as in the last proposition

$$PN^2 - NT^2 > db^2 \quad ba^2$$

whence

$$co \quad NT > \frac{1}{2}co \quad ab,$$

and therefore

$$AN < ab$$

It follows that

$$NO > bo$$

$$> X$$

Again since the specific gravity of the solid is to that of the fluid as the immersed portion of the solid to the whole

$$AM^2 - (Y + Y)^2 \quad AM^2 = AM^2 - PV^2 \quad AM^2,$$

$$\text{or} \quad (\lambda + Y)^2 \quad AM^2 = PV^2 \quad AM^2$$

$$\text{That is} \quad Y + Y = PV$$

$$\text{And} \quad PL \text{ (or } NO) > Y$$

$$> \frac{1}{2}PV$$

so that


$$PL > 2LV$$

Take Γ on PL so that $PF = 2FV$. Then F is the centre of gravity of the portion of the solid above the surface.

Also C is the centre of gravity of the whole solid. Join FC and produce it to H , the centre of gravity of the immersed portion.

Then since $CO = \frac{1}{2}p$, CL is perpendicular to PT and to the surface of the fluid, and

the force acting on the immersed portion of the solid acts upwards along the



$$\begin{aligned}
 &= CM + CK \quad AC - CK \\
 &= \left(\frac{3}{8} + \frac{1}{8}\right) AM \quad \left(\frac{3}{8} - \frac{1}{8}\right) AM \\
 &= 9 \quad 6 \\
 &= MA \quad AC
 \end{aligned} \tag{\beta}$$

Thus C is seen to be on the parabola BA_1B_2 by the converse of Prop 4 of the *Quadrature of the Parabola*]

Also if a perpendicular to AM be drawn from O it will meet the parabola BA_1B_2 in two points, as $Q_2 \ P$. Let $Q_1Q \ Q_2D$ be drawn through Q_2 parallel to AM meeting the parabolas $BAB_1 \ BA_1M$ respectively in $Q_1 \ Q_3$ and BM in D , and let $P_1P_2P_3$ be the corresponding parallel to AM through P_2 . Let the tangents to the outer parabola at $P_1 \ Q_1$ meet MA produced in $T_1 \ U$ respectively

Then, since the three parabolic segments are similar and similarly situated with their bases in the same straight line and having one common extremity, and since $Q_1Q_2 \ Q_2Q_3$ is a diameter common to all three segments, it follows that

$$Q_1Q_2 \ Q_2Q_3 = (B_2B_1 \ B_1B) \ (BM \ MB_2)$$

$$\text{Now} \quad B_2B_1 \ B_1B = MM_2 \ BM \quad \begin{array}{l} \text{(dividing by 2)} \\ \text{by means of } (\beta) \text{ above} \end{array}$$

$$= 2 \quad 5$$

$$\text{And} \quad BM \ MB_2 = BM \ (2BM_2 - BM)$$

$$= 5 \quad (6 - 5)$$

$$= 5 \quad 1 \quad \text{by means of } (\beta),$$

$$\begin{array}{l} \text{It follows that} \\ \text{or} \end{array} \quad \begin{array}{l} Q_1Q \quad Q_2Q_3 = 2 \quad 1 \\ Q_1Q_2 = 2Q_2Q_3 \} \\ \text{Similarly} \quad P_1P_2 = 2P_2P_3 \} \\ \text{Also, since} \quad MR = \frac{3}{2}CO = \frac{3}{2}p \\ \quad AR = AM - MR \\ \quad \quad = AM - \frac{3}{2}p \end{array}$$

(ENUNCIATION)

If the segment of the paraboloid be placed in the fluid with its base entirely above the surface then

(I) if

$$(\text{spec gr of solid}) \ (\text{spec gr of fluid}) < AR^2 \ AM^2$$

$$[< (4M - \frac{3}{2}p)^2 \ AM^2]$$

the solid will rest in the position in which its axis AM is vertical,

(II) if

$$(\text{spec gr of solid}) \ (\text{spec gr of fluid}) < AR^2 \ AM^2$$

$$\text{but} > Q_1Q_2^2 \ AM^2$$

the solid will not rest with its base touching the surface of the fluid in one point only but in such a position that its base does not touch the surface at any point and its axis makes with the surface an angle greater than U

(III a) if

$$(\text{spec gr of solid}) \ (\text{spec gr of fluid}) = Q_1Q_2^2 \ AM^2,$$

the solid will rest and remain in the position in which the base touches the surface of the fluid at one point only and the axis makes with the surface an angle equal to U

(III b) if

$$(\text{spec gr of solid}) \ (\text{spec gr of fluid}) = P_1P_2^2 \ AM^2,$$

the solid will rest with its base touching the surface of the fluid at one point only and with its axis inclined to the surface at an angle equal to T_1 ,

(IV) if

$$(\text{spec gr of solid}) (\text{spec gr of fluid}) > P_1 P_2 \cdot AM^2 \\ \text{but} < Q_1 Q_2 \cdot 1M^2,$$

the solid will rest and remain in a position with its base more submerged,

(V) if

$$(\text{spec gr of solid}) (\text{spec gr of fluid}) < P_1 P_2 \cdot AM^2,$$

the solid will rest in a position in which its axis is inclined to the surface of the fluid at an angle less than T_1 , but so that the base does not even touch the surface at one point

(PROOF)


(I) Since $AM > \frac{1}{2}p$ and

$$(\text{spec gr of solid}) (\text{spec gr of fluid}) < (AM - \frac{1}{2}p)^2 \cdot AM^2$$

it follows by Prop 4 that the solid will be in stable equilibrium with its axis vertical

(II) In this case

$$(\text{spec gr of solid}) (\text{spec gr of fluid}) < AR^2 \cdot AM^2 \\ \text{but} > Q_1 Q_2 \cdot AM^2$$



Suppose the ratio of the specific gravities to be equal to $P^2 \cdot AM^2$, so that
 $\{ < AR^2 \text{ but } > Q_1 Q_2 \cdot$

Place PV' between the two parabolas BAB_1 BP_2Q_2M equal to l and parallel to AM and let PV' meet the intermediate parabola in F

Then by the same proof as before we obtain

$$P'F' = 2FV$$

Let PT the tangent at P' to the outer parabola meet MA in T , and let $P'N'$ be the ordinate at P

Join BV and produce it to meet the outer parabola in Q . Let OQ P_2 meet PV' in I

Now, since in two similar and similarly situated parabolic segments with bases BM BB_1 in the same straight line BV BQ are drawn making the same angle with the bases,

$$BV : BQ = BM : BB_1$$

$$= 1 : 2$$

so that

$$BV = \frac{1}{2} BQ$$

Suppose the segment of the paraboloid placed in the fluid as described with its axis inclined at an angle to the vertical, and with its base touching the surface at one point B only. Let the solid be cut by a plane through the axis and perpendicular to the surface of the fluid and let the plane intersect the solid in the parabolic segment BAB and the plane of the surface of the fluid in BO .

Take the points C, O on AM as before described. Draw the tangent parallel to BQ touching the parabola in P and meeting AM in T and let PI be the diameter bisecting BQ (i.e. the axis of the immersed portion of the solid).

Then $\rho = \frac{4M}{V} = \frac{\text{(spec gr of solid)} \times \text{(spec gr of fluid)} \times \text{(portion immersed)}}{\text{(whole solid)}}$
 $= \frac{PV}{AV} = \frac{P}{A}$

whence

Thus the segments in the two figures namely BPQ BPQ are equal and similar

Therefore

$$\angle PTV = \angle PTN$$

Also

$$AT = AT \quad AN = AN, PV = P\Delta$$

Now, in the first figure $PI < 2IV$

Therefore if OL be perpendicular to PV in the second figure

$$PL < 2LV$$

Take F on LI so that $PF = 2PV$, i.e. so that F is the centre of gravity of the immersed portion of the solid. And C is the centre of gravity of the whole solid. Join FC and produce it to H the centre of gravity of the portion above the surface.

Now since $CO = \frac{1}{2}p$ CL is perpendicular to the tangent at P and to the surface of the fluid. Thus as before we prove that the solid will not rest with B touching the surface but will turn in the direction of increasing the angle PTV .

Hence in the position of rest the axis AM must make with the surface of the fluid an angle greater than the angle ψ which the tangent at Q_1 makes with AM .

(III a) In this case

$$(\text{spec gr of solid}) (\text{pec gr of fluid}) = Q_1 Q_2 \cdot A W^2$$

Let the segment of the paraboloid be placed in the fluid so that its base nowhere touches the surface of the fluid and its axis is inclined at an angle to the vertical

Let the plane through AU perpendicular to the surface of the fluid cut the paraboloid in the parabola BAB and the plane of the surface of the fluid QQ . Let PT be the tangent parallel to QQ . II the diameter bisecting PV the ordinate at P .

(III b) In the case where


$$(\text{spec gr of solid}) \quad (\text{spec gr of fluid}) = P_1 P_2, \quad AM^2$$

we can prove in the same way that if the solid be placed in the fluid so that its axis is inclined to the vertical and its base does not anywhere touch the surface of the fluid the solid will take up and rest in the position in which one point only of the base touches the surface and the axis is inclined to it at an angle equal to T_1 (in the figure on p. 552)

(IV) In this case

$$(\text{spec gr of solid}) (\text{spec gr of fluid}) > P_1 P_2^2 A^2 / \mu^2$$
$$\text{but } \angle O_1 O_2 P = \angle AM^2$$

Suppose the ratio to be equal to l AM^2 , so that l is greater than P_1P_2 but less than O_1O_2 .



Place PV' between the parabolas BP_1Q_1 , BP_2Q_2 , so that PV is equal to l and parallel to AM , and let PV' meet the intermediate parabola in F and OQ_1P_1 in I .

Join BI and produce it to meet the outer parabola in g

Then as before $BV = Vq$ and accordingly the tangent PT at P' is parallel to Bq . Let PN be the ordinate of P

1 Now let the segment be placed in the fluid *first*, with its axis so inclined to the vertical that its base does not

anywhere touch the surface of the fluid

Let the plane through AM perpendicular to the surface of the fluid cut the paraboloid in the parabola BAB' and the plane of the surface of the fluid in

QQ Let PT be the tangent parallel to QQ PV the diameter bisecting QQ Divide AM at C O as before and draw OL perpendicular to PV

Then as before we have PL
 $\text{and } l \text{ and } PV$

Thus the segments BP and QPQ of the paraboloid are equal in volume and it follows that the angle between QQ and BB is less than the angle B_1B_0 .

Therefore


$$\angle PTN < \angle PTN$$

and hence $AN > AN$

so that $\Lambda O > \Lambda O$

$$1e \quad PL > PI$$

$\geq P F$ a fortiori



Thus F again lies between P and L and as before the paraboloid will turn in the direction of diminishing the angle PTN , i.e. so that the base will be more submerged.


(V) In this case

$$(\text{spec gr of solid}) \quad (\text{spec gr of fluid}) < P_1 P_2^2 \quad A M^2$$

If then the ratio is equal to l^2/AM^2 , $l < P_1P_3$. Place $P'V'$ between the parabolas BP_1Q_1 and BP_3Q_3 , equal in length to l and parallel to AM . Let $P'V'$ meet the intermediate parabola in F' and OP_3 in I .

Join BV' and produce it to meet the outer parabola in q . Then, as before, $BV' = Vq$ and the tangent PT' is parallel to Bq .

1 Let the paraboloid be so placed in the fluid that its base touches the surface at one point only



Let the plane through A V perpendicular to the surface of the fluid cut the paraboloid in the parabolic section BAB and the plane of the surface of the fluid in BQ

Making the usual construction, we find

$$PV = l = P' V'$$

and the segments BPQ BP_1q are equal and similar

Therefore

$$\angle PTN = \angle P T N'$$

and

$$AN=AN \quad NO=NO$$

Therefore

$$PL \approx P'I$$

whence it follows that

$$PL < 2LV$$


Thus F , the centre of gravity of the immersed portion of the solid, lies between L and V while CL is perpendicular to the surface of the fluid.

Producing FC to H the centre of gravity of the portion above the surface we prove as usual that there will not be rest but the solid will turn in the direction of increasing the angle PTN so that the base will not anywhere touch the surface

BOOK OF LEMMAS

PROPOSITION 1

If two circles touch at A , and if BD EF be parallel diameters in them ADF is a straight line



Let O & C be the centres of the circles and let OC be joined and produced to A . Draw DH parallel to AO meeting OF in H .

Then since $OH = CD = CA$,
and $OF = OA$

we have by subtraction $HF=CO=DH$

Therefore $\angle HDF = \angle NFD$

Thus both the triangles CAD HDF are isosceles and the third angles ACD DHF in each are equal. Therefore the equal angles in each are equal to one another and

$$\angle ADC = \angle DFH$$

Add to each the angle CDF and it follows that

$$\begin{aligned}\angle ADC + \angle CDF &= \angle CDF + \angle DFH \\ &= (\text{two right angles})\end{aligned}$$


Hence ADF is a straight line

The same proof applies if the circles touch externally

PROPOSITION 2

Let AB be the diameter of a semicircle and let the tangents to it at B and at any other point D on it meet in T . If now DE be drawn perpendicular to AB and $\sphericalangle AT$ DE meet in F .

$DF = FE$



Produce AD to meet BT produced in H . Then the angle ADB in the semicircle is right, therefore the angle BDH is also right. And TB , TD are equal. Therefore T is the centre of the semicircle on BH as diameter, which passes through D .

Hence

$HT \approx TB$

And since $DE \parallel B$ are parallel it follows that $DF = FE$

PROPOSITION 3

Let P be any point on a segment of a circle whose base is AB , and let PN be perpendicular to AB . Take D on AB so that $AN = ND$. If now PQ be an arc equal to the arc PA , and BQ be joined,

BQ BD shall be equal

Join PA , PQ , PD , DQ .

Then since the arcs PA , PQ are equal,

$$PA = PQ$$

But, since $AN = ND$, and the angles at N are right,

$$PA = PD$$

$$PQ = PD$$

Therefore $\angle PQD = \angle PDQ$

and Now, since A , P , Q , B are concyclic,

$$\angle PAD + \angle PQB = (\text{two right angles}),$$

$$\angle PDA + \angle PQB = (\text{two right angles})$$


$$= \angle PDA + \angle PDB$$

$$\text{Therefore } \angle PQB = \angle PDB,$$

and since the parts, the angles PQD , PDQ are equal,

$$\angle BQD = \angle BDQ$$

$$\text{and } BQ = BD$$



PROPOSITION 4

If AB be the diameter of a semicircle and N any point on AB , and if semicircles be described within the first semicircle and having AN , NB as diameters respectively, the figure included between the circumference of the three semicircles is what Archimedes called an $\alpha\rho\beta\eta\lambda\omicron\varsigma$ ¹ and its area is equal to the circle on PN as diameter, where PN is perpendicular to AB and meets the original semicircle in P .

For

$$AB^2 = AN^2 + NB^2 + 2AN \cdot NB$$


$$= AN^2 + NB^2 + 2PN^2$$

But circles (or semicircles) are to one another as the squares of their radii (or diameters)

Hence

$$\begin{aligned} (\text{semicircle on } AB) &= (\text{sum of} \\ &\text{semicircles on } AN, NB) \\ &+ 2(\text{semicircle on } PN) \end{aligned}$$

That is the circle on PN as diameter is equal to the difference between the semicircle on AB and the sum of the semicircles on AN , NB , i.e. is equal to the area of the $\alpha\rho\beta\eta\lambda\omicron\varsigma$.



PROPOSITION 5

Let AB be the diameter of a semicircle C any point on AB and CD perpendicular to it and let semicircles be described within the first semicircle and having AC , CB as diameters. Then if two circles be drawn touching CD on different sides and each touching two of the semicircles the circles so drawn will be equal.

Let one of the circles touch CD at E the semicircle on AB in F , and the semicircle on AC in G .

¹ $\alpha\rho\beta\eta\lambda\omicron\varsigma$ is literally a shoemaker's knife


Draw the diameter EH of the circle which will accordingly be perpendicular to CD and therefore parallel to AB

Join FH , HA , and FE , EB Then, by Prop 1, FHA , FEB are both straight lines, since EH , AB are parallel

For the same reason AGE CGH are straight lines

Let AF produced meet CD in D , and let AE produced meet the outer semi circle in I Join BI , ID

Then since the angles AFB ACD are right, the straight lines ID , AB are such that the perpendiculars on each from the extremity of the other meet in the point E Therefore by the properties of triangles AE is perpendicular to the line joining B to D



But AE is perpendicular to BI

Therefore BID is a straight line

Now, since the angles at G I are right CH is parallel to BD

Therefore

$$\begin{aligned} AB \cdot BC &= AD \cdot DH \\ &= AC \cdot HE \end{aligned}$$

so that

$$AC \cdot CB = AB \cdot HE$$

In like manner if d is the diameter of the other circle, we can prove that

$$AC \cdot CB = AB \cdot d$$

Therefore $d = HE$ and the circles are equal

PROPOSITION 6

Let AB the diameter of a semicircle be divided at C so that $AC = \frac{2}{3}CB$ [or in any ratio] Describe semicircles within the first semicircle and on AC CB as diameters and suppose a circle drawn touching all three semicircles If GH be the diameter of this circle to find the relation between GH and AB

Let GH be that diameter of the circle which is parallel to AB , and let the circle touch the semicircles on AB , AC CB in D E , F respectively


Join AG , GD and BH HD Then, by Prop 1, AGD , BHD are straight lines

For a like reason AEH , BFG are straight lines as also are CEG CFH

Let AD meet the semicircle on AC in I and let BD meet the semicircle on CB in K Join CI CK meeting AE , BF respectively in L M , and let GL , HM produced meet AB in N P respectively

$$(\text{arc } AD) + (\text{arc } CB) = (\text{arc } AC) + (\text{arc } DB)$$

Let the chords intersect at O , and draw the diameter EF parallel to AB intersecting CD in H . EF will thus bisect CD at right angles in H , and



$$(\text{arc } ED) = (\text{arc } EC)$$

Also EDF ECF are semicircles while

$$(\text{arc } ED) = (\text{arc } EA) + (\text{arc } AD)$$

Therefore

$$(\text{sum of arcs } CF, EA, AD) = (\text{arc of a semicircle})$$

And the arcs AE, BF are equal

Therefore


$$(\text{arc } CB) + (\text{arc } AD) = (\text{arc of a semicircle})$$

Hence the remainder of the circumference the sum of the arcs AC, DB is also equal to a semicircle and the proposition is proved.

PROPOSITION 10

Suppose that TA, TB are two tangents to a circle, while TC cuts it. Let BD be the chord through B parallel to TC and let AD meet TC in E . Then, if EH be drawn perpendicular to BD it will bisect it in H .

Let AB meet TC in F , and join BE .



Now the angle TAB is equal to the angle in the alternate segment i.e.

$$\angle TAB = \angle ADB$$

$$= \angle AET, \text{ by parallels}$$

Hence the triangles EAT, AFT have one angle equal and another (at T) common. They are therefore similar and

$$FT \cdot AT = AT \cdot ET$$

$$\text{Therefore } ET \cdot TT = TA^2 = TB^2$$

It follows that the triangles EBT, BFT are similar

$$\angle TEB = \angle TBF$$

$$= \angle TAB$$

But the angle TEB is equal to the angle EBD and the angle TAB was proved equal to the angle EDB .

Therefore

$$\angle EDB = \angle EBD$$

And the angles at H are right angles

It follows that

$$BH = HD$$

PROPOSITION 11

If two chords AB, CD in a circle intersect at right angles in a point O not being the centre then

$$AO^2 + BO^2 + CO^2 + DO^2 = (\text{diameter})^2$$


Draw the diameter CE and join AC, CB, AD, BE .

Then the angle CAO is equal to the angle CEB in the same segment and the angles AOC , EBC are right, therefore the triangles AOC , EBC are similar and

$$\angle ACO = \angle ECB$$

It follows that the subtended arcs and therefore the chords AD , BE are equal

$$\begin{aligned}\text{Thus } (AO^2 + DO^2) + (BO^2 + CO) &= AD^2 + BC^2 \\ &= BE^2 + BC^2 \\ &= CE^2\end{aligned}$$



PROPOSITION 12

If AB be the diameter of a semicircle and TP , TQ the tangents to it from any point T and if AQ , BP be joined meeting in R , then TR is perpendicular to AB

Let TR produced meet AB in M and join PA , QB

Since the angle APB is right,

$$\begin{aligned}\angle PAB + \angle PBA &= (\text{a right angle}) \\ &= \angle AQB\end{aligned}$$

Add to each side the angle RBQ and

$$\angle PAB + \angle QBA = (\text{exterior}) \angle PRQ$$

But

$\angle TPR = \angle PAB$, and $\angle TQR = \angle QBA$,
in the alternate segments

therefore $\angle TPR + \angle TQR = \angle PRQ$

It follows from this that

$$TP = TQ = TR$$

[For if PT be produced to O so that $TO = TQ$ we have

$$\angle TOQ = \angle TQO$$

And by hypothesis,

$$\angle PRQ = \angle TPR + \angle TQR$$

By addition

$$\angle POQ + \angle PRQ = \angle TPR + \angle OQR$$

It follows that, in the quadrilateral $OPRQ$ the opposite angles are together equal to two right angles. Therefore a circle will go round $OPQR$ and T is its centre because $TP = TO = TQ$. Therefore $TR = TP$]

Thus

$$\angle TRP = \angle TPR = \angle PAM$$

Adding to each the angle PRM ,


$$\begin{aligned}\angle PAM + \angle PRM &= \angle TRP + \angle PRM \\ &= (\text{two right angles})\end{aligned}$$

Therefore

$$\angle APR + \angle AMR = (\text{two right angles}),$$

whence

$$\angle AMR = (\text{a right angle})$$



PROPOSITION 13


If a diameter AB of a circle meet any chord CD not a diameter in E and if AM , BN be drawn perpendicular to CD then

$$CN = DM$$

Let O be the centre of the circle and OH perpendicular to CD . Join BM and produce HO to meet BM in K

Then

$$CH = HD$$



And by parallels, since

$$BO = OA,$$

$$BK = KM$$

Therefore

$$NH = HM$$


Accordingly

$$CN = DM$$

PROPOSITION 14

Let ACB be a semicircle on AB as diameter and let AD, BE be equal lengths measured along AB from A, B respectively. On AD, BE as diameters describe semicircles on the side towards C , and on DE as diameter a semicircle on the opposite side. Let the perpendicular to AB through O , the centre of the first semicircle, meet the opposite semicircles in C, F respectively.

Then shall the area of the figure bounded by the circumferences of all the semicircles be equal to the area of the circle on CF as diameter.



By Eucl II 10 since ED is bisected at O and produced to A

$$EA^2 + AD^2 = 2(EO^2 + OA^2)$$

and $CF = OA + OE = EA$

Therefore

$$AB^2 + DE^2 = 4(EO^2 + OA^2) = 2(CF^2 + 4D^2)$$

But circles (and therefore semicircles) are to one another as the squares on their radii (or diameters)

Therefore

$$\begin{aligned} & (\text{sum of semicircles on } AB, DE) \\ &= (\text{circle on } CF) + (\text{sum of semicircles on } AD, BE) \end{aligned}$$

Therefore

$$(\text{area of 'sahnun'}) = (\text{area of circle on } CF \text{ as diam})$$

PROPOSITION 15

Let AB be the diameter of a circle. AC a side of an inscribed regular pentagon. D the middle point of the arc AC . Join CD and produce it to meet BA produced in E . Join AC, DB meeting in F and draw FM perpendicular to AB . Then

$$EM = (\text{radius of circle})$$

Let O be the centre of the circle and join DA, DM, DO, CB

Now

$$\angle ABC = \frac{1}{2}(\text{right angle}),$$

and

$$\angle ABD = \angle DBC = \frac{1}{2}(\text{right angle}),$$

whence

$$\angle AOD = \frac{1}{2}(\text{right angle})$$

Further the triangles FCB, FMB are equal in all respects

Therefore in the triangles DCB, DMB the sides CB, MB being equal and BD common while the angles CBD, MBD are equal

$$\angle BCD = \angle BMD = \frac{1}{2}(\text{right angle})$$

But

$$\begin{aligned} \angle BCD + \angle BAD &= (\text{two right angles}) \\ &= \angle BAD + \angle DAE \\ &= \angle BMD + \angle DMA \end{aligned}$$

so that

$$\angle DAE = \angle BCD$$

and

$$\angle BAD = \angle AMD$$


Therefore

$$AD = MD$$

Now, in the triangle DMO ,

$$\angle MOD = \frac{2}{3}(\text{right angle}),$$

$$\angle DMO = \frac{1}{3}(\text{right angle})$$



Therefore
whence
Again

$$\angle ODM = \frac{2}{3}(\text{right angle}) = \angle AOD,$$

$$OM = MD$$

$$\angle EDA = (\text{supplement of } ADC)$$

$$= \angle CBA$$

$$= \frac{2}{3}(\text{right angle})$$

$$= \angle ODM$$

Therefore, in the triangles EDA , ODM ,

$$\angle EDA = \angle ODM,$$

$$\angle EAD = \angle OMD,$$

and the sides AD MD are equal

Hence the triangles are equal in all respects, and

$$EA = MO$$

Therefore

$$EM = AO$$

Moreover $DE = DO$ and it follows that since DE is equal to the side of an inscribed hexagon and DC is the side of an inscribed decagon EC is divided at D in extreme and mean ratio [i.e. $EC \cdot ED = ED \cdot DC$], "and this is proved in the book of the *Elements*" [Eucl. *III* 9 "If the side of the hexagon and the side of the decagon inscribed in the same circle be put together, the whole straight line is divided in extreme and mean ratio and the greater segment is the side of the hexagon"]

THE METHOD TREATING OF MECHANICAL PROBLEMS

"Archimedes to Eratosthenes greeting

"I sent you on a former occasion some of the theorems discovered by me merely writing out the enunciations and inviting you to discover the proofs which at the moment I did not give. The enunciations of the theorems which I sent are as follows

1 'If in a right prism with a parallelogrammic base a cylinder be inscribed which has its bases in the opposite parallelograms,¹ and its sides [i.e. four generators] on the remaining planes (faces) of the prism, and if through the centre of the circle which is the base of the cylinder and (through) one side of the square in the plane opposite to it a plane be drawn the plane so drawn will cut off from the cylinder a segment which is bounded by two planes and the surface of the cylinder, one of the two planes being the plane which has been drawn and the other the plane in which the base of the cylinder is and the surface being that which is between the said planes and the segment cut off from the cylinder is one sixth part of the whole prism.

2 "If in a cube a cylinder be inscribed which has its bases in the opposite parallelograms² and touches with its surface the remaining four planes (faces), and if there also be inscribed in the same cube another cylinder which has its bases in other parallelograms and touches with its surface the remaining four planes (faces), then the figure bounded by the surfaces of the cylinders which is within both cylinders is two-thirds of the whole cube.

Now these theorems differ in character from those communicated before for we compared the figures then in question conoids and spheroids and segments of them in respect to size, with figures of cones and cylinders but none of those figures have yet been found to be equal to a solid figure bounded by planes whereas each of the present figures bounded by two planes and surfaces of cylinders is found to be equal to one of the solid figures which are bounded by planes. The proofs then of these theorems I have written in this book and now send to you. Seeing moreover in you as I say an earnest student, a man of considerable eminence in philosophy and an admirer [of mathematical inquiry] I thought fit to write out for you and explain in detail in the same book the peculiarity of a certain method by which it will be possible for you to get a start to enable you to investigate some of the problems in mathematics by means of mechanics. This procedure is I am persuaded no less useful even for the proof of the theorems themselves for certain things first became clear to me by a mechanical method although they had to be demonstrated by geom-

¹The parallelograms are apparently squares

²i.e. squares

etry afterwards because their investigation by the said method did not furnish an actual demonstration. But it is of course easier, when we have previously acquired, by the method some knowledge of the questions to supply the proof than it is to find it without any previous knowledge. This is a reason why in the case of the theorems the proof of which Eudoxus was the first to discover, namely that the cone is a third part of the cylinder and the pyramid of the prism, having the same base and equal height, we should give no small share of the credit to Democritus who was the first to make the assertion with regard to the said figure though he did not prove it. I am myself in the position of having first made the discovery of the theorem now to be published [by the method indicated], and I deem it necessary to expound the method partly because I have already spoken of it and I do not want to be thought to have uttered vain words but equally because I am persuaded that it will be of no little service to mathematics for I apprehend that some either of my contemporaries or of my successors, will by means of the method when once established be able to discover other theorems in addition which have not yet occurred to me.

'First then I will set out the very first theorem which became known to me by means of mechanics, namely that

Any segment of a section of a right-angled cone (i.e. a parabola) is four thirds of the triangle which has the same base and equal height

and after this I will give each of the other theorems investigated by the same method. Then at the end of the book, I will give the geometrical [proofs of the propositions]

[I premise the following propositions which I shall use in the course of the work.]

1 'If from [one magnitude] another magnitude be subtracted which has not the same centre of gravity the centre of gravity of the remainder is found by] producing [the straight line joining the centres of gravity of the whole magnitude and of the subtracted part in the direction of the centre of gravity of the whole] and cutting off from it a length which has to the distance between the said centres of gravity the ratio which the weight of the subtracted magnitude has to the weight of the remainder. [*On the Equilibrium of Planes* 1 8]

2 'If the centres of gravity of any number of magnitudes whatever be on the same straight line the centre of gravity of the magnitude made up of all of them will be on the same straight line' [*Cf Ibid* 1 5]

3 The centre of gravity of any straight line is the point of bisection of the straight line [*Cf Ibid* 1 4]

4 'The centre of gravity of any triangle is the point in which the straight lines drawn from the angular points of the triangle to the middle points of the (opposite) sides cut one another' [*Ibid* 1 13 14]

5 The centre of gravity of any parallelogram is the point in which the diagonals meet' [*Ibid* 1 10]

6 The centre of gravity of a circle is the point which is also the centre [of the circle]'

7 'The centre of gravity of any cylinder is the point of bisection of the axis'

8 The centre of gravity of any cone is [the point which divides its axis so that] the portion [adjacent to the vertex is] triple [of the portion adjacent to the base]

[All the e propositions have already been] proved¹ [Besides these I require also the following proposition which is easily proved]

If in two series of magnitudes those of the first series are in order proportional to those of the second series and further, "the magnitudes [of the first series] either all or some of them, are in any ratio whatever [to those of a third series] and if the magnitudes of the second series are in the same ratio to the corresponding magnitudes [of a fourth series], then the sum of the magnitudes of the first series has to the sum of the selected magnitudes of the third series the same ratio which the sum of the magnitudes of the second series has to the sum of the (correspondingly) selected magnitudes of the fourth series" [*On Conoids and Spheroids* Prop 1]

PROPOSITION I

Let ABC be a segment of a parabola bounded by the straight line AC and the parabola ABC , and let D be the middle point of AC Draw the straight line DBE parallel to the axis of the parabola and join AB BC

Then shall the segment ABC be $\frac{2}{3}$ of the triangle ABC

From A draw AKF parallel to DE and let the tangent to the parabola at C meet DBE in E and AKF in F Produce CB to meet AF in K and again produce CK to H making KH equal to CK

Consider CH as the bar of a balance K being its middle point

Let MO be any straight line parallel to ED and let it meet CF CK AC in M N , O and the curve in P

Now since CE is a tangent to the parabola and CD the semi ordinate

$$EB=BD,$$

"for this is proved in the Elements [of Conics]"²

Since FA MO are parallel to ED it follows that

$$FK=KA \quad MN=NO$$

Now by the property of the parabola proved in a lemma'

$$MO \cdot OP = CA \cdot AO \quad [\text{Cf Quadrature of Parabola Prop 5}]$$

$$= CK \cdot KN$$

$$[\text{Eucl vi 2}]$$

$$= HK \cdot KN$$


Take a straight line TG equal to OP and place it with its centre of gravity at H , so that $TH=HG$ then since N is the centre of gravity of the straight line MO and

$$MO \cdot TG = HK \cdot KN$$

it follows that TG at H and MO at N will be in equilibrium about K [*On the Equilibrium of Planes* 1 6 7]

¹The problem of finding the centre of gravity of a cone is not solved in any extant work of Archimedes

² i.e. the works on conics by Aristaeus and Euclid



Similarly, for all other straight lines parallel to DE and meeting the arc of the parabola, (1) the portion intercepted between FC , AC with its middle point on AC and (2) a length equal to the intercept between the curve and AC placed with its centre of gravity at H will be in equilibrium about K .

Therefore K is the centre of gravity of the whole system consisting (1) of all the straight lines as MO intercepted between FC , AC and placed as they actually are in the figure and (2) of all the straight lines placed at H equal to the straight lines as PO intercepted between the curve and AC .

And, since the triangle CFA is made up of all the parallel lines like MO , and the segment CBA is made up of all the straight lines like PO within the curve

it follows that the triangle placed where it is in the figure, is in equilibrium about K with the segment CBA placed with its centre of gravity at H .

Divide AC at W so that $CA = 3AW$, then W is the centre of gravity of the triangle ACF , "for this is proved in the books on equilibrium (*ἐν τοῖς ἰσορροπικοῖς*)"

[Cf. *On the Equilibrium of Planes* 1.15]

Therefore $\triangle ACF$ (segment ABC) $= HA \cdot AW$
 $= 3 \cdot 1$

Therefore segment $ABC = \frac{1}{3} \triangle ACF$

But $\triangle ACF = 4 \triangle ABC$

Therefore segment $ABC = \frac{1}{3} \triangle ABC$

"Now the fact here stated is not actually demonstrated by the argument used but that argument has given a sort of indication that the conclusion is true. Seeing then that the theorem is not demonstrated but at the same time suspecting that the conclusion is true, we shall have recourse to the geometrical demonstration which I myself discovered and have already published."

PROPOSITION 2

We can investigate by the same method the propositions that

(1) Any sphere is (in respect of solid content) four times the cone with base equal to a great circle of the sphere and height equal to its radius and

(2) the cylinder with base equal to a great circle of the sphere and height equal to the diameter is $1\frac{1}{2}$ times the sphere

(1) Let $ABCD$ be a great circle of a sphere and AC , BD diameters at right angles to one another

Let a circle be drawn about BD as diameter and in a plane perpendicular to AC and on this circle as base let a cone be described with A as vertex. Let the surface of this cone be produced and then cut by a plane through C parallel to its base. The section will be a circle on EF as diameter. On this circle as base let a cylinder be erected with height and axis AC , and produce CA to H , making AH equal to CA .

Let CH be regarded as the bar of a balance A being its middle point

Draw any straight line MN in the plane of the circle $ABCD$ and parallel to BD . Let MN meet the circle in O , P the diameter AC in S , and the straight lines AE , AF in Q , R respectively. Join AO .

Through MN draw a plane at right angles to AC ,

thus plane will cut the cylinder in a circle with diameter MN , the sphere in a circle with diameter OP , and the cone in a circle with diameter QR

Now, since $MS=AC$, and $QS=AS$

$$\begin{aligned} MS \cdot SQ &= CA \cdot AS \\ &= 10 \\ &= OS + SQ^2 \end{aligned}$$

And, since $HA=AC$,

$$\begin{aligned} HA \cdot AS &= CA \cdot AS \\ &= MS \cdot SQ \\ &= MS \cdot MS \cdot SQ \\ &= MS \cdot (OS^2 + SQ^2), \\ &\quad \text{from above} \\ &= MN^2 \cdot (OP + QR^2) \\ &= (\text{circle, diam } MN) \cdot (\text{circle diam } OP \\ &\quad + \text{circle, diam } QR) \end{aligned}$$

That is

$$HA \cdot AS = (\text{circle in cylinder}) \cdot (\text{circle in sphere} + \text{circle in cone})$$

Therefore the circle in the cylinder placed where it is is in equilibrium about A , with the circle in the sphere together with the circle in the cone if both the latter circles are placed with their centres of gravity at H

Similarly for the three corresponding sections made by a plane perpendicular to AC and passing through any other straight line in the parallelogram LF parallel to EF

If we deal in the same way with all the sets of three circles in which planes perpendicular to AC cut the cylinder the sphere and the cone, and which make up those solids respectively it follows that the cylinder, in the place where it is will be in equilibrium about A with the sphere and the cone together

when both are placed with their centres of gravity at H

Therefore since A is the centre of gravity of the cylinder

$$HA \cdot AK = (\text{cylinder}) \cdot (\text{sphere} + \text{cone } AEF)$$

$$\text{But } HA = 2AK$$

$$\text{therefore cylinder} = 2(\text{sphere} + \text{cone } AEF)$$

$$\text{Now cylinder} = 3(\text{cone } AEF),$$

[Eucl vii 10]

$$\text{therefore cone } AEF = 2(\text{sphere})$$


$$\text{But since } EF = 2BD$$

$$\text{cone } AEF = 8(\text{cone } ABD)$$

$$\text{therefore sphere} = 4(\text{cone } ABD)$$

(2) Through B D draw $\downarrow BW \downarrow DY$ parallel to AC

and imagine a cylinder which has AC for axis and the circles on VX , WY as diameters for bases



Then

$$\begin{aligned}
 \text{cylinder } VY &= 2(\text{cylinder } VD) \\
 &= 6(\text{cone } ABD) \\
 &= \frac{3}{2}(\text{sphere}), \text{ from above}
 \end{aligned}$$

[Eucl XII 10]

QED

"From this theorem to the effect that a sphere is four times as great as the cone with a great circle of the sphere as base and with height equal to the radius of the sphere I conceived the notion that the surface of any sphere is four times as great as a great circle in it, for, judging from the fact that any circle is equal to a triangle with base equal to the circumference and height equal to the radius of the circle, I apprehended that, in like manner any sphere is equal to a cone with base equal to the surface of the sphere and height equal to the radius

PROPOSITION 3

By this method we can also investigate the theorem that

A cylinder with base equal to the greatest circle in a spheroid and height equal to the axis of the spheroid is $1\frac{1}{2}$ times the spheroid,
and when this is established, it is plain that

If any spheroid be cut by a plane through the centre and at right angles to the axis the half of the spheroid is double of the cone which has the same base and the same axis as the segment (i.e. the half of the spheroid)

Let a plane through the axis of a spheroid cut its surface in the ellipse $ABCD$ the diameters (i.e. axes) of which are AC BD , and let A be the centre


Draw a circle about BD as diameter and in a plane perpendicular to AC imagine a cone with this circle as base and A as vertex produced and cut by a plane through C parallel to its base the section will be a circle in a plane at right angles to AC and about EF as diameter

Imagine a cylinder with the latter circle as base and axis AC , produce CA to H making AH equal to CA

Let HC be regarded as the bar of a balance A being its middle point

In the parallelogram LF draw any straight line MN parallel to EF meeting the ellipse in O P and AE AF , AC in Q R , S respectively

If now a plane be drawn through MN at right angles to AC , it will cut the cylinder in a circle with diameter MN the spheroid in a circle with diameter OP and the cone in a circle with diameter QR



Since $HA = AC$

$$\begin{aligned} HA \quad AS &= CA \quad AS \\ &= EA \quad AQ \\ &= MS \quad SQ \end{aligned}$$

Therefore

$$HA \quad AS = MS \quad MS \quad SQ$$

But, by the property of the ellipse,

$$\begin{aligned} AS \quad SC \quad SO &= AK^2 \quad KB^2 \\ &= AS^2 \quad SQ^2 \end{aligned}$$

therefore

$$\begin{aligned} SQ \quad SO^2 &= AS^2 \quad AS \quad SC \\ &= SQ \quad SQ \quad QM, \end{aligned}$$

and accordingly

$$SO^2 = SQ \quad QM$$

Add SQ^2 to each side and we have

$$SO + SQ^2 = SQ \quad SM$$

Therefore from above we have

$$\begin{aligned} HA \quad AS &= MS^2 \quad (SO + SQ^2) \\ &= MN^2 \quad (OP^2 + QR^2) \\ &= (\text{circle diam } MN) \quad (\text{circle diam } OP + \text{circle diam } QR) \end{aligned}$$

That is

$$HA \quad AS = (\text{circle in cylinder}) \quad (\text{circle in spheroid} + \text{circle in cone})$$

Therefore the circle in the cylinder in the place where it is in equilibrium about A , with the circle in the spheroid and the circle in the cone together if both the latter circles are placed with their centres of gravity at H

Similarly for the three corresponding sections made by a plane perpendicular to AC and passing through any other straight line in the parallelogram LF parallel to EF

If we deal in the same way with all the sets of three circles in which planes perpendicular to AC cut the cylinder the spheroid and the cone and which make up those figures respectively, it follows that the cylinder in the place where it is will be in equilibrium about A with the spheroid and the cone together when both are placed with their centres of gravity at H

Therefore since K is the centre of gravity of the cylinder

$$HA \quad 4K = (\text{cylinder}) \quad (\text{spheroid} + \text{cone } AEF)$$

But $HA = 2AK$

$$\text{therefore} \quad \text{cylinder} = 2(\text{spheroid} + \text{cone } AEF)$$

$$\text{And} \quad \text{cylinder} = 3(\text{cone } AEF)$$

$$\text{therefore} \quad \text{cone } AEF = 2(\text{spheroid})$$

[Eucl XII 10]

But since $EF = 2BD$

$$\text{cone } AEF = 8(\text{cone } ABD)$$

$$\text{therefore} \quad \text{spheroid} = 4(\text{cone } ABD)$$

$$\text{and} \quad \text{half the spheroid} = 2(\text{cone } ABD)$$

Through $B \quad D$ draw $VBW \quad \backslash DY$ parallel to AC and imagine a cylinder which has AC for axis and the circles on VY, WY as diameters for bases

$$\text{Then} \quad \text{cylinder } \backslash Y = 2(\text{cylinder } \backslash D)$$

$$= 6(\text{cone } ABD)$$

$$= \frac{3}{2}(\text{spheroid}) \text{ from above}$$

Q E D

PROPOSITION 4

Any segment of a right-angled conoid (i.e. a paraboloid of revolution) cut off by a plane at right angles to the axis is $1\frac{1}{2}$ times the cone which has the same base and the same axis as the segment

This can be investigated by our method, as follows

Let a paraboloid of revolution be cut by a plane through the axis in the parabola BAC , and let it also be cut by another plane at right angles to the axis and intersecting the former plane in BC . Produce DA , the axis of the segment to H , making HA equal to AD .

Imagine that HD is the bar of a balance, A being its middle point

The base of the segment being the circle on BC as diameter and in a plane perpendicular to AD ,

imagine (1) a cone drawn with the latter circle as base and A as vertex and (2) a cylinder with the same circle as base and AD as axis

In the parallelogram EC let any straight line MN be drawn parallel to BC , and through MN let a plane be drawn at right angles to AD . This plane will cut the cylinder in a circle with diameter MN and the paraboloid in a circle with diameter OP .

Now BAC being a parabola and BD , OS ordinates

$$\begin{array}{l} DA \quad AS = BD^2 \quad OS^2 \\ \text{or} \quad HA \quad AS = MS^2 \quad SO^2 \end{array}$$

Therefore

$$\begin{aligned} HA \quad AS &= (\text{circle, rad } MS) (\text{circle rad } OS) \\ &= (\text{circle in cylinder}) (\text{circle in paraboloid}) \end{aligned}$$

Therefore the circle in the cylinder in the place where it is, will be in equilibrium about A with the circle in the paraboloid, if the latter is placed with its centre of gravity at H .

Similarly for the two corresponding circular sections made by a plane perpendicular to AD and passing through any other straight line in the parallelogram which is parallel to BC .

Therefore as usual if we take all the circles making up the whole cylinder and the whole segment and treat them in the same way we find that the cylinder in the place where it is is in equilibrium about A with the segment placed with its centre of gravity at H .

If K is the middle point of AD K is the centre of gravity of the cylinder, therefore

$$HA \quad AK = (\text{cylinder}) (\text{segment})$$


$$\text{Therefore} \quad \text{cylinder} = 2(\text{segment})$$

$$\text{And} \quad \text{cylinder} = 3(\text{cone } ABC),$$

therefore

$$\text{segment} = \frac{2}{3}(\text{cone } ABC)$$

[Eucl. XII 10]



PROPOSITION 5


The centre of gravity of a segment of a right-angled conoid (i.e. a paraboloid of revolution) cut off by a plane at right angles to the axis is on the straight line which is the axis of the segment and divides the said straight line in such a way that the portion of it adjacent to the vertex is double of the remaining portion

This can be investigated by the method as follows

Let a paraboloid of revolution be cut by a plane through the axis in the parabola BAC , and let it also be cut by another plane at right angles to the axis and intersecting the former plane in BC

Produce DA the axis of the segment, to H making HA equal to AD , and imagine DH to be the bar of a balance its middle point being A

The base of the segment being the circle on BC as diameter and in a plane perpendicular to AD , imagine a cone with this circle as base and A as vertex so that AB AC are generators of the cone



In the parabola let any double ordinate OP be drawn meeting AB , AD AC in Q , S , R respectively

Now, from the property of the parabola,

$$\begin{aligned} BD^2 \cdot OS &= DA \cdot AS \\ &= BD \cdot QS \\ &= BD^2 \cdot BD \cdot QS \end{aligned}$$

Therefore $OS^2 = BD \cdot QS$,

or $BD \cdot OS = OS^2 \cdot QS$,

whence $BD \cdot QS = OS^2 \cdot QS$

$$\begin{aligned} \text{But } BD \cdot QS &= AD \cdot AS \\ &= HA \cdot AS \end{aligned}$$

$$\begin{aligned} \text{Therefore } HA \cdot AS &= OS^2 \cdot QS^2 \\ &= OP^2 \cdot QR^2 \end{aligned}$$

If now through OP a plane be drawn at right angles to AD , this plane cuts the paraboloid in a circle with diameter OP and the cone in a circle with diameter QR

We see therefore that $HA \cdot AS = (\text{circle, diam } OP) \cdot (\text{circle diam } QR)$
 $= (\text{circle in paraboloid}) \cdot (\text{circle in cone}),$

and the circle in the paraboloid in the place where it is is in equilibrium about A with the circle in the cone placed with its centre of gravity at H

Similarly for the two corresponding circular sections made by a plane perpendicular to AD and passing through any other ordinate of the parabola

Dealing therefore in the same way with all the circular sections which make up the whole of the segment of the paraboloid and the cone respectively we see that the segment of the paraboloid in the place where it is is in equilibrium about A with the cone placed with its centre of gravity at H

Now since A is the centre of gravity of the whole system as placed and the centre of gravity of part of it namely the cone as placed is at H , the centre of gravity of the rest namely the segment is at a point K on HA produced such that

$$HA \cdot AK = (\text{segment}) \cdot (\text{cone})$$

But $\text{segment} = 2(\text{cone})$

[Prop 4]

H , balances the cone ABD (alone) in the place where it is therefore the portion N of the cylinder placed with its centre of gravity at H balances the hemisphere (alone) in the place where it is

Now the centre of gravity of the cone is at a point V such that $AG=4GV$, therefore, since M at H is in equilibrium with the cone

$$M \text{ (cone)} = \frac{3}{4}AG \quad HA = \frac{3}{4}AC \quad AC,$$

whence $M = \frac{3}{8}(\text{cone})$

But $M+N=(\text{cone})$ therefore $N=\frac{5}{8}(\text{cone})$

Now let the centre of gravity of the hemisphere be at W which is somewhere on AG

Then, since N at H balances the hemisphere alone,

$$(\text{hemisphere}) \quad N=HA \quad AW$$

But the hemisphere BAD =twice the cone ABD ,

[On the Sphere and Cylinder I 34 and Prop 2 above]

and $N=\frac{5}{8}(\text{cone})$, from above

$$\text{Therefore} \quad 2 \frac{5}{8} = HA \quad AW \\ = 2AG \quad AW$$

whence $AW=\frac{5}{8}AG$, so that W divides AG in such a way that

$$AW \quad WG=5 \quad 3]$$

PROPOSITION 7

We can also investigate by the same method the theorem that


[Any segment of a sphere has] to the cone [with the same base and height the ratio which the sum of the radius of the sphere and the height of the complementary segment has to the height of the complementary segment]

[There is a lacuna here but all that is missing is the construction and the construction is easily understood by means of the figure BAD is of course the

segment of the sphere the volume of which is to be compared with the volume of a cone with the same base and height]

The plane drawn through MN and at right angles to AC will cut the cylinder in a circle with diameter MN the segment of the sphere in a circle with diameter OP and the cone on the base EF in a circle with diameter QR

In the same way as before [cf Prop 2] we can prove that the circle with diameter MN in the place where it is is in equilibrium about A with the two circles with diameters $OP \quad QR$ if these circles are both moved and placed with their centres of gravity at H



The same thing can be proved of all sets of three circles in which the cylinder

der, the segment of the sphere, and the cone with the common height $4G$ are all cut by any plane perpendicular to AC

Since then the sets of circles make up the whole cylinder, the whole segment of the sphere and the whole cone respectively, it follows that the cylinder in the place where it is, is in equilibrium about A with the sum of the segment of the sphere and the cone if both are placed with their centres of gravity at H

Divide AG at W , V in such a way that

$$AW = WG, AV = 3VG$$

Therefore W will be the centre of gravity of the cylinder, and V will be the centre of gravity of the cone

Since, now the bodies are in equilibrium as described,

$$(\text{cylinder}) (\text{cone } AEF + \text{segment } BAD \text{ of sphere}) = HA \quad 4W$$

[The rest of the proof is lost, but it can easily be supplied thus

We have

$$\begin{aligned} (\text{cone } AEF + \text{segmt } BAD) (\text{cylinder}) &= AW \quad AC \\ &= AW \quad AC \quad AC^2 \end{aligned}$$

But

$$\begin{aligned} (\text{cylinder}) (\text{cone } AEF) &= 4C^2 \quad \frac{1}{3}EG^2 \\ &= AC^2 \quad \frac{1}{3}AG^2 \end{aligned}$$

Therefore, *ex aequali*

$$\begin{aligned} (\text{cone } AEF + \text{segmt } BAD) (\text{cone } AEF) &= AW \quad AC \quad \frac{1}{3}AG^2 \\ &= \frac{1}{3}AC \quad \frac{1}{3}AG, \end{aligned}$$

whence

$$\begin{aligned} (\text{segmt } BAD) (\text{cone } AEF) &= (\frac{1}{3}AC - \frac{1}{3}AG) \quad \frac{1}{3}AG \\ (\text{cone } AEF) (\text{cone } ABD) &= EG^2 \quad DG^2 \\ &= AG^2 \quad AG \quad GC \\ &= AG \quad GC \\ &= \frac{1}{3}AG \quad \frac{1}{3}GC \end{aligned}$$

Therefore, *ex aequali*

$$\begin{aligned} (\text{segment } BAD) (\text{cone } ABD) &= (\frac{1}{3}AC - \frac{1}{3}AG) \quad \frac{1}{3}GC \\ &= (\frac{1}{3}AC - AG) \quad GC \\ &= (\frac{1}{3}AC + GC) \quad GC \quad \text{Q.E.D.} \end{aligned}$$

PROPOSITION 8

[The enunciation the setting out, and a few words of the construction are missing

The enunciation however can be supplied from that of Prop 9 with which it must be identical except that it cannot refer to any segment, and the presumption therefore is that the proposition was enunciated with reference to one kind of segment only i.e. either a segment greater than a hemisphere or a segment less than a hemisphere

Heiberg's figure corresponds to the case of a segment greater than a hemisphere The segment investigated is of course the segment BAD The setting-out and construction are self-evident from the figure }

Produce AC to H , O , making HA equal to AC and CO equal to the radius of the sphere

and let HC be regarded as the bar of a balance the middle point being A

In the plane cutting off the segment describe a circle with G as centre and radius (GE) equal to $1G$ and on this circle as base and with A as vertex, let a cone be described AE AF are generators of this cone

Draw KL through any point Q on AG , parallel to EF and cutting the segment in K , L and AE AF in R , P respectively Join AK

Now

$$\begin{aligned}
 HA \cdot AQ &= CA \cdot AQ \\
 &= AK^2 - AQ^2 \\
 &= (KQ^2 + QA^2) - QA^2 \\
 &= (KQ^2 + PQ^2) - PQ^2 \\
 &= (\text{circle, diam } KL + \text{circle diam } PR) - (\text{circle diam } PR)
 \end{aligned}$$

Imagine a circle equal to the circle with diameter PR placed with its centre of gravity at H therefore the circles on diameters KL PR in the places where they are are in equilibrium about A with the circle with diameter PR placed with its centre of gravity at H

Similarly for the corresponding circular sections made by any other plane perpendicular to AG

Therefore, taking all the circular sections which make up the segment ABD of the sphere and the cone AEF respectively, we find that the segment ABD of the sphere and the cone AEF in the places where they are are in equilibrium with the cone AEF assumed to be placed with its centre of gravity at H

Let the cylinder $M+N$ be equal to the cone AEF which has A for vertex and the circle on EF as diameter for base

Divide AG at V so that

$$4G = 4VG,$$

therefore V is the centre of gravity of the cone AEF , for this has been proved before

Let the cylinder $M+N$ be cut by a plane perpendicular to the axis in such a way that the cylinder M (alone) placed with its centre of gravity at H is in equilibrium with the cone AEF


Since $M+N$ suspended at H is in equilibrium with the segment ABD of the sphere and the cone AEF in the places where they are while M , also at H is in equilibrium with the cone AEF in the place where it is, it follows that

N at H is in equilibrium with the segment ABD of the sphere in the place where it is

Now (segment ABD of sphere) (cone ABD) = $OG \cdot GC$
for this is already proved [Cf *On the Sphere and Cylinder* II 2 Cor as well as Prop 7 ante]

And

$$\begin{aligned}
 &(\text{cone } ABD) \cdot (\text{cone } AEF) \\
 &= (\text{circle diam } BD) \cdot (\text{circle diam } EF) \\
 &= BD^2 \cdot EF^2 \\
 &= BG^2 \cdot GE^2 \\
 &= CG \cdot GA \cdot GA^2 \\
 &= CG \cdot GA
 \end{aligned}$$



Therefore, *ex aequali*,

(segment ABD of sphere) (conc AEF) = $OG \cdot GA$

Take a point W on AG such that

$$AW \cdot WG = (GA + 4GC) \cdot (GA + 2GC)$$

We have then inversely,

$$GW \cdot WA = (2GC + GA) \cdot (4GC + GA),$$

and *componendo*,

$$GA \cdot AW = (6GC + 2GA) \cdot (4GC + GA)$$

But $GO = \frac{1}{2}(6GC + 2GA)$, [for $GO - GC = \frac{1}{2}(CG + GA)$]

and $CV = \frac{1}{2}(4GC + GA)$,

therefore $GA \cdot AW = OG \cdot CV$,

and alternately and inversely,

$$OG \cdot GA = CV \cdot WA$$

It follows, from above that

$$(\text{segment } ABD \text{ of sphere}) (\text{cone } AEF) = CV \cdot WA$$

Now since the cylinder M with its centre of gravity at H is in equilibrium about A with the cone AEF with its centre of gravity at V ,

$$\begin{aligned} (\text{cone } AEF) (\text{cylinder } M) &= HA \cdot AV \\ &= CA \cdot AV, \end{aligned}$$

and since the cone AEF = the cylinder $M + N$ we have *dividendo* and *inter tendo*,

$$(\text{cylinder } M) (\text{cylinder } N) = AV \cdot CV$$

Hence *componendo*

$$\begin{aligned} (\text{cone } AEF) (\text{cylinder } N) &= CA \cdot CV \\ &= HA \cdot CV \end{aligned}$$

But it was proved that

$$(\text{segment } ABD \text{ of sphere}) (\text{cone } AEF) = CV \cdot WA,$$

therefore, *ex aequali*,

$$(\text{segment } ABD \text{ of sphere}) (\text{cylinder } N) = HA \cdot AW$$

And it was above proved that the cylinder N at H is in equilibrium about A with the segment ABD in the place where it is
therefore since H is the centre of gravity of the cylinder N , W is the centre of gravity of the segment ABD of the sphere

PROPOSITION 9

In the same way we can investigate the theorem that

The centre of gravity of any segment of a sphere is on the straight line which is the axis of the segment and divides this straight line in such a way that the part of it adjacent to the vertex of the segment has to the remaining part the ratio which the sum of the axis of the segment and four times the axis of the complementary segment has to the sum of the axis of the segment and double the axis of the complementary segment

[Is this theorem related to 'any segment' but states the same result as that proved in the preceding proposition, it follows that Prop 8 must have related to one kind of segment either a segment greater than a semicircle (as in Heiberg's figure of Prop 8) or a segment less than a semicircle, and the present proposition completed the proof for both kinds of segments. It would only require a slight change in the figure in any case.]

PROPOSITION 10

By this method too we can investigate the theorem that

[A segment of an obtuse angled conoid (i.e. a hyperboloid of revolution) has to the cone which has] the same base [as the segment and equal height the same ratio as the sum of the axis of the segment and three times] the "annex to the axis" (i.e. half the transverse axis of the hyperbolic section through the axis of the hyperboloid, or in other words the distance between the vertex of the segment and the vertex of the enveloping cone) has to the sum of the axis of the segment and double of the 'annex' [this is the theorem proved in *On Conoids and Spheroids*, Prop 25] "and also many other theorems which as the method has been made clear by means of the foregoing examples I will omit in order that I may now proceed to compass the proofs of the theorems mentioned above."

PROPOSITION 11

If in a right prism with square bases a cylinder be inscribed having its bases in opposite square faces and touching with its surface the remaining four parallelogrammic faces and if through the centre of the circle which is the base of the cylinder and one side of the opposite square face a plane be drawn, the figure cut off by the plane so drawn is one sixth part of the whole prism

'This can be investigated by the method and when it is set out I will go back to the proof of it by geometrical considerations'


[The investigation by the mechanical method is contained in the two Propositions, 11, 12 Prop 13 gives another solution which, although it contains no mechanics is still of the character which Archimedes regards as inconclusive since it assumes that the solid is actually made up of parallel plane sections

and that an auxiliary parabola is actually made up of parallel straight lines in it Prop 14 added the conclusive geometrical proof]

Let there be a right prism with a cylinder inscribed as stated

Let the prism be cut through the axis of the prism and cylinder by a plane perpendicular to the plane which cuts off the portion of the cylinder let this plane make as section the parallelogram AB , and let it cut the plane cutting off the portion of the cylinder (which plane is perpendicular to AB) in the straight line BC

Let CD be the axis of the prism and cylinder let EF bisect it at right angles and through EF



let a plane be drawn at right angles to CD this plane will cut the prism in a square and the cylinder in a circle

Let MN be the square and $OPQR$ the circle and let the circle touch the sides of the square in O P Q R [F E in the first figure are identical with O Q respectively] Let H be the centre of the circle


Let KL be the intersection of the plane through EF perpendicular to the axis of the cylinder and the plane cutting off the portion of the cylinder KL is bisected by OHQ [and passes through the middle point of HQ]

Let any chord of the circle, as ST , be drawn perpendicular to HQ , meeting HQ in W ,

and through ST let a plane be drawn at right angles to OQ and produced on both sides of the plane of the circle $OPQR$

The plane so drawn will cut the half cylinder having the semicircle PQR for section and the axis of the prism for height in a parallelogram, one side of which is equal to ST and another is a generator of the cylinder, and it will also cut the portion of the cylinder cut off in a parallelogram, one side of which is equal to ST and the other is equal and parallel to UV (in the first figure)

UV will be parallel to BY and will cut off, along EG in the parallelogram DE , the segment EI equal to QV



Now, since EC is a parallelogram, and VI is parallel to GC ,

$$EG \quad GI = EC \quad CV$$

$$= BY \quad UV$$

$$= (\square \text{ in half cyl}) \quad (\square \text{ in portion of cyl})$$

And $EG = HQ$ $GI = HW$, $QH = OH$,

therefore $OH \quad HW = (\square \text{ in half cyl}) \quad (\square \text{ in portion})$

Imagine that the parallelogram in the portion of the cylinder is moved and placed at O so that O is the centre of gravity, and that OQ is the bar of a balance H being its middle point

Then, since W is the centre of gravity of the parallelogram in the half cylinder, it follows from the above that the parallelogram in the half cylinder, in the place where it is, with its centre of gravity at W is in equilibrium about H with the parallelogram in the portion of the cylinder when placed with its centre of gravity at O

Similarly for the other parallelogrammic sections made by any plane perpendicular to OQ and passing through any other chord in the semicircle PQR perpendicular to OQ

If then we take all the parallelograms making up the half cylinder and the portion of the cylinder respectively, it follows that the half cylinder, in the place where it is is in equilibrium about H with the portion of the cylinder cut off when the latter is placed with its centre of gravity at O

PROPOSITION 12

Let the parallelogram (square) MN perpendicular to the axis with the circle $OPQR$ and its diameters $OQ \quad PR$ be drawn separately


Join $HG \quad HM$, and through them draw planes at right angles to the plane of the circle, producing them on both sides of that plane

This produces a prism with triangular section GHM and height equal to the axis of the cylinder, this prism is $\frac{1}{4}$ of the original prism circumscribing the cylinder

Let LK , UT be drawn parallel to OQ and equidistant from it, cutting the circle in K , T , RP in $S \quad P$ and $GH \quad HM$ in $W \quad V$ respectively

Through LK , UT draw planes at right angles to PR , producing them on both sides of the plane of the circle,

these planes produce as sections in the half cylinder PQR and in the prism GHM four parallelograms in which the heights are equal to the axis of the cylinder, and the other sides are equal to KS , TF , LW , UV respectively



[The rest of the proof is missing but, as Zeuthen says the result obtained and the method of arriving at it are plainly indicated by the above

Archimedes wishes to prove that the half cylinder PQR , in the place where it is balances the prism GHM , in the place where it is, about H as fixed point

He has first to prove that the elements (1) the parallelogram with side = KS and (2) the parallelogram with side = LW , in the places where they are, balance about S , or, in other words that the straight lines SK , LW , in the places where they are, balance about S

$$(\text{radius of circle } OPQR)^2 = SK^2 + SH^2,$$

$$SL^2 = SK^2 + SW^2$$

$$LS^2 - SW^2 = SK^2,$$

$$(LS + SW) LW = SK^2,$$

$$\frac{1}{2}(LS + SW) \cdot \frac{1}{2}SK = SK \cdot LW$$

Now
or

Therefore
and accordingly
whence

And $\frac{1}{2}(LS + SW)$ is the distance of the centre of gravity of LW from S while $\frac{1}{2}SK$ is the distance of the centre of gravity of SK from S

Therefore SK and LW , in the places where they are, balance about S

Similarly for the corresponding parallelograms

Taking all the parallelogrammic elements in the half cylinder and prism respectively we find that the half cylinder PQR and the prism GHM in the places where they are respectively, balance about H

From this result and that of Prop 11 we can at once deduce the volume of the portion cut off from the cylinder For in Prop 11 the portion of the cylinder, placed with its centre of gravity at O , is shown to balance (about H) the half cylinder in the place where it is By Prop 12 we may substitute for the half cylinder in the place where it is the prism GHM of that proposition turned the opposite way relatively to RP The centre of gravity of the prism as thus placed is at a point (say Z) on HQ such that $HZ = \frac{2}{3}HQ$

Therefore assuming the prism to be applied at its centre of gravity we have

$$(\text{portion of cylinder}) (\text{prism}) = \frac{2}{3}HQ \cdot OH$$

$$= \frac{2}{3} \cdot \frac{1}{3}HQ^2$$

therefore

$$(\text{portion of cylinder}) = \frac{2}{3}(\text{prism } GHM) \\ = \frac{2}{3}(\text{original prism})$$

PROPOSITION 13

Let there be a right prism with square bases one of which is $ABCD$ in the prism let a cylinder be inscribed the base of which is the circle $EFGH$ touching the sides of the square $ABCD$ in E , F , G , H

Through the centre and through the side corresponding to CD in the square

face *opposite* to $ABCD$ let a plane be drawn, this will cut off a prism equal to $\frac{1}{2}$ of the original prism and formed by three parallelograms and two triangles, the triangles forming opposite faces

In the semicircle EPG describe the parabola which has FA for axis and passes through E, G , draw MN parallel to AF meeting GE in M , the parabola in L , the semicircle in O and CD in N

Then $MN \cdot NL = NF^2$,

"for this is clear"


[Cf Apollonius, *Conics* I 11]

[The parameter is of course equal to GA or AF]

Therefore $MN \cdot NL = GA^2 = LS^2$

Through MN draw a plane at right angles to EG ,

this will produce as sections (1) in the prism cut off from the whole prism a right angled triangle, the base of which is MN , while the perpendicular is perpendicular at N to the plane $ABCD$ and equal to the axis of the cylinder, and the hypotenuse is in the plane cutting the cylinder and (2) in the portion of the cylinder cut off a right angled triangle the base of which is MO while the perpendicular is the generator of the cylinder perpendicular at O to the plane AN and the hypotenuse is



[There is a lacuna here to be supplied as follows

Since $MN \cdot NL = GA^2 = LS^2$

$$= MN^2 \cdot LS$$

it follows that

$$MN \cdot ML = MN^2 \cdot (MN^2 - LS^2)$$

$$= MN^2 \cdot (MN^2 - MA^2)$$

$$= MN^2 \cdot MO^2$$

But the triangle (1) in the prism is to the triangle (2) in the portion of the cylinder in the ratio of $MN^2 : MO^2$

Therefore (Δ in prism) (Δ in portion of cylinder)

$$= MN \cdot ML$$

$$= (\text{straight line in rect } DG) (\text{straight line in parabola})$$

We now take all the corresponding elements in the prism, the portion of the cylinder the rectangle DG and the parabola EFG respectively], and it will follow that

$$(\text{all the } \Delta\text{s in prism}) (\text{all the } \Delta\text{s in portion of cylinder})$$

$$= (\text{all the str lines in } \square DG) (\text{all the straight lines between parabola and } EG)$$

But the prism is made up of the triangles in the prism [the portion of the cylinder is made up of the triangles in it] the parallelogram DG of the straight lines in it parallel to AF and the parabolic segment of the straight lines parallel to AF intercepted between its circumference and EG , therefore

$$(\text{prism}) (\text{portion of cylinder})$$

$$= (\square DG) (\text{parabolic segment } EFG)$$

But $\square DG = 2(\text{parabolic segment } EFG)$,

"for this is proved in my earlier treatise

[*Quadrature of Parabola*]

Therefore prism = $\frac{2}{3}$ (portion of cylinder)

If then we denote the portion of the cylinder by 2 the prism is 3 and the original prism circumscribing the cylinder is 12 (being 4 times the other prism), therefore the portion of the cylinder = $\frac{1}{6}$ (original prism) Q E D

[The above proposition and the next are peculiarly interesting for the fact that the parabola is an auxiliary curve introduced for the sole purpose of analytically reducing the required cubature to the known quadrature of the parabola]

PROPOSITION 14

Let there be a right prism with square bases [and a cylinder inscribed therein having its base in the square $ABCD$ and touching its sides at E, F, G, H let the cylinder be cut by a plane through EG and the side corresponding to CD in the square face opposite to $ABCD$]

This plane cuts off from the prism a prism, and from the cylinder a portion of it

It can be proved that the portion of the cylinder cut off by the plane is $\frac{1}{2}$ of the whole prism

But we will first prove that it is possible to inscribe in the portion cut off from the cylinder, and to circumscribe about it solid figures made up of prisms which have equal height and similar triangular bases in such a way that the circumscribed figure exceeds the inscribed by less than any assigned magnitude

But it was proved that

(prism cut off by oblique plane) $< \frac{2}{3}$ (figure inscribed in portion of cylinder)

Now (prism cut off) (inscribed figure)

= $\square DG$ (\square s inscribed in parabolic segment)

therefore $\square DG < \frac{2}{3}$ (\square s in parabolic segment)

which is impossible since it has been proved elsewhere' that the parallelogram DG is $\frac{2}{3}$ of the parabolic segment

Consequently

not greater

And (all the prisms in prism cut off)

(all prisms in circumscr figure)

= (all \square s in $\square DG$) (all \square s in fig circumscr about parabolic segmt), therefore

(prism cut off) (figure circumscr about portion of cylinder)


= ($\square DG$) (figure circumscr about parabolic segment)

But the prism cut off by the oblique plane is $> \frac{2}{3}$ of the solid figure circumscribed about the portion of the cylinder

[There are large gaps in the exposition of this geometrical proof but the way in which the method of exhaustion was applied and the parallelism between this and other applications of it are clear The first fragment shows that solid figures made up of prisms were circumscribed and inscribed to the portion of the cylinder The parallel triangular faces of these prisms were perpendicular to GE in the figure of Prop 13 they divided GE into equal portions of the requisite smallness each section of the portion of the cylinder by such a

plane was a triangular face common to an inscribed and a circumscribed right prism. The planes also produced prisms in the prism cut off by the same oblique plane as cuts off the portion of the cylinder and standing on GD as base.

The number of parts into which the parallel planes divided GE was made great enough to secure that the circumscribed figure exceeded the inscribed figure by less than a small assigned magnitude.



The second part of the proof began with the assumption that the portion of the cylinder is $> \frac{1}{2}$ of the prism cut off, and this was proved to be impossible, by means of the use of the auxiliary parabola and the proportion

$$MN : ML = MN^2 : MO^2$$


which are employed in Prop. 13.

We may supply the missing proof as follows:

In the accompanying figure are represented (1) the first element-prism circumscribed to the portion of the cylinder (2) two element prisms adjacent to the ordinate OM of which that on the left is circumscribed and that on the right (equal to the other) inscribed (3) the corresponding element-prisms forming part of the prism cut off ($CC'GEDD'$) which is $\frac{1}{2}$ of the original prism.

In the second figure are shown element rectangles circumscribed and inscribed to the auxiliary parabola, which rectangles correspond exactly to the circumscribed and inscribed element prisms represented in the first figure (the length of GM is the same in both figures and the breadths of the element-rectangles are the same as the heights of the element-prisms), the corresponding element-rectangles forming part of the rectangle GD are similarly shown.

For convenience we suppose that GE is divided into an even number of equal parts so that GA contains an integral number of these parts.



For the sake of brevity we will call each of the two element prisms of which OM is an edge 'el prism (O)' and each of the element-prisms of which MNN' is a common face 'el prism (N)'. Similarly we will use the corresponding abbreviations 'el rect (L)' and 'el rect (N)' for the corresponding elements in relation to the auxiliary parabola as shown in the second figure

Now it is easy to see that the figure made up of all the inscribed prisms is less than the figure made up of the circumscribed prisms by twice the final circumscribed prism adjacent to FK , i.e. by twice 'el prism (N)' and as the height of this prism may be made as small as we please by dividing GK into sufficiently small parts it follows that inscribed and circumscribed solid figures made up of element prisms can be drawn differing by less than any assigned solid figure

(1) Suppose if possible that

$$(\text{portion of cylinder}) > \frac{3}{2}(\text{prism cut off}),$$

or $(\text{prism cut off}) < \frac{2}{3}(\text{portion of cylinder})$

Let $(\text{prism cut off}) = \frac{2}{3}(\text{portion of cylinder} - Y)$, say

Construct circumscribed and inscribed figures made up of element-prisms such that

$$(\text{circumscr fig}) - (\text{inscr fig}) < X$$

Therefore $(\text{inscr fig}) > (\text{circumscr fig} - X)$
and a fortiori $> (\text{portion of cyl} - X)$

It follows that

$$(\text{prism cut off}) < \frac{2}{3}(\text{inscribed figure})$$

Considering now the element-prisms in the prism cut off and those in the inscribed figure respectively we have

$$\begin{aligned} \text{el prism } (N) - \text{el prism } (O) &= MN^2 - MO^2 \\ &= MN \cdot ML \quad [\text{as in Prop 13}] \\ &= \text{el rect } (N) - \text{el rect } (L) \end{aligned}$$

It follows that

$$\Sigma\{\text{el prism } (N)\} - \Sigma\{\text{el prism } (O)\} = \Sigma\{\text{el rect } (N)\} - \Sigma\{\text{el rect } (L)\}$$

(There are really two more prisms and rectangles in the first and third than there are in the second and fourth terms respectively but this makes no difference because the first and third terms may be multiplied by a common factor as $n/(n-2)$ without affecting the truth of the proportion Cf the proposition from *On Conoids and Spheroids* quoted on p 571 above)

Therefore

$$\begin{aligned} (\text{prism cut off}) - (\text{figure inscr in portion of cyl}) \\ = (\text{rect } GD) - (\text{fig inscr in parabola}) \end{aligned}$$

But it was proved above that

$$(\text{prism cut off}) < \frac{2}{3}(\text{fig inscr in portion of cyl})$$

therefore $(\text{rect } GD) < \frac{2}{3}(\text{fig inscr in parabola})$,

and a fortiori $(\text{rect } GD) < \frac{2}{3}(\text{parabolic segmt})$

which is impossible since

$$(\text{rect } GD) = \frac{2}{3}(\text{parabolic segmt})$$

Therefore (portion of cyl) is *not* greater than $\frac{2}{3}(\text{prism cut off})$

(2) In the second lacuna must have come the beginning of the next *reductio ad absurdum* demolishing the other possible assumption that the portion of the cylinder is $< \frac{2}{3}$ of the prism cut off

In this case our assumption is that

(prism cut off) $> \frac{2}{3}$ (portion of cylinder),

and we circumscribe and inscribe figures made up of element prisms, such that

(prism cut off) $> \frac{2}{3}$ (fig circumscriber about portion of cyl)

We now consider the element prisms in the prism cut off and in the circumscribed figure respectively, and the same argument as above gives

(prism cut off) (fig circumscriber about portion of cyl)

= (rect GD) (fig circumscriber about parabola),

whence it follows that

(rect GD) $> \frac{2}{3}$ (fig circumscribed about parabola),

and, *a fortiori*,

(rect GD) $> \frac{2}{3}$ (parabolic segment)

which is impossible since

(rect GD) = $\frac{2}{3}$ (parabolic segmt)

Therefore

(portion of cyl) is *not* less than $\frac{2}{3}$ (prism cut off)

But it was also proved that neither is it greater,

therefore (portion of cyl) = $\frac{2}{3}$ (prism cut off)
= $\frac{1}{4}$ (original prism)]

[PROPOSITION 15]

[This proposition, which is lost would be the mechanical investigation of the second of the two special problems mentioned in the preface to the treatise, namely that of the cubature of the figure included between two cylinders, each of which is inscribed in one and the same cube so that its opposite bases are in two opposite faces of the cube and its surface touches the other four faces

Zeuthen has shown how the mechanical method can be applied to this case

In the accompanying figure VII $Y \Lambda$ is a section of the cube by a plane (that of the paper) passing through the axis BD of one of the cylinders inscribed in the cube and parallel to two opposite faces

The same plane gives the circle $ABCD$ as the section of the other inscribed cylinder with axis perpendicular to the plane of the paper and extending on each side of the plane to a distance equal to the radius of the circle or half the side of the cube


AC is the diameter of the circle which is perpendicular to BD

Join AB AD and produce them to meet the tangent at C to the circle in E F

Then $EC = CF = CA$

Let LG be the tangent at A and complete the rectangle $EFGL$

Draw straight lines from A to the four corners of the section in which the plane through BD perpendicular to AA cuts the cube These straight lines if



produced, will meet the plane of the face of the cube opposite to A in four points forming the four corners of a square in that plane with sides equal to EF or double of the side of the cube, and we thus have a pyramid with A for vertex and the latter square for base

Complete the prism (parallelepiped) with the same base and height as the pyramid

Draw in the parallelogram LF any straight line MN parallel to EF and through MN draw a plane at right angles to AC

This plane cuts—

- (1) the solid included by the two cylinders in a square with side equal to OP ,
- (2) the prism in a square with side equal to MN , and
- (3) the pyramid in a square with side equal to QR

Produce CA to H , making HA equal to AC , and imagine HC to be the bar of a balance

Now, as in Prop 2 since $MS=AC$, $QS=AS$,

$$\begin{aligned} MS \cdot SQ &= CA \cdot AS \\ &= AO^2 \\ &= OS^2 + SQ^2 \end{aligned}$$

$$\begin{aligned} \text{Also } HA \cdot AS &= CA \cdot AS \\ &= MS \cdot SQ \\ &= MS \cdot MS \cdot SQ \\ &= MS^2 (OS + SQ) \text{ from above,} \\ &= MN^2 (OP^2 + QR^2) \\ &= (\text{square side } MN) (\text{sq, side } OP + \text{sq, side } QR) \end{aligned}$$

Therefore the square with side equal to MN , in the place where it is is in equilibrium about A with the squares with sides equal to OP , QR respectively placed with their centres of gravity at H

Proceeding in the same way with the square sections produced by other planes perpendicular to AC we finally prove that the prism in the place where it is is in equilibrium about A with the solid included by the two cylinders and the pyramid, both placed with their centres of gravity at H

Now the centre of gravity of the prism is at K

Therefore $HA \cdot AK = (\text{prism}) (\text{solid} + \text{pyramid})$

or $2 \cdot 1 = (\text{prism}) (\text{solid} + \frac{1}{3} \text{ prism})$

Therefore $2 (\text{solid}) + \frac{2}{3} (\text{prism}) = (\text{prism})$

It follows that

$$\begin{aligned} (\text{solid included by cylinders}) &= \frac{1}{3} (\text{prism}) \\ &= \frac{2}{3} (\text{cube}) \end{aligned} \quad \text{Q E D}$$

There is no doubt that Archimedes proceeded to and completed the rigorous geometrical proof by the method of exhaustion

As observed by Prof C Juel (Zeuthen *l c*) the solid in the present proposition is made up of 8 pieces of cylinders of the type of that treated in the preceding proposition. As however the two propositions are separately stated, there is no doubt that Archimedes' proofs of them were distinct

In this case AC would be divided into a very large number of equal parts and planes would be drawn through the points of division perpendicular to AC . These planes cut the solid and also the cube VY in square sections. Thus we can inscribe and circumscribe to the solid the requisite solid figures made up of element prisms and differing by less than any assigned solid magnitude the

prisms have square bases and their heights are the small segments of AC . The element prism in the inscribed and circumscribed figures which has the square equal to OP^2 for base corresponds to an element prism in the cube which has for base a square with side equal to that of the cube, and as the ratio of the element prisms is the ratio $OS^2 : BA^2$, we can use the same auxiliary parabola, and work out the proof in exactly the same way, as in Prop. 14.]

CONICS

BIOGRAPHICAL NOTE

APOLLONIUS, c 262-c 200 B C

APOLLONIUS was born at Perga in Pamphylia Asia Minor some twenty five years after the birth of Archimedes which would place his birth around the year 262 B C He seems to have gone when quite young to Alexandria where, according to Pappus the fourth century mathematician he was attracted by the reputation of the astronomer, Aristarchus of Samos Apollonius studied under the successors of Euclid at Alexandria and continued to reside there during the reigns of Ptolemy Euergetes and of Ptolemy Philopator (247-203 B C) He was also for some time in Pergamum, where he made the acquaintance of the mathematician, Eudemus to whom he dedicated the first three books of his *Conics* and of King Attalus I (269-197 B C), to whom the remaining five books of the *Conics* were dedicated

Apollonius appears to have been associated with the leading mathematicians of his day In the dedicatory epistles of the *Conics* he records that he met Philonides while on a trip to Ephesus and that he undertook the composition of this work in the first instance for Naucrates, who was staying in Alexandria Speaking in the same place of the preceding writers on conics Apollonius points out their limitations and inadequacies in such a way that some of his readers such as Pappus have considered him boastful and envious but it would seem that Apollonius is only trying to explain the appearance of a new text book on the elements of conics (Books I-IV) and the publication of his own original and more advanced investigations (Books V-VIII)

The *Conics* were at once recognized as the authoritative treatise on the subject winning for their author the name of the great geometer They are regularly cited by later writers Pappus added a group of lemmas and Eutocius (fl 500 A D) edited and commented on the first four books These books are extant in the original Greek the fifth sixth and seventh books exist in an Arabic translation, the eighth book is known only indirectly

Although the titles and a general indication of the contents of other works by Apollonius are given by later writers especially by Pappus only one the *Cutting of a Ratio* has survived and that like parts of the *Conics* only in an Arabic version All of the original work with the exception of the second half of the *Conics* has perished Books not extant but known through Pappus are *Cutting of an Area* *Determinate Section* *Tangencies* *Inclinations* and *Plane Loci* He wrote on irrationals and like Archimedes devised a system of multiplication for counting large numbers and calculated an approximate value for the ratio of the circumference of a circle to the diameter The ancient writers also record that Apollonius wrote *On the Burning Glass* in which he probably treated the properties of the parabola a work comparing the dodecahedron and the icosahedron inscribed in the same sphere and a book perhaps on the general principles of mathematics in which he criticized and suggested improvements for Euclid's *Elements* Lastly in astronomy he is credited by Ptol-

emy with an explanation of the motion of the planets by means of epicycles and eccentric circles. He seems to have been especially interested in the theory of the moon, and the Alexandrians are said to have called him Epsilon from the resemblance of that Greek letter to the lunar crescent.

CONTENTS

BIOGRAPHICAL NOTE p 595

BOOK I, p 603 BOOK II, p 682 BOOK III, p 731

TRANSLATOR'S NOTE

If on first appearance this treatise should seem to the reader a jumble of propositions rigorous indeed, but without much rhyme or reason in their sequence, then he can be sure he has not read aright, and as with the planets, he must look further to save the appearances. There are one or two hypotheses at least that can order the apparent wanderings of parabolas, hyperbolas, and ellipses through the first four books. Such hypotheses are the analogies between the three sections, and especially the development of the analogy between the hyperbola and the ellipse reaching its culmination, in the first book, with the final theorem, the construction of conjugate opposite sections.

In First Definitions I 5, Apollonius innocently defines two kinds of diameters, the transverse and the upright. Each one, in a conic section, bisects all the straight lines parallel to the other. But the upright diameter, defined here only as to position, has, in the case of the ellipse, natural bounds fixed by the section itself, and in Proposition I 15 we find it is the mean proportional between the corresponding transverse diameter (or conjugate diameter) and its parameter. The transverse diameter, in turn, is the mean proportional between the upright diameter (or conjugate) and its parameter, so "upright" and "transverse" become meaningless terms in the case of the ellipse, for something better expressed by the symmetrical relation "conjugate" (First Def I 6). Immediately in Proposition I 16, as if arbitrarily, the upright diameter of the hyperbola is bounded in the same way, given a definite magnitude and becomes "the second diameter." But so far transverse and upright diameters, or transverse and second diameters, are distinct things in the case of the hyperbola and there seems to be little reason for giving this second diameter in magnitude *νομος* has not yet become *φύσις*. That the upright diameter should be given even in position for the hyperbola becomes only very significant with two pairs of propositions—Propositions I 37 and 38 and I 39 and 40—where it is shown that certain properties holding for ordinates to the transverse diameter of the hyperbola and ellipse hold also for the ordinates to their conjugates. But it is only with the final proposition of the first book (I 60) that the magnitude of the hyperbola's second diameter is justified in magnitude as well as position. It is the corresponding diameter of the opposite sections conjugate to the first. And this analogy between the hyperbol-

and ellipse now stands on the threshold of a vast development. For this theorem, coming as a climax to the first book, makes possible the main theme of the second book—the asymptotes, those strange lines all but touching each opposite section (II 2, 13, 14) and forming a single bound between each adjacent pair (II 15, 17), so making the hyperbola an all but closed section, a puckered ellipse, a mouth turned inside out. And in the third book, the fruits of this analogy are gathered as in the especially nice case of Proposition III 15.

Although this translation is literal, we have not hesitated to use such symbols and abbreviations as, without prejudicing any Greek number theory or introducing any modern theory of symbols, would yet make the reading and the mechanic of study easier and at the same time preserve all the rigor of Greek mathematics.

As for the Greek text, we have used Heiberg, and have constantly referred to the *editio princeps* of Halley. In certain instances we have been glad to consult the very excellent French translation of Paul Ver Eecke (Desclée de Brouwer, Bruges, 1923). We have also deferred, at all relevant points, to the English usage of T. L. Heath's translation of Euclid's *Elements*.

EXAMPLES OF ABBREVIATIONS AND SYMBOLS USED

$A=B$ for A is equal to B

$A+B$ for A added to B

$A-B$ for B subtracted from A

$A \ B \ C \ D$ for A is to B as C is to D

rect AB, BC for rectangle AB, BC

sq AB for square on AB

ar for area

pllg for parallelogram

trgl for triangle

quadr for quadrilateral

rect $AB \ BC$ rect $CD \ DE$ comp $AB \ CD \ BC \ DE$
for ratio of rectangle $AB \ BC$ to rectangle CD, DE is
compounded of the ratio of AB to CD and of BC to
 DE

ratio comp $AB \ BC, CD \ DE$ =ratio comp $XY \ YZ,$
 $ZW \ WV$ for ratio compounded of AB to BC and of
 CD to DE is the same as the ratio compounded of XY
to YZ and of ZW to WV

$A>B$ for A is greater than B

$A<B$ for A is less than B

rt angle for right angle

BOOK ONE

APOLLONIUS to EUDEMUS greetings

If you are restored in body and other things go with you to your mind well and good and we too fare pretty well At the time I was with you in Pergamum I observed you were quite eager to be kept informed of the work I was doing in conics And so I have sent you this first book revised and we shall dispatch the others when we are satisfied with them For I don't believe you have forgotten hearing from me how I worked out the plan for these conics at the request of Naucrates the geometer at the time he was with us in Alexandria lecturing and how on arranging them in eight books we immediately communicated them in great haste because of his near departure not revising them but putting down whatever came to us with the intention of a final going over And so finding now the occasion of correcting them one book after another we publish them And since it happened that some others among those frequenting us got acquainted with the first and second books before the revision don't be surprised if you come upon them in a different form

Of the eight books the first four belong to a course in the elements The first book contains the generation of the three sections and of the opposite branches and the principal properties (τα αρχικά συμπεράσματα) in them worked out more fully and universally than in the writings of others The second book contains the properties (τα συμβαίνοντα) having to do with the diameters and axes and also the asymptotes and other things of a general and necessary use for limits of possibility (πρὸς τοὺς διορισμούς) And what I call diameters and what I call axes you will know from this book The third book contains many incredible theorems of use for the construction of solid loci and for limits of possibility of which the greatest part and the most beautiful are new And when we had grasped these we knew that the three line and four line locus had not been constructed by Euclid but only a chance part of it and that not very happily For it was not possible for this construction to be completed without the additional things found by us The fourth book shows in how many ways the sections of a cone intersect with each other and with the circumference of a circle and contains other things in addition none of which has been written up by our predecessors that is in how many points the section of a cone or the circumference of a circle and the opposite branches meet the opposite branches The rest of the books are fuller in treatment For there is one dealing more fully with maxima and minima and one with equal and similar sections of a cone and one with limiting theorems and one with determinate conic problems And so indeed with all of them published those happening upon them can judge them as they see fit Good by e

FIRST DEFINITIONS

1 If from a point a straight line is joined to the circumference of a circle which is not in the same plane with the point, and the line is produced in both directions, and if, with the point remaining fixed, the straight line being rotated about the circumference of the circle returns to the same place from which it began then the generated surface composed of the two surfaces lying vertically opposite one another, each of which increases indefinitely as the generating straight line is produced indefinitely, I call a conic surface, and I call the fixed point the vertex, and the straight line drawn from the vertex to the center of the circle the axis

2 And the figure contained by the circle and by the conic surface between the vertex and the circumference of the circle I call a cone, and the point which is also the vertex of the surface I call the vertex of the cone, and the straight line drawn from the vertex to the center of the circle the axis and the circle the base of the cone

3 I call right cones those having axes perpendicular to their bases and oblique those not having axes perpendicular to their bases

4 Of any curved line which is in one plane I call that straight line the diameter which drawn from the curved line, bisects all straight lines drawn to this curved line parallel to some straight line and I call the end of that straight line (the diameter) situated on the curved line the vertex of the curved line, and I say that each of these parallels is drawn ordinatewise to the diameter (*τεταγμένως ἐπὶ τὴν διάμετρον κατὰ ῥαθίαι*)¹

5 Likewise of any two curved lines lying in one plane I call that straight line the transverse diameter (*διάμετρος πλαγία*) which cuts the two curved lines and bisects all the straight lines drawn to either of the curved lines parallel to some straight line and I call the ends of the diameter situated on the curved lines the vertices of the curved lines and I call that straight line the upright diameter (*διάμετρος ὀρθία*) which lying between the two curved lines, bisects all the straight lines intercepted between the curved lines and drawn parallel to some straight line and I say that each of the parallels is drawn ordinatewise to the diameter

6 The two straight lines each of which being a diameter bisects the straight lines parallel to the other I call the conjugate diameters (*συνζυγεῖς διαμέτροι*) of a curved line and of two curved lines

7 And I call that straight line the axis of a curved line and of two curved lines which being a diameter of the curved line or lines cuts the parallel straight lines at right angles

8 And I call those straight lines the conjugate axes of a curved line and of two curved lines which being conjugate diameters cut the straight lines parallel to each other at right angles


PROPOSITION 1

The straight lines drawn from the vertex of the conic surface to points on the surface are on that surface

Let there be a conic surface whose vertex is the point Λ and let there be

¹We shall follow modern usage and generally call these parallels ordinates

taken some point B on the conic surface and let a straight line ACB be joined
I say that the straight line ACB is on the conic surface



For if possible, let it not be, and let the straight line DE be the line generating the surface, and EF be the circle along which ED is moved. Then if, the point A remaining fixed, the straight line DE is moved along the circumference of the circle EF it will also go through the point B (Def 1), and two straight lines will have the same ends. And this is absurd.

Therefore the straight line joined from A to B cannot not be on the surface. Therefore it is on the surface.


PORISM

It is also evident that if a straight line is joined from the vertex to some point among those within the surface it will fall within the conic surface, and if it is joined to some point among those without it will be outside the surface.

PROPOSITION 2

If on either one of the two vertically opposite surfaces two points are taken and the straight line joining the points does not verge to the vertex then it will fall within the surface and produced it will fall outside.

Let there be a conic surface whose vertex is the point A , and a circle BC



along whose circumference the generating straight line is moved and let two points D and E be taken on either one of the two vertically opposite

and let the joining straight line DE not verge to the point A

I say that the straight line DE will be within the surface, and produced will be without

Let AF and AD be joined and produced Then they will fall on the circumference of the circle (r 1) Let them fall to the points B and C , and let BC be joined Therefore the straight line BC will be within the circle, and so too within the conic surface

Then let a point F be taken at random on DE , and let the straight line AF be joined and produced Then it will fall on the straight line BC , for the triangle BCA is in one plane (Eucl r 2) Let it fall to the point G Since then the point G is within the conic surface, therefore the straight line AG is also within the conic surface (r 1, porism), and so too the point F is within the conic surface Then likewise it will be shown that all the points on the straight line DE are within the surface Therefore the straight line DE is within the surface

Then let DE be produced to H I say then it will fall outside the conic surface

For if possible let there be some point H of it not outside the conic surface and let AH be joined and produced Then it will fall either on the circumference of the circle or within (r 1 and porism) And this is impossible, for it falls on BC produced as for example to the point K Therefore the straight line EH is outside the surface

Therefore the straight line DE is within the conic surface and produced is outside

PROPOSITION 3

If a cone is cut by a plane through the vertex the section is a triangle

Let there be a cone whose vertex is the point A and whose base is the circle BC and let it be cut by some plane through the point A and let it make as sections lines AB and AC on the surface and the straight line BC in the base

I say that ABC is a triangle

For since the line joined from A to B is the common section of the cutting plane and of the surface of the cone therefore AB is a straight line And likewise also AC And BC is also a straight line Therefore ABC is a triangle

If then a cone is cut by some plane through the vertex, the section is a triangle




PROPOSITION 4

If either one of the vertically opposite surfaces is cut by some plane parallel to the circle along which the straight line generating the surface is moved, the plane cut off within the surface will be a circle having its center on the axis, and the figure contained by the circle and the conic surface intercepted by the cutting plane on the side of the vertex will be a cone

Let there be a conic surface whose vertex is the point A and whose circle along which the straight line generating the surface is moved is BC and let it be cut by some plane parallel to the circle BC , and let it make on the surface as a section the line DE

I say that the line DE is a circle having its center on the axis

For let the point F be taken as the center of the circle BC , and let AF be joined. Therefore AF is the axis (Def. 1) and meets the cutting plane. Let it meet it at the point G , and let some plane be produced through AF . Then the



section will be the triangle ABC (1. 3). And since the points D, G, E are points in the cutting plane, and are also in the plane of the triangle ABC , therefore DGE is a straight line (Eucl. xi. 3).

Then let some point H be taken on the line DE and let AH be joined and produced. Then it falls on the circumference BC (1. 1). Let it meet it at K and let GH and FK be joined. And since two parallel planes DE and BC are cut by a plane ABC , their common sections are parallel (Eucl. xi. 16). Therefore the straight line DE is parallel to the straight line BC . Then for the same reason the straight line GH is also parallel to the straight line KF . Therefore

$$FA : AG = FB : DG = FC : GE = FK : GH \quad (\text{Eucl. vi. 4})$$

$$\text{And} \quad BF = KF = FC$$

$$\text{Therefore also} \quad DG = GH = GE \quad (\text{Eucl. v. 9})$$

Then likewise we could show also that all the straight lines falling from the point G on the line DE are equal to each other.

Therefore the line DE is a circle having its center on the axis.

And it is evident that the figure contained by the circle DE and the conic surface cut off by it on the side of the point A is a cone.

And it is therewith proved that the common section of the cutting plane and of the axial triangle (triangle through the axis) is a diameter of the circle.

PROPOSITION 5

If an oblique cone is cut by a plane through the axis at right angles to the base, and is also cut by another plane on the one hand at right angles to the axial and on the other cutting off on the side of the vertex a triangle similar to the

triangle and lying subcontrariwise then the section is a circle, and let such a section be called subcontrary

Let there be an oblique cone whose vertex is the point A and whose base is the circle BC , and let it be cut by a plane through the axis perpendicular to the circle BC , and let it make as a section the triangle ABC (1 3). Then let it also be cut by another plane perpendicular to the triangle ABC and cutting off on the side of the point A the triangle AKG similar to the triangle ABC and lying subcontrariwise, that is, so that the angle AKG is equal to the angle ABC . And let it make as a section on the surface, the line GHA .

I say that the line GHA is a circle.

For let any points H and L be taken on the lines GHA and BC , and from the points H and L let perpendiculars be dropped to the plane through the triangle ABC . Then they will fall to the common sections of the planes (Eucl xi def 6). Let them fall as for example FH and LM . Therefore FH is parallel to LM (Eucl xi 6).

Then let the straight line DFE be drawn through F parallel to BC , and FH is also parallel to LM . Therefore the plane through FH and DE is parallel to the base of the cone (Eucl xi 15). Therefore it is a circle whose diameter is the straight line DE (1 1).

Therefore

$$\text{rect } DF \cdot FF = \text{sq } FH \text{ (Eucl iii 31 and vi 8, porism)}$$

And since ED is parallel to BC angle ADE is equal to angle ABC . And angle AKG is supposed equal to angle ABC . And therefore angle AKG is equal to angle ADE . And the vertical angles at the point F are also equal. Therefore triangle DFG is similar to triangle AKE and therefore

$$EF \cdot FK = GF \cdot FD \text{ (Eucl vi 4)}$$

Therefore

$$\text{rect } EF \cdot FD = \text{rect } AF \cdot FG \text{ (Eucl vi 16)}$$

But it has been shown that

$$\text{sq } FH = \text{rect } FF \cdot FD,$$

and therefore


$$\text{rect } AF \cdot FC = \text{sq } FH$$

Likewise then all the perpendiculars drawn from the line GHA to the straight line GA could also be shown to be equal in square to the rectangle, in each case contained by the segments of the straight line GK .


Therefore the section is a circle whose diameter is the straight line GA .

PROPOSITION 6

If a cone is cut by a plane through the axis and some point is taken on the surface of the cone which is not on a side of the axial triangle and from it is drawn a straight line parallel to some straight line which is a perpendicular from the circumference of the circle to the base of the triangle then it meets the axial triangle, and on being produced to the other side of the surface it will be bisected by the triangle.



Let there be a cone whose vertex is the point A and whose base is the circle BC , and let the cone be cut by a plane through the axis and let it make a common section the triangle ABC (1 3), and from some point M of those on the



circumference let the straight line MN be drawn perpendicular to the straight line BC . Then let some point D be taken on the surface of the cone and through D let the straight line DE be drawn parallel to MN .

I say that the straight line DE produced will meet the plane of the triangle ABC and, if further produced toward the other side of the cone until it meet its surface, will be bisected by the triangle ABC .

Let the straight line AD be joined and be produced. Therefore it will meet the circumference of the circle BC (1 1). Let it meet it at K and from K let the straight line KHL be drawn perpendicular to the straight line BC . Therefore KH is parallel to MN , and therefore to DE (Eucl. 1 9).

Let the straight line AH be joined from A to H . Since then in the triangle AHK the straight line DE is parallel to the straight line HK , therefore DE produced will meet AH . But AH is in the plane of ABC , therefore DE will meet the plane of the triangle ABC .

For the same reasons it also meets AH , let it meet it at F and let DF be produced in a straight line until it meet the surface of the cone. Let it meet it at G .

I say that DF is equal to FG .

For since A, G, L are points on the surface of the cone but also in the plane extended through the straight lines AH, AK, DG, KL which is a triangle through the vertex of the cone (1 3), therefore A, C, L are points on the common section of the cone's surface and of the triangle. Therefore the line through A, G, L is a straight line. Since then in the triangle ALK the straight line DG has been drawn parallel to the base KHL and some straight line AFH has been drawn across them from the point A , therefore


$$KH : HL = DF : FG \text{ (Eucl. VI 2)}$$

But KH is equal to HL since KL is a chord in circle BC perpendicular to the diameter (Eucl. III. 3). Therefore DF is equal to FG .

PROPOSITION 7

If a cone is cut by a plane through the axis and if it is also cut by another plane cutting the plane the base of the cone is in, in a straight line perpendicular either to the base of the axial triangle or to it produced, then the straight lines drawn from the resulting section on the cone's surface, made by the cutting plane, parallel to the straight line perpendicular to the base of the triangle will fall on the common section of the cutting plane and of the axial triangle and further produced to the other side of the section, are bisected by the common section, and if it is a right cone the straight line in the base will be perpendicular to the common section of the cutting plane and of the axial triangle and if oblique, it will not always be perpendicular, but whenever the plane through the axis is perpendicular to the base of the cone

Let there be a cone whose vertex is the point A and whose base is the circle BC and let it be cut by a plane through the axis and let it make as a section




the triangle ABC (I. 3). And let it also be cut by another plane cutting the plane the circle BC is in, in the straight line DE perpendicular either to the straight line BC or to it produced and let it make as a section on the surface of the cone the line DFE . Then the straight line FG is the common section of the cutting plane and of the triangle ABC . And let any point H be taken on the section DFE and let the straight line HK be drawn through H parallel to the straight line DE .

I say that the straight line HK meets the straight line FG , and on being produced to the other side of the section DFE , will be bisected by FG .

For since a cone whose vertex is the point A and whose base is the circle BC has been cut by a plane through its axis, and makes as a section the triangle ABC and some point H on the surface, not on a side of the triangle ABC has been taken and since the straight line DG is perpendicular to the straight line BC therefore the straight line drawn through H parallel to DG that is HK , meets the triangle ABC and if further produced to the other side of the surface, will be bisected by the triangle (I. 6).

Then since the straight line drawn through H parallel to the straight line DE meets the triangle ABC and is in the plane of the section DFE , therefore



it will fall on the common section of the cutting plane and of the triangle ABC . But the straight line FG is the common section of the planes. Therefore the straight line drawn through H parallel to DE will fall on FG , and, if further produced to the other side of the section DFE , will be bisected by the straight line FG .

Then either the cone is a right cone or the axial triangle ABC is perpendicular to the circle BC , or neither.

First let the cone be a right cone. Then the triangle ABC would be perpendicular to the circle BC (Def 3 Eucl xi 18). Since then the plane ABC is perpendicular to the plane BC and the straight line DE has been drawn in one of the planes BC perpendicular to their common section the straight line BC therefore the straight line DE is perpendicular to the triangle ABC (Eucl xi def 4) and therefore to all the straight lines touching it and in the triangle ABC (Eucl xi def 3). And so it is also perpendicular to the straight line FG .

Then let the cone not be a right cone. If now the axial triangle is perpendicular to the circle BC , we could likewise show that DE is perpendicular to FG .


Then let the axial triangle ABC not be perpendicular to the circle BC — I say that DE is not perpendicular to FG . For if possible let it be. And it is also perpendicular to the straight line BC . Therefore DE is perpendicular to both BC and FG , and therefore it will be perpendicular to the plane through BC and FG . But the plane through BC and GF is the triangle ABC and therefore DE is perpendicular to the triangle ABC . And therefore all the planes through it are perpendicular to the triangle ABC . But one of the planes through DE is the circle BC therefore the circle BC is perpendicular to the triangle ABC . And so the triangle ABC will also be perpendicular to the circle BC . And this is not supposed. Therefore the straight line DE is not perpendicular to the straight line FG .

PORISM

Then from this it is evident that the straight line FG is the diameter of the section DFE , since it bisects the straight lines drawn parallel to some straight line DE and that it is possible for some parallels to be bisected by the diameter FG and not be perpendicular

PROPOSITION 8

If a cone is cut by a plane through its axis and is cut by another plane cutting the base of the cone in a straight line perpendicular to the base of the axial triangle, and if the diameter of the resulting section on the surface is either parallel to one of the sides of the triangle or meets one of them beyond the vertex of the cone and the surface of the cone and the cutting plane are produced indefinitely then the section will also increase indefinitely, and some straight line drawn from the section of the cone parallel to the straight line in the base of the cone will cut off from the diameter on the side of the vertex a straight line equal to any given straight line



Let there be a cone whose vertex is the point A and whose base is the circle BC and let it be cut by a plane through its axis and let it make as a section the triangle ABC (1 3). And let it be cut also by another plane cutting the

circle BC in a straight line DE perpendicular to the straight line BC , and let it make as a section on the surface the line DFE . And let the diameter FG of the section DFE be either parallel to the straight line AC or on being produced meet it beyond the point A (r 7 and porism)

I say that, if both the surface of the cone and the cutting plane are produced indefinitely, the section DFE also will increase indefinitely


For let both the surface of the cone and the cutting plane be produced. Then it is evident that also the straight lines AB , AC , FG will be therewith produced. Since the straight line FG is either parallel to AC or produced meets it beyond the point A , therefore the straight lines FG and AC on being produced in the direction of C and G will never meet. Then let them be produced and let some point H be taken at random on the straight line FG , and let the straight line KHL be drawn through the point H parallel to the straight line BC and MHN parallel to DE . Therefore the plane through KL and MN is parallel to the plane through BC and DE (Eucl. XI 15). Therefore the plane $KLMN$ is a circle (r 4).

And since the points D , E , M , N are in the cutting plane and also on the surface of the cone, therefore they are on the common section. Therefore the section DFE has increased to the points M and N . Therefore, with the surface of the cone and the cutting plane increased to the circle $KLMN$, the section DFE has also increased to the points M and N . Then likewise we could show also that if the surface of the cone and the cutting plane are extended indefinitely, the section $MDFEN$ will also increase indefinitely.

And it is evident that some straight line will cut off on straight line FH on the side of point F a straight line equal to any given straight line. For if we lay down the straight line FY equal to the given straight line, and draw a parallel to DE through X , it will meet the section just as the straight line through H was also proved to meet the section in the points M and N . And so some straight line is drawn meeting the section parallel to DE , and cutting off on FG on the side of point H a straight line equal to the given straight line.

PROPOSITION 9

If a cone is cut by a plane meeting both sides of the axial triangle and neither parallel to the base nor situated subcontrariwise, then the section will not be a circle.



Let there be a cone whose vertex is the point A and whose base is the circle BC and let it be cut by some plane neither parallel to the base nor situated subcontrariwise, and let it make as a section on the surface the line DKE .

I say that the line DKE will not be a circle.

For if possible let it be and let the cutting plane meet the base and let the straight line FG be the common section of the planes, and let the point H be the center of the circle BC and let the straight line HG be

drawn from it perpendicular to the straight line FG . And let a plane be extended through GHI and the axis and let it make as sections on the conic surface the straight lines BA and AC (I 1). Since then D, E, G are points in the plane through the line $D\Lambda E$ and also in the plane through the points A, B, C , therefore D, E, G are points on the common section of the planes. Therefore GED is a straight line (Eucl VI 3).

Then let some point Λ be taken on the line $D\Lambda E$, and through Λ let the straight line KL be drawn parallel to the straight line FG , then KM will be equal to ML (I 7). Therefore the straight line DE is the diameter of the circle $D\Lambda LE$ (Def 4). Then let the straight line $NM\Lambda$ be drawn through M parallel to the straight line BC . But ΛL is also parallel to FG . And so the plane through the straight lines NV and ΛM is parallel to the plane through the straight lines BC and FG , that is to the base (Eucl VI 15), and the section will be a circle (I 4). Let it be the circle NKX .

And since the straight line FG is perpendicular to the straight line BG , the straight line KM is also perpendicular to the straight line NX (Eucl VI 10). And so

$$\text{rect } NM \cdot M\Lambda = \text{sq } KM \text{ (Eucl III 31 VI 8, porism)}$$

But

$$\text{rect } DM \cdot ME = \text{sq } \Lambda M,$$

for the line $D\Lambda EL$ is supposed a circle, and the straight line DE is its diameter. Therefore

$$\text{rect } NM \cdot M\Lambda = \text{rect } DM \cdot ME$$

Therefore

$$MN \cdot MD = EM \cdot M\Lambda \text{ (Eucl VI 16)}$$

Therefore triangle DMN is similar to triangle ΛME (Eucl VI 6, VI def 1) and angle DNM is equal to angle $ME\Lambda$. But angle DNM is equal to angle ABC , for the straight line $N\Lambda$ is parallel to the straight line BC . And therefore angle ABC is equal to angle $M\Lambda X$. Therefore the section is subcontrary (I 5). And this is not supposed. Therefore the line $D\Lambda E$ is not a circle.


PROPOSITION 10

If two points are taken on the section of a cone the straight line joining the two points will fall within the section, and produced in a straight line it will fall outside.

Let there be a cone whose vertex is the point A and whose base is the circle BC , and let it be cut by a plane through the axis and let it make as a section the triangle ABC (I 3). Then let it also be cut by another plane, and let it make as a section on the surface of the cone the line DEF , and let two points G and H be taken on the line DEF .

I say that the straight line joining the two points G and H will fall within the line DEF , and produced in a straight line it will fall outside.

For since a cone, whose vertex is the point A and whose base is the circle BC , has been cut by a plane through the axis and some points G and H have been taken on its surface which are not on a side of the axial



triangle and since the straight line joining G and H does not verge to the point A therefore the straight line joining G and H will fall within the cone and produced in a straight line it will fall outside (1 2), consequently also outside the section DFE

PROPOSITION 11

If a cone is cut by a plane through its axis and also cut by another plane cutting the base of the cone in a straight line perpendicular to the base of the axial triangle, and if further the diameter of the section is parallel to one side of the axial triangle then any straight line which is drawn from the section of the cone to its diameter parallel to the common section of the cutting plane and of the cone's base will equal in square the rectangle contained by the straight line cut off by it on the diameter beginning from the section's vertex and by another straight line which has the ratio to the straight line between the angle of the cone and the vertex of the section that the square on the base of the axial triangle has to the rectangle contained by the remaining two sides of the triangle And let such a section be called a parabola (παράβολη)

Let there be a cone whose vertex is the point A , and whose base is the circle BC , and let it be cut by a plane through its axis and let it make as a section the triangle ABC (1 3) And let it also be cut by another plane cutting the base of the cone in the straight line DE perpendicular to the straight line BC and let it make as a section on the surface of the cone the line DFE , and let the diameter of the section FG (1 7 and def 4) be parallel to one side AC of the axial triangle And let the straight line FH be drawn from the point F perpendicular to the straight line FG and let it be contrived that

$$\text{sq } BC \text{ rect } BA \ AC \quad FH \ FA$$

And let some point K be taken at random on the section and through K let the straight line KL be drawn parallel to the straight line DE

I say that $\text{sq } KL = \text{rect } HF \ FL$

For let the straight line MN be drawn through L parallel to the straight line BC And the straight line DE is also parallel to the straight line KL Therefore the plane through KL and MN is parallel to the plane through BC and DE (Eucl xi 15) that is to the base of the cone Therefore the plane through KL and MN is a circle whose diameter is MN (1 4) And KL is perpendicular to MN since DE is also perpendicular to BC (Eucl xi 10) Therefore

$$\text{rect } ML \ LN = \text{sq } KL \text{ (Eucl iii 31 vi 8 porism)}$$

And since

$$\text{sq } BC \text{ rect } BA \ AC \quad HF \ FA$$

and

therefore $\text{sq } BC \text{ rect } BA \ AC \text{ comp } BC \ CA \ BC \ BA$ (Eucl vi 23),


$$HF \ FA \text{ comp } BC \ CA \ BC \ BA$$

But

$$BC \ CA \ MN \ NA \ ML \ LF \text{ (Eucl vi 4),}$$

and

$$BC \ BA \ MN \ MA \ LM \ MF \ AL \ FL \text{ (Eucl vi 2)}$$



Therefore

$$HF \text{ FA comp } ML \text{ LF, NL FA}$$

But

$$\text{rect } ML, LN \text{ rect } LF, FA \text{ comp } ML \text{ LF, LN FA (Eucl vi 23)}$$

Therefore

$$HF \text{ FA rect } ML, LN \text{ rect } LF, FA$$

But with the straight line FL taken as common height

$$HF \text{ FA rect } HF, FL \text{ rect } LF, FA \text{ (Eucl vi 1),}$$

therefore

$$\text{rect } ML, LN \text{ rect } LF, FA \text{ rect } HF, FL \text{ rect } LF, FA \text{ (Eucl v 11)}$$

Therefore

$$\text{rect } ML, LN = \text{rect } HF, FL \text{ (Eucl v 9)}$$

But

$$\text{rect } ML \text{ LN} = \text{sq } KL$$

therefore also

$$\text{sq } KL = \text{rect } HF \text{ FL}$$

And let such a section be called a parabola, and let HIF be called the straight line to which the straight lines drawn ordinatewise to the diameter FG are applied in square ($\pi\alpha\rho\ \eta\nu\ \delta\upsilon\nu\alpha\tau\alpha\iota\ \alpha\iota\ \kappa\alpha\tau\alpha\gamma\omicron\mu\epsilon\nu\alpha\iota\ \tau\epsilon\tau\alpha\gamma\mu\epsilon\nu\omega\varsigma\ \epsilon\pi\iota\ \tau\eta\nu\ ZH\ \delta\iota\alpha\mu\epsilon\tau\rho\omega\nu$), and let it also be called the upright side ($\delta\omicron\rho\theta\iota\alpha$)¹

PROPOSITION 12

If a cone is cut by a plane through its axis and also by another plane cutting the base of the cone in a straight line perpendicular to the base of the axial triangle, and if the diameter of the section produced meets one side of the axial triangle beyond the vertex of the cone then any straight line which is drawn from the section to its diameter parallel to the common section of the cutting plane and of the cone's base will equal in square some area applied to a straight line to which the straight line added along the diameter of the section and subtending the exterior angle of the triangle has the ratio that the square on the straight line drawn from the cone's vertex to the triangle's base parallel to the section's diameter has to the rectangle contained by the sections of the base which this straight line makes when drawn, this area having as breadth the straight line cut off on the diameter beginning from the section's vertex by this straight line from the section to the diameter and exceeding ($\upsilon\pi\epsilon\rho\beta\acute{\alpha}\lambda\lambda\omicron\nu$) by a figure ($\epsilon\iota\delta\omicron\varsigma$) similar and similarly situated to the rectangle contained by the straight line subtending the exterior angle of the triangle and by the parameter. And let such a section be called an hyperbola ($\upsilon\pi\epsilon\rho\beta\omicron\lambda\eta$).

Let there be a cone whose vertex is the point A and whose base is the circle BC , and let it be cut by a plane through its axis, and let it make as a section the triangle ABC (1 3). And let it also be cut by another plane cutting the base of the cone in the straight line DE perpendicular to BC the base of the triangle ABC , and let it make as a section on the surface of the cone the line DFE , and

¹The Greek of the phrase the straight line to which the straight lines drawn ordinatewise to the diameter are applied in square that is $\eta\ \pi\alpha\rho\ \eta\ \delta\ \alpha\upsilon\tau\alpha\iota\ \alpha\iota\ \kappa\alpha\tau\alpha\gamma\omicron\mu\epsilon\nu\alpha\iota\ \tau\epsilon\tau\alpha\gamma\mu\epsilon\nu\omega\varsigma\ \epsilon\pi\iota\ \tau\eta\nu\ \delta\ \delta\iota\alpha\mu\epsilon\tau\rho\omega$ soon becomes abbreviated to $\eta\ \pi\alpha\rho\ \eta\ \delta\epsilon\ \alpha\upsilon\tau\alpha\ \alpha\iota\ \kappa\alpha\tau\alpha\gamma\omicron\mu\epsilon\nu\alpha\iota$ and to $\eta\ \pi\alpha\rho\ \eta\ \delta\epsilon\ \alpha\upsilon\tau\alpha$. We shall translate these abbreviations by the word parameter. And we shall later on after proposition XIV shorten the long expression to the parameter of the ordinates to the diameter.

The Latin translation of $\delta\omicron\rho\theta\iota\alpha$ ($\tau\lambda\epsilon\nu\rho\acute{\alpha}$) is *latus rectum* which has become an English term too.

let FG the diameter of the section (1 7 and def 4) when produced meet AC one side of the triangle ABC beyond the vertex of the cone at the point H . And let the straight line AA be drawn through A parallel to the diameter of the section FG , and let it cut BC . And let the straight line FL be drawn from F perpendicular to FG , and let it be contrived that

$$\text{sq } AA \text{ rect } BK \text{ } KC \text{ } FH \text{ } FL$$

And let some point M be taken at random on the section, and through M let the straight line MN be drawn parallel to DE , and through N let the straight line VOX be drawn parallel to FL . And let the straight line HL be joined and produced to Y , and let the straight lines LO and XP be drawn through L and X parallel to FN .

I say that MN is equal in square to the parallelogram $F\Lambda$ which is applied to FL having FN as breadth and exceeding by a figure LY similar to the rectangle contained by HF and FL .

For let the straight line RNS be drawn through N parallel to BC and NM is also parallel to DE . Therefore the plane through MN and RS is parallel to the plane through BC and DE that is to the base of the cone (Eucl xi 15). Therefore if the plane is produced through MN and RS , the section will be a circle whose diameter is the straight line RNS (1 4). And MN is perpendicular to it. Therefore

$$\text{rect } RN, NS = \text{sq } MN$$

And since

$$\text{sq } AA \text{ rect } BK \text{ } KC \text{ } FH \text{ } FL,$$

and

$\text{sq } AA \text{ rect } BK \text{ } KC \text{ comp } AK \text{ } KC \text{ } AK \text{ } KB$ (Eucl vi 23),
therefore also

$$FH \text{ } FL \text{ comp } AK \text{ } KC \text{ } AK \text{ } KB$$

But

$$AK \text{ } KC \text{ } HG \text{ } GC \text{ } HN \text{ } NS \text{ (Eucl vi 4),}$$

and

$$AA \text{ } KB \text{ } FG \text{ } GB \text{ } FN \text{ } NR$$

Therefore

$$HF \text{ } FL \text{ comp } HN \text{ } NS \text{ } FN \text{ } NR$$

And

$$\text{rect } HN \text{ } NF \text{ rect } SN \text{ } NR \text{ comp } HN \text{ } NS \text{ } FN \text{ } NR \text{ (Eucl vi 23)}$$

Therefore also

$$\text{rect } HN \text{ } NF \text{ rect } SN \text{ } NR \text{ } HF \text{ } FL \text{ } HN \text{ } NY \text{ (Eucl vi 4)}$$

But, with the straight line FN taken as common height

$$HN \text{ } NY \text{ rect } HN \text{ } NF \text{ rect } FN \text{ } NY \text{ (Eucl vi 1)}$$

Therefore also


$$\text{rect } HN \text{ } NF \text{ rect } SN \text{ } NR \text{ rect } HN \text{ } NF \text{ rect } XN \text{ } NF \text{ (Eucl v 11)}$$

Therefore

$$\text{rect } SN \text{ } NR = \text{rect } XN \text{ } NF \text{ (Eucl v 9)}$$

But it was shown

$$\text{sq } MN = \text{rect } SN \text{ } NR,$$



therefore also

$$\text{sq } MN = \text{rect } XN \text{ } NF$$


But the rectangle contained by YN and NF is the parallelogram YF . Therefore the straight line MN is equal in square to YF which is applied to the straight line FL , having FN as breadth and exceeding by the parallelogram LX similar to the rectangle contained by HF and FL (Eucl vi 24)

And let such a section be called an hyperbola, and let LF be called the straight line to which the straight lines drawn ordinatewise to FG are applied in square, and let the same straight line also be called the upright side, and the straight line FH the transverse side

PROPOSITION 13

If a cone is cut by a plane through its axis, and is also cut by another plane on the one hand meeting both sides of the axial triangle and on the other extended neither parallel to the base nor subcontrariwise, and if the plane the base of the cone is in and the cutting plane meet in a straight line perpendicular either to the base of the axial triangle or to it produced then any straight line which is drawn from the section of the cone to the diameter of the section parallel to the common section of the planes will equal in square some area applied to a straight line to which the diameter of the section has the ratio that the square on the straight line drawn from the cone's vertex to the triangle's base parallel to the section's diameter has to the rectangle contained by the intercepts of this straight line (on the base) from the sides of the triangle an area having as breadth the straight line cut off on the diameter beginning from the section's vertex by this straight line from the section to the diameter and deficient (ἐλλείπων) by a figure similar and similarly situated to the rectangle contained by the diameter and parameter. And let such a section be called an ellipse (ἐλλειψις)

Let there be a cone whose vertex is the point A and whose base is the circle BC and let it be cut by a plane through its axis, and let it make as a section



the triangle ABC . And let it also be cut by another plane on the one hand meeting both sides of the axial triangle and on the other extended neither parallel to the base of the cone nor subcontrariwise, and let it make as a section on the surface of the cone the line DE . And let the common section of the cut

ting plane and of the plane the base of the cone is in, be the straight line FG perpendicular to the straight line BC , and let the diameter of the section be the straight line ED (i 7 and Def 4) And let the straight line EH be drawn from E perpendicular to ED and let the straight line IA be drawn through A parallel to ED and let it be contrived that

$$\text{sq } AA \text{ rect } BK \ AC \ DE \ EH$$

And let some point L be taken on the section and let the straight line LM be drawn through L parallel to FG

I say that the straight line LM is equal in square to some area which is applied to EH , having EM as breadth and deficient by a figure similar to the rectangle contained by DE and EH

For let the straight line DH be joined and on the one hand let the straight line MAN be drawn through M parallel to HE and on the other let the straight lines HN and VO be drawn through H and X parallel to EM and let the straight line PMR be drawn through M parallel to BC

Since then PR is parallel to BC and LM is also parallel to FG therefore the plane through LM and PR is parallel to the plane through FG and BC that is to the base of the cone (Eucl xi 15) If therefore a plane is extended through LM and PR the section will be a circle whose diameter is PR (i 4) And LM is perpendicular to it Therefore

$$\text{rect } PM \ MR = \text{sq } LM$$

And since

$$\text{sq } AA \text{ rect } BK \ AC \ DE \ EH$$

and

$$\text{sq } AK \text{ rect } BK \ AC \text{ comp } AA \ KB \ AA \ AC \text{ (Eucl vi 23),}$$

but

$$AA \ KB \ EG \ GB \ EM \ MP \text{ (Eucl vi 4)}$$

and

$$AK \ AC \ DG \ GC \ DV \ MR$$

therefore

$$DE \ EH \text{ comp } EM \ MP \ DV \ MR$$

But

$$\text{rect } EM \ MD \text{ rect } PM \ MR \text{ comp } EM \ MP \ DV \ MR \text{ (Eucl vi 23)}$$

Therefore

$$\text{rect } EM \ MD \text{ rect } PM \ MR \ DE \ EH \ DV \ MX \text{ (Eucl vi 4)}$$

And with the straight line ME taken as common height

$$DV \ MX \text{ rect } DV \ ME \text{ rect } XM \ ME \text{ (Eucl vi 1)}$$

Therefore also

$$\text{rect } DV, ME \text{ rect } PM \ MR \text{ rect } DV \ ME \text{ rect } XM \ ME \text{ (Eucl v 11)}$$

Therefore

$$\text{rect } PM \ MR = \text{rect } VM \ ME \text{ (Eucl v 9)}$$

But it was shown

$$\text{rect } PM \ MR = \text{sq } LM$$

therefore also

$$\text{rect } VM \ ME = \text{sq } LM$$

Therefore the straight line LM is equal in square to the parallelogram MO which is applied to the straight line HE having EM as breadth and deficient by the figure OV similar to the rectangle contained by DE and EH (Eucl vi 24)

And let such a section be called an ellipse, and let EH be called the straight line to which the straight lines drawn ordinatewise to DE are applied in square, and let the same straight line also be called the upright side, and the straight line ED the transverse side

PROPOSITION 14

If the vertically opposite surfaces are cut by a plane not through the vertex the section on each of the two surfaces will be that which is called the hyperbola and the diameter of the two sections will be the same straight line and the straight lines to which the straight lines drawn to the diameter parallel to the straight line in the cone's base are applied in square, are equal and the transverse side of the figure, that between the vertices of the sections, is common And let such sections be called opposite (*ἀντικείμεναι*)

Let there be the vertically opposite surfaces whose vertex is the point A and let them be cut by a plane not through the vertex and let it make as sections on the surface the lines DEF and GHA

I say that each of the two sections DEF and GHA is the so-called hyperbola

For let there be the circle $BDCF$ along which the line generating the surface moves and let the plane λGOK be extended parallel to it on the vertically opposite surface and the straight lines FD and GA are common sections of the sections GHA and FED , and of the circles (1 4) Then they will be parallel (Eucl xi 16) And let the straight line LAU be the axis of the conic surface and the points L and U be the centers of the circles and let a straight line drawn from L perpendicular to the straight line FD be produced to the points B and C and let a plane be produced through the straight line BC and the axis Then it will make as sections in the circles the parallel straight lines λO and BC (Eucl xi 16) and on the surface the straight lines BAO and CAY (1 1 and Def 4)


Then the straight line λO will be perpendicular to the straight line GK , since the straight line BC is also perpendicular to the straight line FD and each of the two is parallel to the other (Eucl xi 10) And since the plane through the axis meets the sections in the points M and N within the lines it is clear that the plane also cuts the lines Let it cut them at H and E , therefore $M E H$ and N are points on the plane through the axis and in the plane the lines are in therefore the line $MEHN$ is a straight line (Eucl xi 3) It is also evident both that $\lambda H A$, and C are in a straight line and B, E, A , and O also For they are both on the conic surface and in the plane through the axis (1 1)

Let then the straight lines HR and EP be drawn from H and E perpendicular to HE and let the straight line SAT be drawn through A parallel to $MEHN$ and let it be contrived that

$$HE \cdot EP = \text{sq } AS = \text{rect } BS \cdot SC,$$

and

$$EH \cdot HR = \text{sq } AT = \text{rect } OT, T\lambda$$



Since then a cone whose vertex is the point A and whose base is the circle BC , has been cut by a plane through its axis and it has made as a section the triangle ABC and it has also been cut by another plane cutting the base of the cone in the straight line DUF perpendicular to the straight line BC and it has made as a section on the surface the line DEF , and the diameter ME produced has met one side of the axial triangle beyond the vertex of the cone and through the point A the straight line AS has been drawn parallel to the diameter of the section EM , and from E the straight line EP has been drawn perpendicular to the straight line EM , and

$$EH \cdot EP = \text{sq } AS \text{ rect } BS, SC,$$

therefore the section DEF is an hyperbola (r 12) and EP is the straight line to which the straight lines drawn ordinatewise to EM are applied in square and the straight line HE is the transverse side of the figure. And likewise GHA is also an hyperbola whose diameter is the straight line HN and whose straight line to which the straight lines drawn ordinatewise to HN are applied is HR , and the transverse side of whose figure is HE

I say that the straight line HR is equal to the straight line EP

For since BC is parallel to AO ,

$$AS : SC = AT : TX$$

and

$$AS : SB = AT : TO$$

But

$$\text{sq } AS \text{ rect } BS, SC \text{ comp } AS : SC = AS : SB \text{ (Eucl vi 23)}$$

and

$$\text{sq } AT \text{ rect } XT, TO \text{ comp } AT : TX = AT : TO,$$

therefore

$$\text{sq } AS \text{ rect } BS : SC = \text{sq } AT \text{ rect } XT, TO$$

Also

$$\text{sq } AS \text{ rect } BS : SC = HE : EP,$$

and

$$\text{sq } AT \text{ rect } XT : TO = HE : HR$$

Therefore also

$$HE : EP = EH : HR \text{ (Eucl v 11)}$$

Therefore

$$EP = HR \text{ (Eucl v 9)}$$

PROPOSITION 15


If in an ellipse a straight line drawn ordinatewise from the midpoint of the diameter, is produced both ways to the section and if it is contrived that as the straight line so produced is to the diameter so is the diameter to some straight line then any straight line which is drawn, from the section to the straight line produced parallel to the diameter, will equal in square the area applied to this third proportional and having as breadth the straight line cut off by it beginning from the section and deficient by a figure similar to the rectangle contained by the straight line to which the straight lines are drawn and by the parameter and if further produced to the other side of the section will be bisected by the straight line to which it has been drawn

Let there be an ellipse whose diameter is the straight line AB , and let AB be bisected at the point C and through C let the straight line DCE be drawn

ordinatewise and produced both ways to the section and from the point D let the straight line DF be drawn perpendicular to DE And let it be contrived that

$DE \cdot AB = AB \cdot DF$

And let some point G be taken on the section, and through G let the straight line GH be drawn parallel to AB , and let FF be joined and through H let the straight line HL be drawn parallel to DF , and through F and L let the straight lines FK and LM be drawn parallel to HD



I say that the straight line GH is equal in square to the area DL

which is applied to the straight line DF having as breadth the straight line DH and deficient by a figure LF similar to the rectangle contained by FD and DF

For let AV be the parameter of the ordinates to AB , and let BV be joined, and through G let the straight line GY be drawn parallel to DE , and through X and C let the straight lines XO and CP be drawn parallel to AN , and through N , O and P let the straight lines NV , OS and TP be drawn parallel to AB Therefore

$$\text{sq } DC = \text{ar } AP, \text{ sq } GX = \text{ar } AO \text{ (I 13)}$$

And since

$$BA \cdot AN = BC \cdot CP = PT \cdot TN \text{ (I ucl vi 4),}$$

and

$$BC = CA = TP,$$

and

$$CP = TA,$$

therefore

$$\text{ar } AP = \text{ar } TR,$$

and

$$\text{ar } VT = \text{ar } TU$$

Since also

$$\text{ar } OT = \text{ar } OR \text{ (Eucl I 43),}$$

and area NO is common therefore

$$\text{ar } TU = \text{ar } NS$$

But

$$\text{ar } TU = \text{ar } TX$$

and TS is common Therefore

$$\text{ar } NP = \text{ar } PA = \text{ar } AO + \text{ar } PO$$

and so

$$\text{ar } PA - \text{ar } AO = \text{ar } PO$$

Also

$$\text{ar } AP = \text{sq } CD, \text{ ar } NO = \text{sq } XG,$$

and

$$\text{ar } OP = \text{rect } OS, SP$$

therefore

$$\text{sq } CD - \text{sq } GX = \text{rect } OS, SP$$

Since also the straight line DE has been cut into equal parts at C , and into unequal parts at H therefore

$$\text{rect } EH, HD + \text{sq } CH = \text{sq } CD \text{ (Eucl II 5),}$$

or

$$\text{rect } EH, HD + \text{sq } XG = \text{sq } CD$$

Therefore

$$\text{sq } CD - \text{sq } XG = \text{rect } EH, HD,$$

but

$$\text{sq } CD - \text{sq } XG = \text{rect } OS, SP,$$

therefore

$$\text{rect } EH, HD = \text{rect } OS, SP$$

And since

$$DE \text{ } 4B \text{ } AB \text{ } DF,$$

therefore

$$DE \text{ } DF \text{ } \text{sq } DE \text{ } \text{sq } AB \text{ (Eucl VI 20),}$$

that is

$$DE \text{ } DF \text{ } \text{sq } CD \text{ } \text{sq } CB \text{ (Eucl V 15),}$$

And

$$\text{rect } PC, CA = \text{rect } PC, CB = \text{sq } CD \text{ (I 13),}$$

and since

$$DE \text{ } DF \text{ } EH \text{ } HL \text{ (Eucl VI 4),}$$

or

$$DE \text{ } DF \text{ } \text{rect } EH, HD \text{ } \text{rect } DH, HL \text{ (Eucl VI 1),}$$

and since

$$DE \text{ } DF \text{ } \text{rect } PC, CB \text{ } \text{sq } CB,$$

and

$$\text{rect } PC, CB \text{ } \text{sq } CB \text{ } \text{rect } OS, SP \text{ } \text{sq } OS^1$$

therefore also

$$\text{rect } EH, HD \text{ } \text{rect } DH, HL \text{ } \text{rect } OS, SP \text{ } \text{sq } OS$$

And

$$\text{rect } EH, HD = \text{rect } OS, SP$$

therefore

$$\text{rect } DH, HL = \text{sq } OS = \text{sq } GH$$

Therefore the straight line GH is equal in square to the area DL which is applied to the straight line DF deficient by a figure FL similar to the rectangle contained by ED and DF (Eucl VI 24)

I say then that also if produced to the other side of the section the straight line GH will be bisected by the straight line DE

For let it be produced and let it meet the section at W , and let the straight line WY be drawn through Y parallel to GX , and through Y let the straight line YZ be drawn parallel to AN . And since

$$GX = WY,$$

therefore also

$$\text{sq } GX = \text{sq } WY$$

¹This follows from the proportions

$$\frac{PC}{PC} = \frac{CB}{CB} = \frac{PS}{rect \ PC \ CB} = \frac{OS}{\text{sq } OS} \text{ (Eucl VI 4)}$$

and

$$\frac{PC}{PS} = \frac{CB}{OS} = \frac{rect \ PC \ CB}{\text{sq } OS} \text{ (Eucl VI 1)}$$

and

$$\frac{PS}{PS} = \frac{OS}{OS} = \frac{\text{rect } PS \ OS}{\text{sq } OS} \text{ (Eucl VI 1)}$$

But

$$\text{sq } GX = \text{rect } A\lambda, \lambda O \text{ (I 13),}$$

and

$$\text{sq } W\lambda = \text{rect } A\gamma, \gamma Z \text{ (I 13)}$$

Therefore

$$O\gamma \ Z\lambda \quad \gamma A \ A\gamma \text{ (Eucl vi 16)}$$

And

$$O\lambda \ Z\lambda \quad \lambda B \ B\gamma \text{ (Eucl vi 4),}$$

therefore also

$$\gamma A \ A\lambda \quad \lambda B \ B\gamma$$

And *separando*

$$\gamma\lambda \ A\lambda \quad \lambda X \ B\gamma \text{ (Eucl v 17)}$$

Therefore

$$A\gamma = \lambda B$$

And also

$$AC = CB,$$

therefore also the remainders

$$XC = C\gamma,$$

and so also


$$GH = HW$$

Therefore the straight line HG , produced to the other side of the section, is bisected by the straight line DH

PROPOSITION 16

If through the midpoint of the transverse side of the opposite sections a straight line be drawn parallel to a straight line drawn ordinatewise it will be a diameter of the opposite sections conjugate to the diameter just mentioned

Let there be the opposite sections whose diameter is the straight line AB ,



and let AB be bisected at C and through C let the straight line CD be drawn parallel to a straight line drawn ordinatewise

I say that the straight line CD is a diameter conjugate to AB

For let the straight lines AE and BF be the parameters and let the straight

lines AF and BE be joined and produced, and let some point G be taken at random on either section and through G let the straight line GH be drawn parallel to AB , and from G and H let the straight lines GK and HL be drawn ordinatewise, and through K and L let the straight lines KM and LN be drawn parallel to AE and BF . Since then

$$GA = HL \text{ (Eucl. I 34),}$$

therefore also

$$\text{sq } GA = \text{sq } HL,$$

But

$$\text{sq } GA = \text{rect } AK, KM \text{ (I 12),}$$

and

$$\text{sq } HL = \text{rect } BL, LN \text{ (I 12),}$$

therefore

$$\text{rect } AK, KM = \text{rect } BL, LN$$

And since

$$AE = BF,$$

therefore

$$AE \cdot AB = BF \cdot BA \text{ (Eucl. V 7)}$$

But

$$AE \cdot AB = MA \cdot AB \text{ (Eucl. VI 4),}$$

and as

$$BF \cdot BA = NL \cdot LA \text{ (Eucl. VI 4)}$$

And therefore

$$MA \cdot AB = NL \cdot LA$$

But, with AA taken as common height,

$$MA \cdot AB = \text{rect } MA, AB = \text{rect } BK, KA,$$

and, with BL taken as common height

$$NL \cdot LA = \text{rect } NL, LB = \text{rect } AL, LB$$

And therefore

$$\text{rect } MA, AB = \text{rect } BK, KA = \text{rect } NL, LB = \text{rect } AL, LB$$

And alternately

$$\text{rect } MA, KA = \text{rect } NL, LB = \text{rect } BK, KA = \text{rect } AL, LB \text{ (Eucl. V 16)}$$

And

$$\text{rect } AK, KM = \text{rect } BL, LN,$$

therefore

$$\text{rect } BK, KA = \text{rect } AL, LB,$$

therefore

$$AK = LB^1$$

But also

$$AC = CB,$$

and therefore

$$KC = CL,$$

and so also

$$GX = XH$$

¹The intermediary steps to this conclusion are as follows. If

$$\text{rect } BK, KA = \text{rect } AL, LB$$

then

$$BK \cdot LB = AL \cdot KA$$

or

$$BA + AK \cdot LB = BA + LB \cdot AK$$

and componendo

$$BA + AK + LB \cdot LB = BA + LB + AK \cdot AK$$

Therefore the straight line GH has been bisected by the straight line XCD and is parallel to the straight line AB . Therefore the straight line XCD is a diameter and conjugate to the straight line AB (Defs. 4, 6)

SECOND DEFINITIONS

9 Let the midpoint of the diameter of both the hyperbola and the ellipse be called the center of the section, and let the straight line drawn from the center to meet the section be called the radius of the section.

10 And likewise let the midpoint of the transverse side of the opposite sections be called the center.

11 And let the straight line drawn from the center parallel to an ordinate, being a mean proportional to the sides of the figure ($\tau\delta$ $\epsilon\iota\delta\omicron\varsigma$) and bisected by the center, be called the second diameter.


PROPOSITION 17

If in a section of a cone a straight line is drawn from the vertex of the line, and parallel to an ordinate, it will fall outside the section (Cf. Eucl. III. 16)

Let there be a section of a cone whose diameter is the straight line AB .

I say that the straight line drawn from the vertex that is from the point A parallel to an ordinate will fall outside the section.

For if possible let it fall within as AC . Since then a point C has been taken at random on a section of a cone, therefore the straight line drawn from the point C within the section parallel to an ordinate will meet the diameter AB and will be bisected by it (17). Therefore the straight line AC produced will be bisected by the straight line AB . And this is absurd. For the straight line AC if produced will fall outside the section (10). Therefore the straight line drawn from the point A parallel to an ordinate will not fall within the line; therefore it will fall outside and therefore it is tangent to the section.




PROPOSITION 18

If a straight line meeting a section of a cone and produced both ways falls outside the section and some point is taken within the section and through it a parallel to the straight line meeting the section is drawn, the parallel so drawn if produced both ways will meet the section.

Let there be a section of a cone and the straight line APB meeting it and let it fall when produced both ways outside the section. And let some point C be taken within the section and through C let the straight line CD be drawn parallel to the straight line AB .

I say that the straight line CD produced both ways will meet the section.

For let some point E be taken on the



ection and let the straight line EF be joined. And since the straight line AB is parallel to CD and some straight line EF meets AB , therefore CD produced will also meet EF . And if it meets EF between the points E and F , it is evident that it also meets the section, but if beyond the point F , that it will first meet the section. Therefore CD produced to the side of points D and E meets the section. Then likewise we could show that, produced to the side of points F and B it also meets it. Therefore the straight line CD produced both ways will meet the section.


PROPOSITION 19

In every section of a cone, any straight line drawn from the diameter parallel to the ordinate, will meet the section.

Let there be a section of a cone whose diameter is the straight line AB , and let some point B be taken on the diameter, and through B let the straight line BC be drawn parallel to an ordinate.

I say that the straight line BC produced will meet the section.

For let some point D be taken on the section. But A is also on the section, therefore the straight line joined from A to D will fall within the section (i 10). And since the straight line drawn from A parallel to an ordinate falls outside the section (i 17), and the straight line AD meets it and the straight line BC is parallel to the ordinate therefore BC will also meet AD . And if it meets AD between the points A and D it is evident that it will also meet the section, but if beyond point D as at E that it will first meet the section. Therefore the straight line drawn from B parallel to an ordinate will meet the section.



PROPOSITION 20

If in a parabola two straight lines are dropped ordinatewise to the diameter, the squares on them will be to each other as the straight lines cut off by them on the diameter beginning from the vertex¹ are to each other.

Let there be a parabola whose diameter is the straight line AB and let some points C and D be taken on it and from the points C and D let the straight lines CE and DF be dropped ordinatewise to AB .

I say that

$$\text{sq } DF : \text{sq } CE :: FA : AE$$

For let AG be the parameter therefore

$$\text{sq } DF = \text{rect } FA, AG,$$

and


$$\text{sq } CE = \text{rect } EA, AG \text{ (i 11)}$$

Therefore

$$\text{sq } DF : \text{sq } CE :: \text{rect } FA, AG : \text{rect } EA, AG$$

But

$$\text{rect } FA, AG : \text{rect } EA, AG :: FA : AE \text{ (Eucl vi 1),}$$



¹These are usually called *abscissas* from the Latin *abscindere*, to cut off


and therefore

$$\text{sq } DF \text{ sq } CE \quad FA \quad AE$$


PROPOSITION 21

If in an hyperbola or ellipse or in the circumference of a circle straight lines are dropped ordinatewise to the diameter the squares on them will be to the areas contained by the straight lines cut off by them beginning from the ends of the transverse side of the figure, as the upright side of the figure is to the transverse and to each other as the areas contained by the straight lines cut off (abscissas), as we have said

Let there be an hyperbola or ellipse or circumference of a circle whose diam



eter is AB and whose parameter is the straight line AC and let the straight lines DE and FG be dropped ordinatewise to the diameter



I say that

$$\text{sq } FG \text{ rect } AG \quad GB \quad AC \quad AB$$

and

$$\text{sq } FG \text{ sq } DE \text{ rect } AG \quad GB \text{ rect } AE \quad EB$$

For let the straight line BC determining the figure be joined and through E and G let the straight lines EH and GK be drawn parallel to the straight line AC . Therefore

$$\begin{aligned} \text{sq } FG &= \text{rect } KG \text{ } GA \\ \text{sq } DE &= \text{rect } HE \text{ } EA \text{ (I' 12, 13)} \end{aligned}$$

And since

$$KG \text{ } GB \text{ } CA \text{ } AB,$$

and with AG taken as common height

$$KG \text{ } GB \text{ } \text{rect } KG \text{ } GA \text{ } \text{rect } BG \text{ } GA,$$

therefore

$$CA \text{ } AB \text{ } \text{rect } KG \text{ } GA \text{ } \text{rect } BG \text{ } GA$$

or

$$CA \text{ } AB \text{ } \text{sq } FG \text{ } \text{rect } BG \text{ } GA$$

Then also for the same reasons

$$CA \text{ } AB \text{ } \text{sq } DE \text{ } \text{rect } BE \text{ } EA$$

And therefore



$$\text{sq } FG \text{ } \text{rect } BG \text{ } GA \text{ } \text{sq } DE \text{ } \text{rect } BE \text{ } EA$$

alternately

$$\text{sq } FG \text{ } \text{sq } DE \text{ } \text{rect } BG \text{ } GA \text{ } \text{rect } BE \text{ } EA^1$$

PROPOSITION 22

If a straight line cuts a parabola or hyperbola in two points not meeting the diameter inside it will, if produced meet the diameter of the section outside the section. Let there be a parabola or hyperbola whose diameter is the straight line AB , and let some straight line cut the section in two points C and D .



I say that the straight line DC , if produced, will meet the straight line AB outside the section.

For let the straight lines CE and DB be dropped ordinatewise from C and D , and first let the section be a parabola. Since then in the parabola

$$\text{sq } CE \text{ } \text{sq } DB \text{ } EA \text{ } AB \text{ (I 20)}$$

and

$$EA > AB,$$

¹Eutocius commenting says It is to be noted that the parameter that is the upright side in the case of the circle is equal to the diameter. For if

$$\text{sq } DE \text{ } \text{rect } AE \text{ } EB \text{ } CA \text{ } AB$$

and only in the case of the circle

$$\text{sq } DE = \text{rect. } AE \text{ } EB$$

therefore also

$$CA = AB$$

And this must also be noted that the ordinates on the circumference of the circle are in every case perpendicular to the diameter and are in a straight line with the parallels to AC (Eucl. III 3 4)

therefore also

$$\text{sq } CE > \text{sq } DB \text{ (Eucl v 14)}$$

And so also

$$CE > DB$$

And they are parallel, therefore CD produced will meet the diameter AB outside the section (I 10, Eucl I 33)

But then let it be an hyperbola. Since then in the hyperbola

$$\text{sq } CE \text{ sq } DB \text{ rect } FE, EA \text{ rect } FB, BA \text{ (I 21),}$$

therefore also

$$\text{sq } CE > \text{sq } DB$$

And they are parallel, therefore the straight line CD produced will meet the diameter of the section outside the section

PROPOSITION 23

If a straight line lying between the two (conjugate) diameters¹ cuts the ellipse, it will when produced, meet each of the diameters outside the section

Let there be an ellipse whose diameters are the straight lines AB and CD (I 15), and let some straight line EF lying between the diameters AB and CD cut the section:

I say that the straight line EF , when produced will meet each of the straight lines AB and CD outside the section

For let the straight lines GE and FH be dropped ordinately from E and F to AB and the straight lines EK and FL ordinately to CD . Therefore

$$\text{sq } EG \text{ sq } FH \text{ rect } BG, GA \text{ rect } BH, HA \text{ (I 21)}$$

and

$$\text{sq } FL \text{ sq } EK \text{ rect } DL, LC \text{ rect } DK, KC \text{ (I 21)}$$

And

$$\text{rect } BG, GA > \text{rect } BH, HA$$

for the point G is nearer the midpoint (Eucl VI 27 II 5), and

$$\text{rect } DL, LC > \text{rect } DK, KC,$$

therefore also

$$\text{sq } GE > \text{sq } FH$$

and

$$\text{sq } FL > \text{sq } EK,$$


therefore also

$$GE > FH,$$

and

$$FI > EK$$


And GE is parallel to FH and FL to EK therefore the straight line EF produced will meet each of the diameters AB and CD outside the section (I 10 Eucl I 33)



¹So far Apollonius by theorems I 6 13 15 has shown for every ellipse the existence of at least one diameter and of one set of conjugate diameters but of no more. He can therefore now speak of the two diameters. Later on he will show the existence of an infinite number of such sets. The same is true of hyperbolas.

PROPOSITION 24

If a straight line, meeting a parabola or hyperbola at a point when produced both ways, falls outside the section, then it will meet the diameter



Let there be a parabola or hyperbola whose diameter is the straight line AB , and let the straight line CDE meet it at D and when produced both ways, let it fall outside the section


I say that it will meet the diameter AB

For let some point F be taken on the section, and let the straight line DF be joined therefore DF produced will meet the diameter of the section (I 22) Let it meet it at A and the straight line CDE lies between the section and the straight line FDA . And therefore the line CDE produced will meet the diameter outside the section

PROPOSITION 25

If a straight line, meeting an ellipse between the two (conjugate) diameters and produced both ways, falls outside the section, it will meet each of the diameters

Let there be an ellipse whose diameters are the straight lines AB and CD (I 15) and let EF , some straight line between the two diameters meet it at G , and produced both ways fall outside the section




I say that the straight line EF will meet each of the straight lines AB and CD

Let the straight lines GH and GK be dropped ordinately to the straight lines AB and CD respectively. Since GK is parallel to AB (I 15) and some straight line GF has met GK therefore it will also meet AB . Then likewise EF will also meet CD

PROPOSITION 26

If in a parabola or hyperbola a straight line is drawn parallel to the diameter of the section it will meet the section in one point only



Let there first be a parabola whose diameter is the straight line ABC , and whose upright side is the straight line AD and let the straight line LE be drawn parallel to AB

I say that the straight line LE produced will meet the section

For let some point E be taken on LE and from E let the straight line EG be drawn parallel to an ordinate, and let

$$\text{rect } DA, AC > \text{sq } GE,$$

and from C let CH be erected ordinatewise (I 19) Therefore
 $\text{sq } HC = \text{rect } DA, AC$ (I 11)

But

$$\text{rect } DA, AC > \text{sq } EG,$$

therefore

$$\text{sq } HC > \text{sq } EG,$$

therefore

$$HC > EG$$


And they are parallel, therefore the straight line EF produced cuts the straight line HC , and so it will also meet the section

Let it meet it at the point K

Then I say also that it will meet it in the one point K only

For if possible let it also meet it in the point L Since then a straight line cuts a parabola in two points, if produced it will meet the diameter of the section (I 22) And this is absurd for it is supposed parallel Therefore the straight line EF produced meets the section in only one point

Next let the section be an hyperbola and the straight line AB the transverse



side of the figure and the straight line AD the upright side and let the straight line DB be joined and produced Then with the same things being constructed, let the straight line CM be drawn from C parallel to AD Since then
 $\text{rect } MC, CA > \text{rect } DA, AC,$

and

$$\text{sq } CH = \text{rect } MC, CA,$$

and

$$\text{rect } DA, AC > \text{sq } GE,$$

therefore also

$$\text{sq } CH > \text{sq } GE$$

And so also


$$CH > GE$$

and the same things as in the first case will come to pass

PROPOSITION 27

If a straight line cuts the diameter of a parabola, then produced both ways it will meet the section

Let there be a parabola whose diameter is the straight line AB , and let some straight line CD cut it within the section



I say that the straight line CD produced both ways will meet the section

For let some straight line AE be drawn from A parallel to an ordinate, therefore the straight line AE will fall outside the section (I 17)

Then either the straight line CD is parallel to the straight line AE or not

If now it is parallel to it it has been dropped ordinatewise so that produced both ways it will meet the section (I 18)

Next let it not be parallel to AE but produced let it meet AE at E . Then it is evident that it meets the section the side the point E is on, for if it meets AE , *a fortiori* it cuts the section

I say that, produced the other way it also meets the section. For let the straight line MA be the parameter and the straight line GF an ordinate and let

$$\text{sq } AD = \text{rect } BA \cdot AF \text{ (Eucl vi 11)}$$

and let the straight line BK , parallel to the ordinate, meet the straight line DC at C . Since

$$\text{rect } BA \cdot AF = \text{sq } AD$$

hence

$$AB \cdot AD = AD \cdot AF$$

and therefore

$$BD \cdot DF = 4B \cdot AD \text{ (Eucl v 19)}$$

Therefore also

$$\text{sq } BD + \text{sq } DF = \text{sq } AB + \text{sq } AD$$

But since

$$\text{sq } AD = \text{rect } BA \cdot AF$$

hence

$$AB \cdot AF + \text{sq } AB + \text{sq } AD = \text{sq } BD + \text{sq } FD$$

But

$$\text{sq } BD + \text{sq } DF = \text{sq } BC + \text{sq } FG$$

and

$$AB \cdot AF = \text{rect } BA \cdot AM = \text{rect } FA \cdot 4M$$

Therefore

$$\text{sq } BC + \text{sq } FG = \text{rect } BA \cdot AM + \text{rect } FA \cdot 4M$$

and alternately

$$\text{sq } BC + \text{rect } BA \cdot AM = \text{sq } FG + \text{rect } FA \cdot 4M$$

But

$$\text{sq } FG = \text{rect } FA \cdot 4M$$

because of the section (1 11) Therefore also

$$\text{sq } BC = \text{rect } BA, AM$$


But the straight line AM is the upright side,¹ and the straight line BC is parallel to an ordinate Therefore the section passes through the point C (1 20), and the straight line CD meets the section at the point C

PROPOSITION 28

If a straight line touches one of the opposite sections, and some point is taken within the other section and through it a straight line is drawn parallel to the tangent then produced both ways it will meet the section

Let there be opposite sections whose diameter is the straight line AB , and let some straight line CD touch the section A , and let some point E be taken within the other section and through E let the straight line EF be drawn parallel to the straight line CD

I say that the straight line EF produced both ways will meet the section



Since then it has been proved that the straight line CD produced will meet the diameter AB (1 24), and EF is parallel to it therefore EF produced will meet the diameter I let it meet it at G , and let AH be made equal to GB , and through H let HK (1 18) be drawn parallel to EF , and let the straight line KL be dropped ordinately and let GM be made equal to LH , and let the straight line MN be drawn parallel to an ordinate and let GN be further produced in the same straight line And since KI is parallel to MN and KH to GN and LV is one straight line triangle KHL is similar to triangle HMN And

$$LH = GM$$

therefore

$$KL = MN$$

And so also

$$\text{sq } KL = \text{sq } MN$$

And since

$$LH = GM$$

and

$$AH = BG,$$

and AB is common therefore

$$BL = AM$$

¹The text reads $\kappa\lambda\gamma\iota\alpha$ which is impossible I have corrected to $\alpha\beta\theta\iota\alpha$

therefore

$$\text{rect } BL, LA = \text{rect } AM MB$$

Therefore

$$\text{rect } BL, LA \text{ sq } LK \quad \text{rect } AM MB \text{ sq } MN$$

And

$$\text{rect } BL, LA \text{ sq } LK \quad \text{the transverse the upright (I 21),}$$

therefore also

$$\text{rect } AM MB \text{ sq } MN \quad \text{the transverse the upright}$$


Therefore the point N is on the section Therefore the straight line EF produced will meet the section at the point N (I 21)

Likewise then it could be shown that produced to the other side it will meet the section

PROPOSITION 29

If in opposite sections a straight line is drawn through the center to meet either of the sections, then produced it will cut the other section

Let there be opposite sections whose diameter is the straight line AB , and whose center is the point C , and let the straight line CD cut the section AD



I say that it will also cut the other section

For let the straight line ED be dropped ordinatewise and let the straight line BF be made equal to the straight line AE and let the straight line FG be drawn ordinatewise (I 19) And since

$$EA = BF$$

and AB is common therefore

$$\text{rect } BE EA = \text{rect } BF FA$$

And since

$$\text{rect } BE, EA \text{ sq } DE \quad \text{the transverse the upright (I 21)}$$

but also

$$\text{rect } BF FA \text{ sq } FG \quad \text{the transverse the upright (I 21),}$$

therefore also

$$\text{rect } BE, EA \text{ sq } DE \quad \text{rect } BF FA \text{ sq } FG \text{ (I 14)}$$

But

$$\text{rect } BE EA = \text{rect } BF FA$$

therefore also

$$\text{sq } DE = \text{sq } FG$$

Since then

$$EC = CF,$$

and


$$DF = FG,$$

and EF is a straight line, and ED is parallel to FG , therefore DG is also a straight line (Eucl vi 32) And therefore CD will also cut the other section

PROPOSITION 30

If in an ellipse or in opposite sections a straight line is drawn in both directions from the center meeting the section, it will be bisected at the center

Let there be an ellipse or opposite sections and their diameter the straight line AB , and their center C , and through C let some straight line DCE be drawn (I 29)



I say that the straight line CD is equal to the straight line CE

For let the straight lines DF and EG be drawn ordinatewise And since
rect BF, FA sq FD the transverse the upright (I 21),
but also

rect AG, GB sq GE the transverse the upright (I 21),
therefore also

$$\text{rect } BF, FA \text{ sq } FD = \text{rect } AG, GB \text{ sq } CE \text{ (I 14)}$$

And alternately

$$\text{rect } BF, FA \text{ rect } AG, GB \text{ sq } FD \text{ sq } GE$$

But

$$\text{sq } FD \text{ sq } GE = \text{sq } FC \text{ sq } CG \text{ (Eucl vi 4),}$$

therefore alternately

$$\text{rect } BF, FA \text{ sq } FC = \text{rect } AG, GB \text{ sq } CG$$

Therefore also *componendo* in the case of the ellipse and inversely and *convertendo* in the case of the opposite sections (Eucl v Defs 14, 13, 16),

$$\text{sq } FC \text{ sq } CG = \text{sq } BC \text{ sq } CG \text{ (Eucl II 5 6),}$$

and alternately But

$$\text{sq } CB = \text{sq } AC$$

therefore also

$$\text{sq } CG = \text{sq } CF$$

Therefore

$$CG = CF$$


And the straight lines DF and GE are parallel, therefore also

$$DC = CE$$

PROPOSITION 31

If on the transverse side of the figure of an hyperbola some point be taken cutting off from the vertex of the section not less than half of the transverse side of the figure and a straight line be drawn from it to meet the section, then, when further produced it will fall within the section on the near side of the section

Let there be an hyperbola whose diameter is the straight line AB and let C some point on the diameter be taken cutting off the straight line CB not less



than half of AB and let some straight line CD be drawn to meet the section

I say that the straight line CD produced will fall within the section

For if possible let it fall outside the section as the line CDF (1. 24), and from E a point at random let the straight line EG be dropped ordinatewise also DH and first let

$$AC = CB$$

And since

$$\text{sq } EG \text{ sq } DH > \text{sq } FG \text{ sq } DH \text{ (Eucl v 8),}$$

but

$$\text{sq } EG \text{ sq } DH \text{ sq } CG \text{ sq } CH$$

because of EG 's being parallel to DH and

$$\text{sq } FG \text{ sq } DH \text{ rect } AG \text{ GB rect } AH \text{ HB}$$

because of the section (1. 21),

therefore

$$\text{sq } CG \text{ sq } CH > \text{rect } AG \text{ GB rect } AH \text{ HB}^1$$

Alternately therefore

$$\text{sq } CG \text{ rect } AG \text{ GB} > \text{sq } CH \text{ rect } AH, HB$$

Therefore *separando*

$$\text{sq } CB \text{ rect } AG \text{ GB} > \text{sq } CB \text{ rect } AH \text{ HB}$$

and this is impossible (Eucl v 8) Therefore the straight line CDE will not fall outside the section therefore inside And for this reason the straight line from some one of the points on the straight line AC will a *fortiori* fall inside, since it will also fall inside CD

¹The rules governing operations on inequalities in proportions are not developed by Euclid in Book V of the *Elements*. But they can be deduced on Euclid's principles.

PROPOSITION 32

If a straight line is drawn through the vertex of a section of a cone parallel to an ordinate, then it touches the section, and another straight line will not fall into the space between the conic section and this straight line

Let there be a section of a cone, first the so called parabola whose diameter is the straight line AB , and from A let the straight line AC be drawn parallel to an ordinate

Now it has been shown that it falls outside the section (1 17)

Then I say that also another straight line will not fall into the space between the straight line AC and the section

For if possible, let it fall in, as the straight line AD and let some point D be taken on it at random and let the straight line DE be dropped or dinatewise, and let the straight line AF be the parameter of the ordinates And since

$\text{sq } DE \text{ sq } EA > \text{sq } GE \text{ sq } EA$ (Eucl v 8),
and

$$\text{sq } GE = \text{rect } FA \cdot 4E \text{ (1 11),}$$

therefore also

$$\text{sq } DE \text{ sq } EA > \text{rect } FA \cdot AE \text{ sq } EA$$

or

$$> FA \cdot EA$$

Let it be contrived then that

$$\text{sq } DE \text{ sq } EA = FA \cdot HA \text{ (Eucl vi 20 11),}$$

and through the point H let the straight line HLA be drawn parallel to ED
Since then

$$\text{sq } DE \text{ sq } EA = FA \cdot AH = \text{rect } FA \cdot AH \text{ sq } AH$$

and

$$\text{sq } DE \text{ sq } EA = \text{sq } KH \text{ sq } HH \text{ (Eucl vi 22)}$$

and

$$\text{sq } HL = \text{rect } FA \cdot AH \text{ (1 11)}$$

therefore also

$$\text{sq } KH \text{ sq } HA = \text{sq } LH \text{ sq } HA$$

Therefore

$$KH = HL$$


and this is absurd Therefore another straight line will not fall into the space between the straight line $1C$ and the section

Next let the section be an hyperbola or ellipse or circumference of a circle whose diameter is the straight line AB and whose upright side is the straight line AF and let the straight line BF be joined and produced, and from the point A let the straight line AC be drawn parallel to an ordinate


Now it has been shown that it falls outside the section (1 17)

Then I say that also another straight line will not fall into the space between the straight line AC and the section

For if possible let it fall as the straight line AD and let some point D be taken at random on it and from it let the straight line DE be dropped ordi



parallel and through E let the straight line EM be drawn parallel to the straight line AF



And since

$$\text{sq } GE = \text{rect } AE \cdot EM \quad (\text{r } 12 \text{ } 13)$$

let it be contrived that

$$\text{rect } AE \cdot EN = \text{sq } DE,$$

and let the straight line joining AN cut the straight line FM at X and through X let the straight line XH be drawn parallel to FA and through H HLA parallel to AC Since then

$$\text{sq } DE = \text{rect } AM \cdot EN,$$

hence

$$NE \cdot ED = DE \cdot EA$$

and therefore

$$NE \cdot EA = \text{sq } DE = \text{sq } EA \quad (\text{Eucl vi } 20)$$

But

$$NE \cdot EA = XH \cdot HA$$

and

$$\text{sq } DE = \text{sq } EA = \text{sq } AH = \text{sq } HA$$

Therefore

$$XH \cdot HA = \text{sq } AH = \text{sq } HA$$

therefore

$$\sqrt{AH} \cdot HK = KH \cdot HA \text{ (Eucl vi 20)}$$

Therefore

$$\text{sq } KH = \text{rect } AH \cdot HX,$$

but also

$$\text{sq } LH = \text{rect } AH \cdot HX$$

because of the section (i 12, 13),

therefore

$$\text{sq } KH = \text{sq } HL$$

and this is absurd. Therefore another straight line will not fall into the space between the straight line AC and the section.

PROPOSITION 33

If in a parabola some point is taken, and from it an ordinate is dropped to the diameter and to the straight line cut off by it on the diameter from the vertex a straight line in the same straight line from its extremity is made equal then the straight line joined from the point thus resulting to the point taken will touch the section.

Let there be a parabola whose diameter is the straight line AB and let the straight line CD be dropped ordinately and let the straight line AE be made equal to the straight line ED and let the straight line AC be joined.

I say that the straight line AC produced will fall outside the section.

For if possible let it fall within as the straight line CF and let the straight line GB be dropped ordinately. And since

$$\text{sq } BG = \text{sq } CD > \text{sq } FB = \text{sq } CD$$

but

$$\text{sq } FB = \text{sq } CD = \text{sq } BA + \text{sq } AD$$

and

$$\text{sq } BG = \text{sq } CD = BE \cdot DE \text{ (i 20),}$$

therefore

$$BE \cdot DE > \text{sq } BA + \text{sq } AD$$

But

$$BE \cdot DE = 4 \text{ rect } BE \cdot EA = 4 \text{ rect } DE \cdot EA$$

therefore also

$$4 \text{ rect } BE \cdot EA = 4 \text{ rect } DE \cdot EA > \text{sq } AB + \text{sq } AD$$

Therefore alternately

$$4 \text{ rect } BE \cdot EA = \text{sq } AB > 4 \text{ rect } DE \cdot EA = \text{sq } AD$$

and this is absurd for since

$$DE,$$

hence


But

$$4$$

for E is not the π
 AC does not fall

$$\text{of } A$$




line



PROPOSITION 34

If on an hyperbola or ellipse or circumference of a circle some point is taken and from it a straight line is dropped ordinatewise to the diameter, and whatever ratio the straight lines cut off by the ordinate from the ends of the figure's transverse side have to each other, that ratio have the segments of the transverse side to each other so that the segments from the vertex arc corresponding then the straight line joining the point taken on the transverse side and that taken on the section will touch the section

Let there be an hyperbola or ellipse or circumference of a circle whose diam-



eter is the straight line AB and let some point C be taken on the section and from C let the straight line CD be drawn ordinatewise and let it be contrived that

$BD \quad DA \quad BE \quad EA'$

and let the straight line EC be joined

I say that the straight line CE touches the section

¹This construction is easy. In the case of the hyperbola *componendo*

BD+DA DA BA EA

and in the case of the ellipse *sepa ando*

BD-DA DA BA EA

This proportion is the same as the harmonic proportion defined by Nicomachus in his *Introduction to Arithmetic*. For if

$BD \quad D^4 \quad BE \quad EA$

$BD+D4 \quad BD \quad BA \quad BE$

$$BA \quad BD \quad BE-EA \quad BE$$

$$BD+D4 \quad B4 \quad BA \quad BE-E4$$

$$DA = BD - BA \quad EA = BA - BE$$

2BD-B1 B4 BA 2BE-B4

And so $B4$ is the harmonic mean between BD and BE

For if possible, let it cut it, as the straight line ECF , and let some point F be taken on it, and let the straight line GFH be dropped ordinatewise and let the straight lines AL and BA be drawn through A and B parallel to the straight line EC , and let the straight lines DC , BC , and GC be joined and produced to the points M , Λ , and K . And since

$$BD \ DA \ BE \ EA$$

but

$$BD \ DA \ BA \ AN,$$

and

$$BE \ AE \ BC \ CY \ BA \ \Lambda N \text{ (Eucl vi 4),}$$

therefore

$$BA \ AN \ BA \ \Lambda N,$$

therefore

$$AN = N\Lambda$$

Therefore

$$\text{rect } \Lambda N \ N\Lambda > \text{rect } AO \ O\Lambda \text{ (Eucl vi 27 ii 5)}$$

Therefore

$$NY \ \Lambda O > OA \ AN^1$$

But

$$N\Lambda \ \Lambda O \ \Lambda B \ BM \text{ (Eucl vi 4)}$$

therefore

$$\Lambda B \ BM > OA \ AN$$

Therefore

$$\text{rect } \Lambda B \ AN > \text{rect } BM, OA$$

And so

$$\text{rect } \Lambda B \ \Lambda N \text{ sq } CE > \text{rect } BM \ OA \text{ sq } CE \text{ (Eucl v 8)}$$

But

$$\text{rect } \Lambda B \ AN \text{ sq } CE \text{ rect } BD \ DA \text{ sq } DE$$

through the similarity of the triangles $B\Lambda D$, ECD and $N\Lambda D$ and

¹Eutocius commenting says For since

$$\text{rect } AN \ NY > \text{rect } AO \ O\Lambda$$

let $\text{rect } \Lambda N \ NY = \text{rect } AO \ XP$

where XP is some line such that $XP > \Lambda C$

therefore $OA \ AN \ N\Lambda \ XP$

But $NY \ \Lambda O > N\Lambda \ \Lambda P$ (Eucl v 8)

and therefore $NY \ XO > OA \ AN$

Then the converse is also evident that if

$$NY \ \Lambda O > OA \ AN$$

then $\text{rect } \Lambda N \ NA > \text{rect } AO \ O\Lambda$

For let it be that $OA \ AN \ \Lambda Y \ \Lambda P$

where $XP > \Lambda O$

therefore $\text{rect } \Lambda Y \ NA = \text{rect } AO \ XP$

and so $\text{rect } \Lambda N \ NA > \text{rect } AO \ O\Lambda$

²Eutocius commenting, says Since then because AN , EC and ΛB are parallel

$$AN \ EC \ AD \ DE$$

and $EC \ \Lambda B \ ED \ DB$

therefore ex aequali $AN \ \Lambda B \ AD \ DB$

therefore also $\text{sq } AN \text{ rect } AN \ \Lambda B \text{ sq } AD \text{ rect } AD \ DB$

But $\text{sq } EC \text{ sq } AN \text{ sq } ED \text{ sq } AD$

therefore ex aequali $\text{sq } EC \text{ rect } AN \ \Lambda B \text{ sq } ED \text{ rect } AD \ DB$

and inversely $\text{rect } \Lambda B \ \Lambda N \text{ sq } EC \text{ rect } AD \ DB \text{ sq } ED$

A similar proof holds for the proportion following

rect BM, OA sq CE rect BG, GA sq GE ,

therefore

rect BD, DA sq $DE >$ rect BG, GA sq GE

Therefore alternately

rect BD, DA rect $BG, GA >$ sq DE sq GE

But

rect BD, DA rect AG, GB sq CD sq GH (I 21),

and

sq DE sq EG sq CD sq FG (Eucl vi 4),

therefore also

sq CD sq $HG >$ sq CD sq FG

Therefore


$HG < FG$ (Eucl v 10)

and this is impossible Therefore the straight line EC does not cut the section therefore it touches it

PROPOSITION 35

If a straight line touches a parabola meeting the diameter outside the section, the straight line drawn from the point of contact ordinatewise to the diameter will cut off on the diameter beginning from the vertex of the section a straight line equal to the straight line between the vertex and the tangent, and no straight line will fall into the space between the tangent and the section

Let there be a parabola whose diameter is the straight line AB , and let the straight line BC be erected ordinatewise and let the straight line AC be tangent to the section



I say that the straight line AG is equal to the straight line GB

For if possible, let it be unequal to it, and let the straight line GE be made equal to AG and let the straight line EF be erected ordinatewise and let the straight line AF be joined Therefore AF produced will meet the straight line AC (I 33) and this is impossible For two straight lines will have the same ends Therefore the straight line AG is not unequal to the straight line GB , therefore it is equal

Then I say that no straight line will fall into the space between the straight line AC and the section

For if possible let the straight line CD fall in between and let GE be made equal to GD and let the straight line EF be erected ordinatewise Therefore the straight line joined from D to F touches the section (I 33) therefore produced it will fall outside it And so it will meet DC and two straight lines will have the same ends and this is impossible Therefore a straight line will not fall into the space between the section and the straight line AC

PROPOSITION 36

If some straight line meeting the transverse side of the figure touches an hyperbola or ellipse or circumference of a circle and a straight line is dropped from the

of contact ordinatewise to the diameter then as the straight line cut off by the tangent from the end of the transverse side is to the straight line cut off by the tangent from the other end of that side so will the straight line cut off by the ordinate from the end of the side be to the straight line cut off by the ordinate from the other end of the side in such a way that the corresponding straight lines are continuous and another straight line will not fall into the space between the tangent and the section of the cone

Let there be an hyperbola or ellipse or circumference of a circle whose diameter is the straight line AB and let the straight line CD be tangent, and let the straight line CE be dropped ordinatewise


I say that

$$BE \quad EA \quad BD \quad DA$$

For if it is not, let it be

$$BD \quad DA \quad BG \quad GA,$$

and let the straight line GF be erected ordinatewise therefore the straight line



joined from D to F will touch the section (1 31) therefore produced it will meet CD Therefore two straight lines will have the same ends, and this is impossible

I say that no straight line will fall between the section and the straight line CD

For if possible let it fall between as the straight line CH , and let it be contrived that

$$BH \quad HA \quad BG \quad GA,$$

and let the straight line CF be erected ordinatewise therefore the straight line joined from H to F when produced will meet HC (1 31) Therefore two straight lines will have the same ends and thus is impossible Therefore a straight line will not fall into the space between the section and the straight line CD

PROPOSITION 37

If a straight line touching an hyperbola or ellipse or circumference of a circle meets the diameter, and from the point of contact to the diameter a straight line is dropped ordinatewise, then the straight line cut off by the ordinate from the center of the section with the straight line cut off by the tangent from the center of the section will contain an area equal to the square on the radius of the section, and with the straight line between the ordinate and the tangent will contain an area having the ratio to the square on the ordinate which the transverse has to the upright

Let there be an hyperbola or ellipse or circumference of a circle whose diameter is the straight line AB , and let the straight line CD be drawn tangent and let the straight line CE be dropped ordinatewise, and let the point F be the center

I say that

$$\text{rect } DF, FE = \text{sq } FB,$$

and $\text{rect } DE, EF = \text{sq } EC$ the transverse the upright


For since CD touches the section, and CE has been dropped ordinatewise hence

$$AD \cdot DB = AE \cdot EB \quad (\text{I 36})$$

Therefore *componendo*

$$AD + DB \cdot DB = AE + EB \cdot EB$$

And let the halves of the antecedents be taken (Eucl v 15) in the case of the



hyperbola we shall say but

$$\text{half } (AE + EB) = FE,$$

and

$$\text{half } AB = FB,$$

therefore

$$FE \cdot EB = FB \cdot BD$$

Therefore *conuertendo*

$$FE \cdot FB = FB \cdot FD,$$

therefore

$$\text{rect } EF, FD = \text{sq } FB$$

And since

$$FE \text{ EB } FB \text{ BD } AF \text{ BD,}$$

alternately

$$AF \text{ FE } DB \text{ BE,}$$

componendo

$$AE \text{ EF } DE \text{ EB}$$

and so

$$\text{rect } AE \text{ EB} = \text{rect } FE, ED$$

But

rect $AE \text{ EB}$ sq CE the transverse the upright (I 21),
therefore also

$$\text{rect } FE \text{ ED} = \text{sq } CE \text{ the transverse the upright}$$

And in the case of the ellipse and of the circle we shall say but
half $(AD + DB) = DF$,

and

$$\text{half } AB = FB$$

therefore

$$FD \text{ DB } FB \text{ BE}$$

Therefore *convertendo*

$$DF \text{ FB } BF \text{ FE}$$

Therefore

$$\text{rect } DF \text{ FE} = \text{sq } BF$$

But

$$\text{rect } DF \text{ FE} = \text{rect } DE \text{ EF} + \text{sq } FE \text{ (Eucl II 3),}$$

and

$$\text{sq } BF = \text{rect } AE \text{ EB} + \text{sq } FE \text{ (Eucl II 5)}$$

Let the common square on EF be subtracted therefore

$$\text{rect } DE \text{ EF} = \text{rect } AE \text{ EB}$$

Therefore

$$\text{rect } DE \text{ EF} \text{ sq } CE \text{ rect } AE \text{ EB} \text{ sq } CE$$

But

$$\text{rect } AE \text{ EB} \text{ sq } CE \text{ the transverse the upright (I 21)}$$

Therefore

$$\text{rect } DE \text{ EF} \text{ sq } CE \text{ the transverse the upright}$$

PROPOSITION 38

If a straight line touching an hyperbola or ellipse or circumference of a circle meets the second diameter and from the point of contact a straight line is dropped to the same diameter parallel to the other diameter, then the straight line cut off by the dropped straight (*καταμύων*)¹ line from the center of the section with the straight

¹When this word *καταμύων* is used in connection with the first diameter we translate it as ordinate but we have preferred to stick more closely to the original when it is referred to the second diameter. For although it is certainly an ordinate in the case of the ellipse yet in the case of the hyperbola it is only analogically an ordinate. This analogy however becomes stronger and stronger as the treatise moves on. It is therefore no accident that *καταμύων* is used in both cases. On the other hand in First Definitions 1-5 Apollonius definitely calls both cases ordinates as if announcing the culmination of an analogy to be worked out in the course of the treatise.

line cut off by the tangent from the center of the section will contain an area equal to the square on the half of the second diameter and with the straight line between the dropped straight line and the tangent will contain an area having a ratio to the square on the dropped straight line which the upright side of the figure has to the transverse

Let there be an hyperbola or ellipse or circumference of a circle whose diameter is the straight line AGB and whose second diameter is the straight line CQD and let the straight line ELF meeting CD at F , be a tangent to the section and let the straight line HE be parallel to AB

I say that



$$\text{rect } FG \cdot GH = \text{sq } GC$$

and

$$\text{rect } GH, HF \text{ sq } HE \quad \text{the upright the transverse}$$

Let the straight line ME be drawn ordinatewise therefore

$$\text{rect } GM \cdot ML \text{ sq } ME \quad \text{the transverse the upright (1 37)}$$



But

$$\text{the transverse } BG \cdot CD \quad CD \text{ the upright (see Def 11)}$$

and therefore

$$\text{the transverse the upright sq } BA \text{ sq } CD \text{ (Eucl vi 20)}$$

and as the quarters of them that is

$$\text{the transverse the upright sq } GA \text{ sq } GC$$

therefore also

$$\text{rect } GM \cdot ML \text{ sq } ME \text{ sq } GA \text{ sq } GC$$

But

$$\text{rect } GM \cdot ML \text{ sq } ME \text{ comp } GM \cdot ME \cdot LM \cdot MF$$

or

$$\text{rect } GM \cdot ML \text{ sq } ME \text{ comp } CM \cdot CH \cdot LM$$

Therefore inversely

sq CG sq GA comp EM MG or HG GM EM ML or FG GL

Therefore

sq GC sq GA comp HG GM, FG GL ,

which is the same as

rect FG GH rect MG GL

Therefore

rect FG, GH rect MG, GL sq CG sq GA

And alternately therefore

rect FG GH sq CG rect MG GL sq GA

But

rect $MG, GL = \text{sq } GA$ (1 37),

therefore also

rect $FG, GH = \text{sq } CG$

Again since

the upright the transverse sq EM rect CV VL (1 37),

and

sq EM rect GM ML comp EM GM EM ML

or

sq EM rect GM ML comp HG HE FG GL or FH HE

which is the same as

rect FH HG sq HE

therefore

rect FH HG sq HE the upright the transverse

With the same things supposed it remains to be shown that as the straight line between the tangent and the end of the (second) diameter on the same side with the dropped straight line is to the straight line between the tangent and the second diameter so is the straight line between the other end and the dropped straight line to the straight line between the first end and the dropped straight line

For since

rect FG $GH = \text{sq } GC = \text{rect } CG$ GD (2 para above),

for

$CG = GD$

therefore

rect FG $GH = \text{rect } CG$ GD

therefore

FG GD CG GH

And *convertendo*

GF FD GC CH

And let the doubles of the antecedents be taken, but

$2GF = CF + FD$

because

$CG = GD$

and

$2GC = CD$

therefore

$CF + FD$ GD DC CH

And *separando*

$CF \quad FD \quad DH \quad HC,$

and this was to be shown

Then it is clear from what has been said that the straight line EF touches the section either if

$$\text{rect } FG, GH = \text{sq } GC,$$

or if

$$\text{rect } FH \quad HG \quad \text{sq } GC$$

in the ratio we said, for it could be shown conversely

PROPOSITION 39

If a straight line touching an hyperbola or ellipse or circumference of a circle meets the diameter, and from the point of contact a straight line is dropped ordinatewise to the diameter, then whichever of the two straight lines is taken of which one is the straight line between the ordinate and the center of the section and the other is between the ordinate and the tangent then the ordinate will have to it the ratio compounded of the ratio of the other of the two straight lines to the ordinate and of the ratio of the upright side of the figure to the transverse

Let there be an hyperbola or ellipse or circumference of a circle whose diameter is the straight line AB and let the center of it be the point F , and let the straight line CD be drawn tangent to the section, and the straight line CE be dropped ordinatewise

I say that


$$CE \quad FE \text{ comp the upright the transverse } ED \quad EC,$$

and

$$CE \quad ED \text{ comp the upright the transverse } FE \quad EC$$

For let

$$\text{rect } FE \quad ED = \text{rect } EC \quad G$$



And since

$$\text{rect } FE \quad ED \quad \text{sq } CE \quad \text{the transverse the upright (1 37),}$$

650 APOLLONIUS OF PERGA

and

rect FE LD = rect CE , G ,

therefore

rect CE G sq CE G CE the transverse the upright

And since

rect FE LD = rect CE , G ,

hence

FE EC G ED

And since

CE ED comp CE G , G ED ,

but



CE G the upright the transverse

therefore

CE ED comp the upright the transverse, FE EC

PROPOSITION 40

If a straight line touching an hyperbola or ellipse or circumference of a circle meets the second diameter, and from the point of contact a straight line is dropped to the same diameter parallel to the other diameter, then whichever of the two straight lines is taken of which one is the straight line between the dropped straight line and the center of the section and the other is between the dropped straight line and the



tangent the dropped straight line will have to it the ratio compounded of the ratio of the transverse side to the upright and of the ratio of the other of the two straight lines to the dropped straight line

Let there be an hyperbola or ellipse or circumference of a circle AB and its diameter the straight line BFC and its second diameter the straight line DFE and let the straight line HLA be drawn tangent and the straight line AG parallel to the straight line BC

I say that

AG HG comp the transverse the upright FG GA ,
and

AG FG comp the transverse the upright, HG GA
Let

$$\text{rect } GA \cdot K = \text{rect } HG \cdot GF$$

And since

the upright the transverse $\text{rect } HG, GF$ sq GA (r 38),
and

$$\text{rect } GA, K = \text{rect } HG, GF,$$

therefore also

$\text{rect } GA, K$ sq GA K AG the upright the transverse
And since

$$AG \cdot GF \text{ comp } AG \cdot K \cdot K \cdot GF,$$

but

$$AG \cdot K \text{ the transverse the upright,}$$

and

$$K \cdot GF \cdot HG \cdot GA$$

because

$$\text{rect } HG \cdot GF = \text{rect } AG \cdot K$$

therefore

$$AG \cdot GF \text{ comp the transverse the upright } GH \cdot GA$$

PROPOSITION 41

If in an hyperbola or ellipse or circumference of a circle a straight line is dropped ordinatewise to the diameter and equiangular parallelogrammic figures are described both on the ordinate and on the radius and the ordinate side has to the remaining side of the figure the ratio compounded of the ratio of the radius to the remaining side of its figure, and of the ratio of the upright side of the section's figure to the transverse then the figure on the straight line between the center and the ordinate similar to the figure on the radius is in the case of the hyperbola greater than the figure on the ordinate by the figure on the radius and, in the case of the ellipse and circumference of a circle, together with the figure on the ordinate is equal to the figure on the radius

Let there be an hyperbola or ellipse or circumference of a circle whose diameter is the straight line AB and center the point E and let the straight line CD be dropped ordinatewise and on the straight lines EA and CD let the equiangular figures AF and DG be described and let

$$CD \cdot CG \text{ comp } AE \cdot EF \text{ the upright the transverse}$$

I say that with the figure on ED similar to AF , in the base of the hyperbola
figure on $ED = AF + GD$

and in the case of the ellipse and circle

$$\text{figure on } ED + GD = AF$$

For let it be contrived that

$$\text{the upright the transverse } DC \cdot CH$$

And since


$$DC \cdot CH \text{ the upright the transverse}$$

but

$$DC \cdot CH \text{ sq } DC \text{ rect } DC \cdot CH$$

and

the upright the transverse sq DC rect $BD DA$ (1 21),



therefore

$$\text{rect } BD DA = \text{rect } DC CH$$

And since

DC CG comp AE EF the upright the transverse

or

$$DC \text{ } CG \text{ comp } AE \text{ } EF \text{ } DC \text{ } CH,$$

and further

$$DC \text{ } CG \text{ comp } DC \text{ } CH \text{ } CH \text{ } CG$$

therefore

$$\text{ratio comp } AE \text{ } EF \text{ } DC \text{ } CH = \text{ratio comp } DC \text{ } CH \text{ } CH \text{ } CG$$

Let the common ratio DC CH be taken away therefore

$$AE \text{ } EF \text{ } CH \text{ } CG$$

But

$$HC \text{ } CG \text{ rect } HC \text{ } CD \text{ rect } GC \text{ } CD,$$

and

$$AL \text{ } EF \text{ sq } AE \text{ rect } AE \text{ } EF$$

therefore

$$\text{rect } HC, CD \text{ rect } GC \text{ } CD \text{ sq } AE \text{ rect } AE \text{ } EF$$

And it has been shown that

$$\text{rect } HC, CD = \text{rect } BD \text{ } D1,$$

therefore

$$\text{rect } BD, DA \text{ rect } GC \text{ } CD \text{ sq } AE \text{ rect } AE \text{ } EF$$

Alternately

$$\text{rect } BD, DA \text{ sq } AE \text{ rect } GC \text{ } CD \text{ rect } AE \text{ } EF$$

And

$$\text{rect } GC, CD \text{ rect } AE \text{ } EF \text{ pll } g \text{ } DG \text{ pll } g \text{ } F4$$

for they are equiangular and have to one another the ratio compounded of their sides GC AE and CD EF (Eucl vi 23) and therefore

$$\text{rect } BD \text{ } DA \text{ sq } EA \text{ pll } g \text{ } DG \text{ pll } g \text{ } FA$$

Moreover in the case of the hyperbola we are to say *componendo*

$$\text{rect } BD \text{ } DA + \text{sq } AE \text{ sq } AE \text{ pll } g \text{ } GD + \text{pll } g \text{ } AF \text{ pll } g \text{ } 4F,$$

or

$$\text{sq } DE \text{ sq } EA \text{ pll } g \text{ } GD + \text{pll } g \text{ } 4F \text{ pll } g \text{ } AF \text{ (Eucl ii 6)}$$

And as the square on DE is to the square on EA so is the figure described on ED similar and similarly situated to the parallelogram AF , to the parallelogram AF (Eucl vi 20 porism) therefore, with the figure on ED similar to the parallelogram AF ,

$$\text{pll } g \text{ } GD + \text{pll } g \text{ } AF \text{ pll } g \text{ } AF \text{ figure on } ED \text{ pll } g \text{ } AF$$

Therefore

$$\text{figure on } ED = \text{pll } g \text{ } GD + \text{pll } g \text{ } AF$$

the figure on ED being similar to the parallelogram AF

And in the case of the ellipse and of the circumference of a circle we shall say since then

$$\text{whole sq } AE \text{ whole pll } g \text{ } AF$$

$$\text{rect } AD \text{ } DB \text{ subtracted pll } g \text{ } DG \text{ subtracted,}$$

also remainder is to remainder as whole to whole (Eucl v 19)

And

$$\text{sq } AE - \text{rect } BD \text{ } DA = \text{sq } DE \text{ (Eucl ii 5),}$$

therefore

$$\text{sq } DE \text{ pll } g \text{ } AF - \text{pll } g \text{ } DG \text{ sq } AE \text{ pll } g \text{ } AF$$

But

$$\text{sq } AE \text{ pll } g \text{ } 4F \text{ sq } DE \text{ figure on } DE \text{ (Eucl vi 20 porism)}$$

the figure on DE being similar to the parallelogram AF Therefore the figure on DE being similar to the parallelogram AF

$$\text{sq } DE \text{ pll } g \text{ } AF - DG \text{ sq } DE \text{ figure on } DE$$

Therefore the figure on DE being similar to the parallelogram AF ,

$$\text{figure on } DE = \text{pll } g \text{ } AF - \text{pll } g \text{ } DG$$

Therefore

$$\text{figure on } DE + \text{pll } g \text{ } DG = \text{pll } g \text{ } AF$$

PROPOSITION 42

If a straight line touching a parabola meets the diameter, and from the point of contact a straight line is dropped ordinately to the diameter and, some point being taken on the section two straight lines are dropped to the diameter, one of them parallel to the tangent and the other parallel to the straight line dropped from

the point of contact then the triangle resulting from them is equal to the parallelogram contained by the straight line dropped from the point of contact and by the straight line cut off by the parallel from the vertex of the section

Let there be a parabola, whose diameter is the straight line AB and let the straight line AC be drawn tangent to the section, and let the straight line CH be dropped ordinatewise and from some point at random let the straight line DF be dropped ordinatewise and through the point D let the straight line DE be drawn parallel to the straight line AC , and through the point C the straight line CG parallel to the straight line BF and through the point B the straight line BG parallel to the straight line HC

I say that

$$\text{trgl } DEF = \text{pllg } GF$$

For since the straight line AC touches the section and the straight line CH has been dropped ordinatewise

$$AB = BH \text{ (I 35),}$$

therefore

$$AH = 2BH$$

Therefore

$$\text{trgl } AHC = \text{pllg } BC \text{ (Eucl I 41)}$$

And since

$$\text{sq } CH \text{ sq } DF \text{ } HB \text{ } BF$$

because of the section (I 20) but

$$\text{sq } CH \text{ sq } DF \text{ } \text{trgl } ACH \text{ } \text{trgl } EDF \text{ (Eucl VI 19)}$$

and

$$HB \text{ } BF \text{ } \text{pllg } GH \text{ } \text{pllg } GF \text{ (Eucl VI 1)}$$

therefore

$$\text{trgl } ACH \text{ } \text{trgl } EDF \text{ } \text{pllg } HG \text{ } \text{pllg } FG$$

Therefore alternately

$$\text{trgl } AHC \text{ } \text{pllg } BC \text{ } \text{trgl } EDF \text{ } \text{pllg } GF$$

But

$$\text{trgl } ACH = \text{pllg } GH$$


therefore

$$\text{trgl } EDF = \text{pllg } GF$$

PROPOSITION 43


If a straight line touching an hyperbola or ellipse or circumference of a circle meets the diameter and from the point of contact a straight line is dropped ordinatewise to the diameter and a parallel to it is drawn through the vertex meeting the straight line drawn through the point of contact and the center, and some point being taken on the section two straight lines are drawn to the diameter one of which is parallel to the tangent and the other parallel to the straight line dropped from the point of contact then the triangle resulting from them in the case of the hyperbola will be less than the triangle the straight line through the center and the point of contact cuts off, by the triangle on the radius similar to the triangle cut off and in the case of the ellipse and the circumference of the circle together with the triangle cut off from the center, will be equal to the triangle on the radius similar to the triangle cut off

Let there be an hyperbola or ellipse or circumference of a circle whose diam



eter is the straight line AB , and center the point C , and let the straight line DE be drawn tangent to the section, and let the straight line CE be joined and let the straight line EF be dropped ordinatewise, and let some point G be taken on the section and let the straight line GH be drawn parallel to the tangent, and let the straight line GK be dropped ordinatewise, and through B let the straight line BL be erected ordinatewise

I say that triangle AMC differs from triangle CLB by triangle GKH



For since the straight line ED touches and the straight line EF has been dropped hence

$EF \cdot FD$ comp $CF \cdot FE$ the upright the transverse (I 39)

But

$EF \cdot FD = GK \cdot KH$

and

$CF \cdot FE = CB \cdot BL$ (Eucl VI 4)

therefore

$GK \cdot KH$ comp $BC \cdot BL$ the upright the transverse

And through those things shown in the forty first theorem (I 41), triangle CAM differs from triangle BCL by triangle GKH for the same things have also been shown in the case of the parallelograms their doubles

PROPOSITION 44


If a straight line touching one of the opposite sections meets the diameter, and from the point of contact some straight line is dropped ordinatewise to the diameter and a parallel to it is drawn through the vertex of the other section meeting the straight line drawn through the point of contact and the center and some point being taken at random on the section let straight lines be dropped to the diameter, one of which

is parallel to the tangent and the other parallel to the straight line dropped ordinate-wise from the point of contact then the triangle resulting from them will be less than the triangle the dropped straight line cuts off from the center of the section, by the triangle on the radius similar to the triangle cut off

Let there be the opposite sections AF and BE , and let their diameter be the straight line AB and center the point C , and from some point F of those on the section FA let the straight line FG be drawn tangent to the section, and the straight line FO ordinate-wise, and let the straight line CF be joined and produced, as CE (1 29), and through B let the straight line BL be drawn parallel to the straight line FO and let some point N be taken on the section BE , and from N let the straight line NH be dropped ordinate-wise, and let the straight line NA be drawn parallel to the straight line FG

I say that

$$\text{trgl } HAN + \text{trgl } CBL = \text{trgl } CMH$$



For through E let the straight line ED be drawn tangent to the section BE , and let the straight line $E\lambda$ be drawn ordinate-wise. Since then FA and BE are opposite sections whose diameter is AB and the straight line through whose center is C and FG and ED are tangents to the section hence DE is parallel to FG ¹ And the straight line NA is parallel to FG , therefore NA is also parallel to ED , and the straight line MH to BL . Since then BE is an hyperbola, whose diameter is the straight line AB , and whose center is C and the straight line DE is tangent to the section, and $E\lambda$ drawn ordinate-wise and BL is parallel to $E\lambda$ and N has been taken on the section as the point from which NH has

¹Futocius commenting says For since AF is an hyperbola and BE a tangent and FO an ordinate
 (1 37) likewise then also
 Therefore
 and alternately
 But
 therefore also
 And
 and therefore
 and also
 therefore
 And they contain equal angles at the point C for they are vertical And so also
 and
 And they are alternate therefore the straight line FG is parallel to the straight line ED


$\text{rect } OC \cdot CC = \text{sq } CA$
 $\text{rect } \lambda C \cdot CD = \text{sq } CB$
 $\text{rect } OC \cdot CC \text{ sq } AC = \text{rect } \lambda C \cdot CD \text{ sq } BC$
 $\text{rect } OC \cdot CC \text{ rect } \lambda C \cdot CD \text{ sq } AC \text{ sq } CB$
 $\text{sq } \lambda C = \text{sq } CB$
 $\text{rect } OC \cdot CG = \text{rect } \lambda C \cdot CD$
 $OC = C\lambda$ (1 14 30)
 $GC = CD$
 $FC = CE$ (1 30)
 $FC = FC \quad CG = CD$
 $FG = FD$
 $\text{angle } CFG = \text{angle } CFD$

been dropped ordinatewise and AN has been drawn parallel to DE , therefore
 $\text{trgl } NHA + \text{trgl } BCL = \text{trgl } HMC$,
 for this has been shown in the forty third theorem (1 43)

PROPOSITION 45

If a straight line touching an hyperbola or ellipse or circumference of a circle meets the second diameter, and from the point of contact some straight line is dropped to the same diameter parallel to the other diameter, and through the point of contact and the center a straight line is produced, and some point being taken at random on the section two straight lines are drawn to the second diameter one of which is parallel to the tangent and the other parallel to the dropped straight line, then the triangle resulting from them is greater in the case of the hyperbola, than the triangle the dropped straight line cuts off from the center, by the triangle whose base is the tangent and vertex is the center of the section, and in the case of the ellipse and circle, together with the triangle cut off will be equal to the triangle whose base is the tangent and whose vertex is the center of the section

Let there be an hyperbola or ellipse or circumference of a circle ABC , whose



diameter is the straight line AH and second diameter HD , and center H and let the straight line CML touch it at C , and let the straight line CD be drawn parallel to AH and let the straight line HC be joined and produced and let some point B be taken at random on the section, and from B let the straight lines BE and BF be drawn from B parallel to the straight lines LC and CD

I say that in the case of the hyperbola

$$\text{trgl } BEF = \text{trgl } GHF + \text{trgl } LCH,$$


and in the case of the ellipse and circle

$$\text{trgl } BEF + \text{trgl } FGH = \text{trgl } CLH$$

For let the straight lines CA and BA be drawn parallel to DH Since then the straight line CM is tangent and the straight line CA has been dropped ordinatewise hence

CA AH comp MA AC the upright the transverse (1 39)

$MK \ AC \ CD \ DL$ (Eucl vi 4),



therefore

$CK \ KH$ comp $CD \ DL$ the upright the transverse
And triangle CDL is the figure on KH and triangle CAH , that is triangle CDH is the figure on CK that is on DH therefore, in the case of the hyperbola,

$\text{trgl } CDL = \text{trgl } CAH + \text{trgl on } AH \text{ similar to trgl } CDL$,
and in the case of the ellipse and the circle

$\text{trgl } CDH + \text{trgl } CDL = \text{trgl on } AH \text{ similar to trgl } CDL$,
for this was also shown in the case of their doubles in the forty first theorem (i 41)

Since then triangle CDL differs either from triangle CAH or from triangle CDH by the triangle on AH similar to triangle CDL , and it also differs by triangle CHL therefore

$\text{trgl } CHL = \text{trgl on } AH \text{ similar to trgl } CDL$

Since then triangle BFE is similar to triangle CDL , and triangle GFH to triangle CDH therefore they have the same ratio¹ And triangle BFE is described on NH between the ordinate and the center and triangle GFH on the ordinate BN , that is on FH and by things already shown (i 41) triangle BFE differs from triangle GFH by the triangle on AH similar to CDL , and so also by triangle CHL

¹That is (Eucl vi 4)


$$\frac{BF}{GF} = \frac{FE}{FH} = \frac{CD}{CD} = \frac{DL}{DH} = \frac{CK}{CK} = \frac{KH}{KH}$$

and
Therefore these first ratios can be substituted in the central proportion of the theorem
 $CK \ KH$ comp $CD \ DL$ the upright the transverse
and so satisfy i 41

PROPOSITION 46

If a straight line touching a parabola meets the diameter, the straight line drawn through the point of contact parallel to the diameter in the direction of the section bisects the straight lines drawn in the section parallel to the tangent

Let there be a parabola whose diameter is the straight line ABD , and let the



straight line AC touch the section (1 24) and through C let the straight line HCM be drawn parallel to the straight line AD (1 26), and let some point L be taken at random on the section, and let the straight line LNF (1 18, 22) be drawn parallel to AC

I say that

$$LN = NF$$

Let the straight lines BH , KFG , and LMD be drawn ordinately. Since then by the things already shown in the forty-second theorem (1 42)

$$\text{trgl } ELD = \text{pllg } BM,$$

and

$$\text{trgl } EFG = \text{pllg } BK$$

therefore the remainders

$$\text{pllg } GM = \text{quadr } LFGD$$

Let the common pentagon $MDGFN$ be subtracted therefore the remainders

$$\text{trgl } KFN = \text{trgl } LMN$$

And KF is parallel to LM therefore

$$FN = LN \text{ (Eucl vi 22 lemma)}$$

PROPOSITION 47

If a straight line touching an hyperbola or ellipse or circumference of a circle meets the diameter and through the point of contact and the center a straight line is drawn in the direction of the section it bisects the straight lines drawn in the section parallel to the tangent

Let there be an hyperbola or ellipse or circumference of a circle whose center is the straight line AB and center C and let the straight line DE tangent to the section and let the straight line CE be joined

let a point N be taken at random on the section, and through N let the straight line $HNOG$ be drawn parallel

I say that




$$NO = OG$$

For let the straight lines ANF , BL and GMA be dropped ordinates. Therefore by things already shown in the forty third theorem (143)

$$\text{trgl } HNF = \text{quadr } LBF\lambda,$$

and

$$\text{trgl } GHK = \text{quadr } LBA M$$



Therefore the remainders

$$\text{quadr } NGKF = \text{quadr } MKFA,$$

Let the common pentagon $ONFAM$ be subtracted

therefore the remainders

$$\text{trgl } OMG = \text{trgl } NXO$$

And the straight line MG is parallel to the straight line NX , therefore


$$NO = OG \text{ (Eucl vi 22, lemma)}$$

PROPOSITION 48

If a straight line touching one of the opposite sections meets the diameter, and through the point of contact and the center a straight line produced cuts the other section then whatever line is drawn in the other section parallel to the tangent, will be bisected by the straight line produced

Let there be opposite sections whose diameter is the straight line AB and center C and let the straight line AI touch the section A and let the straight line LC be joined and produced (129), and let some point V be taken on the

section B , and through N let the straight line NG be drawn parallel to the straight line LA .



I say that

$$NO = OG$$

For let the straight line ED be drawn through E tangent to the section therefore ED is parallel to LA (1 44 note) And so also to NG . Since then BNG is an hyperbola whose center is C and tangent DE and since CE has been joined and a point N has been taken on the section and through it NG has been drawn parallel to DE , by a theorem already shown (1 47) for the hyperbola

$$NO = OG$$

PROPOSITION 49

If a straight line touching a parabola meets the diameter and through the point of contact a parallel to the diameter is drawn, and from the vertex a straight line is drawn parallel to an ordinate and it is contrived that as the segment of the tangent between the erected straight line and the point of contact is to the segment of the parallel between the point of contact and the erected straight line so is some straight line to the double of the tangent, then whatever straight line is drawn [parallel to the tangent] from the section to the straight line drawn through the point of contact parallel to the diameter will equal in square the rectangle contained by the straight line found and by the straight line cut off by it from the point of contact

Let there be a parabola whose diameter is the straight line MBC and CD its tangent and through D let the straight line FDN be drawn parallel to the straight line BC , and let the straight line FB be erected ordinatewise (1 17),

and let it be contrived that

$$ED : DF :: \text{some straight line} : G \cdot 2CD$$


and let some point K be taken on the section and let the straight line KLP be drawn through K parallel to CD

I say that

$$\text{sq } KL = \text{rect } G \cdot DL$$

that is that with the straight line DL as diameter the straight line G is the upright side

For let the straight lines DY and KNM be dropped ordinate wise And since the straight line



CD touches the section, and the straight line $D\lambda$ has been dropped ordinate-wise, then

$$CB = B\lambda \quad (1 \ 35)$$

But

$$B\lambda = FD$$

and therefore

$$CB = FD$$

And so also

$$\text{trgl } ECB = \text{trgl } EFD$$

Let the common figure $DEBMN$ be added, therefore

$$\begin{aligned} \text{quadr } DCMN &= \text{pllg } FM \\ &= \text{trgl } APM \quad (1 \ 42) \end{aligned}$$

Let the common quadrilateral $LPMN$ be subtracted therefore the remainders

$$\text{trgl } ALN = \text{pllg } LC$$

And

$$\text{angle } DLP = \text{angle } ALN,$$

therefore

$$\text{rect } AL \ LN = 2 \text{ rect } LD, DC^1$$

And since

$$ED \ DF \ G \ 2CD$$

and

$$ED \ DF \ AL \ LN,$$

therefore also

$$G \ 2CD \ AL \ LN$$

But

$$AL \ LN \ \text{sq} \ AL \ \text{rect } AL \ LN,$$

and

$$G \ 2CD \ \text{rect } G \ DL \ 2 \text{rect } LD \ DC,$$

therefore

¹Eutocius commenting says For let the triangle ALN and the parallelogram $DLPC$ be set out And since

$$\text{trgl } ALN = \text{pllg } DP$$

let the straight line NR be drawn through N parallel to LA and through A AR parallel to LN therefore LR is a parallelogram and

$$\text{pllg } LR = 2 \text{ trgl } ALN$$

and so also

$$\text{pllg } LR = 2 \text{ pllg } DP$$

Then let the straight lines DC and LP be produced to S and T and let CS be made equal to DC and PT to LP and let ST be joined therefore

$$\text{pllg } DT = 2 \text{ pllg } DP$$

and so

$$\text{pllg } LR = \text{pllg } LS$$

But it is also equiangular with it because of the angles at L being vertical but in equal and equiangular parallelograms the sides about the equal angles are reciprocally proportional therefore

$$AL \ LT \ \text{or} \ DS \ DL \ LN$$

and

$$\text{rect } AL \ LN = \text{rect } LD \ DS$$

And since

$$DS = 2DC$$

hence

$$\text{rect } KL \ LN = 2 \text{ rect } LD \ DC$$


And if DC is parallel to LP and CP is not parallel to LD it is clear $DCPL$ is a trapezoid and so I say that

$$\text{rect } KL \ LN = \text{rect } DL \ CD + LR$$

For if LP is filled out as we have said before and the straight lines DC and LP are produced and CS is made equal to LP and PT to DC and the straight line ST is joined then

$$\text{pllg } DT = 2DP$$

and the same demonstration will fit And this will be useful in what follows (1 50)




sq KL rect KL, LN rect $G DL$ 2rect $CD DL$
 And alternately but
 rect $KL LN = 2\text{rect } CD DL$,
 therefore a.l.o

$$\text{sq } KL = \text{rect } G, DL$$

PROPOSITION 50

If a straight line touching an hyperbola or ellipse or circumference of a circle meets the diameter, and a straight line is produced through the point of contact and the center and from the vertex a straight line erected parallel to an ordinate meets the straight line drawn through the point of contact and the center, and if it is con-
 trued that as the segment of the tangent between the point of contact and the straight line erected is to the segment of the straight line, drawn through the point of contact and the center, between the point of contact and the straight line erected, so some straight line is to the double of the tangent, then whatever straight line is drawn from the section to the straight line drawn through the point of contact and the center parallel to the tangent will equal in square a rectangular area applied to the straight line found having as breadth the straight line cut off by it from the

CASES I




point of contact, and exceeding, in the case of the hyperbola, by a figure similar to the rectangle contained by the double of the straight line between the center and the point of contact and by the straight line found but in the case of the ellipse and circle, defectue by it

Let there be an hyperbola or ellipse or circumference of a circle whose diameter is the straight line AB , and center C , and let the straight line DE be a tangent, and let the straight line CE be joined and produced both ways, and let the straight line CA be made equal to the straight line EC , and through B let the straight line BFG be erected ordinally and through E let the straight line EH be drawn perpendicular to FC and let it be that

$$FF \text{ EG } EH \text{ } 2FD,$$

CASES II



and let the straight line HA be joined and produced and let some point L be taken on the section and through it let the straight line LMN be drawn parallel to ED , and the straight line LRN parallel to BG , and the straight line MP parallel to EH

I say that

$$\text{sq } LM = \text{rect } EM \ MP$$

For let the straight line CSO be drawn through C parallel to AP And since $EC = CA$,

and

Cases I

$$EC \ AC \ ES \ SH$$

therefore also

$$ES = SH$$

Cases II

And since

$$FE \ EG \ HE \ 2ED,$$

and

$$2ES = EH,$$

therefore also

$$FE \ EG \ SE \ ED$$

And

$$FE \ EG \ LM \ MR$$

therefore

$$LM \ MR \ SE \ ED$$

And since it was shown (I 43) that, in the case of the hyperbola,

$$\begin{aligned} \text{trgl } RNC &= \text{trgl } LNX + \text{trgl } GBC, \\ &= \text{trgl } LNY + \text{trgl } CDE^1 \end{aligned}$$

and, in the case of the ellipse and circle

$$\begin{aligned} \text{trgl } RNC + \text{trgl } LNY &= \text{trgl } GBC \\ &= \text{trgl } CDE \end{aligned}$$

therefore in the case of the hyperbola with the common triangle ECD and the common quadrilateral $NRMN$ subtracted and in the case of the ellipse and circle with the common triangle MAC subtracted ²

¹That

$$\text{trgl } GBC = \text{trgl } CDE$$

is proved by Apollonius in the course of another proof of I 43 reported by Eutocius It is also proved in III 1 without the help of intervening propositions

²The position of point L furnishes different cases which at times as in the present theorem require a change in the course of the proof The figures marked Cases I are drawn to fit the proof as set down but we have added figures marked Cases II as an example of the possible differences

For the hyperbola of Case II instead of the subtraction in the theorem above we have

$$\begin{aligned} \text{trgl } RVC &= \text{trgl } LNY + \text{trgl } CDE \\ \text{quadr } MCNL &= \text{quadr } MCNL \end{aligned}$$

Subtracting the first equals from the second identity we have

$$\text{trgl } LMR = \text{quadr } MEDY$$

The rest of the proof is the same

For the ellipse and circle of Case II we have as in the theorem above

$$\text{trgl } PAC + \text{trgl } LNY = \text{trgl } CDE$$

and subtracting the common triangle CMY

$$\text{trgl } LMP = \text{trgl } CDE - \text{trgl } CMY$$

therefore

$$\text{rect } LM \ MR = \text{rect } EM \ LD + MY$$

For let CM be made equal to CM and CA to CA Then

trgl $LMR = \text{quadr } MED\lambda$

And $M\lambda$ is parallel to DE , and

angle $LMR = \text{angle } EM\lambda$,

therefore

rect $LM, MR = \text{rect } EM ED + M\lambda$ (1.49 note, para 2)

And since

$MC CE M\lambda ED$

and

$MC CE MO FS$,

therefore

$MO ES MX ED$

And componendo

$MO + ES ES MX + ED ED$,

alternately

$MO + ES MX + ED ES ED$

But

$MO + ES MX + ED \text{ rect } MO + ES EM \text{ rect } M\lambda + ED EM$,

and

$ES ED IM MR FE EG$ (Eucl vi 4)

or

$ES ED \text{ sq } LM \text{ rect } LM, MR$

therefore

rect $MO + ES ME \text{ rect } M\lambda + ED EM \text{ sq } LM \text{ rect } LM MR$

And alternately

rect $MO + ES ME \text{ sq } LM \text{ rect } M\lambda + ED EM \text{ rect } LM, MR$

But

rect $LM MR = \text{rect } ME M\lambda + ED$ (above),

therefore

sq $LM = \text{rect } EM MO + ES$

And

$SE = SH$

and

$SH = OP$,

therefore

sq $LM = \text{rect } EM MP$

PROPOSITION 51

If a straight line touching either of the opposite sections meets the diameter and through the point of contact and the center some straight line is produced to the other section and from the vertex a straight line is erected parallel to an ordinate and meets the straight line drawn through the point of contact and the center and if it is contrived that as the segment of the tangent between the straight line erected


trgl $CDF - \text{trgl } CM\lambda = \text{quadr } MFDX$
and $M\lambda$ is parallel to FD and
angle $F M \lambda = \text{angle } RMI$



These cases will come up again in Book III and in general it is convenient to think of quadrilateral $MFDX$ as standing for the difference of the two triangles when one pair of its sides cross each other

and the point of contact is to the segment of the straight line, drawn through the point of contact and the center, between the point of contact and the straight line erected, so is some straight line to the double of the tangent, then whatever straight line in the other of the sections is drawn to the straight line through the point of contact and the center, parallel to the tangent, will equal in square the rectangle applied to the straight line found having as breadth the straight line cut off by it from the point of contact and exceeding by a figure similar to the rectangle contained by the straight line between the opposite sections and the straight line found

Let there be opposite sections whose diameter is the straight line AB and



center E , and let the straight line CD be drawn tangent to the section B and the straight line CE joined and produced (1 29) and let the straight line BLG be drawn ordinatewise (1 17) and let it be contrived that

$$LC \cdot CG \text{ some straight line } \Lambda \cdot 2CD$$

Now it is evident that the straight lines in the section BC , parallel to CD and drawn to EC produced are equal in square to the areas applied to Λ having as breadths the straight line cut off by them from the point of contact and exceeding by a figure similar to the rectangle $CF \cdot K$, for

$$GC = 2CE$$

I say then that in section FA the same thing will come about

For let the straight line MF be drawn through F tangent to the section AF , and let the straight line $\Lambda \cdot \Lambda N$ be erected ordinatewise. And since BC and AF are opposite sections and CD and MF are tangents to them therefore CD is equal and parallel to MF (1 44 note). But also

$$CE = EF$$

therefore also

$$ED = EM$$

And since

$$LC \cdot CG \text{ } \Lambda \cdot 2CD \text{ or } 2MF$$

therefore also

$$\Lambda F \cdot FN \text{ } \Lambda \cdot 2MF$$

Since then AF is an hyperbola whose diameter is AB and tangent MF , and AN has been drawn ordinatewise and

$$\Lambda F \cdot FN \text{ } \Lambda \cdot 2FM,$$

hence any lines drawn from the section to EF produced parallel to FM will equal in square the rectangle contained by the straight line Λ and the line cut off by them from F exceeding by a figure similar to the rectangle $CF \cdot \Lambda$ (1 50)


And with these things shown it is at once evident that in the parabola each of the straight lines drawn off parallel to the original diameter is a diameter

(I 46), but in the hyperbola and ellipse and opposite sections each of the straight lines drawn through the center is a diameter (I 47-48), and that in the parabola the straight lines dropped to each of the diameters parallel to the tangents will equal in square the rectangles applied to it (I 49), but in the hyperbola and opposite sections they will equal in square the areas applied to it and exceeding by the same figure (I 50-51), but in the ellipse the areas applied to it and defective by the same figure (I 50) and that all the things which have been already proved about the sections as following when the principal diameters are used,¹ will also, those very same things follow when the other diameters are taken

PROPOSITION 52 (PROBLEM)

Given a straight line in a plane bounded at one point, to find in the plane the section of a cone called parabola, whose diameter is the given straight line, and whose vertex is the end of the straight line and where whatever straight line is dropped from the section to the diameter at a given angle will equal in square the rectangle contained by the straight line cut off by it from the vertex of the section and by some other given straight line

Let there be the straight line AB given in position and bounded at the point A , and another straight line CD given in magnitude and first let the given



angle be a right angle it is required then to find a parabola in the plane of reference whose diameter is the straight line AB and whose vertex is the point A and whose upright side is the straight line CD , and where the straight lines dropped ordinatewise will be dropped at a right angle that is so that AB is the axis (First Def 1 7)

Let AB be produced to E and let CG be taken as the fourth part of CD , and let

$$EA > CG$$

and let

$$CD : H :: H : EA$$

Therefore

$$CD : EA :: \text{sq } H : \text{sq } EA$$

and

$$CD < 4EA$$

therefore also

$$\text{sq } H < 4 \text{ sq } EA$$

¹The principal diameter ($\delta \mu \epsilon \tau \rho \iota \varsigma \alpha \rho \chi \epsilon \varsigma$) is that whose being is established in I 7 porism.

Therefore

$$H < 2EA,$$

and so the two straight lines EA are greater than H . It is therefore possible for a triangle to be constructed from H and two straight lines EA . Then let the triangle EAF be constructed on EA at right angles to the plane of reference so that

$$EA = AF,$$

and

$$H = FE$$

and let the straight line AK be drawn parallel to FE and FK to EA and let a cone be conceived whose vertex is the point F and whose base is the circle about diameter KA , at right angles to the plane through AFX . Then the cone will be a right cone (First Def 1 3) for

$$AF = FK$$

And let the cone be cut by a plane parallel to the circle KA , and let it make as a section the circle MNX (1 4) at right angles clearly to the plane through MFN , and let the straight line MN be the common section of the circle MNX and of the triangle MFN , therefore it is the diameter of the circle. And let the straight line XL be the common section of the plane of reference and of the circle. Since then circle MNX is at right angles to triangle MFN , and the plane of reference also at right angles to triangle MFN , therefore the straight line LX , their common section, is at right angles to triangle MFN , that is to triangle KFA (Eucl XI 19), and therefore it is perpendicular to all the straight lines touching it and in the triangle, and so it is perpendicular to both MN and AB .

Again since a cone, whose base is the circle MNX and whose vertex is the point F , has been cut by a plane at right angles to the triangle MFN and makes as a section circle MNX and since it has also been cut by another plane, the plane of reference, cutting the base of the cone in a straight line XL at right angles to MN which is the common section of the circle MNX and the triangle MFN , and the common section of the plane of reference and of the triangle MFN the straight line AB , is parallel to the side of the cone FA , therefore the resulting section of the cone in the plane of reference is a parabola and its diameter AB (1 11), and the straight lines dropped ordinately from the section to AB will be dropped at right angles for they are parallel to XL which is perpendicular to AB . And since

$$CD = H = EA,$$

and

$$EA = AF = FK$$

and

$$H = EF = AK$$

therefore

$$CD = AK = AK = AF$$

And therefore

$$CD = AF = \text{sq } AK = \text{sq } AF \text{ or rect } AF, FK$$

Therefore CD is the upright side of the section, for this has been shown in the eleventh theorem (1 11)

PROPOSITION 53 (PROBLEM)

With the same things supposed let the given angle not be right, and let the angle HAE be made equal to it and let

$$AH = \text{half } CD,$$

and from H let the straight line HE be drawn perpendicular to AE , and through E let the straight line EL be drawn parallel to BH , and from A let the straight line AL be drawn perpendicular to EL , and let EL be bisected at K , and from A let the straight line AM be drawn perpendicular to EL and produced to F and G and let rect $LH \cdot AM = \text{sq } AL$. And given the two straight lines LH and AM in position and bounded at A and AM in magnitude and let a parabola be described with a right angle whose diameter is the straight line AL , and whose vertex is the point A and whose upright side is the straight line AM , as has been shown before (I 52), and it will pass through the point A because

$$\text{sq } AL = \text{rect } LH, AM \text{ (I 11),}$$

and the straight line EA will touch the section because

$$EA = AL \text{ (I 33)}$$

And HA is parallel to EKL , therefore HAB is the diameter of the section and the straight lines dropped to it parallel to AE will be bisected by AB (I 46) And they will be dropped at angle HAE . And since

$$\text{angle } AEH = \text{angle } AGF,$$

and angle at A is common therefore triangle AHE is similar to triangle AGF . Therefore

$$HA : EA :: FA : AG,$$

therefore

$$2AH : 2AE :: FA : AG$$

But

$$CD = 2AH,$$

therefore


$$FA : AG :: CD : 2AE$$

Then by things already shown in the forty ninth theorem (I 49) the straight line CD is the upright side


PROPOSITION 54

Given two bounded straight lines perpendicular to each other one of them being produced on the side of the right angle to find on the straight line produced the section of a cone called hyperbola in the same plane with the straight lines, so that the straight line produced is a diameter of the section and the point at the angle is the vertex, and where whatever straight line is dropped from the section to the diameter making an angle equal to the given angle will equal in square the rectangle applied to the other straight line having as breadth the straight line cut off by the dropped straight line beginning with the vertex and exceeding by a figure similar and similarly situated to that contained by the original straight lines

Let there be the two bounded straight lines AB and BC perpendicular to




each other and let AB be produced to D it is required then to find in the plane through the lines AB, BC an hyperbola whose diameter will be the straight



line ABD and vertex B and upright side the straight line BC and where the straight lines dropped from the section to BD at the given angle will equal in square the rectangles applied to BC having as breadths the straight lines cut off by them from B and exceeding by a figure similar and similarly situated to the rectangle $AB BC$

First let the given angle be a right angle and on AB let a plane be erected at right angles to the plane of reference and let the circle $AEBF$ be described in it about AB so that the segment of the circle's diameter within the sector AEB has to the segment of the diameter within the sector AFB a ratio not greater than that of AB to BC ¹ and let AEB be bisected at E and let the straight line

¹Eutocius commenting, adds Let there be two straight lines AB and BC and let it be required to describe a circle on AB so that its diameter is cut by AB in such a way that the part of it on the side of C has to the remainder a ratio not greater than that of AB to BC



Now let it be supposed that they have the same ratio and let AB be bisected at D and through it let the straight line EDF be drawn perpendicular to AB and let it be contrived that $AB BC ED DF$

and let EF be bisected then it is clear that if

$$AB = BC$$

$$ED = DF$$

and the point D will be the midpoint of EF and if

$$AB > BC$$

$$ED > DF$$

and the midpoint will be below D and if

EA be drawn perpendicular from E to the straight line AB and let it be produced to L , therefore the straight line EL is a diameter (Eucl III 1) If then

$$AB \ BC \ EA \ AL$$

we use point L , but if not, let it be contrived that

$$AB \ BC \ EA \ AM$$

with

$$AM < AL \text{ (Eucl V 8),}$$

and through M let MF be drawn parallel to AB , and let AF , EF , and FB be joined, and through B let $B\chi$ be drawn parallel to FE Since then

$$\text{angle } AFE = \text{angle } EFB,$$

but

$$\text{angle } AFE = \text{angle } A\chi B,$$

and

$$\text{angle } EFB = \text{angle } \chi BF,$$

therefore also

$$\text{angle } \chi BF = \text{angle } F\chi B,$$

therefore also

$$FB = F\chi$$

Let a cone be conceived whose vertex is the point F and whose base is the circle about diameter $B\chi$ at right angles to triangle BFX Then the cone will be a right cone for

$$FB = F\chi$$

Then let the straight lines BF , $F\chi$ and MF be produced, and let the cone be cut by a plane parallel to the circle $B\chi$ then the section will be a circle (I 4) Let it be the circle GPR and so GH will be the diameter of the circle (I 4 end) And let the straight line PDR be the common section of circle GH and of the plane of reference then PDR will be perpendicular to both of the straight lines GH and DB for both of the circles AB and HG are perpendicular to triangle FGH and the plane of reference is perpendicular to triangle FGH , and therefore their common section the straight line PDR is perpendicular to triangle FGH therefore it makes right angles also with all the straight lines touching it and in the same plane

$$AB < BC$$

it will be above D

And now let it be below as G and with center O and radius OF let a circle be described, then it will have to pass either within or without the points A and B And if it should pass through the points A and B what was enjoined would be done but let it fall beyond the points A and B and let the straight line AB produced both ways meet the circumference at H and K and let FH , HF , EA and AF be joined and let MB be drawn through B parallel to FA and BL parallel to AF and let MA and AL be joined then these will also be parallel to FH and HE because

$$AD = DB$$

$$DH = DA$$

and FDE is perpendicular to HA And since the angle at A is a right angle and MB and BL are parallel to FA and AF therefore the angle at B is a right angle then for the same reasons also the angle at A And so the circle described on ML will pass through the points A and B (Eucl III 31) Let the circle $MALB$ be described And since MB is parallel to FA ,

$$FD \ DM \ AD \ DB$$

$$\text{Then likewise also} \quad AD \ DB \ FD \ DL$$

$$\text{And therefore} \quad FD \ DM \ ED \ DL$$

$$\text{And alternately} \quad ED \ DF \ AB \ BC \ LD \ DM$$

And likewise if the circle described on FE cuts AB the same thing could be shown.

And since a cone whose base is circle GH and vertex F , has been cut by a plane perpendicular to triangle FGH , and has also been cut by another plane the plane of reference, in the straight line PDR perpendicular to the straight line GDH and the common section of the plane of reference and of triangle GFH , that is the straight line DB , produced in the direction of B , meets the straight line GF at A , therefore by things already shown before (1 12) the section PBR will be an hyperbola whose vertex is the point B , and where the straight lines dropped ordinatewise to BD will be dropped at a right angle, for they are parallel to straight line PDR And since

$$AB \cdot BC = EA \cdot KM,$$

and

$$EA \cdot KM = EN \cdot NF \quad \text{rect } EN \cdot NF = \text{sq } NF,$$

therefore

$$AB \cdot BC = \text{rect } EN \cdot NF = \text{sq } NF$$

And

$$\text{rect } EN \cdot NF = \text{rect } AN \cdot NB,$$

therefore

$$AB \cdot CB = \text{rect } AN \cdot NB = \text{sq } NF$$

But

$$\text{rect } AN \cdot NB = \text{sq } NF \text{ comp } AN \cdot NF \text{ } BN \cdot NF,$$

but

$$AN \cdot NF = AD \cdot DG = FO \cdot OG,$$

and

$$BN \cdot NF = FO \cdot OH,$$

therefore

$$AB \cdot BC \text{ comp } FO \cdot OG, FO \cdot OH,$$

that is

$$\text{sq } FO = \text{rect } OG \cdot OH$$

Therefore

$$AB \cdot BC = \text{sq } FO = \text{rect } OG \cdot OH$$

And the straight line FO is parallel to the straight line AD , therefore the straight line AB is the transverse side and BC the upright side, for these things have been shown in the twelfth theorem (1 12)

PROPOSITION 55 (PROBLEM)

Then let the given angle not be a right angle and let there be the two given straight lines AB and AC , and let the given angle be equal to angle BAH then it is required to describe an hyperbola whose diameter will be the straight line AB , and upright side AC and where the ordinates will be dropped at angle HAB

Let the straight line AB be bisected at D and let the semicircle AFD be described on AD and let some straight line FG parallel to AH , be drawn to the semicircle making

$$\text{sq } FG = \text{rect } DG \cdot GA = AC \cdot AB^1$$

¹Cutocius, commenting gives this construction. Let there be the semicircle ABC on the diameter AC and the given ratio EF to FG and let it be required to do what is proposed.

Let FH be made equal to EF and let HG be bisected at A and let the straight line CB be drawn in the semicircle at angle ACB (the required angle) and from the center L let the straight line LS be drawn perpendicular to it and produced let it meet the circumference at

And $4H$ will touch it, for

$$\text{rect } FD, DH = \text{sq } DL \text{ (I 37)}$$

And so $4B$ is a diameter of the section (I 51) And since

$$CA \cdot 2AD \text{ or } AB \cdot \text{sq } FG \text{ rect } DG, GA,$$

but

$$CA \cdot 2AD \text{ comp } CA \cdot 2AH, 2AH \cdot 2AD$$

or

$$CA \cdot 2AD \text{ comp } CA \cdot 2AH, AH \cdot AD$$

and

$$AH \cdot AD = FG \cdot GD,$$

therefore

$$CA \cdot AB \text{ comp } CA \cdot 2AH, FG \cdot GD$$

But also

$$\text{sq } FG \text{ rect } DG, GA \text{ comp } FG \cdot GD, FG \cdot GA,$$

therefore

$$\text{ratio comp } CA \cdot 2AH, FG \cdot GD = \text{ratio comp } FG \cdot GA, FG \cdot GD$$

Let the common ratio $FG \cdot GD$ be taken away, therefore

$$CA \cdot 2AH = FG \cdot GA$$

But

$$FG \cdot GA = OA \cdot AX$$

therefore


$$CA \cdot 2AH = OA \cdot AX$$

But whenever this is so the straight line AC is a parameter for this has been shown in the fiftieth theorem (I 50)

PROPOSITION 56 (PROBLEM)

Given two bounded straight lines perpendicular to each other to find about one of them as diameter and in the same plane with the two straight lines the section of a cone called ellipse, whose vertex will be the point at the right angle and where the straight lines dropped ordinatewise from the section to the diameter at a given angle will equal in square the rectangles applied to the other straight line having as breadth the straight line cut off by them from the vertex of the section and defective by a figure similar and similarly situated to the rectangle contained by the given straight lines

Let there be two given straight lines AB and AC perpendicular to each other,



of which the greater is the straight line AB , then it is required to describe in the plane of reference an ellipse whose diameter will be the straight line AB and vertex A , and upright side AC , and where the ordinates will be dropped from the section to the diameter at a given angle and will equal in square the rectangles applied to AC having as breadths the straight lines cut off by them from A and defective by a figure similar and similarly situated to rectangle $B\Lambda AC$

And first let the given angle be a right angle, and let a plane be erected from AB at right angles to the plane of reference and in it on AB , let the sector of a circle ADB be described and its midpoint be D and let the straight lines DA and DB be joined and let the straight line $A\Lambda$ be made equal to AC , and through Λ let the straight line ΛO be drawn parallel to DB , and through O let OF be drawn parallel to AB , and let DF be joined and let it meet AB produced at E , then we will have

$$AB \ AC \ AB \ A\Lambda \ DA \ AO \ DE \ FF$$

And let the straight lines AF and FB be joined and produced, and let some point G be taken at random on FA , and through it let the straight line GL be drawn parallel to DE and let it meet AB produced at K , then let FO be produced and let it meet GK at L Since then

$$\begin{aligned} \text{arc } AD &= \text{arc } DB, \\ \text{angle } ABD &= \text{angle } DFB \text{ (Eucl iii 27)} \end{aligned}$$

And since

$$\text{angle } EFA = \text{angle } FDA + \text{angle } FAD,$$

but

$$\text{angle } FAD = \text{angle } FBD,$$

and

$$\text{angle } FDA = \text{angle } FBA,$$

therefore also

$$\text{angle } EFA = \text{angle } DBA = \text{angle } DFB$$

And also DE is parallel to LG therefore

$$\text{angle } EFA = \text{angle } FGH,$$

and

$$\text{angle } DFB = \text{angle } FHG$$

And so also

$$\text{angle } FGH = \text{angle } FHG,$$

and

$$FG = FH$$

Then let circle GHN be described about HG at right angles to triangle HGF , let a cone be conceived whose base is the circle GHN , and whose vertex is the point F , then the cone will be a right cone because

$$FG = FH$$

And since the circle GHN is at right angles to plane HGF and the plane of reference is also at right angles to the plane through GH and HF therefore their common section will be at right angles to the plane through GH and HF Then let their common section be the straight line KM therefore the straight line KM is perpendicular to both of the straight lines AK and KG

And since a cone whose base is the circle GHN and whose vertex is the point F has been cut by a plane through the axis and makes as a section triangle GHF , and has been cut also by another plane through AK and KM ,

which is the plane of reference in the straight line KM which is perpendicular to GA and the plane meets the sides of the cone FG and IH , therefore the resulting section is an ellipse whose diameter is AB and where the ordinates will be dropped at a right angle (I 13), for they are parallel to KM And since

$$DE \cdot EF \text{ rect } DE \cdot EF \text{ or rect } BE, EA \text{ sq } EF,$$

and

$$\text{rect } BE, EA \text{ sq } EF \text{ comp } BE \cdot EF, AE \cdot EF,$$

but

$$BE \cdot EF = BK \cdot KH,$$

and

$$AE \cdot EF = AK \cdot KG \quad FL \cdot LG,$$

therefore

$$BA \cdot AC \text{ comp } FL \cdot LG \quad FL \cdot LH \text{ (see above),}$$

which is the same as

$$\text{sq } FL \text{ rect } GL, LH$$

therefore

$$BA \cdot AC \text{ sq } FL \text{ rect } GL \cdot LH$$

And whenever this is so, the straight line AC is the upright side of the figure as has been shown in the thirteenth theorem (I 13)

PROPOSITION 57 (PROBLEM)

With the same things supposed let the straight line AB be less than AC , and let it be required to describe an ellipse about diameter AB so that AC is the upright

Let AB be bisected at D and from D let the straight line EDF be drawn perpendicular to AB , and let

$$\text{sq } FE = \text{rect } BA \cdot AC$$

so that

$$FD = DE,$$

and let FG be drawn parallel to AB , and let it be contrived that

$$AC \cdot AB = EF \cdot FG,$$

therefore also

$$EF > FG$$

And since

$$\text{rect } CA, AB = \text{sq } EF,$$

hence

$$CA \cdot AB \text{ sq } FE \text{ sq } AB \text{ sq } DF \text{ sq } DA$$

But

$$CA \cdot AB = EF \cdot FG$$

therefore

$$EF \cdot FG \text{ sq } FD \text{ sq } DA$$


But

$$\text{sq } FD = \text{rect } FD \cdot DE,$$

therefore

$$EF \cdot FG \text{ rect } ED \cdot DF \text{ sq } AD$$

Then with two bounded straight lines situated at right angles to each other and with EF greater let an ellipse be described whose diameter is EF and upright side FG (I 56) then the section will pass through A because



and let

$$EA = EH$$

and let it be contrived that

$$\text{rect } HF \text{ } FL = \text{sq } AT,$$

and let the straight line AL be joined and from H let the straight line HM be drawn perpendicular to HF and so parallel to the straight line AFL for the angle at F is right. And with the two given bounded straight lines KH , and HM perpendicular to each other, let an ellipse be described whose transverse diameter is KH , and the upright side of whose figure is HM , and where the ordinates to HK will be dropped at right angles (156-57) then the section will pass through A because sq $FA = \text{rect } HF, FL$ (113). And since

$$HE = EK.$$

and

$$AE = EB,$$

the section will also pass through B , and E will be the center, and the straight line AEB the diameter. And the straight line DA will touch the section because

$$\text{rect } DE \cdot EF = \text{sq } EH$$

And since

$CA \quad AB \quad \text{sq} \quad FG \quad \text{rect} \quad AG, GE$

but

$$CA \quad AB \text{ comp } CA \quad 2AD \quad 2AD \quad AB \text{ or } DA \quad AE,$$

and

$$\text{sq } FG = \text{rect } AG, GE \text{ comp } FG \quad GE \quad FG \quad GA,$$

therefore

$$\text{ratio comp } CA \quad 2AD \quad DA \quad AE = \text{ratio comp } FG \quad GE \quad FG \quad GA$$

But

$$D4 \quad AE \quad FG \quad GE$$


and the common ratio being taken away we will have

CA 2AD FG GA

or

CA 2AD YA 4N

And whenever this is so the straight line AC is the upright side of the figure (r. 50)




PROPOSITION 59 (PROBLEM)

Guen two bounded straight lines perpendicular to each other to find opposite sections whose diameter is one of the guen straight lines and whose vertex is the ends of the straight line and where the straight lines dropped in each of the sections at a guen angle will equal in square the rectangles applied to the other of the straight lines and exceeding by a figure similar to the rectangle contained by the guen straight lines

Let there be the two given bounded straight lines BE and BH perpendicular to each other and let the given angle be G then it is required to describe

opposite sections about one of the straight lines BE and BH , so that the ordinates are dropped at an angle G



And given the two straight lines BE and BH let an hyperbola be described whose transverse diameter will be the straight line BE and the upright side of whose figure will be BH and where the ordinates to BE produced will be at an angle G , and let it be the line ABC for we have already described how this must be done (1 55) Then let the straight line EA be drawn through E perpendicular to BE and equal to BH , and let another hyperbola DEF be likewise described whose diameter is BE and the upright side of whose figure is EA and where the ordinates from the section will be dropped at a same angle G Then it is evident that B and E are opposite sections, and there is one diameter for them, and their uprights are equal

PROPOSITIO 60 (PROBLEM)

Given two straight lines bisecting each other to describe about each of them opposite sections so that the straight lines are their conjugate diameters and the diameter of one pair of opposite sections is equal in square to the figure of the other pair and likewise the diameter of the second pair of opposite sections is equal in square to the figure of the first pair

Let there be the two given straight lines AC and DE bisecting each other then it is required to describe opposite sections about each of them as a diameter so that the straight lines AC and DE are conjugates in them and DE is equal in square to the figure about AC and AC is equal in square to the figure about DE


Let

$$\text{rect } AC \cdot CL = \text{sq } DE$$

and let LC be perpendicular to CA And given two straight lines AC and CL perpendicular to each other let the opposite sections RAG and HCA be described whose transverse diameter will be CA and whose upright side will be CL and where the ordinates from the sections to C will be dropped at the given angle (1 59) Then the straight line DE will be a second diameter of the opposite sections (See Def 1 11), for it is the mean proportion between the sides of the figure, and parallel to an ordinate it has been bisected at B

Then again let

$$\text{rect } DE, DF = \text{sq } AC.$$



and let DF be perpendicular to DE . And given two straight lines ED and DF lying perpendicular to each other, let the opposite sections MDN and OEX be described whose transverse diameter will be DE and the upright side of whose figure will be DF , and where the ordinates from the sections will be dropped to DE at the given angle (159) then the straight line AC will also be a second diameter of the sections MDN and XEO . And so AC bisects the parallels to DE between the sections RAG and HCK and DE the parallels to AC , and thus it was required to do

And let such sections be called conjugate

BOOK TWO

APOLLONIUS TO EUDEMUS


If you are well well and good and I too fare pretty well

I have sent you my son Apollonius bringing you the second book of the conics as arranged by us Go through it then carefully and acquaint those with it worthy of sharing in such things And Philonides the geometer, I introduced to you in Ephesus, if ever he happen about Pergamum, acquaint him with it too And take care of yourself, to be well Good bye

PROPOSITION 1

If a straight line touch an hyperbola at its vertex and from it on both sides of the diameter a straight line is cut off equal to the straight line equal in square to the fourth of the figure, then the straight lines drawn from the center of the section to the ends thus taken on the tangent will not meet the section

Let there be an hyperbola whose diameter is the straight line AB and center C and upright the straight line BF , and let the straight line DE touch the



section at B and let the squares on BD and BE each be equal to the fourth of the figure $AB \cdot BF$ and the straight lines CD and CE be joined and produced

I say that they will not meet the section

For if possible, let CD meet the section at G and from G let the straight line GH be dropped ordinatewise therefore it is parallel to DB (1 17) Since then

$$AB \cdot BF = \text{sq } AB + \text{rect } AB \cdot BF,$$

therefore also

$$GL = LM$$

And since

$$GH = DB,$$

therefore

$$GK > DB$$

And also

$$AM > BE,$$

since also

$$LM > BE$$

therefore

$$\text{rect } MA, AG > \text{rect } DB BE \\ > \text{sq } DB$$

Since then

$$AB \ BF \ \text{sq } CB \ \text{sq } BD \text{ (II 1),}$$

but

$$AB \ BF \ \text{rect } AL \ LB \ \text{sq } LA \text{ (I 21),}$$

and

$$\text{sq } CB \ \text{sq } BD \ \text{sq } CL \ \text{sq } LG,$$

therefore also

$$\text{sq } CL \ \text{sq } LG \ \text{rect } AL \ LB \ \text{sq } LK$$

Since then

$$\begin{array}{l} \text{whole sq } LC \ \text{whole sq } LG \\ \text{part subtr rect } AL, LB \ \text{part subtr sq } LA, \end{array}$$

therefore also

$$\text{sq } LC \ \text{sq } LG \ \text{remainder sq } CB \ \text{remainder rect } MA \ AG$$

that is

$$\text{sq } CB \ \text{rect } MA \ AG \ \text{sq } CB \ \text{sq } DB$$

Therefore

$$\text{sq } DB = \text{rect } MA \ AG,$$

and this is absurd for it has been shown to be greater than it. Therefore the straight line CH is not an asymptote to the section.

PROPOSITION 3


If a straight line touches an hyperbola it will meet both of the asymptotes and it will be bisected at the point of contact, and the square on each of its segments will be equal to the fourth of the figure resulting on the diameter drawn through the point of contact

Let there be the hyperbola ABC and its center E , and asymptotes FE and EG and let some straight line HA touch it at B

I say that the straight line HA produced will meet the straight lines FE and EG

For if possible, let it not meet them and let EB be joined and produced and let ED be made equal to EB therefore the straight line BD is a diameter. Then let the squares on HB and BA each be made equal to the fourth of the figure on BD , and let HI and FA be joined. Therefore they are asymptotes (II 1) and this is absurd (II 2) for FE and EG are supposed asymptotes. Therefore HA produced will meet the asymptotes EF and EC at F and C

I say then also that the squares on BF and BG will each be equal to the fourth of the figure on BD




For let it not be, but if possible let the squares on BH and BK each be equal to the fourth of the figure. Therefore HE and EA are asymptotes (II 1), and this is absurd (II 2). Therefore the squares on BF and BG will each be equal to the fourth of the figure on BD .

PROPOSITION 4 (PROBLEM)

Given two straight lines containing an angle and a point within the angle to describe through the point the section of a cone called hyperbola so that the given straight lines are its asymptotes

Let there be the two straight lines AC and AB containing a chance angle at A and let some point D be given and let it be required to describe through D an hyperbola to the asymptotes CA and AB



Let the straight line AD be joined and produced to E and let AE be made equal to DA and let the straight line DF be drawn through D parallel to AB , and let FC be made equal to AF and let CD be joined and produced to B and let it be contrived that
 $\text{rect } DE, G = \text{sq } CB$
 and with AD extended let an hyperbola be described about it through D so that the

ordinates equal in square the areas applied to G and exceeding by a figure similar to rectangle DE, G . Since then DF is parallel to BA , and

$$CF = FA$$

therefore

$$CD = DB$$

and so

$$\text{sq } CB = 4 \text{ sq } CD$$

And

$$\text{sq } CB = \text{rect } DE, G$$

therefore the squares on CD and DB are each equal to the fourth part of the figure DE, G . Therefore the straight lines AB and AC are asymptotes to the hyperbola described.

PROPOSITION 5

If the diameter of a parabola or hyperbola bisects some straight line, the tangent to the section at the end of the diameter will be parallel to the bisected straight line.

Let there be the parabola or hyperbola ABC whose diameter is the straight line DBE and let the straight line FBG touch the section and let some straight line AEC be drawn in the section making AE equal to EC .

I say that AC is parallel to FG .


For if not let the straight line CH be drawn through C parallel to FG and let HA be joined. Since then ABC is a parabola or hyperbola whose diameter is DE , and tangent FG , and CH is parallel to it therefore

$$CK = KH \quad (\text{I } 46, 47)$$

But also

$$CE = EA$$

Therefore AH is parallel to KE , and this is absurd, for produced it meets BD (I 22).



PROPOSITION 6

If the diameter of an ellipse or circumference of a circle bisects some straight line not through the center, the tangent to the section at the end of the diameter will be parallel to the bisected straight line.

Let there be an ellipse or circumference of a circle whose diameter is the straight line AB and let AB bisect CD a straight line not through the center at the point E .

I say that the tangent to the section at A is parallel to CD .

For let it not be but if possible, let DF be parallel to the tangent at A therefore

$$DG = FG$$

But also

$$DE = EC$$

therefore CF is parallel to GE and this is absurd.


For if G is the center of the section AB the straight line CF will meet the straight line AB (I 23), and if it is not suppose it to be K and let DK be joined and produced to H and let CH be joined. Since then

$$DK = KH,$$

and also

$$DE = EC$$

therefore CH is parallel to AB . But also CF and this is absurd. Therefore the tangent at A is parallel to CD .



PROPOSITION 7

If a straight line touches a section of a cone or circumference of a circle and a parallel to it is drawn in the section and bisected the straight line joined from the point of contact to the midpoint will be a diameter of the section

Let there be a section of a cone or circumference of a circle ABC , and FG tangent to it, and AC parallel to FG and bisected at E , and let BE be joined


I say that BE is a diameter of the section

For let it not be, but if possible, let BH be a diameter of the section Therefore

$AH = HC$ (First Def 1 4),
and thus is absurd, for

$$AE = FC$$


Therefore BH will not be a diameter of the section Then likewise we could show that there is no other than BE



PROPOSITION 8

If a straight line meets an hyperbola in two points produced both ways it will meet the asymptotes and the straight lines cut off on it by the section from the asymptotes will be equal

Let there be the hyperbola ABC and the asymptotes ED and DF , and let some straight line AC meet ABC



I say that produced both ways it will meet the asymptotes

Let AC be bisected at G and let DG be joined Therefore it is a diameter of the section (II 7) therefore the tangent at B is parallel to AC (II 5 6) Then let HBA be the tangent (I 32), then it will meet ED and DF (II 3) Since then AC is parallel to HA and HA meets DK and DH therefore also AC will meet DE and DF

Let it meet them at E and F and

$$HB = BA \text{ (II 3),}$$

therefore also

$$FG = GE$$

And so also

$$CF = AE$$

therefore

$$\text{sq } EG \text{ sq } GD \text{ rect } HG, GB \text{ sq } GA$$

Since then

$$\begin{array}{l} \text{whole sq } EG \text{ whole sq } GD \\ \text{part subtr rect } HG, GB \text{ part subtr sq } AG, \end{array}$$

therefore also

$$\text{remainder sq } EB \text{ remainder rect } DA, AF \text{ sq } EG \text{ sq } GD,$$

or

$$\text{remainder sq } EB \text{ remainder rect } DA, AF \text{ sq } EB \text{ sq } BK$$

Therefore

$$\text{rect } FA, AD = \text{sq } BK$$

Then likewise it could be shown also that

$$\text{rect } DC, CF = \text{sq } BL,$$

therefore also

$$\text{rect } FA, AD = \text{rect } DC, CF$$

PROPOSITION 11

If some straight line cut each of the straight lines containing the angle adjacent to the angle containing the hyperbola, it will meet the section in one point only and the rectangle contained by the straight lines cut off between the containing straight lines and the section will be equal to the fourth part of the square on the diameter drawn parallel to the cutting straight line

Let there be an hyperbola whose asymptotes are CA, AD and let DA be produced to E and through some point E let EF be drawn cutting EA and AC


Now it is evident that it meets the section in one point only for the straight line drawn through A parallel to EF as AB will cut angle CAD and will meet the section (II 2) and be its diameter (I 50), therefore EF will meet the section in one point only (I 26)

Let it meet it at G

I say then also that

$$\text{rect } EG, GF = \text{sq } AB$$

For let the straight line $HGLK$ be drawn ordinatewise through G , therefore



the tangent through B is parallel to GH (II 5) Let it be CD Since then $CB = BD$ (II 3),

therefore

$$\text{sq } CB \text{ or rect } CB, BD \text{ sq } BA \text{ comp } CB \text{ } BA \text{ } DB \text{ } BA$$

But

$$CB \ BA \ HG \ GT,$$

and

$$DB \ BA \ GK \ GE$$

therefore

$$\text{sq } CB \ \text{sq } B1 \ \text{comp } HG \ GT, \ \Lambda G \ GE$$

But also

$$\text{rect } \Lambda G, GH \ \text{rect } EG \ GF \ \text{comp } HG \ GF, KG \ GE,$$

therefore

$$\text{rect } \Lambda G \ GH \ \text{rect } EG \ GF \ \text{sq } CB \ \text{sq } BA$$

Alternately

$$\text{rect } \Lambda G, GH \ \text{sq } CB \ \text{rect } EG, GF \ \text{sq } BA$$

But it was shown

$$\text{rect } \Lambda G, GH = \text{sq } CB \ (\text{II } 10),$$

therefore also

$$\text{rect } EG \ GF = \text{sq } AB$$

PROPOSITION 12

If two straight lines at chance angles are drawn to the asymptotes from some point of those on the section and parallels are drawn to them from some point of those on the section then the rectangle contained by the parallels will be equal to that contained by those straight lines to which they were drawn parallel

Let there be an hyperbola whose asymptotes are AB and BC and let some point D be taken on the section, and from it let DE and DF be dropped to AB and BC and let some other point on the section G be taken and through G let GH and GK be drawn parallel to ED and DF

I say that

$$\text{rect } ED \ DF = \text{rect } HG, GK$$

For let DG be joined and produced to A and C Since then

$$\text{rect } AD \ DC = \text{rect } AG, GC \ (\text{II } 8)$$

therefore

$$AG \ AD \ DC \ CG$$

But

$$AG \ AD \ GH \ ED,$$

and

$$DC \ CG \ DF \ GK$$

therefore


$$GH \ DE \ DF \ GK$$

Therefore


$$\text{rect } ED \ DF = \text{rect } HG \ GK$$

PROPOSITION 13

If in the place bounded by the asymptotes and the section some straight line is drawn parallel to one of the asymptotes, it will meet the section in one point only



Let there be an hyperbola whose asymptotes are CA and AB , and let some point E be taken and through it let EF be drawn parallel to AB



I say that it will meet the section

For if possible, let it not meet it and let some point G on the section be taken and through G let GC and GH be drawn parallel to CA and AB and let

$$\text{rect } CG, GH = \text{rect } AE, EF$$

and let AF be joined and produced, then it will meet the section (1 2) Let it meet it at K , and through K parallel to CA and AB let KL and KD be drawn, therefore

$$\text{rect } CG, GH = \text{rect } LA, KD \text{ (II 12)}$$

And it is supposed that also

$$\text{rect } CG, GH = \text{rect } AE, EF$$

therefore

$$\text{rect } LA, KD \text{ or } \text{rect } KL, LA = \text{rect } AE, EF$$

and thus is impossible for both

$$KL > EF$$

and

$$LA > AE$$

Therefore EF will meet the section

Let it meet it at M

I say then that it will not meet it at any other point

For if possible let it also meet it at N and through M and N let MX and NB be drawn parallel to CA Therefore


$$\text{rect } EM, MX = \text{rect } EN, NB \text{ (II 12),}$$

and this is impossible Therefore it will not meet the section in another point

PROPOSITION 14

The asymptotes and the section if produced indefinitely draw nearer to each other and they reach a distance less than any given distance

Let there be an hyperbola whose asymptotes are AB and AC and a given distance K



I say that AB and AC and the section if produced, draw nearer to each other and will reach a distance less than K

For let EHF and CGD be drawn parallel to the tangent and let AH be joined and produced to Y Since then $\text{rect } CG, GD = \text{rect } FH, HE$ (II 10), therefore

$$DG \cdot FH = HE \cdot CG$$

But

$$DG > FH \text{ (I 8, 26),}$$

therefore also

$$HE > CG$$

Then likewise we could show that the succeeding straight lines are less

Then let the distance EL be taken less than K and through L let LN be drawn parallel to AC therefore it will meet the section (II 13) Let it meet it at N , and through N let MNB be drawn parallel to EF Therefore

$$MN = EL$$

and so

$$MN < K$$

PROBLEM

Then from this it is evident that the straight lines AB and AC are never than all the asymptotes to the section and the angle contained by BA, AC is clearly less than that contained by other asymptotes to the section


PROPOSITION 15

The asymptotes of opposite sections are common

Let there be opposite sections whose diameter is AB and center C

I say that the asymptotes of the sections A and B are common

Let the straight lines DAE and FBG be drawn tangent to the sections through the points A and B they are therefore parallel (I 44, note) Then let



each of the straight lines DA, AE, FB and BG be cut off equal in square to the fourth of the figure applied to AB therefore

$$DA = AE = FB = BG$$

Then let CD, CE, CF and CG be joined. Then it is evident that DC is in a straight line with CG and CE with CF because of the parallels. Since then it is an hyperbola whose diameter is AB and tangent DE and DA and AE are each equal in square to the fourth of the figure applied to AB therefore DC and CE are asymptotes (II 1). For the same reasons then FC and CG are also asymptotes to section B . Therefore the asymptotes of opposite sections are common

PROPOSITION 16

If in opposite sections some straight line is drawn cutting each of the straight lines containing the angle adjacent to the angles containing the sections, it will meet

each of the opposite sections in one point only, and the straight lines cut off on it by the sections from the asymptotes will be equal


For let there be the opposite sections A and B whose center is C and asymptotes DCG and ECF , and let some straight line HA be drawn through cutting each of the straight lines DC and CF

I say that produced it will meet each of the sections in one point only

For since DC and CE are asymptotes of section A , and some straight line HK has been drawn across cutting both of the straight lines containing the adjacent angle DCF , therefore HA produced will meet the section (II 11) Then likewise also B

Let it meet them at L and M

Let the straight line ACB be drawn through C parallel to LM , therefore



$$\text{rect } KL LH = \text{sq } AC \text{ (II 11)}$$

and

$$\text{rect } HM MK = \text{sq } CB \text{ (II 11)}$$

And so also

$$\text{rect } KL LH = \text{rect } HM, MK,$$


and

$$LH = KM$$

PROPOSITION 17

The asymptotes of conjugate opposite sections are common

Let there be conjugate opposite sections whose conjugate diameters are AB and CD , and whose center is E



I say that their asymptotes are common

For let the straight lines FAG , GDH , HBA , and KCF be drawn through the points A , B , C , and D touching the sections, therefore $FGHK$ is a parallelogram (I 44, note) Then let FEH and KEG be joined therefore they are straight lines (II 15) and diagonals of the parallelogram, and they are all bisected at the point E And since the figure on AB is equal to the square on CD (I 60), and

$$CE = ED,$$

therefore each of the squares on FA , AG , KB , and BH is equal to a fourth of the figure on AB Therefore the straight lines FEH and KEG are asymptotes of the sections A and B (II 1) Then likewise we could show that the same straight lines are also asymptotes of the sections C and D Therefore the asymptotes of conjugate opposite sections are common


PROPOSITION 18

If a straight line meeting one of the conjugate opposite sections when produced both ways, falls outside the section, it will meet both of the adjacent sections in one point only

Let there be the conjugate opposite sections A , B , C , and D , and let some straight line EF meet the section C and produced both ways fall outside the section

I say that it will meet both of the sections A and B in one point only

For let GH and KL be asymptotes of the sections Therefore EF meets both



GH and KL (II 3) Then it is evident that it will also meet the sections A and B in one point only (II 16)

PROPOSITION 19

If some straight line is drawn touching some one of the conjugate opposite sections at random, it will meet the adjacent sections and will be bisected at the point of contact


Let there be the conjugate opposite sections A , B , C and D , and let some straight line ECF touch it at C

I say that produced it will meet sections A and B and will be bisected at C

It is evident now that it will meet sections A and B (II 18), let it meet them at G and H

I say that

$$CG = CH$$



For let the asymptotes of the sections KL and MN be drawn. Therefore $EG = FH$ (II 16),

and

$$CE = CF \text{ (II 3),}$$


and

$$CG = CH$$

PROPOSITION 20

If a straight line touches one of the conjugate opposite sections, and two straight lines are drawn through their center, one through the point of contact and one parallel to the tangent until it meet one of the adjacent sections then the straight line touching the section at the point of meeting will be parallel to the straight line drawn through the point of contact and the center, and those through the points of contact and the center will be conjugate diameters of the opposite sections

Let there be conjugate opposite sections whose conjugate diameters are the



straight lines AB and CD and center Y , and let EF be drawn touching the section A , and produced let it meet CY at T , and let EY be joined and produced to λ and through Y let YG be drawn parallel to EF , and through G let HG be drawn touching the section

I say that HG is parallel to YE and GO and $E\lambda$ are conjugate diameters

For let the straight lines KE , GL and CRP be drawn ordinatewise, and let AM and CN be the parameters Since then

$$BA \cdot AM = NC \cdot CD \text{ (I 60),}$$

but

$$BA \cdot AM = \text{rect } YA \cdot AF = \text{sq } KE \text{ (I 37),}$$

and

$$NC \cdot CD = \text{sq } GL = \text{rect } YL \cdot LH \text{ (I 37),}$$

therefore also

$$\text{rect } YA \cdot AF = \text{sq } EA = \text{sq } GL = \text{rect } YL \cdot LH$$

But

$$\text{rect } YA \cdot AF = \text{sq } EK \text{ comp } YK = KE \cdot FA = KE,$$

and

$$\text{sq } GL = \text{rect } YL \cdot LH \text{ comp } GL = LY \cdot GL = LH,$$

therefore

$$\text{ratio comp } YK = KE \cdot FA = KE = \text{ratio comp } GL = LY \cdot GL = LH$$

and of these

$$FA \cdot AE = GL \cdot LY$$

for each of the straight lines EK , AF and FE is parallel to each of the straight lines YL , LG and GY respectively Therefore as remainder

$$YA \cdot AE = GL \cdot LH$$

Also the sides about the equal angles at A and L are proportional therefore triangle EAY is similar to triangle GHL and will have equal the angles the corresponding sides subtend Therefore

$$\text{angle } EYA = \text{angle } LGH$$

But also

$$\text{angle } AYG = \text{angle } LGY$$

and therefore

$$\text{angle } EYG = \text{angle } HGY$$

Therefore EY is parallel to GH

Then let it be contrived that

$$PG \cdot GR = HG \cdot S$$

therefore S is a half of the parameter of the ordinates to the diameter GO in sections C and D (I 51) And since CD is the second diameter of the sections A and B and ET meets it therefore

$$\text{rect } TY \cdot EA = \text{sq } CY$$

for if we draw from E a parallel to AY the rectangle contained by TY and the straight line cut off by the parallel will be equal to the square on CY (I 38)

And therefore

$$TY \cdot EA = \text{sq } TY = \text{sq } TC \text{ (Eucl vi 20)}$$

But

$$TY \cdot EA = TF \cdot FE$$

or

$$TY \cdot FA = \text{trgl } TYF = \text{trgl } EYF \text{ (Eucl vi 1),}$$

and

sq TY sq CY trgl YTF trgl YCP (Eucl vi 19)

or

sq TY sq CY trgl YTF trgl GHY (iii 1)

Therefore

trgl TYF trgl EFY trgl TFY trgl YGH

Therefore

trgl $GHY = \text{trgl } YEF$

But they also have

angle $HGY = \text{angle } YEF$

for EY is parallel to GH , and EF to GY . Therefore the sides about the equal angles are reciprocally proportional (Eucl vi 15) Therefore

$GH : EY = EF : GY$

therefore

rect $HG : GY = \text{rect } YE : EF$

And since

$S : HG = RG : GP$,

and

$RG : GP = YE : EF$,

for they are parallel therefore also

$S : HG = YE : EF$

But with YG taken as common height,

$S : HG = \text{rect } SYG : \text{rect } HGY$,

and

$YE : EF = \text{sq } YE : \text{rect } YEF$

And therefore

$\text{rect } SYG : \text{rect } HGY = \text{sq } YE : \text{rect } YEF$

Alternately

$\text{rect } SYG : \text{sq } EY = \text{rect } HGY : \text{rect } FEY$

But

$\text{rect } HGY = \text{rect } YEF$ (above),

therefore also

$\text{rect } SYG = \text{sq } EY$

And rectangle SYG is a fourth of the figure on GO for

$GY = \text{half } GO$

and S is the parameter and

$\text{sq } EY = \text{fourth sq } EX$

for

$EY = YX$

Therefore the square on EY is equal to the figure on GO . Then likewise we could show also that GO is equal in square to the figure on EX . Therefore EX and GO are conjugate diameters of the opposite sections A, B, C , and D .

PROPOSITION 21

The same things being supposed it is to be shown that the point of meeting of the tangents is on one of the asymptotes

Let there be conjugate opposite sections whose diameters are the straight lines AB and CD and let the straight lines AE and EC be drawn tangent

I say that the point E is on the asymptote

PROPOSITION 23


If in conjugate opposite sections some radius is drawn to any one of the sections, and a parallel is drawn to it meeting the three adjacent sections, then the rectangle contained by the segments produced between the three sections on the straight line drawn is twice the square on the radius

Let there be the conjugate opposite sections A, B, C , and D and let the center of the section be Y , and from the point Y let some straight line CY be drawn to meet any one of the sections, and let KL be drawn parallel to CY cutting the three adjacent sections

I say that

$$\text{rect } KM, ML = 2 \text{ sq } CY$$

Let the asymptotes to the sections, EF and GH , be drawn therefore



$$\text{sq } CY = \text{rect } HM, ME \text{ (II 22)} = \text{rect } HK, KE \text{ (II 11)}$$


And

$$\text{rect } HM, ME + \text{rect } HK, KE = \text{rect } LM, MK$$

because of the straight lines on the ends being equal (II 8 16) Therefore also
 $\text{rect } LM, MK = 2 \text{ sq } CY$

PROPOSITION 24

If two straight lines meet a parabola each at two points and if a point of meeting of neither one of them is contained by the points of meeting of the other, then the straight lines will meet each other outside the section



Let there be the parabola $ABCD$ and let the two straight lines AB and CD meet $ABCD$ and let a point of meeting of neither of them be contained by the points of meeting of the other

I say that the straight lines produced will meet each other

Let the diameters of the section EBF and GCH be drawn through the points B and C therefore they are parallel (I 51, end) and each one cuts the section in one point only (I 26) Then let BC be joined therefore

$$\text{angle } EBC + \text{angle } BCG = 2 \text{ rt angles}$$

and DC and BA produced make angles less than two right angles. Therefore they will meet each other outside the section (I 10, Eucl Post 5)

PROPOSITION 25


If two straight lines meet an hyperbola each at two points and if a point of meeting of neither of them is contained by the points of meeting of the other, then the straight lines will meet each other outside the section, but within the angle containing the section

Let there be an hyperbola, whose asymptotes are AB and AC , and let the two straight lines EF and GH cut the section, and let a point of meeting of neither of them be contained by the points of meeting of the other

I say that the straight lines EF and GH produced will meet outside the section, but within the angle CAB

For let the straight lines AF and AH be joined and produced and let FH be joined. And since the straight lines EF and GH produced cut the angles AFH and AHF , and the said angles are less than two right angles (Eucl I 17), the straight lines EF and GH produced will meet each other outside the section but within the angle BAC

Then we could likewise show it even if the straight lines EF and GH are tangents to the sections




PROPOSITION 26

If in an ellipse or circumference of a circle two straight lines not through the center cut each other then they do not bisect each other

For if possible, in the ellipse or circumference of a circle let the two straight lines CD and EF not through the center bisect each other at G and let the point H be the center of the section and let GH be joined and produced to A and B



Since then the straight line AB is a diameter bisecting EF , therefore the tangent at A is parallel to EF (II 6). We could then likewise show that it is also parallel to CD . And so also EF is parallel to CD . And this is impossible. Therefore CD and EF do not bisect each other



PROPOSITION 27

If two straight lines touch an ellipse or circumference of a circle, and if the straight line joining the points of contact is through the center of the section the tangents will be parallel but if not they will meet on the same side of the center

Let there be the ellipse or circumference of a circle AB , and let the straight lines CD and EBF touch it, and AB be joined, and first let it be through the center



I say that CD is parallel to EF

For since AB is a diameter of the section, and CD touches it at A , therefore CD is parallel to the ordinates to AB (I 17). Then for the same reasons BF is also parallel to the same ordinates. Therefore CD is also parallel to EF .

Then let AB not be through the center, as in the second drawing, and let the diameter AH be drawn, and let KHL be drawn tangent through H . therefore KL is parallel to CD . Therefore EF produced will meet CD on the same side of the center as AB .

PROPOSITION 28

If in a section of a cone or circumference of a circle some straight line bisects two parallel straight lines, then it will be a diameter of the section

For let AB and CD two parallel straight lines in a conic section, be bisected at E and F , and let EF be joined and produced

I say that it is a diameter of the section

For if not let the straight line GFH be so if possible. Therefore the tangent at G is parallel to AB (II 5 6). And so the same straight line is parallel to CD . And GH is a diameter. therefore

$$CH = HD \text{ (First Def 1 4)}$$

and this is impossible for it is supposed


$$CE = ED$$

Therefore GH is not a diameter. Then likewise we could show that there is no other except EF . Therefore EF will be a diameter of the section.

PROPOSITION 29

If in a section of a cone or circumference of a circle two tangents meet the straight line drawn from their point of meeting to the midpoint of the straight line joining the points of contact is a diameter of the section

Let there be a section of a cone or circumference of a circle to which let the straight lines AB and AC meeting at A be drawn tangent and let BC be joined and bisected at D , and let AD be joined



I say that it is a diameter of the section

For if possible let DE be a diameter, and let EC be joined, then it will cut the section (1 35 36) Let it cut it at F , and through F let FKG be drawn parallel to CDB Since then

$$CD = DB$$


also

$$FH = HG$$

And since the tangent at L is parallel to BC (11 5, 6), and FG is also parallel to BC , therefore also FG is parallel to the tangent at L Therefore

$$FH = HK \text{ (1 46, 47)}$$

and this is impossible Therefore DF is not a diameter Then likewise we could show that there is no other except AD



PROPOSITION 30

If two straight lines tangent to a section of a cone or to a circumference of a circle meet the diameter drawn from the point of meeting will bisect the straight line joining the points of contact

Let there be the section of a cone or circumference of a circle BC , and let two tangents BA and AC be drawn to it meeting at A and let BC be joined and let AD be drawn through A as a diameter of the section


I say that

$$DB = DC$$

For let it not be but if possible let

$$BE = EC$$

and let BE be joined therefore AF is a diameter of the section (11 29) But AD is also a diameter and this is absurd For if the section is an ellipse the point A at which the diameters meet each other, will be a center outside the section and this is impossible and if the section is a parabola, the diameters meet each other (1 51 end) and if it is an hyperbola and the straight lines BA and AC meet the section without containing one another's points of meeting then the center is within the angle containing the hyperbola (11 25), but it is also on it, for it has been supposed a center since DA and AE are diameters (1 51 end) and this is absurd Therefore BE is not equal to EC



PROPOSITION 31


If two straight lines touch each of the opposite sections then if the straight line joining the points of contact falls through the center the tangents will be parallel, but if not, they will meet on the same side as the center

Let there be the opposite sections 1 and B , and let the straight lines CAD and EBF be tangent to them at A and B and let the straight line joined from

A to B fall first through the center of the sections

I say that CD is parallel to EF

For since they are opposite sections of which AB is a diameter and CD touches one of them at A , therefore the straight line drawn through B parallel




to CD touches the section (1 44 note) But EF also touches it therefore CD is parallel to EF

Then let the straight line from A to B not be through the center of the sections and let AG be drawn as a diameter of the sections and let HA be drawn tangent to the section therefore HK is parallel to CD and since the straight lines EF and HK touch an hyperbola therefore they will meet (11 25 end) And HK is parallel to CD therefore also the straight lines CD and EF produced will meet And it is evident they are on the same side as the center

PROPOSITION 32

If straight lines meet each of the opposite sections in one point when touching or in two points when cutting, and when produced the straight lines meet then their point of meeting will be in the angle adjacent to the angle containing the section



Let there be opposite sections and the straight lines AB and CD

touching the opposite sections in one point or cutting them in two points, and let them meet when produced


I say that their point of meeting will be in the angle adjacent to the angle containing the section

Let FG and HA be asymptotes to the sections, therefore AB produced will meet the asymptotes (II 8) Let it meet them at H and G And since FA and HG are supposed as meeting it is evident that either they will meet in the place under angle HLF or in that under angle ALG And likewise also, if they touch (II 3)

PROPOSITION 33

If a straight line meeting one of the opposite sections, when produced both ways, falls outside the section, it will not meet the other section, but will fall through the three places of which one is that contained by the angle containing the section, and two are those contained by the angle adjacent to the angle containing the section

Let there be the opposite sections A and B , and let some straight line CD



cut A , and when produced both ways, let it fall outside the section

I say that the straight line CD does not meet the section B

For let EF and GH be drawn as asymptotes to the sections therefore CD produced will meet the asymptotes (II 8) And it only meets them in the points E and H And so it will not meet the section B

And it is evident that it will fall through the three places For if some straight line meets both of the opposite sections it will meet neither of the opposite sections in two points For if it meets it in two points by what has just been proved it will not meet the other section

PROPOSITION 34


If some straight line touch one of the opposite sections and a parallel to it be drawn in the other section then the straight line drawn from the point of contact to the midpoint of the parallel will be a diameter of the opposite sections

Let there be the opposite sections A and B , and let some straight line CD touch one of them A at A , and let EF be drawn parallel to CD in the other section, and let it be bisected at G and let AG be joined

I say that AG is a diameter of the opposite sections

For if possible, let AHA be Therefore the tangent at H is parallel to CD

(II.31) But CD is also parallel to EF , and therefore the tangent at H is parallel to EF . Therefore



$$EK = KF \text{ (I. 47),}$$

and this is impossible, for

$$EG = GF$$

Therefore AH is not a diameter of the opposite sections

Therefore AB is

PROPOSITION 35

If a diameter in one of the opposite sections bisects some straight line the straight line touching the other section at the end of the diameter will be parallel to the bisected straight line

Let there be the opposite sections A and B and let their diameter AB bisect the straight line CD in section B at E



I say that the tangent to the section at A is parallel to CD

For if possible let DF be parallel to the tangent to the section at A , therefore

$$DG = GF \text{ (I. 48)}$$

But also


$$DE = EC$$

Therefore CF is parallel to EG , and this is impossible, for produced it meets it (I. 22). Therefore DF is not parallel to the tangent to the section at A nor is any other straight line except CD

PROPOSITION 36

If parallel straight lines are drawn one in each of the opposite sections then the straight line joining their midpoints will be a diameter of the opposite sections

Let there be the opposite sections A and B , and let the straight lines CD and EF be drawn one in each of them, and let them be parallel, and let them both



be bisected at points G and H and let GH be joined

I say that GH is a diameter of the opposite sections

For if not, let GA be one. Therefore the tangent to A is parallel to CD (II 5), and so also to EF . Therefore

$$EA = AF \text{ (I 48)}$$

and this is impossible since also

$$EH = HF$$

Therefore GA is not a diameter of the opposite sections. Therefore GH is

PROPOSITION 37

If a straight line not through the center cuts the opposite sections then the straight line joined from its midpoint to the center is a so-called upright diameter of the opposite sections and the straight line drawn from the center parallel to the bisected straight line is a transverse diameter conjugate to it.

Let there be the opposite sections A and B , and let some straight line CD not through the center cut the sections A and B and let it be bisected at E , and let Y be the center of the sections, and let YE be joined, and through Y let AB be drawn parallel to CD .


I say that the straight lines AB and CD are conjugate diameters of the sections.

For let DY be joined and produced to F , and let CF be joined. Therefore

$$DY = YF \text{ (I 30)}$$

But also

$$DE = EC,$$



therefore EY is parallel to FC . Let BA be produced to G . And since

$$D\mathbf{y} = \mathbf{y}F,$$

therefore also

$$EY = FG,$$

and so also

$$CG=FG$$

Therefore the tangent at A is parallel to CF (II 5), and so also to EY . Therefore EY and AB are conjugate diameters (I 16).


PROPOSITION 38

If two straight lines meeting touch opposite sections the straight line joined from the point of meeting to the midpoint of the straight line joining the points of contact will be a so-called upright diameter of the opposite sections, and the straight line drawn through the center parallel to the straight line joining the points of contact is a transverse diameter conjugate to it

Let there be the opposite sections A and B , and CY and $1 D$ touching the sections and let CD be joined and bisected at E , and let EY be joined

I say that the diameter EY is a so-called upright and the straight line drawn through the center parallel to CD is a transverse diameter conjugate to it

For if possible, let EF be a diameter and let F be a point taken at random, therefore DY will meet EF . Let it meet it at F , and let CF be joined, therefore



CF will hit the section (1 32) Let it hit it at *A*, and through *A* let *AB* be drawn parallel to *CD* Since then *EF* is a diameter, and bisects *CD*, it also bisects the parallels to it (First Def 1 4) Therefore

$$AG = GB$$

And since

$$CE = ED$$

and is on triangle CFD therefore also

$$AG = GK$$

And so also


$$GK = GB$$

and thus is impossible. Therefore EF will not be a diameter

PROPOSITION 39

If two straight lines meeting touch opposite sections the straight line drawn through the center and the point of meeting of the tangents bisects the straight line joining the points of contact

Let there be the opposite sections A and B , and let two straight lines CE and



ED be drawn touching A and B , and let CD be joined, and let EF be drawn as a diameter

I say that

$$CF = FD$$

For if not, let CD be bisected at G , and let GE be joined, therefore GE is a diameter (II 38). But EF is also, therefore E is the center (I 31, end). Therefore the point of meeting of the tangents is at the center of the sections and this is absurd (II 32). Therefore CF is not unequal to FD . Therefore equal.


PROPOSITION 40

If two straight lines touching opposite sections meet and through the point of meeting a straight line is drawn, parallel to the straight line joining the points of contact and meeting the sections then the straight lines drawn from the points of meeting to the midpoint of the straight line joining the points of contact touch the sections

Let there be the opposite sections A and B , and let two straight lines CE and ED be drawn touching A and B , and let CD be joined and through E let FEG be drawn parallel to CD , and let CD be bisected at H , and let FH and HG be joined

I say that FH and HG touch the sections

Let EH be joined, therefore EH is an upright diameter and the straight line drawn through the center parallel to CD a transverse diameter conjugate to it (II 38). And let the center Y be taken and let AYB be drawn parallel to CD therefore HE and AB are conjugate diameters. And CH has been drawn



ordinatewise to the second diameter and CE has been drawn touching the section and meeting the second diameter. Therefore the rectangle EY, YH is

equal to the square on the half of the second diameter (I 38); that is to the fourth part of the figure on AB (Second Def I 10) And since FE has been drawn ordinatewise and FH joined, therefore FH touches the section A (I 38) Likewise then also GH touches section B Therefore FH and GH touch sections A and B


PROPOSITION 41

If in opposite sections two straight lines not through the center cut each other, then they do not bisect each other

Let there be the opposite sections A and B , and in A and B let the two straight lines CB and AD not through the center cut each other at E

I say that they do not bisect each other

For if possible, let them bisect each other, and let Y be the center of the sections and let EY be joined, therefore EY is a diameter (II 37) Let YF be drawn through Y parallel to BC , therefore YF is a diameter and conjugate to




EY (II 37) Therefore the tangent at F is parallel to EY (First Def I 6) Then for the same reasons with HK drawn parallel to AD the tangent at H is parallel to EY and so also the tangent at F is parallel to the tangent at H , and this is absurd for it has been shown it also meets it (II 31) Therefore the straight lines CB and AD not being through the center do not bisect each other

PROPOSITION 42

If in conjugate opposite sections two straight lines not through the center cut each other, they do not bisect each other

Let there be the conjugate opposite sections A , B , C and D , and in A , B , C



and D let the two straight lines not through the center EF and GH , cut each other at K

I say that they do not bisect each other

For if possible, let them bisect each other, and let the center of the sections be Y , and let AB be drawn parallel to EF and CD to HG , and let AY be joined, therefore AY and AB are conjugate diameters (II 37) Likewise YH and CD are also conjugate diameters. And so also the tangent at A is parallel to the tangent at C , and this is impossible, for it meets it, since the tangent at C cuts the sections A and B (II 19), and the tangent at A sections C and D , it is evident also that their point of meeting is in the place under angle AYC (II 21). Therefore the straight lines EF and GH not being through the center do not bisect each other


PROPOSITION 43

If a straight line cuts one of the conjugate opposite sections in two points, and through the center one straight line is drawn to the midpoint of the cutting straight line and another straight line is drawn parallel to the cutting straight line, they will be conjugate diameters of the opposite sections

Let there be the conjugate opposite sections A , B , C and D , and let some straight line cut section A at the two points E and F , and let EF be bisected at G , and let Y be center, and let YG be joined, and let CY be drawn parallel to EF .

I say that AY and YC are conjugate diameters

For since AY is a diameter and bisects EF , the tangent at A is parallel to



EF (II 5) and so also to CY . Since then they are opposite sections and a tangent has been drawn to one of them A at A and from the center Y one straight line YG is joined to the point of contact and another CY has been drawn parallel to the tangent therefore YA and YC are conjugate diameters for this has been shown before (II 20)

PROPOSITION 44 (PROBLEM)

Given a section of a cone to find a diameter

Let there be the given conic section on which are the points A , B , C , D and E . Then it is required to find a diameter

Let it have been done and let it be CH . Then with DF and EH drawn or drawnwise and produced


$$DF = FB,$$

and

$$EH = HA \text{ (First Def 1 4)}$$

If then we fix the straight lines BD and EA in position to be parallel, the points H and F will be given. And so HFC will be given in position.


Then it will be constructed (*συμβησεται*) thus let there be the given conic section on which are the points A, B, C, D and E , and let the straight lines BD and AE be drawn parallel and be bisected at F and H . And the straight line FH joined will be a diameter of the section (First Def 1 4). And in the same way we could also find an indefinite number of diameters.



PROPOSITION 45 (PROBLEM)

Given an ellipse or hyperbola to find the center

And this is evident, for if two diameters of the section, AB and CD are



drawn through (II 44) the point at which they cut each other will be the center of the section, as indicated below.

PROPOSITION 46 (PROBLEM)

Given a section of a cone to find the axis

Let the given section of a cone first be a parabola on which are the points F, C and E . Then it is required to find its axis.

For let AB be drawn as a diameter of it (I 44). If then AB is an axis, what was enjoined would have been done; but if not, let it have been done, and let CD be the axis, therefore the axis CD is parallel to AB (I 51, end) and bisects the straight lines drawn perpendicular to it (First Def 1 7). And the perpendiculars to CD are also perpendiculars to AB , and so CD bisects the perpendic-

ulars to AB . If then I fix EF the perpendicular to AB , it will be given in position, and therefore

$$ED = DF,$$

therefore the point D is given. Therefore through the given point D , CD has been drawn parallel in position to AB , therefore CD is given in position.

Then it will be constructed thus: let there be the given parabola on which are the points F , E and A , and let AB , a diameter of it, be drawn (I 44), and let BE be drawn perpendicular to it and let it be produced to F . If then

$$EB = BF$$

it is evident that AB is the axis (First Def 1 7), but if not, let EF be bisected by D , and let CD be drawn parallel to AB . Then it is evident that CD is the axis of the section, for being parallel to a diameter that is being a diameter (I 51, end), it bisects EF at right angles. Therefore CD has been found as the axis of the given parabola (First Def 1 7).

And it is evident that the parabola has only one axis. For if there is another, as AB , it will be parallel to CD (I 51, end) and it cuts EF , and so it also bisects it (First Def 1 4). Therefore

$$BE = BF$$



and this is absurd.

PROPOSITION 47 (PROBLEM)

Given an hyperbola or ellipse to find the axis

Let there be the hyperbola or ellipse ABC then it is required to find its axis.

Let it have been found and let it be AD , and A the center of the section, therefore AD bisects the ordinates to itself and at right angles (First Def 1 7).



Let the perpendicular CDA be drawn and let KA and AC be joined. Since then

$$CD = DA,$$

therefore

$$CA = KA$$

If then we fix the given point C , CA will be given. And so the circle described with center A and radius AC will also pass through A and will be given in position. And the section ABC is also given in position, therefore the point A

is given. But the point C is also given, therefore CA is given in position. Also

$$CD = D - 1$$

therefore the point D is given. But also K is given therefore DK is given in position.

Then it will be constructed thus let there be the given hyperbola or ellipse ABC , and let K be taken as its center and let a point C be taken at random on the section and let the circle CEA with center K and radius KC be described, and let CA be joined and bisected at D , and let AC , KD and KA be joined and let KD be drawn through to B

Since then

$AD=DC$

and DK is common, therefore the two straight lines CD and DK are equal to the two straight lines AD and DK and

$$\text{base } K \cdot 1 = \text{base } KC$$

Therefore KBD bisects ADC at right angles. Therefore AD is an axis (First Def. 17).


Let MKN be drawn through K parallel to CA therefore MA is the axis of the section conjugate to BK (First Def 1.8)

PROPOSITION 48 (PROBLEM)

Then with these things shown, let it be next in order to show that there are no other axes of the same sections

For if possible let there also be another axis KG . Then in the same way as before with AH drawn perpendicular

$AH = HL$ (First Def 1 4),



and so also

$$AK=KL$$

But also

$$AK = KC$$

therefore

$$AL = KC,$$

and this is absurd

Now that the circle AEC does not hit the section also in another point between the points A, B and C , is evident in the case of the hyperbola, and in the case of the ellipse let the perpendiculars CR and LS be drawn Since then

$$KC = KL,$$

for they are radii, also

$$\text{sq } KC = \text{sq } AL$$

But

$$\text{sq } CR + \text{sq } RK = \text{sq } CK,$$

and

$$\text{sq } AS + \text{sq } SL = \text{sq } LA,$$

therefore

$$\text{sq } CR + \text{sq } RA = \text{sq } AS + \text{sq } SL$$

Therefore

$$\begin{aligned} &\text{difference between sq } CR \text{ and sq } SL = \\ &\text{difference between sq } AS \text{ and sq } RA \end{aligned}$$

Again since

$$\text{rect } MR, RN + \text{sq } RA = \text{sq } AM,$$

and also

$$\text{rect } MS, SN + \text{sq } SA = \text{sq } AM \text{ (Eucl II 5)}$$

therefore

$$\text{rect } MR, RN + \text{sq } RK = \text{rect } MS, SN + \text{sq } SA$$

Therefore

$$\begin{aligned} &\text{difference between sq } SA \text{ and sq } AR = \\ &\text{difference between rect } MR, RN \text{ and rect } MS, SN \end{aligned}$$

And it was shown that

$$\begin{aligned} &\text{difference between sq } SA \text{ and sq } AR = \\ &\text{difference between sq } CR \text{ and sq } SL \end{aligned}$$

therefore

$$\begin{aligned} &\text{difference between sq } CR \text{ and sq } SL = \\ &\text{difference between rect } MR, RN \text{ and rect } MS, SN \end{aligned}$$

And since CR and LS are ordinates

$$\text{sq } CR \text{ rect } MR, RN \text{ sq } SL \text{ rect } MS, SN \text{ (I 21)}$$

But the same difference was also shown for both therefore

$$\text{sq } CR = \text{rect } MR, RN$$

and

$$\text{sq } SL = \text{rect } MS, SN \text{ (Eucl I 16, 17 9)}$$

Therefore the line ICM is a circle, and this is absurd, for it is supposed an ellipse

PROPOSITION 49 (PROBLEM)

Given a section of a cone and a point not within the section to draw from the point a straight line touching the section in one point

Let the given section of a cone first be a parabola whose axis is BD Then it is

required to draw a straight line as prescribed from the given point which is not within the section

Then the given point is either on the line or on the axis or somewhere else outside

Now let it be on the line, and let it be A , and let it have been done, and let it be AE , and let AD be drawn perpendicular, then it will be given in position And

$$BE = BD \text{ (I 35)}$$

and BD is given, therefore BE is also given And the point B is given, therefore E is also given But A also therefore AE is given in position


Then it will be constructed thus let AD be drawn perpendicular from A , and let BE be made equal to BD , and let AE be joined Then it is evident that it touches the section (I 33)

Again let the given point E be on the axis and let it have been done and let AE be drawn tangent, and let AD be drawn perpendicular, therefore

$$BE = BD \text{ (I 35)}$$

And BE is given therefore also BD is given And the point B is given, therefore D is also given And DA is perpendicular, therefore DA is given in position Therefore the point A is given But also E , therefore AE is given in position

Then it will be constructed thus let BD be made equal to BE , and from D let DA be drawn perpendicular to ED , and let AE be joined Then it is evident that AE touches (I 33)



And it is evident also that even if the given point is the same as B the straight line drawn from B perpendicular touches the section (I 17)

Then let C be the given point and let it have been done and let CA be it and through C let CF be drawn parallel to the axis that is to BD therefore CF is given in position And from A let AF be drawn ordinatewise to CF then


$$CG = FG \text{ (I 35)}$$

And the point G is given therefore F is also given And FA has been

erected ordinatewise that is parallel to the tangent at G (I 32) therefore FA is given in position Therefore A is also given but also C Therefore CA is given in position

It will be constructed thus let CF be drawn through C parallel to BD , and let FG be made equal to CG , and let FA be drawn parallel to the tangent at G (above) and let AC be joined It is evident then that this will do the problem (I 33)

Again let it be an hyperbola whose axis is DBC and center H , and asymptotes HE and HF . Then the given point will be given either on the section or



on the axis or within angle EHF or in the adjacent place or on one of the asymptotes containing the section or in the place between the straight lines containing the angle vertical to angle EHF .

Let the point A first be on the section and let it have been done, and let AG be tangent and let AD be drawn perpendicular and let BC be the transverse side of the figure, then

$$CD : DB :: CG : GB \quad (1.36)$$

And the ratio of CD to DB is given, for both the straight lines are given therefore also the ratio of CG to GB is given. And BC is given, therefore point G is given. But also A therefore AG is given in position.

It will be constructed thus let AD be drawn perpendicular from A , and let

$$CG : GB :: CD : DB$$

and let AG be joined. Then it is evident that AG touches the section (1.34).

Then again let the given point G be on the axis and let it have been done, and let AG be drawn tangent, and let AD be drawn perpendicular. Then for the same reasons

$$CG : GB :: CD : DB \quad (1.36)$$

And BC is given therefore the point D is given. And DA is perpendicular, therefore DA is given in position. And also the section is given in position therefore the point A is given. But also G therefore AG is given in position.

Then it will be constructed thus let the other things be supposed the same, and let it be contrived that


$$CG : GB :: CD : DB$$

and let DA be drawn perpendicular and let AC be joined. Then it is evident that AG does the problem (1.34) and that from G another tangent to the section could be drawn on the other side.

With the same things supposed let the given point A be in the place inside angle EHF , and let it be required to draw a tangent to the section from A . Let it have been done and let it be AK and let AH be joined and produced, and let HN be made equal to LH therefore they are all given. Then also LN will be given. Then let AM be drawn ordinately to MN then also

$$AK : AI :: AM : MI$$

And the ratio of NA to AL is given, therefore also the ratio of NM to ML is given. And the point L is given, therefore also M is given. And MA has been erected parallel to the tangent at L therefore MA is given in position. And also the section ALB is given in position, therefore the point A is given. But K is also given, therefore AK is given.



Then it will be constructed thus: let the other things be supposed the same and the given point K , and AH be joined and produced, and

let HN be made equal to HL and let it be contrived that

$$NK : AL = NM : ML$$

and let MA be drawn parallel to the tangent at L (above), and let KA be joined, therefore KA touches the section (I 34)

And it is evident that a tangent to the section could also be drawn to the other side.


With the same things supposed let the given point F be on one of the asymptotes containing the section, and let it be required to draw from F a tangent to the section. And let it have been done, and let it be FAE , and through A let AD be drawn parallel to EH then

$$DH = DF$$

since also

$$FA = AE \text{ (II 3)}$$

And FH is given, therefore also point D is given. And through the given point D DA has been drawn parallel in position to EH , therefore DA is given in position. And the section is also given in position, therefore the point A is given. But F is also given, therefore the straight line FAE is given in position.



Then it will be constructed thus: let there be the section AB and asymptotes EH and HF , and the given point F on one of the asymptotes containing the section and let FH be bisected at D and through D

let DA be drawn parallel to HE and let FA be joined. And since

$$FD = DH,$$


therefore also

$$FA = AE$$

And so by things shown before, the straight line FAE touches the section (II 9)

With the same things supposed let the given point be in the place under the

Then it will be constructed thus let AD be drawn perpendicular, and let CG GB CD DE ,




and let AG be joined Then it is evident that AG touches, as also in the case of the hyperbola (I 34)

Then again let the given point be K , and let it be required to draw a tangent Let it have been done, and let it be KA , and let the straight line KLH be joined to the center H and produced to N , then

it will be given in position And if AM is drawn ordinatewise, then

NA KI NM ML (I 36)



And the ratio of NK KL is given, therefore the ratio of MN to LM is also given Therefore the point M is given And MA has been erected ordinatewise, for it is parallel to the tangent at L , therefore MA is given in position Therefore the point A is given But also K , therefore KA is given in position


And the construction (*σινθεσις*) is the same as for the preceding

PROPOSITION 50 (PROBLEM)

Given the section of a cone to draw a tangent which will make with the axis, on the same side as the section, an angle equal to a given acute angle

Let the section of a cone first be a parabola whose axis is AB then it is required to draw a tangent to the section which will make with the axis AB , on the same side as the section an angle equal to the given acute angle

Let it have been done and let it be CD therefore angle BDC is given Let BC



be drawn perpendicular then the angle at B is also given Therefore the ratio of DB to BC is given But the ratio of BD to BA is given therefore also the ratio of AB to BC is given And the angle at B is given therefore angle BAC

is also given. And it is in position with respect to BA and the given point A , therefore CA is given in position. And the section is also given in position, therefore the point C is given. And CD touches, therefore CD is given in position.

Then the problem will be constructed thus: let the given section of a cone first be a parabola whose axis is AB , and the given acute angle, angle EFG , and let some point E be taken on EF , and let EG be drawn perpendicular, and let FG be bisected by H , and let HE be joined, and let angle BAC be constructed equal to angle GHE , and let BC be drawn perpendicular, and let AD be made equal to BA , and let CD be joined. Therefore CD is tangent to the section (I 33).

I say then that

$$\text{angle } CDB = \text{angle } EFG$$

For since

$$FG \quad GH \quad DB \quad BA$$

and

$$HG \quad GE \quad AB \quad BC,$$

therefore *ex aequali*

$$FG \quad GE \quad DB \quad BC,$$

And the angles at G and B are right angles, therefore

$$\text{angle at } F = \text{angle at } D$$

Let the section be an hyperbola and let it have been done and let CD be tangent and let the center of the section Γ be taken, and let $C\Gamma$ be joined and let CE be perpendicular: therefore the ratio of rectangle $YE \cdot ED$ to the square on CE is given: for it is the same as the transverse to the upright (I 37). And the ratio of the square on CE to the square on ΓD is given, for each of the rectangles $CD \cdot DE$ and $DE \cdot EC$ is given. Therefore the ratio of rectangle $YE \cdot ED$ to the square on ED is given: and so also the ratio of

YE to ED is given. And the angle at E is given, therefore the angle at Γ is also given.

Then some straight line $C\Gamma$ has been drawn across in position with respect to the straight line YE and to the given point Γ at a given angle: therefore $C\Gamma$ is given in position. And the section


is also given in position: therefore the point C is given. And CD has been drawn across as tangent: therefore CD is given in position.

Let the asymptote to the section ΓF be drawn: therefore CD produced will meet the asymptote (II 3). Let it meet it at F . Therefore

$$\text{angle } FDE > \text{angle } \Gamma \Gamma D$$


Therefore, for the construction, the given acute angle will have to be greater than half the angle contained by the asymptotes.

Then the problem will be constructed thus: let there be the given hyperbola whose axis is AB , and asymptote YF , and the given acute angle $A\Gamma G$ greater than angle $A\Gamma F$, and let



angle $\angle HHL = \text{angle } \angle A\gamma F$,

and let AF be drawn from A perpendicular to AB , and let some point G be



taken on GH , and let GK be drawn from it perpendicular to HK . Since then

angle $\angle \gamma A = \text{angle } \angle LHA$,

and also the angles at A and K are right, therefore

$$\frac{\gamma A}{HK} = \frac{AF}{KL} = \frac{HK}{KG}$$

therefore also

$$\gamma A \cdot AF > HK \cdot KG$$

And so also

$$\text{sq } \gamma A + \text{sq } AF > \text{sq } HK + \text{sq } KG$$

But

$$\text{sq } \gamma A + \text{sq } AF = \text{transverse upright (II 1)}$$

therefore also

$$\text{transverse upright} > \text{sq } HK + \text{sq } KG$$

If then we shall contrive that

$$\text{sq } \gamma A + \text{sq } AF = \text{some other sq } KG,$$

it will be greater than the square on HA . Let it be the rectangle MK, KH and let GM be joined. Since then

$$\text{sq } MA > \text{rect } MK, KH,$$

therefore

$$\begin{aligned} \text{sq } MA + \text{sq } KG &> \text{rect } MK, KH + \text{sq } KG \\ &> \text{sq } \gamma A + \text{sq } AF \end{aligned}$$

And if we shall contrive that

$$\text{sq } MA + \text{sq } KG = \text{sq } \gamma A + \text{some other}$$

it will be to a magnitude less than the square on AF and the straight line joined from γ to the point taken will make similar triangles and therefore

$$\text{angle } \angle \gamma A > \text{angle } \angle CMK$$

Let angle $\angle AYC$ be made equal to angle $\angle GMK$ therefore γC will cut the section (II 2). Let it cut it at C , and from C let CD be drawn tangent to the section (II 49) and CE drawn perpendicular, therefore triangle CYE is similar to triangle GMA . Therefore

$$\text{sq } \gamma E + \text{sq } EC = \text{sq } MA + \text{sq } KG$$

But also

transverse upright rect $\angle F E D$ sq EC (i 37),

and

transverse upright rect MA, AH sq AG

And inversely

sq CE rect $\angle E, ED$ sq GA rect MA, AH ,

therefore *ex aequali*

sq $\angle F$ rect $\angle E F D$ sq MA rect MA, AH

And therefore

$\angle E E D$ $MA AH$

But also we had

$CE EY$ $GA AM$,

therefore *ex aequali*


$CE ED$ $GA AH$

And the angles at E and A are right angles, therefore

angle at D = angle GHA

Let the section be an ellipse whose axis is AB . Then it is required to draw a tangent to the section which with the axis will contain, on the same side as the section, an angle equal to the given acute angle

Let it have been done and let it be CD . Therefore angle CDA is given. Let CE be drawn perpendicular, therefore the ratio of the square on DE to the square on EC is given. Let Y be the center of the section, and let CY be joined. Then the ratio of the square on CE to the rect angle $DE EY$ is given for it is the same as the ratio of the upright to the transverse (i 37) and therefore the ratio of the square on DE to rectangle DE, EY is given, and therefore the ratio of DE to EY is given. And of DE to FC therefore also the ratio of CE to FY is given. And the angle at E is right, therefore the angle at Y is given. And it is given with respect to a straight line given in position and to a given point, therefore the point C is given. And from the given point C let CD be drawn tangent therefore CD is given in position.



Then the problem will be constructed thus: let there be the given acute angle FGH and let some point I be taken on FG , and let FH be drawn perpendicular and let it be contrived that

upright transverse sq FH rect $GH HA$,

and let AF be joined and let I be the center of the section and let angle $AIY C$ be constructed equal to angle GAF and let CD be drawn tangent to the section (ii 49)

I say that CD does the problem that is

angle CDE = angle FGH

For since

$$YE \cdot EC = KH \cdot FH,$$

therefore also

$$\text{sq } YE = \text{sq } EC - \text{sq } KH = \text{sq } FH$$

But also

$$\text{sq } EC \cdot \text{rect } DE, EY = \text{sq } FH \cdot \text{rect } KH, HG,$$

for each is the same ratio as that of the upright to the transverse (I 37, and above) And *ex aequali*, therefore

$$\text{sq } YE \cdot \text{rect } DE, EY = \text{sq } KH \cdot \text{rect } KH, HG$$

And therefore

$$YE \cdot ED = KH \cdot HG$$

But also

$$YE \cdot EC = KH \cdot FH,$$

ex aequali, therefore

$$DE \cdot EC = HG \cdot FH$$


And the sides about the right angles are proportional therefore
angle $CDF = \text{angle } FGH$

Therefore CD does the problem

PROPOSITION 51 (PROBLEM)

Given a section of a cone to draw a tangent which with the diameter drawn through the point of contact will contain an angle equal to a given acute angle

Let the given section of a cone first be a parabola whose axis is AB and the given angle H then it is required to draw a tangent to the parabola which with



the diameter from the point of contact will contain an angle equal to the angle at H

Let it have been done and let CD be drawn a tangent making with the diameter EC drawn through the point of contact angle ECD equal to angle H and let CD meet the axis at D (I 24) Since then AD is parallel to EC (I 51, end)

$$\text{angle } ADC = \text{angle } ECD$$

But angle ECD is given for it is equal to angle H , therefore angle ADC is also given

Then it will be constructed thus let there be a parabola whose axis is AB ,

and the given angle H . Let CD be drawn a tangent to the section making with the axis the angle ADC equal to angle H (II 50), and through C let EC be drawn parallel to AB . Since then

$$\text{angle } H = \text{angle } ADC,$$


and

$$\text{angle } ADC = \text{angle } ECD,$$

therefore also

$$\text{angle } H = \text{angle } ECD$$

Let the section be an hyperbola whose axis is AB and center E , and asymptote ET , and the given acute angle Q , and let CD be tangent and let CF be joined doing the problem, and let CG be drawn perpendicular. Therefore the ratio of the transverse to the upright is given, and so also the ratio of rectangle EG, GD to the square on CG (I 37). Then let some given straight line FL be laid out, and on it let there be described a segment of a circle admitting an angle equal to angle Q (Fuel III 33) therefore it will be greater than a semi circle (Fuel III 31). And from some point K of those on the circumference let KL be drawn perpendicular making



$$\text{rect } FL \cdot LH \text{ sq } LK \quad \text{transverse upright}$$

and let FA and KH be joined. Since then

$$\text{angle } FKH = \text{angle } LCD,$$

but also


$$\text{rect } EG \cdot GD \text{ sq } GC \quad \text{transverse upright,}$$

and

$$\text{rect } FL \cdot LH \text{ sq } LK \quad \text{transverse upright,}$$



angle $HFA = \text{angle } CED$

U and let a given straight line *FH* be laid out, and on it let there be described a segment of a circle, greater than semicircle and admitting an angle equal to angle *Q* (Eucl. III 31, 33), and let it be *FAH*, and let the center of the circle *N* be taken, and from *N* let *NO* be drawn perpendicular to *FH* and let *NO* be cut at *P* in the ratio of *UW* to *WY* and through *P* let *PK* be drawn parallel to *FH*, and from *K* let *KL* be drawn perpendicular to *FH* produced and let *FK* and *KH* be joined and let *LK* be produced to *M*, and from *N* let *NA* be drawn perpendicular to it, therefore it is parallel to *FH*. And therefore



NP PO or UW WY XA AL

ZW WY MK KL.

 $ZY \quad YW \quad ML \quad LK$
$$ML \quad LK \quad \text{rect} \quad ML \quad LK \quad \text{sq} \quad LK$$
$$ZY \quad YW \quad \text{rect } ML, LH \quad \text{sq } LK \quad \text{rect } FL, LH \quad \text{sq } LK \quad (\text{Eucl III 36})$$
$$\text{rect } BC \cdot CG \text{ sq } CA \quad \text{rect } EF \cdot FH \text{ sq } DF$$


whole angle A = whole angle D

$$\text{angle } BAG = \text{angle } EDH$$

remaining angle GAC = remaining angle HDF

$$\angle C = \angle F$$
 $GC \quad CA \quad HF \quad FD$ $BC \quad CA \quad EF \quad FD$

ounded ratio is the same with compounded. Th

$$\text{rect } BC \cdot CG \text{ sq } C^4 \quad \text{rect } EF \cdot FH \text{ sq } F^4 D$$

But

$ZI \parallel H$ transverse upright,

therefore also

rect FL, LH sq LK transverse upright

Then let AT be drawn from A perpendicular to AB Since then
sq EA sq AT transverse upright (ii 1),

and also

transverse upright rect FL, LH sq LK ,

and

sq FL sq $LK >$ rect $FI LH$ sq LK ,

therefore also

sq FL sq $IA >$ sq EA sq AT

And the angles at A and L are right angles, therefore

angle $F <$ angle E

Then let angle AEC be constructed equal to angle LFA , therefore EC will meet
the section (ii 2) Let it meet it at C Then let CD be drawn tangent from C
(ii 49), and let CG be drawn perpendicular, then

transverse upright rect FG, GD sq CG (i 37)

Therefore also

rect FL, LH sq LK rect EG, GD sq CG

therefore triangle KFL is similar to triangle ECG and triangle KHL to tri-
angle CGD , and triangle KFH to triangle CED And so

angle $ECD =$ angle $FKH =$ angle Q


And if the ratio of the transverse to the upright is equal to equal, KI
touches the circle FAH (Eucl iii 37), and the straight line joined from the
center to K will be parallel to FH and itself will do the problem

PROPOSITION 52

If a straight line touches an ellipse making an angle with the diameter drawn
through the point of contact it is not less than the angle adjacent to the one con-
tained by the straight lines deflected at the middle of the section

Let there be an ellipse whose axes are AB and CD and center E and let AB
be the major axis and let the straight line
 GFL touch the section and let AC CB and
 FE be joined and let BC be produced to L

I say that angle LFE is not less than angle
 LCA



For FE is either parallel to LB or not

Let it first be parallel and

$AF = EB$,

therefore also

$AH = HC$

And FE is a diameter therefore the tangent at F is parallel to AC (ii 6) But
also FF is parallel to LB therefore $FHCL$ is a parallelogram and therefore
angle $IFH =$ angle ICH

And since AE and FB are each greater than FC angle ACB is obtuse there-
fore angle ICA is acute And so also angle LFE And therefore angle GFE is
obtuse

Then let FE not be parallel to LB , and let FA be drawn perpendicular,

therefore $\angle BE$ is not equal to angle FE 4 But

rt angle at E = rt angle at H ,

therefore it is not true that

$$sq\ BE \quad sq\ EC \quad sq\ EH \quad sq\ HF$$

But

sq BE sq EC rect AE, EB sq EC transverse upright (i 21)


and

transverse upright rect GK, KE sq KF (1 37)

Therefore it is not true that

$$\text{rect } GH, HE \text{ sq } HF \text{ sq } HE \text{ sq } HF$$

Therefore GK is not equal to AE . Let there be laid out a segment of a circle



MUN admitting an angle equal to angle ACB (Eucl III 33) and angle MCB is obtuse therefore MUN is a segment less than a semicircle (Eucl III 31) Then let it be contrived that

GA KE NY XM

and from X let UXY be drawn at right angles and let NU and UM be joined, and let MN be bisected at T and let OTP be drawn at right angles, therefore it is a diameter. Let the center be R and from it let RS be drawn perpendicular, and ON and OM be joined. Since then

angle $MON = \text{angle } ACB$

and AB and MN have been bisected the one at E and the other at T and the angles at E and T are right angles therefore triangles OTN and BEC are similar Therefore

$$\text{sq } TN \quad \text{sq } TO \quad \text{sq } BE \quad \text{sq } EC$$

And since

$TR \approx 5$

and

$RO \geq SU$

therefore

RO TR>SU S\

and *converting*

RO OT SU UY

And doubling the antecedents therefore

$$PO \quad TO < YU \quad UX$$

And separando

PT TO<1 Y UA

But

$PT \perp O$ sq TN sq TO sq BE sq EC transverse upright (i 21),
and

transverse upright rect $GA, \Lambda E$ sq ΛF (i 37),

therefore

$$\begin{aligned} \text{rect } GA, \Lambda E \text{ sq } \Lambda F &< YX \text{ } XU \\ &< \text{rect } YV, XU \text{ sq } XU \\ &< \text{rect } NV, VM \text{ sq } XU \end{aligned}$$

If then we contrive it that

rect $GA, \Lambda E$ sq ΛF rect $M\Lambda, XN$ some other,
it will be greater than the square on ΛU Let it be to the square on ΛW Since
then

$$GK \text{ } KE \text{ } NX \text{ } \Lambda M,$$

and ΛF and YW are perpendicular and

$$\text{rect } GA, \Lambda F \text{ sq } \Lambda F \text{ rect } MY, VN \text{ sq } \Lambda W,$$

therefore

$$\text{angle } GFE = \text{angle } MWN$$

Therefore

$$\text{angle } MUN \text{ or angle } ACB > \text{angle } GFE,$$


and the adjacent angle LFH is greater than angle LCH

Therefore angle LFH is not less than angle LCH

PROPOSITION 53 (PROBLEM)

Given an ellipse to draw a tangent which will make with the diameter drawn through the point of contact an angle equal to a given acute angle then it is required that the given acute angle be not less than the angle adjacent to the angle contained by the straight lines deflected at the middle of the section

Let there be the given ellipse whose major axis is AB and minor axis CD , and center E , and let AC and CB be joined and let angle U be the given angle



not less than angle ACG , and so also angle ACB is not less than angle Y

Therefore angle U is either greater than or equal to angle ACG

Let it first be equal and through E let EA be drawn parallel to BC and through A let AH be drawn tangent to the section (ii 49) Since then

$$AE = EB$$

and

$$AE \parallel B \quad AF \parallel C,$$

therefore

$$AF = FC$$

Then let it be that

transverse upright $QA' \ 1'Y'$,
and let QY' be bisected at Y' Since then
transverse upright $> RO \ ON$,
also

$$QA' \ A'Y' > RO \ ON$$

And *componendo*

$$QY' \ Y'A' > RV \ NO$$

Let the center of the circle be W , and so also

$$Y'Y' \ Y'A' > WN \ NO$$

And *separando*

$$A'Y' \ A'Y' > WO \ ON$$

Then let it be contrived that

$$A'Y' \ A'X' \ WO \text{ less than } ON$$

such as IO , and let YX and XT and WZ be drawn parallel

Therefore

$$A'Y' \ A'X' \ WO \ OI \ ZS \ SY,$$

and *componendo*

$$Y'Y' \ Y'A' \ ZY \ XS$$

And doubling the antecedents

$$QY' \ Y'A' \ TX \ XS$$

And *separando*

$$QA' \ A'X' \text{ or transverse upright } TS \ SX$$

Then let MY and XP be joined, and let angle AEK be constructed on straight line AE at point E equal to angle MPY and through K let KH be drawn touching the section (II 49), and let KL be dropped ordinatewise Since then
angle $MPY = \text{angle } AEK$,

and

$$\text{rt angle at } S = \text{rt angle at } L$$

therefore triangle YSP is equiangular with triangle KEL

And

$$\text{transverse upright } TS \ SY \text{ rect } TS \ SY$$

$$\text{sq } SX \text{ rect } MY \ SP \text{ sq } SY$$

therefore triangle KLE is similar to triangle SPY and triangle MXP to triangle KHE and therefore

$$\text{angle } MXP = \text{angle } HKE$$

But

$$\text{angle } MXP = \text{angle } MNP = \text{angle } Y$$

therefore also

$$\text{angle } HKE = \text{angle } Y$$

And therefore

$$\text{adjacent angle } GKE = \text{adjacent angle } U$$


Therefore GH has been drawn across tangent to the section and making with the diameter AE drawn through the point of contact angle GKE equal to the given angle U and this it was required to do

BOOK THREE

PROPOSITION 1

If straight lines touching a section of a cone or circumference of a circle, meet and diameters are drawn through the points of contact meeting the tangents, the resulting vertically related triangles will be equal

Let there be the section of a cone or circumference of a circle AB and let AC and BD , meeting at E , touch AB , and let the diameters of the section CB and



DA be drawn through A and B meeting the tangents at C and D

I say that

$$\text{trgl } ADE = \text{trgl } EBC$$

For let AF be drawn from A parallel to BD therefore it has been dropped ordinatewise (1 32) Then in the case of the parabola


$$\text{pllg } ADBF = \text{trgl } ACF \text{ (1 42)}$$

and with the common area $AEBF$ subtracted

$$\text{trgl } ADE = \text{trgl } CBE$$

And in the case of the others let the diameters meet at center G Since then

AF has been dropped ordinatewise and AC is a
rect $FG, GC = sq\ BG$ (17)



Therefore

therefore also

$$FG \cdot GB = BG \cdot GC,$$

But

$$FG \cdot GC = sq\ FG = sq\ GB \text{ (Euc vi 29)}$$

and

$$sq\ FG = sq\ GB \implies \text{trgl } AGF = \text{trgl } DGB \text{ (Euc vi 18)}$$

therefore also

$$FG \cdot GC = \text{trgl } AGF = \text{trgl } AGC,$$

Therefore



$$\text{trgl } AGF = \text{trgl } AGC = \text{trgl } AGF = \text{trgl } DGB$$

Let the common area $DGBE$ be subtracted, therefore as re
 $\text{trgl } AGC = \text{trgl } DGB$
 $\text{trgl } AED = \text{trgl } CLB$


PROPOSITION 2

With the same things supposed if some point is taken on the sec-
ence of a circle and through it parallels to the tangents are dr-
diameters then the quadrilateral produced on one of the tangen-
diameters will be equal to the triangle produced on the same tangen-

For let there be a section of a cone or circumference of a circle



AEC and BED be tangents, and AD and BC diameters, and let some point G



be taken on the section and GKL and GMF be drawn parallel to the tangents
I say that

$$\text{trgl } AIM = \text{quadr } CLGI$$

For triangle GKM has been shown equal to quadrilateral AL (1 42, 43), let the common quadrilateral IA be added or subtracted, and

$$\text{trgl } AIM = \text{quadr } CG^1$$

PROPOSITION 3

With the same things supposed, if two points are taken on the section or circumference of a circle and through them parallels to the tangents are drawn as far as the diameters, the quadrilaterals produced by the straight lines drawn and standing on the diameters as bases are equal to each other

For let there be the section and tangents and diameters as said before and let two points at random F and G be taken on the section, and through F let the straight lines $FHAL$ and $NFIM$ be drawn parallel to the tangents and through G the straight lines GYO and HPR

I say that

$$\text{quadr } LG = \text{quadr } MH$$

¹Eutocius commenting gives the proof for another and important case. It must be remarked that if the point G is taken between A and B so that the parallels are for instance

and

$$\text{quadr } LN = \text{quadr } RN$$

For since it has already been shown that

$$\text{trgl } RPA = \text{quadr } CG \text{ (III 2),}$$

and

$$\text{trgl } AMI = \text{quadr } CF \text{ (III 2),}$$


and

$$\text{trgl } RPA = \text{trgl } AMI + \text{quadr } PM,$$

therefore also

$$\text{quadr } CG = \text{quadr } CF + \text{quadr } PM,$$

MIGI and *LGA* one must draw *IA* to the section at *N* for instance and through *N* draw



infer-
the
the

AN parallel to *BD* for by what was said in the forty ninth and fiftieth theorems of the first book (I 49 50) and in the notes to them

$$\text{trgl } ANN = \text{quadr } AC$$

But triangle *ANN* is similar to triangle *AMC* because *MG* is parallel to *AN* but it is also equal to it because *AC* is a tangent and *CV* is parallel to it and *MY* is a diameter and

$$GA = AN$$

Since then

$$\text{trgl } ANN = \text{quadr } AC = \text{trgl } AMG$$


with the common quadrilateral *AC* subtracted as remainders

$$\text{trgl } ANM = \text{quadr } CC$$

It will be noticed that, just as in the second note to I 50 the quadrilateral *CG* is to be considered as the difference between the triangles *CIF* and *ICF*

and so

$$\text{quadr } CG = \text{quadr } CH + \text{quadr } RF$$



be taken

I say

For the common quadrilateral CH be subtracted therefore as remainders

$$\text{quadr } LG = \text{quadr } HM$$

and therefore as wholes

$$\text{quadr } LN = \text{quadr } RN$$

PROPOSITION 4

If two straight lines touching opposite sections meet each other, and diameters are drawn through the points of contact meeting the tangents then the triangles at the tangents will be equal

Let there be the opposite sections A and B and let the tangents to them AC and BC meet at C and let D be the center of the sections and let AB and CD be joined and CD produced to E and let DA and BD also be joined and produced to F and G

I say that

$$\text{trgl } AGD = \text{trgl } BDF,$$


and

$$\text{trgl } ACF = \text{trgl } BCC$$

For let HL be drawn through H tangent to the section, therefore it is parallel to AG (1 44 note) And since

$$AD = DH \text{ (1 30),}$$

$\text{trgl } AGD = \text{trgl } DHL$ (Eucl vi 19)



But

$$\text{trgl } DHL = \text{trgl } BDF \text{ (III 1),}$$

therefore also

$$\text{trgl } AGD = \text{trgl } BDF$$


And so also

$$\text{trgl } ACF = \text{trgl } BCG$$

PROPOSITION 5

If two straight lines touching opposite sections meet, and some point is taken on either of the sections, and from it two straight lines are drawn, the one the tangent, the other parallel to the line joining the points of intersection of the triangle produced by them on the diameter drawn through the point of meeting of the tangent, the angle cut off on the tangent and the diameter drawn through the point of meeting of the tangent differs from the triangle cut off at the point of meeting of the tangent and the diameter drawn through the point of meeting of the tangent.

Let there be the opposite sections A and B whose center is C , and let the tangents ED and DF meet at D , and let EF and CD be joined and let




produced, and let FC and EC be joined and produced and let some point G be taken on the section, and through it let $HGA L$ be drawn parallel to EF , and GM parallel to DF .

I say that triangle GHM differs from triangle KHD by triangle KLF .

For since CD has been shown to be a diameter of the opposite sections (II

39, 38), and EF to be an ordinate to it (II 38, First Def 15), and GH has been




drawn parallel to EF , and MG parallel to DF , therefore triangle GHM differs from triangle CLH by triangle CDF (I 45, or I 44, according to the case) And so triangle GHM differs from triangle KHD by triangle KLF

And it is evident that

trgl $ALF = \text{quadr } MGKD$


PROPOSITION 6

ame things supposed if some point is taken on one of the opposite sec-
from it parallels to the tangents are drawn meeting the tangents and the
then the quadrilateral produced by them on one of the tangents and on
diameters will be equal to the triangle produced on the same tangent and
iameter

re be opposite sections of which AEC and BED are diameters and let $3G$ touch the section AB meeting each other at H and let some point K be on the section and from it let KML and KNX be drawn parallel

$$\text{quadr } KF = \text{trgl } AIN$$

v since AB and CD are opposite sections, and AF , meeting BD , touches



ction AB and AL has been drawn parallel to AF therefore


$\text{trgl } AIN = \text{quadr } AF \text{ (III 2)}^1$

PROPOSITION 7

With the same things supposed, if points are taken on each of the sections, and from them parallels to the tangents are drawn meeting the tangents and the diameters, then the quadrilaterals produced by the straight lines drawn and standing on the diameters as bases will be equal to each other

For let the aforesaid things be supposed and let points K and L be taken on both sections and through them let $MAPI$ and $NTLQ$ be drawn parallel to AF and $NIOK$ and $IWULZ$ parallel to BG

I say that what was said in the enunciation will be so



For since

$$\text{trgl } AOI = \text{quadr } RO \text{ (III 2),}$$


¹ Another and important case where the point K falls between C and D is given by Eutocius in his commentary to this proposition. It is as follows and let CPR be drawn tangent to the section then it is evident that it is parallel to AF and VL (I 44 note) since it has been shown in the second theorem (III 2) in the figure of the hyperbola

$$\text{trgl } PNC = \text{quadr } LP \text{ (III 2 note)}$$

let the common quadrilateral MP be added therefore

$$\text{trgl } MAN = \text{quadr } MLRC$$

Let there be added the common triangle CRE which is equal to triangle AEF by I 44 (and I 30) therefore



$$\text{whole trgl } MEL = \text{trgl } MAN + \text{trgl } AEF$$

With common triangle AMV subtracted as remainders

$$\text{trgl } LEF = \text{quadr } AIFN$$

Let the common quadrilateral $FFNI$ be added therefore in whole

$$\text{trgl } AIN = \text{quadr } AIFI$$

And likewise also

$$\text{trgl } BOL = \text{quadr } A\backslash GO$$

let the quadrilateral EO be added to both therefore
 whole $\text{trgl } AEF = \text{quadr } KE$

But also

$$\text{trgl } BGE = \text{quadr } LE \text{ (III 5, note),}$$

and

$$\text{trgl } AEF = \text{trgl } BGE \text{ (III 1),}$$

therefore

$$\text{quadr } LE = \text{quadr } IKRE$$

Let the common quadrilateral NE be added, therefore as wholes

$$\text{whole quadr } TH = \text{quadr } IL,$$

and also

$$\text{quadr } KU = \text{quadr } RL$$

PROPOSITION 8


With the same things supposed, instead of K and L let there be taken the points C and D at which the diameters hit the sections and through them let the parallels to the tangents be drawn

I say that

$$\text{quadr } DG = \text{quadr } FC$$

and

$$\text{quadr } \lambda I = \text{quadr } OT$$



For since it was shown

$$\text{trgl } AGH = \text{trgl } HBF \text{ (III 1)}$$

and the straight line from A to B is parallel to the straight line from G to F ¹
 therefore

$$AE \cdot EG = BE \cdot EF$$

and *conuertendo*

$$EA \cdot AG = EB \cdot BF$$

And also

$$CA \cdot AE = DB \cdot BE,$$

¹For the point H falls within the angle AEB (II 25) and the straight line drawn from H to the midpoint of AB that is S is a diameter (II 29) and must therefore pass through E (I 51 end). An analogous series of propositions is found for the opposite sections II. 32 38 39

Then, since
 therefore

$$\text{trgl } GHA = \text{trgl } FHB$$

$$\text{trgl } GFB = \text{trgl } GFA$$

Their bases are the same therefore their heights are equal (Eucl. VI 1)

for each is double the other therefore *ex aequali*

$$CA \ AG \ DB \ BF$$

And the triangles are similar because of the parallels, therefore

$$\text{trgl } CTA \ \text{trgl } AHG \ \text{trgl } ABD \ \text{trgl } HBF \text{ (Euel vi 19)}$$

And alternately, but

$$\text{trgl } AHG = \text{trgl } HBF \text{ (iii 1),}$$

therefore

$$\text{trgl } CTA = \text{trgl } ABD$$

As parts of these it was shown

$$\text{trgl } AHG = \text{trgl } HBF,$$

therefore also as remainders

$$\text{quadr } DH = \text{quadr } CH$$

And so also

$$\text{quadr } DG = \text{quadr } CF$$

And since *CO* is parallel to *AF*

$$\text{trgl } COE = \text{trgl } AEF$$

And likewise also

$$\text{trgl } DEI = \text{trgl } BEG$$

But

$$\text{trgl } BEG = \text{trgl } AEF \text{ (iii 1),}$$

therefore also

$$\text{trgl } COE = \text{trgl } DEI$$

And also

$$\text{quadr } DG = \text{quadr } CF \text{ (above)}$$

Therefore as wholes,

$$\text{quadr } VI = \text{quadr } OT$$


PROPOSITION 9

With the same things supposed if one of the points is between the diameters as *K*, and the other is the same with one of the points *C* and *D* for instance *C*, and the parallels are drawn I say that

$$\text{trgl } CEO = \text{quadr } AE$$

and

$$\text{quadr } LO = \text{quadr } LM$$



And this is evident For since it was shown

$$\text{trgl } CEO = \text{trgl } AEF,$$

and

$$\text{trgl } AEF = \text{quadr } \triangle A E \text{ (III 5, note),}$$

therefore also

$$\text{trgl } CEO = \text{quadr } \triangle A E$$

And so also

$$\text{trgl } CRM = \text{quadr } \triangle A O,$$

and

$$\text{quadr } \triangle AC = \text{quadr } \triangle LO$$


PROPOSITION 10

With the same things supposed, let A and L be taken not as points at which the diameters hit the sections

Then it is to be shown that

$$\text{quadr } LTRY = \text{quadr } QYAI$$

For since the straight lines AF and BG touch, and AE and BE are diameters



through the points of contact and LT and KI are parallel to the tangents,

$$\text{trgl } TUE = \text{trgl } UQL + \text{trgl } EFA \text{ (I 44)}$$

And likewise also

$$\text{trgl } \triangle EI = \text{trgl } XRK + \text{trgl } BEG$$

But

$$\text{trgl } EFA = \text{trgl } BEG \text{ (III 1)}$$

therefore

$$\text{trgl } TUE - \text{trgl } UQL = \text{trgl } XEI - \text{trgl } XRK$$

Therefore

$$\text{trgl } TUE + \text{trgl } XRK = \text{trgl } \triangle EI + \text{trgl } UQL$$

Let the common area $\triangle YEULY$ be added therefore

$$\text{quadr } LTRY = \text{quadr } QYAI$$

PROPOSITION 11

With the same things supposed if some point is taken on either of the sections and from it parallels are drawn one parallel to the tangent and the other parallel to the straight line joining the points of contact then the triangle produced by them on the diameter drawn through the point of meeting of the tangents differs from the tri-

for each is double the other, therefore *ex aequali*

$$CA \cdot AG = DB \cdot BF$$

And the triangles are similar because of the parallels, therefore

$$\text{trgl } CTA \sim \text{trgl } AIG \sim \text{trgl } XBD \sim \text{trgl } HBF \text{ (Eucl vi 19)}$$

And alternately, but

$$\text{trgl } AIG = \text{trgl } HBF \text{ (iii 1),}$$

therefore

$$\text{trgl } CTA = \text{trgl } XBD$$

As parts of these it was shown

$$\text{trgl } AIG = \text{trgl } HBF$$

therefore also as remainders

$$\text{quadr } DH = \text{quadr } CH$$

And so also

$$\text{quadr } DG = \text{quadr } CF$$

And since *CO* is parallel to *AF*

$$\text{trgl } COE = \text{trgl } AEF$$

And likewise also

$$\text{trgl } DEI = \text{trgl } BEG$$

But

$$\text{trgl } BEG = \text{trgl } AEF \text{ (iii 1),}$$

therefore also

$$\text{trgl } COE = \text{trgl } DEI$$

And also

$$\text{quadr } DG = \text{quadr } CF \text{ (above)}$$

Therefore, as wholes

$$\text{quadr } XI = \text{quadr } OT$$


PROPOSITION 9

With the same things supposed, if one of the points is between the diameters, as *K*, and the other is the same with one of the points *C* and *D* for instance *C*, and the parallels are drawn, I say that

$$\text{trgl } CEO = \text{quadr } KE$$

and



$$\text{quadr } LO = \text{quadr } LM$$



And this is evident For since it was shown

$$\text{trgl } CEO = \text{trgl } AEF,$$

For let there be the same things as before and let the points B and K be taken at random on section AB and through them let $LBMN$ and $KVOUP$



be drawn parallel to AD and $B\lambda R$ and LKS parallel to AE

I say that

$$\text{quadr } BP = \text{quadr } KR$$

For since it has been shown

trgl AOP=quadr KOES (III 11 end)

and

trgl	$AMN = \text{quadr } BMER$ (in 11 end).

therefore, as remainders either

$$\text{quadr } KR - \text{quadr } BO = \text{quadr } MP$$

OR

$$\text{quadr } AR + \text{quadr } BO = \text{quadr } MP$$

And with the common quadrilateral BO added or subtracted

$$\text{quadr } BP = \text{quadr } XS$$

PROPOSITION 13

If in conjugate opposite sections straight lines tangent to the adjacent sections meet and diameters are drawn through the points of contact then the triangles whose common vertex is the center of the opposite sections will be equal

Let there be conjugate opposite sections on which there are the points A, B, C and D and let BE and AE meeting at E touch the sections A and B and let H be the center and let AH and BH be joined and produced to D and C



angle cut off on the tangent and the diameter drawn through the point of contact by the triangle cut off at the point of meeting of the tangents

Let there be the opposite sections AB and CD , and let the tangents AE and DE meet at E and let the center be H , and let AD and EHG be joined, and let some point B be taken at random on the section AB , and through it let BFL be drawn parallel to AG , and BM parallel to AE

I say that triangle BFM differs from triangle AKL by triangle AEF

For it is evident that AD is bisected by EH (II 39 and II 29), and that EH is a diameter conjugate to the diameter drawn through H parallel to AD (II 38), and so AG is an ordinate to EG (First Def 1 6)

Since then GE is a diameter and AE touches and AG is an ordinate and with point B taken on the section, BF has been dropped to EG parallel to AG ,



and BM parallel to AE , therefore it is clear that triangle BMF differs from triangle LHF by triangle HAE (I 45, I 43)¹ And so also triangle BMF differs from triangle AKL by triangle AEF

And it has been proved at the same time that

$$\text{quadr } BAE M = \text{trgl } LHA$$

PROPOSITION 12

With the same things being so if on one of the sections two points are taken and parallels are drawn from each of them, likewise the quadrilaterals produced by them will be equal

¹That is in the first case

$$\text{trgl } BMF = \text{trgl } LHF + \text{trgl } HAE \text{ (I 45)}$$

In the second case only the more general statement differs but is true (I 43). It will be noticed these are different cases of I 43 and I 45 from those given in the text itself

B and through it let λRS be drawn parallel to AG and λTO parallel to BE

I say that triangle OHT differs from triangle λST by triangle $HB F$

For let AU be drawn from A parallel to BF Since then because of the same things as before, LHM is a diameter of the section AL , and DHB is a second diameter and conjugate to it (II 20), and AG is a tangent at A , and AU has been dropped parallel to LM , therefore

AU UG comp HU UA ,
transverse side of figure on LM upright (I 40)

But

AU UG λT TS ,

and

HU UA HT TO HB BF ,

and

transverse side of figure on LM upright
upright side of figure on BD transverse (I 60)

Therefore

λT TS comp HB BF , upright side of figure on BD transverse

or

λT TS comp HT TO upright side of figure on BD transverse

And by things shown in the forty first theorem of the first book (I 41), triangle THO differs from triangle λTS by triangle BFH

And so also by triangle AGH (III 13)

PROPOSITION 15

*If straight lines touching one of the conjugate opposite sections meet and diameters are drawn through the points of contact and some point is taken on any one of the conjugate sections and from it parallels to the tangents are drawn as far as the diameters then the triangle produced by them at the section is greater than the triangle produced at the center by the triangle having the tangent as base and the center of the opposite sections as vertex*¹

Let there be conjugate opposite sections AB GS T and λ whose center is H and let ADE and BDC touch the section AB and let the diameters $AHFW$ and BHT be drawn through the points of contact A and B and let some point S be taken on the section GS and through it let SFL be drawn parallel to BC and SU parallel to AE

I say that

$\text{trgl } SLU = \text{trgl } HLF + \text{trgl } HCB$

For let λHG be drawn through H parallel to BC and λIG through G parallel to AE and SO parallel to BT then it is evident that λG is a diameter conjugate to BT (II 20) and that SO being parallel to BT has been dropped ordinatewise to HGO (First Def I 6) and that $SLHO$ is a parallelogram

Since then BC touches and BH is through the point of contact and AE is another tangent let it be contrived that

DB BE λN $2BC$,

therefore λN is the so-called upright side of the figure on BT (I 50) Let MN be bisected at P therefore

¹This proposition comes as a climax to a long series and shows that the conjugate opposite sections taken as a unit have the same property as the other conic sections. The conjugate opposite sections seem to be a sort of fifth section.

Again since

$$HB \ BC \text{ comp } HB \ MP, MP \ BC,$$

but

$$HB \ MP \ TB \ MN \ R \ \lambda G \text{ (above } \alpha \text{ and } \gamma),$$

and

$$MP \ BC \ DB \ BE \text{ (above, } \beta)$$

therefore

$$HB \ BC \text{ comp } DB \ BE \ R \ \gamma G$$

And since BC is parallel to SL , and triangle HCB is similar to triangle HLF ,
and

$$HB \ BC \ HL \ LF,$$

therefore

$$HL \ LF \text{ comp } R \ \lambda G \ DB \ BE$$

or

$$HL \ IF \text{ comp } R \ \lambda G \ HG \ HI$$

Since then GS is an hyperbola having λG as a diameter and R as an upright side and from some point S SO has been dropped ordinatewise and figure HIG has been described on radius HG and figure HLF has been described on the ordinate SO or its equal HL , and on HO the straight line between the center and the ordinate or on SL its equal the figure SLU has been described similar to the figure HIG described on the radius, and there are the compounded ratios as already given therefore

$$\text{trgl } SLU = \text{trgl } HLF + \text{trgl } HCB \text{ (I 41)}$$

PROPOSITION 16

If two straight lines touching a section of a cone or circumference of a circle meet and from some point of those on the section a straight line is drawn parallel to one tangent and cutting the section and the other tangent then as the squares on the tangents are to each other so the area contained by the straight lines between the section and the tangent will be to the square cut off at the point of contact

Let there be the section of a cone or circumference of a circle AB and let the straight lines AC and CB meeting at C touch it and let some point D be taken on the section AB and through it let EDF be drawn parallel to CB

I say that

$$\text{sq } BC \text{ sq } AC \text{ rect } FE, ED \text{ sq } EA$$

For let the diameters AGH and ABL be drawn through A and B , and DMN through D parallel to AL , it is at once evident that

$$DA = AF \text{ (I 46 47)}$$

and

$$\text{trgl } AEG = \text{quadr } LD \text{ (III 2),}$$

and

$$\text{trgl } BLC = \text{trgl } ACH \text{ (III 1)}$$

Since then

$$DA = AF$$

and DE is added





$$\text{rect } FE \ ED + \text{sq } DK = \text{sq } KE$$

And since triangle ELK is similar to triangle DNK

$$\text{sq } EK \text{ sq } KD \text{ trgl } ELK \text{ trgl } DNK$$

And alternately

whole sq EK whole trgl ELK
 part subtracted sq DK part subtracted trgl DNK



Therefore also

remainder rect $FE ED$ remainder quadr DL sq EA trgl FLA

But

sq EK trgl ELK sq CB trgl BLC

therefore also

rect $FE ED$ quadr LD sq CB trgl LCB

But

quadr $LD = \text{trgl } AFG$

and

trgl $BLC = \text{trgl } ACH$,

therefore also

rect $FE ED$ trgl AEG sq CB trgl ACH

Alternately

rect $FE ED$ sq CB trgl AFG trgl ACH

But

trgl AEC trgl ACH sq EA sq AC

therefore also

rect $FE ED$ sq CB sq FA sq AC

And alternately

PROPOSITION 17

If two straight lines touching a section of a cone or circumference of a circle meet, and two points are taken at random on the section, and from them in the section are drawn parallel to the tangents straight lines cutting each other and the line of the section, then as the squares on the tangents are to each other, so will the rectangles contained by the straight lines taken similarly

Let there be the section of a cone or circumference of a circle AB , and tangents to AB AC and CB meeting at C , and let points D and E be taken at random on the section and through them at $EFIK$ and $DFGH$ be drawn parallel to AC and CB


I say that

$$\text{sq } CA : \text{sq } CB :: \text{rect } KF : \text{rect } FE :: \text{rect } HF : \text{rect } FD$$

For let the diameters $ALMN$ and $BOXP$ be drawn through A and B , and let the tangents and parallels be produced to the diameters and let DX and EM be drawn from D and E parallel to the tangents, then it is evident that

$$KI = IE, HG = GD \quad (\text{I 46 47})$$

Since then KE has been cut equally at I and unequally at F ,
 $\text{rect } KF, FE + \text{sq } FI = \text{sq } EI$ (Eucl II 5)



And since the triangles are similar because of the parallels
 $\text{whole sq } EI : \text{whole trgl } IME$

part subtracted sq IF part subtracted trgl FIL

Therefore also

remainder rect KF, FE remainder quadr FM whole
sq EI whole trgl IME

But

sq EI trgl IME sq CA trgl CAN

Therefore

rect KF, FE quadr FM sq CA trgl CAN

But

trgl $CAN = \text{trgl } CPB$ (in 1),

and

quadr $FM = \text{quadr } FA$ (in 3)

therefore

rect KF, FE quadr FA sq CA trgl CPB

Then likewise it could be shown that

rect HF, FD quadr FY sq CB trgl CPB

Since then

rect KF, FE quadr FY sq CA trgl CPB

and inversely

quadr FA rect HF, FD trgl CPB sq CB


therefore ex aequali

sq CA sq CB rect KF, FE rect HF, FD

PROPOSITION 18

If two straight lines touching opposite sections meet and some point is taken on either one of the sections and from it some straight line is drawn parallel to one of the tangents cutting the section and the other tangent then as the squares on the tangents are to each other so will the rectangle contained by the straight lines between the section and the tangent be to the square on the straight line cut off at the point of contact

Let there be the opposite sections AB and MN and tangents ACL and BCH and through the points of contact the diameters AM and BN and let some point D be taken at random on the section MN and through it let EDF be drawn parallel to BH



I say that

sq BC sq CA rect FE, FD sq AF

For let DX be drawn through D parallel to AL Since then AB is an hyper

bola and BN its diameter and BH a tangent and DF parallel to BH , therefore $FO=OD$ (I 48)

And ED is added therefore

$$\text{rect } FE, ED + \text{sq } DO = \text{sq } EO \text{ (Eucl II 6)}$$

And since EL is parallel to DY , triangle EOL is similar to triangle $D\Lambda O$
Therefore

$$\begin{array}{ll} \text{whole sq } EO & \text{whole trgl } EOL \\ \text{part subtracted sq } DO & \text{part subtracted trgl } D\Lambda O, \end{array}$$

therefore also

$$\text{remainder rect } DE, EF \quad \text{remainder quadr } DL \quad \text{sq } EO \quad \text{trgl } EOL$$

But

$$\text{sq } OE \quad \text{trgl } EOL \quad \text{sq } BC \quad \text{trgl } BCL$$

therefore also

$$\text{rect } FE, ED \quad \text{quadr } DL \quad \text{sq } BC \quad \text{trgl } BCL$$

And

$$\text{quadr } DL = \text{trgl } AEG \text{ (III 6 note),}$$

and

$$\text{trgl } BCL = \text{trgl } ACH \text{ (III 1),}$$

therefore

$$\text{rect } FE, ED \quad \text{trgl } AEG \quad \text{sq } BC \quad \text{trgl } ACH$$

But also

$$\text{trgl } AEG \quad \text{sq } EA \quad \text{trgl } ACH \quad \text{sq } AC,$$

therefore *ex aequali*


$$\text{sq } BC \quad \text{sq } AC \quad \text{rect } FE, ED \quad \text{sq } EA^1$$

¹Eutocius gives an alternative proof of Apollonius demonstrating another and important case For let there be the opposite sections A and B and tangents to them AC and CB meeting at C and let D be taken on section B and through it let EDF be drawn parallel to AC I say that

$$\text{sq } AC \quad \text{sq } CB \quad \text{rect } EF, FD \quad \text{sq } FB$$

For let AHG be drawn as a diameter through A and through B and G GK and BL parallel to EF Since then BH touches the hyperbola at B and BL has been drawn ordinate-wise

$$AL \quad LG \quad AH \quad HG \text{ (I 36)}$$



But
and
therefore also
And alternately
and
But it was shown
therefore also

$$\begin{array}{llll} AL & LG & CB & BK \\ AH & HG & AC & KG \\ CB & BA & AC & AG \\ AC & CB & KG & KB \\ \text{sq } AC & \text{sq } CB & \text{sq } GK & \text{sq } AB \\ \text{sq } GA & \text{sq } KB & \text{rect } EF, FD & \text{sq } FB \\ \text{sq } AC & \text{sq } CB & \text{rect } EF, FD & \text{sq } FB \end{array}$$


PROPOSITION 19

If two straight lines touching opposite sections meet and parallels to the tangents are drawn cutting each other and the section then, as the squares on the tangents are to each other so will the rectangle contained by the straight lines between the section and the point of meeting of the straight lines be to the rectangle contained by the straight lines taken similarly

Let there be opposite sections whose diameters are AC and BD and center in E , and let the tangents AF and FD meet at F , and let $GHIKL$ and $MNAL$ be drawn from any points parallel to AF and FD

I say that

$$\text{sq } AF \text{ sq } FD \text{ rect } GL LI \text{ rect } ML, LA$$



Let IP and XR be drawn through λ and I parallel to AF and FD
 BH touches the hyperbola at B and BL has been drawn ordinatewise,
 $AL LG \quad AH HG$ (I, 36)

And since

$$\text{sq } AF \text{ trgl } AFS \quad \text{sq } HL \text{ trgl } HLO \quad \text{sq } HI \text{ trgl } HIP$$

therefore

$$\text{remainder rect } GL LI \text{ remainder quadr } IPOL \quad \text{sq } AF \text{ trgl } AFS$$

But

$$\text{trgl } AFS = \text{trgl } DTF \text{ (III 4)}$$

and

$$\text{quadr } IPOL = \text{quadr } KRYL \text{ (III 7)}$$

therefore also

$$\text{sq } AF \text{ trgl } DTF \quad \text{rect } GL LI \text{ quadr } KRYL$$

But

$$\text{trgl } DTF \text{ sq } FD \quad \text{quadr } KRYL \text{ rect } ML LY \text{ (likewise),}$$

and therefore ex aequali

$$\text{sq } AF \text{ sq } FD \quad \text{rect } GL LI \text{ rect } ML LY$$

PROPOSITION 20


If two straight lines touching opposite sections meet and through the point of meeting some straight line is drawn parallel to the straight line joining the points of contact and meeting each of the sections and some other straight line is drawn parallel to the same at right line and cutting the sections and the tangents then as the rectangle contained by the straight lines drawn from the point of meeting to cut the sections is to the square on the tangent so is the rectangle contained by the

straight lines between the sections and the tangent to the square on the straight line cut off at the point of contact

Let there be the opposite sections AB and CD whose center is E and tangents AF and CF and let AC be joined, and let EF and AE be joined and produced and let BFH be drawn through F parallel to AC , and let the point K be taken at random and through it let $KLSMNX$ be drawn parallel to AC

I say that

$$\text{rect } BF, FD \text{ sq } FA \quad \text{rect } KL, LX \text{ sq } AL$$



For let KP and BR be drawn from K and B parallel to AF Since then
 $\text{sq } BF \text{ trgl } BFR \quad \text{sq } KS \text{ trgl } KSP \quad \text{sq } LS \text{ trgl } LSF$
 and

$$\begin{aligned} & \text{sq } AS \text{ trgl } ASP \\ & \text{remainder rect } KL \text{ LX (Eucl II 5)} \\ & \text{remainder quadr } KLFP \text{ (Eucl V 19)} \end{aligned}$$

and
 $\text{sq } BF = \text{rect } BF, FD$ (II 39, 38)

and
 $\text{trgl } BRF = \text{trgl } AFH$ (III 11 and special case)

and
 $\text{quadr } KLFP = \text{trgl } ALN$ (III 5)

therefore

$$\text{rect } BF, FD \text{ trgl } AFH \quad \text{rect } KL \text{ LX trgl } ALN$$

And
 $\text{trgl } AFH \text{ sq } AF \quad \text{trgl } ALN \text{ sq } AL,$

then
 $\text{rect } BF \text{ FD sq } FA \quad \text{rect } KL, LX \text{ sq } AL$

PROPOSITION 21


With the same things supposed if two points are taken on the section and through them straight lines are drawn the one parallel to the tangent, the other parallel to the straight line joining the points of contact and cutting each other and the sections then as the rectangle contained by the straight lines drawn from the point

of meeting to cut the sections as to the square on the tangent, so will the rectangle contained by the straight lines between the sections and the point of meeting be to the rectangle contained by the straight lines between the section and the point of meeting

For let there be the same things as before, and let points G and K be taken and through them let $NAGOPR$ and KST be drawn parallel to AF , and GLV and KOW IZQ parallel to AC

I say that

$$\text{rect } BF, FD \text{ sq } FA \quad \text{rect } KO, OQ \text{ rect } NO, OG$$



For since

$$\text{sq } AF \text{ trgl } AFH \quad \text{sq } AL \text{ trgl } ALM \quad \text{sq } XO \text{ trgl } XOZ$$

and

$$\text{sq } XO \text{ trgl } XOZ \quad \text{sq } YG \text{ trgl } YGM,$$

therefore

$$\text{whole sq } XO \quad \text{whole trgl } XOZ$$

$$\text{part subtracted sq } XG \quad \text{part subtracted trgl } YGM,$$

therefore also

$$\text{remainder rect } NO, OG \quad \text{remainder quadr } GOZM \quad \text{sq } AF \text{ trgl } AFH$$

But

$$\text{trgl } AFH = \text{trgl } BUF \text{ (in 11 end special case),}$$

and

$$\text{quadr } GOZM = \text{quadr } KORT \text{ (in 12),}$$

therefore

$$\text{sq } AF \text{ trgl } BFU \quad \text{rect } NO, OG \quad \text{quadr } KORT$$

But it was shown (in the course of III 20) $\text{trgl } BUF \text{ sq } BF \text{ or rect } BF, FD$

$$\text{(II 33 38) quadr } KORT \text{ rect } KO, OQ$$

therefore ex aequali

$$\text{sq } AF \text{ rect } BF, FD \quad \text{rect } NO, OG \text{ rect } KO, OQ$$

And inversely

$$\text{rect } BF, FD \text{ sq } FA \quad \text{rect } KO, OQ \text{ rect } NO, OG$$

PROPOSITION 22


If two parallel straight lines touch opposite sections and any straight lines are drawn cutting each other and the sections one parallel to the tangent the other parallel to the straight line joining the points of contact, then as the transverse side

of the figure on the straight line joining the points of contact is to the upright, so the rectangle contained by the straight lines between the sections and the point of meeting will be to the rectangle contained by the straight lines between the section and the point of meeting

Let there be the opposite sections A and B , and let AC and BD be parallel and tangent to them, and let AB be joined. Then let EXG be drawn across parallel to AB and $KELM$ parallel to AC

I say that

AB upright side of the figure rect $GE EY$ rect KE, EM
Let λN and GF be drawn through G and λ parallel to AC



For since AC and BD are parallels tangent to the sections AB is a diameter (II 31), and KL , XN and GF are ordinates to it (I 32), then (I 21)

AB upright side
rect BL, LA sq LK rect $BN NA$ sq $N\lambda$ or sq LK

Therefore

whole rect $BL LA$ whole sq LK
part subtracted rect $BN NA$ part subtracted sq LE ,

or

rect BL, LA sq LK rect $FA AN$ sq LE ,

for

$NA = BF$ (I 21),

therefore also

remainder rect $FL LN$ remainder rect KE, EM AB upright

But

rect $FL LN =$ rect $GE EY$,


therefore

AB the transverse side of figure upright
rect $GE EY$ rect KE, EM

PROPOSITION 23

If in conjugate opposite sections two straight lines touching contrary sections meet in any one section at random and any straight lines are drawn parallel to the tangents and cutting each other and the other opposite sections then as the squares on the tangents are to each other so the rectangle contained by the straight lines between the sections and the point of meeting will be to the rectangle contained by the straight lines similarly taken

Let there be the conjugate opposite sections AB CD , EF and GH , and their center K , and let $AWCL$ and $EYDL$, tangents to the sections AB and EF



meet at L and let AK and EK be joined and produced to B and F , and let $GMNXO$ be drawn from G parallel to AL and $HPR\lambda S$ from H parallel to EL

I say that

$$\text{sq } EL \text{ sq } LA \text{ rect } HX, \lambda S \text{ rect } GY \lambda D$$

For let ST be drawn through S parallel to AL , and OU from O parallel to EL . Since then BE is a diameter of the conjugate opposite sections AB CD , EF and GH and EL touches the section and HS has been drawn parallel to it,

$$HP = PS \text{ (II 20, First Def 1 5),}$$

and for the same reasons

$$GM = MO$$

And since

$$\text{sq } EL \text{ trgl } EWL \text{ sq } PS \text{ trgl } PTS \text{ sq } PY \text{ trgl } PNX,$$

also

$$\text{remainder rect } HX, \lambda S \text{ remainder quadr } TN \lambda S \text{ sq } EL \text{ trgl } WLE$$

But

$$\text{trgl } EWL = \text{trgl } ALY \text{ (III 4),}$$

and

$$\text{quadr } TN \lambda S = \text{quadr } XRUO \text{ (III 15) }^1$$

therefore

$$\text{sq } EL \text{ trgl } ALY \text{ rect } HX \lambda S \text{ quadr } \lambda RUO$$

But

$$\text{trgl } ALY \text{ sq } AL \text{ quadr } \lambda RUO \text{ rect } GY \lambda O \text{ (same way),}$$

therefore *ex aequali*

$$\text{sq } EL \text{ sq } AL \text{ rect } HX, \lambda S \text{ rect } GY, XO$$

PROPOSITION 24

If in conjugate opposite sections two straight lines are drawn from the center through to the sections and one of them is taken as the transverse diameter and the other as the upright diameter and any straight lines are drawn parallel to the two diameters and meeting each other and the sections and the point of meeting of the straight lines is the place between the four sections then the rectangle contained by

¹This is the case of III 15 where the tangents are one to each of the opposite sections

Compare with the two cases of III 12 and III 18

For $\text{trgl } TSI - \text{trgl } APR = \text{trgl } ANA \text{ (III 15)}$

and $\text{trgl } MOU - \text{trgl } MVA = \text{trgl } ANA \text{ (III 15)}$

the segments of the parallel to the transverse diameter together with the rectangle to which the rectangle contained by the segment of the parallel to the upright diameter has the ratio which the square on the upright diameter has to the square on the transverse will be equal to twice the square on the half of the transverse


Let there be the conjugate opposite sections A, B, C and D whose center is E , and from E let the transverse diameter ABC and the upright diameter DEB be drawn through and let $FGHIKL$ and $MNXOPR$ be drawn parallel to AC and DB and meeting each other at X , and first let Y be within the angle SEW or the angle UET

I say that the rectangle FX, XL together with the rectangle to which the rectangle RX, XM has the ratio which the square on DB has to the square on AC , is equal to twice the square on AE

For let the asymptotes of the sections SET and UEW be drawn and through A , $SGAW$ tangent to the section

Since then

$$\text{rect } SA, AW = \text{sq } DE \text{ (I 60, II 1),}$$



therefore

$$\text{rect } SA, AW = \text{sq } EA + \text{sq } DE + \text{sq } EA$$

And

$$\text{rect } SA, AW = \text{sq } AE + \text{comp } SA, AE \cdot WA + AE$$

But

$$SA, AE = NX, XH$$

and

$$WA, AE = PY, YK$$

therefore

$$\text{sq } DE + \text{sq } AE + \text{comp } NX, XH, PY, YK$$

But

$$\text{rect } PX, YN = \text{rect } KX, XH + \text{comp } NX, YH, PX, YK,$$

therefore

$$\text{sq } DE + \text{sq } AE = \text{rect } PX, YN = \text{rect } KY, XH$$

Therefore also

$$\text{sq } DL = \text{sq } AE + \text{sq } DE + \text{rect } PY, XN = \text{sq } AE + \text{rect } KY, XH$$

And

$$\text{sq } DE = \text{rect } PM, MN \text{ (II 11)} = \text{rect } RN, NM \text{ (II 16)}$$

and

$$\text{sq } AE = \text{rect } KF, FH \text{ (II 11)} = \text{rect } LH, HF \text{ (II 16),}$$

therefore

$$\text{sq } DE \text{ sq } AE \text{ rect } PY, \lambda N + \text{rect } RN, NM \\ \text{rect } K \lambda, \lambda H + \text{rect } LH, HF$$

And

$$\text{rect } P \lambda \lambda N + \text{rect } RN, NM = \text{rect } RY, XM,^1$$

therefore

$$\text{sq } DE \text{ sq } AE \text{ rect } R \lambda, XM \text{ rect } K \lambda, \lambda H + \text{rect } KF, FH$$

Then it must be shown that

$$\text{rect } F \lambda \lambda L + \text{rect } K \lambda \lambda H + \text{rect } KF, FH = 2 \text{ sq } AE$$

Let the common square AE that is rectangle KF, FH , be subtracted, therefore it remains to be shown that

$$\text{rect } F \lambda \lambda L + \text{rect } K \lambda, \lambda H = \text{sq } AE$$

And this is so, for

$$\text{rect } F \lambda \lambda L + \text{rect } K \lambda \lambda H = \text{rect } LH, HF, \dagger$$

$$\text{rect } F \lambda, \lambda L + \text{rect } K \lambda \lambda H = \text{rect } KF, FH \text{ (II 16)}$$

$$= \text{sq } AE \text{ (II 11)}$$

Then let the straight lines FL and MR meet on one of the asymptotes at H . Then


$$\text{rect } FH, HL = \text{sq } AE,$$

and

$$\text{rect } MH, HR = \text{sq } DE \text{ (II 11 16),}$$

therefore

$$\text{sq } DE \text{ sq } AE \text{ rect } MH, ER \text{ rect } FH, HL$$



And so we want twice rectangle FH, HL to equal twice the square on AE . And it does

¹For

$$RP = \lambda M \text{ (II 8)}$$

and

$$RO = OM \text{ (II 3)}$$

therefore

$$PO = O \lambda$$

But

$$\text{rect } PY, \lambda \lambda + \text{sq } O \lambda = \text{sq } ON \text{ (Eucl II 5)}$$

and for the same reasons

$$\text{rect } RN, \lambda M + \text{sq } O \lambda = \text{sq } OM$$

and

$$\text{rect } R \lambda, \lambda M + \text{sq } O \lambda = \text{sq } OM$$

Hence

$$\text{rect } R \lambda, \lambda M + \text{sq } O \lambda = \text{rect } R \lambda, \lambda M + \text{sq } O \lambda$$

and adding equals to equals,

$$\text{rect } R \lambda, \lambda M + \text{sq } O \lambda + \text{rect } P \lambda, \lambda \lambda + \text{sq } O \lambda = \text{rect } R \lambda, \lambda M + \text{sq } O \lambda + \text{sq } ON$$

Subtracting the common squares

$$\text{rect } R \lambda, \lambda M + \text{rect } P \lambda, \lambda \lambda = \text{rect } RY, XM$$

[†]By the same manner of proof as in the note above but using also Euclid II 6 because of the different position of the point λ .

And let the point Y be within the angle SEA or angle WET Then likewise by the composition of ratios

$$\text{sq } DE \text{ sq } AE \text{ rect } P\lambda, \lambda N \text{ rect } K\lambda \lambda H$$

And


$$\text{sq } DE = \text{rect } PM, RN = \text{rect } RN NM,$$

and

$$\text{sq } AL = \text{rect } FH HL,$$

therefore

$$\text{rect } RN, NM \text{ rect } FH, HL$$



part subtracted rect $P\lambda \lambda N$ part subtracted rect $K\lambda \lambda H$

Therefore also

$$\text{rect } RN NM \text{ rect } FH HL$$

remainder rect $RX XM$ remainder (sq $AE - \text{rect } K\lambda \lambda H$)

Therefore it must be shown that

$$\text{rect } FX \lambda L + (\text{sq } AE - \text{rect } KX \lambda H) = 2 \text{ sq } AE$$

Let the common square on AE that is rectangle $FH HL$ be subtracted therefore it remains to be shown that

$$\text{rect } KX \lambda H + (\text{sq } AE - \text{rect } KX \lambda H) = \text{sq } AE$$

And this is so for

$$\text{rect } K\lambda \lambda H + \text{sq } AE - \text{rect } KX \lambda H = \text{sq } AE$$

PROPOSITION 25

With the same things supposed let the point of meeting of the parallels to AC and BD be within one of the sections D and B as set out below at X

I say that the rectangle contained by the segments of the parallel to the transverse that is rectangle $O\lambda \lambda N$, will be greater than the rectangle to which the rectangle contained by the segments of the parallel to the upright diameter that is rectangle $RX XM$ has the ratio which the square on the upright diameter has to the square on the transverse by twice the square on the half of the transverse

For for the same reasons

$$\text{sq } DE \text{ sq } AE \text{ rect } PX XH \text{ rect } SY \lambda L$$

and

$$\text{sq } DE = \text{rect } PM MH$$

and


$$\text{sq } AE = \text{rect } LO OS \text{ (II 11)}$$

therefore also

$$\text{sq } DE = \text{sq } AE - \text{rect } PM, MH - \text{rect } LO, OS$$

And since

$$\text{whole rect } PX, \backslash H = \text{whole rect } L\lambda, \lambda S$$



$$\text{part subtracted rect } PM, MH = \text{part subtracted rect } LO, OS \\ \text{or rect } ST, TL \text{ (II 22),}$$

therefore also

$$\text{remainder rect } RX, SM = \text{remainder rect } T\lambda, \lambda K \\ (\text{first note to III 24 II 8}) \quad \text{sq } DE = \text{sq } AE$$

Therefore it must be shown that

$$\text{rect } O\backslash, \backslash N = \text{rect } T\lambda, \lambda K + 2 \text{ sq } AE$$

Let the common rectangle $TY, \backslash A$ be subtracted, therefore it must be shown that

$$\text{rect } OT, TN \text{ (first note to III 24)} = 2 \text{ sq } AE$$

And it is (II 23)

PROPOSITION 26

And if the point of meeting of the parallels at Y is within one of the sections A and C as set out below then the rectangle contained by the segments of the parallel to the transverse that is rectangle $LX, \backslash F$ will be less than the rectangle to which the rectangle contained by the segments of the other parallel that is rectangle $R\backslash, \backslash G$ has the ratio which the square on the upright diameter has to the square on the transverse by twice the square on half of the transverse

For, since for the same reasons as before

$$\text{sq } DE = \text{sq } AE - \text{rect } W\lambda, \backslash S - \text{rect } K\backslash, \backslash H$$

therefore also

$$\text{whole rect } R\backslash, \backslash G^1 = \text{whole rect } K\backslash, \backslash H + \text{sq } AE \\ \text{sq upright diameter} = \text{sq transverse}$$

Therefore it must be shown that

$$\text{rect } I\backslash, \backslash F + 2 \text{ sq } AL = \text{rect } K\backslash, \backslash H + \text{sq } AE$$

¹For by II 11

$$\text{rect } WG, GS = \text{sq } DE$$


and

$$IW = GS \text{ (II 16)}$$

Therefore by the first note to III 24 and II 16

$$\text{rect } W\backslash, \backslash S + \text{sq } DE = \text{rect } W\backslash, \backslash S + \text{rect } WG, GS = \text{rect } R\backslash, \backslash G$$

Let the common square on AE be subtracted, therefore it remains to be shown that



$$\text{rect } LX \cdot XF + \text{sq } AE = \text{rect } KY, \backslash H$$

or

$$\text{rect } LX, \backslash F + \text{rect } LH \cdot HF = \text{rect } K\backslash \backslash H \quad (\text{II } 16 \text{ } 11)$$

And it is for

$$\text{rect } LH, HF + \text{rect } LX, \backslash F = \text{rect } K\backslash, \backslash H^1$$

PROPOSITION 27

If the conjugate diameters of an ellipse or circumference of a circle are drawn and one of them is called the upright diameter and the other the transverse and two straight lines meeting each other and the line of the section, are drawn parallel to them then the squares on the straight lines cut off on the straight line drawn parallel to the transverse between the point of meeting of the straight lines and the line of the section plus the figures described on the straight lines cut off on the straight line drawn parallel to the upright diameter between the point of meeting of the straight lines and the line of the section figures similar and similarly situated to the figure on the upright diameter will be equal to the square on the transverse diameter

For let there be the ellipse or circumference of a circle $ABCD$, whose center is E and let two of its conjugate diameters be drawn the upright AFC and the transverse BED and let $NGFH$ and $KFLM$ be drawn parallel to AC and BD

I say that the squares on NF and FI plus the figures described on AF and FM , similar and similarly situated to the figure on AC will be equal to the square on BD

Let NX be drawn from N parallel to AF therefore it has been dropped ordinatewise to BD And let BP be the upright side Now since

$$\frac{BP}{AC} = \frac{AC}{BD} \quad (\text{I } 15),$$

therefore also

$$\frac{BP}{BD} = \frac{\text{sq } AC}{\text{sq } BD}$$

And

$$\text{sq } BD = \text{figure on } AC,$$


therefore

$$\frac{BP}{BD} = \frac{\text{sq } AC}{\text{figure on } AC}$$

¹This is another case of the first note to III 21

And

sq AC figure on AC
sq NY figure on NY similar to the figure on AC (Eucl vi 22),



therefore also

$BP \cdot BD$ sq NY figure on NY similar to the figure on AC

And also

$BP \cdot BD$ sq NY rect $BY \cdot YD$ (i 21)

therefore

figure on NA or FL similar to the figure on AC = rect $BY \cdot YD$

Then likewise we could show that

figure on AL similar to the figure on AC = rect $BL \cdot LD$

And since the straight line AH has been cut equally at G and unequally at F ,

sq HF + sq FN = 2[sq HG + sq GF] = 2[sq NG + sq GF] (Eucl vi 9)

Then for the same reasons also

sq MF + sq FK = 2[sq KL + sq LF]

and the figure on MF and FK similar to the figure on AC are double the similar figures on AL and LF

And

figure on AI + figure on FL = rect $BY \cdot YD$ + rect $BL \cdot LD$ (above),

and

sq NG + sq GF = sq AE + sq EL ,

therefore

sq NI + sq FH + figures on AI and FH similar to the figure on AC =
2[rect $BY \cdot YD$ + rect $BL \cdot LD$ + sq AE + sq EL]

And since the straight line BD has been cut equally at E and unequally at Y ,

rect $BY \cdot YD$ + sq YE = sq BE (Eucl ii 5)

And likewise also

rect $BL \cdot LD$ + sq LE = sq BE ,

and so

rect $BY \cdot YD$ + rect $BL \cdot LD$ + sq AE + sq LE = 2 sq BE

Therefore the squares on NI and FH together with figures on AI and FH similar to the figure on AC are double the square on BE . But also


sq BD = 2 sq BE ,

therefore the squares on NI and FH plus the figures on AI and FH similar to the figure on AC are equal to the square on BD

PROPOSITION 28

If in conjugate opposite sections conjugate diameters are drawn, and one of them is called the upright, and the other the transverse and two straight lines are drawn parallel to them and meeting each other and the sections then the squares on the straight lines cut off on the straight line drawn parallel to the upright between the point of meeting of the straight lines and the sections have to the squares on the straight lines cut off on the straight line drawn parallel to the transverse between the point of meeting of the straight lines and the sections the ratio which the square on the upright diameter has to the square on the transverse diameter

Let there be the conjugate opposite sections A, B, C , and D and let AEC be the upright diameter and BED the transverse and let $FGHK$ and $LGMN$



be drawn parallel to them and cutting each other and the sections

I say that

$$\text{sq } LG + \text{sq } GN : \text{sq } FG + \text{sq } GK :: \text{sq } AC : \text{sq } BD$$

For let LY and FO be drawn ordinatewise from F and L , therefore they are parallel to AC and BD . And from B let the upright side for BD BP be drawn, then it is evident that

$$PB \cdot BD : \text{sq } AC : \text{sq } BD \text{ (I 15)} :: \text{sq } AE : \text{sq } EB$$

$$\text{sq } FO : \text{rect } BO \cdot OD \text{ (I 21)} :: \text{rect } CX, XA : \text{sq } LY \text{ (I 60 21)}$$

Therefore as one of the antecedents is to one of the consequents so are all of the antecedents to all of the consequents (Eucl v 12) therefore

$$\text{sq } AC : \text{sq } BD :: \text{rect } CX, XA + \text{sq } AE + \text{sq } OF : \text{rect } DO, OB + \text{sq } BE, \\ + \text{sq } LX$$

or

$$\text{sq } AC : \text{sq } BD :: \text{rect } CX, XA + \text{sq } AE + \text{sq } EH : \\ \text{rect } DO, OB + \text{sq } BE + \text{sq } ME$$

But

$$\text{rect } CX, XA + \text{sq } AE = \text{sq } YE,$$

and

$$\text{rect } DO, OB + \text{sq } BE = \text{sq } OE \text{ (Eucl II 6),}$$

therefore

$$\text{sq } AC : \text{sq } BD :: \text{sq } YE + \text{sq } EH : \text{sq } OE + \text{sq } EM \\ \text{sq } LM + \text{sq } MG : \text{sq } FH + \text{sq } HG$$

And as has been shown

$$\text{sq } NG + \text{sq } GL = 2[\text{sq } LM + \text{sq } MG]$$

and

$$\text{sq } FG + \text{sq } GK = 2[\text{sq } FH + \text{sq } HG] \text{ (Eucl II 9),}$$

therefore also

$$\text{sq } AC - \text{sq } BD = \text{sq } NG + \text{sq } GL - \text{sq } FG + \text{sq } GK$$

PROPOSITION 29

With the same things supposed, if the parallel to the upright diameter cuts the asymptotes then the squares on the straight lines cut off on the straight line drawn parallel to the upright between the point of meeting of the straight lines and the asymptotes plus the half of the square on the upright diameter has to the squares on the straight lines cut off on the straight line drawn parallel to the transverse between the point of meeting of the straight lines and the sections the ratio which the square on the upright diameter has to the square on the transverse


For let there be the same things as before and let NL cut the asymptotes at V and O

It is to be shown that

$$\text{sq } \lambda G + \text{sq } GO + \text{half sq } AC = \text{sq } FG + \text{sq } GK - \text{sq } AC - \text{sq } BD$$

or

$$\text{sq } \lambda G + \text{sq } GO + 2 \text{ sq } AL = \text{sq } FG + \text{sq } GK - \text{sq } AC - \text{sq } BD$$



For since

$$LV = ON \text{ (II 16),}$$

$$\text{sq } LG + \text{sq } GN + 2 \text{ rect } NV, LV = \text{sq } \lambda G + \text{sq } GO,¹$$

therefore

$$\text{sq } \lambda G + \text{sq } GO + 2 \text{ sq } AE = \text{sq } LG + \text{sq } GN$$

And

$$\text{sq } IG + \text{sq } GN - \text{sq } FG + \text{sq } GK = \text{sq } AC - \text{sq } BD \text{ (III 28),}$$

therefore also

$$\text{sq } \lambda G + \text{sq } GO + 2 \text{ sq } AE - \text{sq } FG + \text{sq } GK = \text{sq } AC - \text{sq } BD$$

¹For

$$OM = MV$$

Therefore as in a lemma of Pappus since

$$2 \text{ rect } NV - \lambda L + 2 \text{ sq } MV = 2 \text{ sq } ML \text{ (Eucl II 5)}$$

adding the common square on CV

$$2 \text{ rect } NV - \lambda L + 2 \text{ sq } MV + 2 \text{ sq } GV = 2 \text{ sq } ML + 2 \text{ sq } GV$$

And

$$2 \text{ sq } \lambda L + 2 \text{ sq } CV = \text{sq } NG + \text{sq } LG$$


and

$$2 \text{ sq } MX + 2 \text{ sq } GV = \text{sq } OG + \text{sq } GX \text{ (Eucl II 9)}$$

Therefore as above

PROPOSITION 30

If two straight lines touching an hyperbola meet and through the points of contact a straight line is produced and through the point of meeting a straight line is drawn parallel to some one of the asymptotes and cutting both the section and the straight line joining the points of contact then the straight line between the point of meeting and the straight line joining the points of contact will be bisected by the section¹



Let there be the hyperbola ABC , and let AD and DC be tangents and EF and FG asymptotes and let AC be joined and through D parallel to FE let DKL be drawn

I say that

$$DK = KL$$

For let $FDBM$ be joined and produced both ways and let FH be made equal to BF and through the points B and K let BE and KN be drawn

parallel to AC therefore they have been dropped ordinatewise (Π 30 5 7) And since triangle BEF is similar to triangle DNK , therefore

$$\text{sq } DN : \text{sq } NK :: \text{sq } BF : \text{sq } BE \quad (\alpha)$$

And

$$\text{sq } BF : \text{sq } BE :: HB : \text{upright } (\Pi 1),$$

therefore also

$$\text{sq } DN : \text{sq } NK :: HB : \text{upright}$$

But

$$HB : \text{upright} :: \text{rect } HN, NB : \text{sq } NK \quad (\text{I } 21),$$

therefore also

$$\text{sq } DN : \text{sq } NK :: \text{rect } MN, NB : \text{sq } NK \quad (\beta)$$

Therefore

$$\text{rect } HN, NB = \text{sq } DN$$

¹The propositions from 30 to 34 inclusive are one special case and propositions 35 and 36 are another special case of proposition 37. The first group takes the line drawn through the intersection of the tangents as parallel to an asymptote. The second group takes one of the tangents as an asymptote. Proposition 34 lying between is special in both these ways.

In proposition 37 we have the line CF divided by the section at D and F and at E by the straight line joining the points of contact in such a way that

$$CF : CD :: FE : ED$$

This is the same form of the harmonic proportion as we found in Π 34 and DF is the harmonic mean between CF and FE .

If we argue by analogy from this proportion treating infinity as a definite magnitude and two such infinities as would occur here as equal and subject to the general laws of magnitudes we can immediately deduce the special cases of propositions 30 to 36. Thus in the case of the first group CF and FE both become infinite therefore CD is equal to ED .

And also

$$\text{rect } MF \cdot FD = \text{sq } FB \text{ (I 37),}$$

because AD touches and AM has been dropped ordinatewise, and so also

$$\text{rect } HN, NB + \text{sq } FB = \text{rect } MF, FD + \text{sq } DN$$

But

$$\text{rect } HN, NB + \text{sq } FB = \text{sq } FN \text{ (Eucl II 6),}$$

and therefore

$$\text{rect } MF \cdot FD + \text{sq } DN = \text{sq } FN$$

Therefore DM has been bisected at N with DF added (Eucl II 6) And KN and LM are parallel, therefore

$$DK = KL$$


PROPOSITION 31

If two straight lines touching opposite sections meet and a straight line is produced through the points of contact, and through the point of meeting a straight line is drawn parallel to the asymptote and cutting both the section and the straight line joining the points of contact then the straight line between the point of meeting and the straight line joining the points of contact will be bisected by the section

Let there be the opposite sections A and B and tangents AC and CB , and let AB be joined and produced, and let FE be an asymptote and through C let CGH be drawn parallel to FE

I say that

$$CG = GH$$



Let CE be joined and produced to D and through E and G let $NEKM$ and $G\lambda$ be drawn parallel to AB , and through G and K let KF and GL be drawn parallel to CD

Since triangle KFF is similar to triangle MLG

$$\text{sq } KE : \text{sq } KF :: \text{sq } ML : \text{sq } LG$$

And it has been shown

$$\text{sq } KEE : \text{sq } KF :: \text{rect } NL : LK :: \text{sq } LG \text{ (}\alpha \text{ and } \beta \text{ of III 30),}$$

therefore

$$\text{rect } NL : LK = \text{sq } MI$$

Let the square on KF be added to each therefore

$$\text{rect } NL : LK + \text{sq } KE = \text{sq } LF = \text{sq } G\lambda = \text{sq } ML + \text{sq } KE$$

And

$$\text{sq } G\lambda : \text{sq } ML :: \text{sq } KE : \text{sq } KC :: \text{sq } LG + \text{sq } KF \text{ (Eucl VI 4, v 12),}$$

therefore

$$\text{sq } KC = \text{sq } IG + \text{sq } KF$$

And

$$\text{sq } LG = \text{sq } YE$$

and

$$\text{sq } KF = \text{sq on half of second diameter (II 1),} \\ = \text{rect } CE, ED \text{ (I 38),}$$

therefore

$$\text{sq } \lambda C = \text{sq } \lambda E + \text{rect } CE ED$$

Therefore the straight line CD has been cut equally at X and unequally at E (Eucl II 5)

And DH is parallel to GY , therefore

$$CG = GH$$

PROPOSITION 32

If two straight lines touching an hyperbola meet, and a straight line is produced through the points of contact and a straight line is drawn through the point of meeting of the tangents parallel to the straight line joining the points of contact and a straight line is drawn through the midpoint of the straight line joining the points of contact parallel to one of the asymptotes, then the straight line cut off between this midpoint and the parallel will be bisected by the section

Let there be the hyperbola ABC , whose center is D , and asymptote DE , and let AF and FC touch and let CA and FD be joined and produced to G and H ,

then it is evident that

$$AH = HC$$

Then let FK be drawn through F parallel to AC and HLK through H parallel to DE

I say that

$$KL = HL$$

Let LM and BE be drawn through B and L parallel to AC then as has been already shown (III 30 α , β and conclusion)

$$\text{sq } DB + \text{sq } BE = \text{sq } HM + \text{sq } ML = \text{rect } BM, MG + \text{sq } ML$$

therefore

$$\text{rect } GM, MB = \text{sq } MH$$

And also

$$\text{rect } HD, DF = \text{sq } DB$$

because AF touches and AH has been dropped ordinatewise (I 37) therefore


$$\text{rect } GM, MB + \text{sq } DB = \text{rect } HD, DF + \text{sq } MH = \text{sq } DM \text{ (Eucl II 6)}$$

Therefore FH has been bisected at M with DF added And KF and LM are parallel, therefore

$$KL = LH$$

PROPOSITION 33

If two straight lines touching opposite sections meet, and one straight line is produced through the points of contact and another straight line is drawn through the point of meeting of the tangents parallel to the straight line joining the points of



PROPOSITION 34

If one point is taken on one of the asymptotes of an hyperbola, and a straight line from it touches the section and through the point of contact a parallel to the asymptote is drawn then the straight line drawn from the point taken parallel to the other asymptote will be bisected by the section

Let there be the hyperbola AB and the asymptotes CD and DE , and let a point C be taken at random on CD , and through it let CBE be drawn touching the section and through B let FBG be drawn parallel to CD and through C let CAG be drawn parallel to DE

I say that

$$CA = AG$$

For let AH be drawn through A parallel to CD , and BK through B parallel to DE Since then


$$CB = BE \text{ (II 3),}$$

therefore also

$$CK = KD$$

and

$$DF = FE$$



And since

$$\text{rect } KB, BF = \text{rect } CA, AH \text{ (II 12),}$$

and

$$BF = DK = CK,$$

and

$$AH = DC,$$

therefore

$$\text{rect } DC, CA = \text{rect } GC, CK$$

Therefore

$$DC \cdot CK = GC \cdot CA$$

And

$$CD = 2CK,$$

therefore also

$$GC = 2CA$$

Therefore

$$CA = AG$$

PROPOSITION 35

With the same things being so if from the point taken some straight line is drawn cutting the section at two points then as the whole straight line is to the straight line cut off outside so will the segments of the straight line cut off inside be to each other

For let there be the hyperbola AB and the asymptotes CD and DE and CBE touching and HB parallel and through C let some straight line $CALFG$ be drawn across cutting the section at A and F

I say that

$$FC \cdot CA = FL \cdot AL$$

For let CNX KAM $OPBR$ and FU be drawn through C A B and F

parallel to DE , and APS and $TFRMX$ through A and F parallel to CD

Since then

$$AC = FG \text{ (ii 8),}$$

therefore also

$$KA = TG \text{ (Eucl vi 4)}$$

But

$$KA = DS,$$

therefore also

$$TG = DS$$

And so also

$$CK = DU$$

And since

$$CK = DU,$$

also

$$DK = CU,$$

therefore

$$DK \quad CK \quad CU \quad CK$$

And

$$CU \quad CK \quad FC \quad AC,$$

and

$$FC \quad AC \quad MK \quad KA,$$

and

$$MK \quad KA \quad \text{p||g } MD \quad \text{p||g } DA \text{ (Eucl vi 1),}$$

and

$$DK \quad CK \quad \text{p||g } HK \quad \text{p||g } KN,$$

therefore also

$$\text{p||g } MD \quad \text{p||g } DA \quad \text{p||g } HK \quad \text{p||g } KN$$

But

$$\text{p||g } DA = \text{p||g } DB \text{ (ii 12)} = \text{p||g } ON,$$

for

$$CB = BE \text{ (ii 3),}$$

and

$$DO = OC,$$

therefore

$$\text{p||g } MD \quad \text{p||g } ON \quad \text{p||g } HK \quad \text{p||g } KN$$

and

$$\text{remainder p||g } MH \quad \text{remainder p||g } BK \quad \text{whole p||g } MD \quad \text{whole p||g } ON$$

And since

$$\text{p||g } DA = \text{p||g } DB$$

let the common parallelogram DP be subtracted,

therefore

$$\text{p||g } AP = \text{p||g } PH$$

Let the common parallelogram AB be added therefore

$$\text{whole p||g } BK = \text{whole p||g } AH$$


Therefore

$$\text{p||g } MD \quad \text{p||g } DA \quad \text{p||g } MH \quad \text{p||g } AH$$

But

$$\text{p||g } MD \quad \text{p||g } DA \quad MK \quad KA \quad FC \quad AC,$$

and



pllg MH pllg AH MW WA FL LA ,

therefore also

FC AC FL LA

PROPOSITION 36


With the same things being so if the straight line drawn across from the point neither cuts the section at two points nor is parallel to the asymptote it will meet the opposite section and as the whole straight line is to the straight line between the section and the parallel through the point of contact so will the straight line between the opposite section and the asymptote be to the straight line between the asymptote and the other section

Let there be the opposite sections A and B whose center is C and asymptotes DE and FG and let some point G be taken on CG and from it let GBE be drawn tangent and GH neither parallel to CE nor cutting the section in two points (I 26)

It has been shown that GH produced meets CD and therefore also section A Let it meet it at A and let KBL be drawn through B parallel to CG

I say that

AK KH AG GH



For let HM and AN be drawn from the points A and H parallel to CG and BX GP and $RHSN$ from B G and H parallel to DE Since then

$AD = GH$ (I 16)
 AG GH DH HG

But

AG GH NS SH

and

DH GH CS SG

And therefore

NS SH CS SG

But

NS SH pllg NC pllg CH

and

CS SG pllg RC pllg RG

therefore also

pllg NC pllg CH pllg RC pllg RG

And as one is to one so are all to all therefore

And since $\text{pllg } NC \text{ pllg } CH \text{ whole pllg } NL \text{ whole pllg } CH + \text{pllg } RG$

$$EB = BG,$$

also

$$LB = BP$$

and

$$\text{pllg } L\lambda = \text{pllg } BG$$

And

$$\text{pllg } L\lambda = \text{pllg } CH \text{ (II 12),}$$

therefore also

$$\text{pllg } BG = \text{pllg } CH$$

Therefore

$$\text{pllg } NC \text{ pllg } CH \text{ whole pllg } NL \text{ whole pllg } BG + \text{pllg } RG$$

or

$$\text{pllg } NC \text{ pllg } CH \text{ pllg } NL \text{ pllg } R\chi$$

But

$$\text{pllg } R\lambda = \text{pllg } LH$$

since also

$$\text{pllg } CH = \text{pllg } BC \text{ (II 12),}$$

and

$$\text{pllg } MB = \text{pllg } \lambda H$$

Therefore

$$\text{pllg } NC \text{ pllg } CH \text{ pllg } NL \text{ pllg } LH$$

But

$$\text{pllg } NC \text{ pllg } CH \text{ } NS \text{ } SH \text{ } AG \text{ } GH,$$

and

$$\text{pllg } NL \text{ pllg } LH \text{ } NR \text{ } RH \text{ } AK \text{ } KH,$$

therefore also

$$AK \text{ } KH \text{ } AG \text{ } GH$$

PROPOSITION 37

If two straight lines touching a section of a cone or circumference of a circle or opposite sections meet and a straight line is joined to their points of contact and from the point of meeting of the tangents some straight line is drawn across cutting the line (of the section) at two points then as the whole straight line is to the straight line cut off outside so will the segments produced by the straight line joining the points of contact be to each other

Let there be the section of a cone AB and tangents AC and CB and let AB be joined and let $CDEF$ be drawn across

I say that

$$CF \text{ } CD \text{ } FE \text{ } FD$$

Let the diameters CH and AK be drawn through C and A and through F and D DP FR LFM and NDO parallel to AH and LC Since then LFM is parallel to λDO

$$FC \text{ } CD \text{ } LF \text{ } \lambda D \text{ } FM \text{ } DO \text{ } LM \text{ } \lambda O$$

and therefore

$$\text{sq } LM \text{ sq } \lambda O \text{ sq } FM \text{ sq } DO$$

But

$$\text{sq } LM \text{ sq } \lambda O \text{ } \text{trgl } LMC \text{ } \text{trgl } \lambda CO \text{ (Eucl vi 19),}$$

and

$$\text{sq } FM \text{ sq } DO \quad \text{trgl } FRM \quad \text{trgl } DPO,$$

therefore also



$$\text{trgl } LMC \quad \text{trgl } \lambda CO \quad \text{trgl } FRM \quad \text{trgl } DPO \\ \text{remainder quadr } LCRF \quad \text{remainder quadr } \lambda CP$$

But

$$\text{quadr } LCRF = \text{trgl } ALK \text{ (III 2, III 11),}$$

and

$$\text{quadr } YCPD = \text{trgl } ANY \text{ (III 2, III 11),}$$



therefore

$$\text{sq } LM \text{ sq } XO \quad \text{trgl } ALK \quad \text{trgl } ANX$$

But

$$\text{sq } LM \text{ sq } XO \quad \text{sq } FC \text{ sq } CD,$$

and

$$\text{trgl } ALK \quad \text{trgl } ANX \quad \text{sq } LA \text{ sq } AX \quad \text{sq } FE \text{ sq } ED,$$




therefore also

$$\text{sq } FC \text{ sq } CD \quad \text{sq } FE \text{ sq } ED$$

And therefore

$$FC \quad CD \quad FE \quad ED$$

For let $LFAM$ and $DHGAN$ be drawn through F and D parallel to AB , and through F and G FR and GP parallel to LC . Then likewise as before (III 37) it will be shown that
 $\text{sq } LM = \text{sq } \backslash H = \text{sq } LA = \text{sq } AX$
 And
 $\text{sq } LM = \text{sq } XH = \text{sq } LC = \text{sq } CX$
 $\text{sq } FO = \text{sq } OD$,
 and
 $\text{sq } LA = \text{sq } AX = \text{sq } FE = \text{sq } ED$,
 therefore
 $\text{sq } FO = \text{sq } OD = \text{sq } FE = \text{sq } ED$,
 and
 $FO = OD = FE = ED$



PROPOSITION 39

If two straight lines touching opposite sections meet and a straight line is produced through the points of contact and a straight line drawn from the point of meeting of the tangents cuts both of the sections and the straight line joining the points of contact then as the whole straight line drawn across is to the straight line

and

$$\text{trgl } FRM = \text{trgl } A \backslash N + \text{trgl } \backslash MD \text{ (nr 11),}$$

therefore

$$\text{trgl } DHS \text{ trgl } \backslash MD \text{ trgl } ASK + \text{trgl } DHS \text{ trgl } A \backslash N + \text{trgl } \backslash MD,$$

and

$$\text{remainder trgl } ASK \text{ remainder trgl } AN \backslash \text{ trgl } DHS \text{ trgl } \backslash MD$$

But

$$\text{trgl } ASK \text{ trgl } AN \backslash \text{ sq } KA \text{ sq } AN \text{ sq } EG \text{ sq } FG,*$$

and

$$\text{trgl } DHS \text{ trgl } \backslash MD \text{ sq } HD \text{ sq } DM \text{ sq } ED \text{ sq } DF$$

Therefore also

$$EG \text{ } FG \text{ } ED \text{ } DF$$

PROPOSITION 40

With the same things being so if a straight line is drawn through the point of meeting of the tangents parallel to the straight line joining the points of contact and if a straight line drawn from the midpoint of the straight line joining the points of contact cuts both of the sections and the straight line parallel to the straight line joining the points of contact then as the whole straight line drawn across is to the straight line cut off outside between the parallel and the section, so will the straight line's segments produced by the sections and the straight line joining the points of contact be to each other

Let there be the opposite sections A and B whose center is C , and tangents AD and DB , and let AB and CDE be joined, therefore

$$AE = EB \text{ (nr 39)}$$

And from D let FDG be drawn parallel to AB , and from E , LE at random

I say that

$$HL \text{ } LA \text{ } HE \text{ } EA$$

From H and A let NMH and KOP be drawn parallel to AB and HR and AS parallel to AD , and let $\backslash ACT$ be drawn through

Since then $\backslash AU$ and MAP have been drawn across the parallels $\backslash M$ and AP

$$XA \text{ } AU \text{ } MA \text{ } AP$$

But

$$XA \text{ } AU \text{ } HE \text{ } EA,$$

and

$$HE \text{ } EA \text{ } HN \text{ } KO$$

because of the similarity of the triangles HEN and KEO , therefore

$$HN \text{ } KO \text{ } MA \text{ } AP,$$

therefore also

$$\text{sq } HN \text{ sq } KO \text{ sq } MA \text{ sq } AP$$

For

$$EG \text{ } TG \text{ } KA \text{ } TA$$

and

$$TG \text{ } TF \text{ } TA \text{ } TN$$

and

$$TG - TF \text{ } TG \text{ } TA - TN \text{ } TA$$

therefore ex aequali


$$EG \text{ } FG \text{ } KA \text{ } AV$$

But

$$\text{sq } HN \text{ sq } KO \quad \text{trgl } HRN \quad \text{trgl } KSO,$$

and

$$\text{sq } MA \text{ sq } AP \quad \text{trgl } \lambda MA \quad \text{trgl } AUP,$$



therefore also

$$\text{trgl } HRN \quad \text{trgl } KSO \quad \text{trgl } \lambda MA \quad \text{trgl } AUP$$

And

$$\text{trgl } HNR = \text{trgl } \lambda MA + \text{trgl } MND \text{ (iii 11),}$$

and

$$\text{trgl } KSO = \text{trgl } AUP + \text{trgl } DOP \text{ (iii 11),}$$

therefore also

$$\text{trgl } \lambda MA + \text{trgl } MND \quad \text{trgl } AUP + \text{trgl } DOP \quad \text{trgl } \lambda MA \quad \text{trgl } AUP,$$

therefore also

$$\text{remainder trgl } NMD \quad \text{remainder trgl } DOP \quad \text{whole whole}$$

But

$$\text{trgl } \lambda MA \quad \text{trgl } AUP \quad \text{sq } \lambda A \quad \text{sq } AU,$$

and

$$\text{trgl } NMD \quad \text{trgl } DOP \quad \text{sq } MN \quad \text{sq } PO,$$

therefore also

$$\text{sq } MN \quad \text{sq } PO \quad \text{sq } \lambda A \quad \text{sq } AU$$

But

$$\text{sq } MN \quad \text{sq } PO \quad \text{sq } ND \quad \text{sq } OD,$$

and

$$\text{sq } \lambda A \quad \text{sq } AU \quad \text{sq } HF \quad \text{sq } EK,$$

and

$$\text{sq } ND \quad \text{sq } DO \quad \text{sq } HL \quad \text{sq } LK,$$

therefore also

$$\text{sq } HE \quad \text{sq } FK \quad \text{sq } HL \quad \text{sq } LK$$

Therefore

$$HE \quad EK \quad HL \quad LK$$

PROPOSITION 11


If three straight lines touching a parabola meet each other they will be cut in the same ratio

Let there be the parabola ABC , and tangents ADE FFC and DBF

I say that

$$CF \quad FE \quad ED \quad DA \quad FB \quad BD$$

For let AC be joined and bisected at G



Then it is evident that the straight line from E to G is a diameter of the section (II 29)

If then it goes through B DF is parallel to AC , (I 5) and will be bisected by EG and therefore

$$AD = DE \quad (\text{I } 35)$$

and

$$CF = FE \quad (\text{I } 35),$$

and what was sought is apparent

Let it not go through B but through H , and let KHL be drawn through H parallel to AC therefore it will touch the section at H (I 32), and because of things already said (I 35),

$$AK = KE$$

and

$$LC = LE$$

Let $MNBX$ be drawn through B parallel to EG and AO and CP through A and C parallel to DF Since then MB is parallel to EH , MB is a diameter (I 40, I 51 end), and DF touches at B , therefore AO and CP have been dropped ordinatewise (II 5, First Def I 4) And since MB is a diameter, and CM a tangent and CP an ordinate

$$MB = BP \quad (\text{I } 35),$$

and so also

$$MF = FC$$

And since

$$MF = FC$$

and

$$EL = LC,$$

$$MC \quad CF \quad EC \quad CL,$$

and alternately

$$MC \quad EC \quad CF \quad CL$$

But

$$MC \quad EC \quad XC \quad CG$$

therefore also

$$CF \quad CL \quad XC \quad CG$$

And

$$CL \quad EC \quad CG \quad CA$$

therefore *ex aequali*

$$CA \quad XC \quad EC \quad CF$$

and *convertendo*

$$EC \quad FE \quad CA \quad AX,$$

separando

$$CF \quad FE \quad XC \quad AX$$

Again since MB is a diameter and AN a tangent and AO an ordinate,

$$NB=BO \text{ (r 35),}$$

and

$$ND=DA$$

And also

$$EK=KA,$$

therefore

$$AE \quad KA \quad NA \quad DA,$$

alternately

$$AE \quad NA \quad KA \quad DA$$

But

$$AE \quad NA \quad GA \quad AX,$$

therefore also

$$KA \quad DA \quad GA \quad AX$$

And also

$$AE \quad KA \quad CA \quad GA,$$

therefore, *ex aequali*,

$$AE \quad DA \quad CA \quad AX,$$

separando

$$ED \quad DA \quad XC \quad AX$$

And it was also shown

$$XC \quad AX \quad CF \quad FE,$$

therefore

$$CF \quad FE \quad FD \quad DA$$

Again since

$$XC \quad AX \quad CP \quad AO,$$

and

$$CP=2 BF,$$

and

$$CM=2 MF,$$

and

$$AO=2 BD,$$

and

$$AN=2 ND,$$

therefore

$$XC \quad AX \quad FB \quad BD \quad CF \quad FE \quad ED \quad DA$$

PROPOSITION 42

If in an hyperbola or ellipse or circumference of a circle or opposite sections straight lines are drawn from the vertex of the diameter parallel to an ordinate and some other straight line at random is drawn tangent it will cut off from them straight lines containing a rectangle equal to the fourth part of the figure to the same diameter


For let there be some one of the aforesaid sections whose diameter is AB and from A and B let AC and DB be drawn parallel to an ordinate, and let some other straight line CED be tangent at E

I say that

$$\text{rect } AC \quad BD = \text{fourth part of figure to } AB$$

For let its center be F and through it let FG be drawn parallel to AC and BD . Since then AC and BD are parallel and FG is also parallel, therefore it is the diameter conjugate to AB (First Def, 16), and so

sq FG = fourth part of figure to AB (Sec Def 13)



If then FG goes through E in the case of the ellipse and circle
 $AC = FG = BD$ (11 7)

and it is immediately evident that

rect $AC \cdot BD =$ q FG or fourth part of figure to AB

Then let it not go through it and let DC and BA produced meet at K and let EI be drawn through E parallel to AC and FM parallel to AB . Since then

rect $KF, FL =$ sq AF (1 37)

$KF \cdot AF = AF \cdot FL$

and

$$KA \cdot AL = KF \cdot AF \text{ or } FB \text{ (Eucl v 19),}$$

inversely

$$FB \cdot KF = AL \cdot KA,$$

componendo or separando

$$BK \cdot KF = LK \cdot KA$$

Therefore also

$$DB \cdot FH = EL \cdot CA$$

Therefore

$$\begin{aligned} \text{rect } DB, CA &= \text{rect } FH, EL, \\ &= \text{rect } HF, FM \end{aligned}$$

But

$$\begin{aligned} \text{rect } HF, FM &= \text{sq } FG \text{ (I 38),} \\ &= \text{fourth figure to } AB \text{ (Sec Def I 11),} \end{aligned}$$

therefore also

$$\text{rect } DB \cdot CA = \text{fourth figure to } AB$$

PROPOSITION 43

If a straight line touch an hyperbola it will cut off from the asymptotes beginning with the center of the section straight lines containing a rectangle equal to the rectangle contained by the straight lines cut off by the tangent at the section's vertex at its axis

Let there be the hyperbola AB , and asymptotes CD and DE , and axis BD , and let FBG be drawn through B tangent, and some other tangent at random CAH

I say that

$$\text{rect } FD \cdot DG = \text{rect } CD \cdot DH$$

For let AK and BL be drawn from A and B parallel to DG and AM and BN parallel to CD . Since then CAH touches

$$CA = AH \text{ (II 3),}$$

and so

$$CH = 2AH$$

and

$$CD = 2AM$$

and

$$DH = 2AK$$

Therefore

$$\text{rect } CD \cdot DH = 4 \text{ rect } KA \cdot AM$$

Then likewise it could be shown

$$\text{rect } FD \cdot DG = 4 \text{ rect } LB \cdot BN$$


But

$$\text{rect } KA \cdot AM = \text{rect } LB \cdot BN \text{ (II 12)}$$

Therefore also


$$\text{rect } CD \cdot DH = \text{rect } FD \cdot DG$$

Then likewise it could be shown even if DB were some other diameter and not the axis



PROPOSITION 44

If two straight lines touching an hyperbola or opposite sections meet the asymptotes then the straight lines drawn to the sections will be parallel to the straight line joining the points of contact



For let there be either the hyperbola or the opposite sections AB and asymptotes CD and DE and tangents $CAHF$ and $EBHG$, and let AB FG , and CE be joined

I say that they are parallel

For since

$$\text{rect } CD \text{ } DF = \text{rect } GD, DE \text{ (III 43),}$$

therefore

$$\frac{CD}{HF} = \frac{DE}{FC} = \frac{GD}{HG} = \frac{DF}{GE},$$

therefore CE is parallel to FG And therefore

And

$$\frac{FC}{HG} = \frac{AF}{GB} = \frac{GE}{HF} = \frac{GB}{FA},$$

for each is double (II 3) therefore *ex aequali*




Therefore FG is parallel to AB

PROPOSITION 45

If in an hyperbola or ellipse or circumference of a circle or opposite sections straight lines are drawn from the vertex of the axis at right angles and a rectangle equal to the fourth part of the figure is applied to the axis on each side and exceed

ing by a square figure in the case of the hyperbola and opposite sections but deficient in the case of the ellipse and some straight line is drawn tangent to the section and meeting the perpendicular straight lines then the straight lines drawn from the points of meeting to the points produced by the application make right angles at the aforesaid points¹

Let there be one of the sections mentioned whose axis is AB , and AC and BD



¹ The points of application are in modern terminology the foci of the conics. The circle is seen here as an ellipse whose two foci or focal points coincide with the center. This theory is of course a special application of Euclid vi. 28 and 29 two theorems on which depends one whole side of Greek geometry.

Apollonius never speaks of the focus of the parabola but it can be found by analogy with the ellipse.

Thus in the ellipse above

$$\text{rect } AF \cdot FB = \text{fourth rect } AB \cdot R$$

where R is the parameter. Or

$$\text{rect } 1F (AB - AF) = \text{fourth rect } AB \cdot R$$

or

$$1F \cdot \text{fourth } R = AB (AB - AF)$$

Then if we consider the ellipse as its axis AB gets as large as we please we can think of it as approaching as near as we please to a parabola with parameter R . The ratio $1B : (1B - 1F)$ approaches as near as we please to equality and hence also the ratio $AF : \text{fourth } R$. At the limit we can think of the ellipse as the parabola its axis AB as infinite and AB as equal to $AB - 1F$. Then $1F$ will be equal to a fourth R . Thus the focus of a parabola will be defined as the point on its axis at a distance from the vertex equal to one quarter of the parameter. Then many of the properties of the foci of the ellipse can be proved analogously for the parabola. Thus in the case of this proposition, FD will become parallel to CE . Hence any straight line from the focus of a parabola parallel to a tangent will make a right angle with the straight line drawn from the focus to the intersection of the tangent and the perpendicular to the axis at the vertex.



at right angles and CED tangent and let the rectangle $AF FB$ and the rectangle $4G GB$ equal to the fourth part of the figure be applied on each side (Eucl vi 28 29), as has been said and let CF , CG , DF , and DG be joined

I say that angle CFD and angle CGD are each a right angle

For since it has been shown

$$\text{rect } AC \cdot BD = \text{fourth figure on } AB \text{ (III 42),}$$

and since also

$$\text{rect } AF \cdot FB = \text{fourth figure on } AB,$$

therefore

$$\text{rect } AC \cdot BD = \text{rect } AF \cdot FB$$

Therefore

$$AC : AF = FB : BD$$

And the angles at points A and B are right therefore

$$\text{angle } ACF = \text{angle } BFD \text{ (Eucl vi 6),}$$

and

$$\text{angle } AFC = \text{angle } FDB$$

And since angle CAF is right, therefore

$$\text{angle } ACF + \text{angle } AFC = 1 \text{ rt. angle}$$

And it has also been shown that

$$\text{angle } ACF = \text{angle } DFB,$$

therefore

$$\text{angle } AFC + \text{angle } DFB = 1 \text{ right angle}$$

Therefore

$$\text{angle } DFC = 1 \text{ right angle}$$

Then likewise it could also be shown

$$\text{angle } CGD = 1 \text{ right angle}$$

PROPOSITION 46

With the same things being so the straight lines joined make equal angles with the tangents

For with the same things supposed I say that

$$\text{angle } ACF = \text{angle } DCG$$

and

$$\text{angle } CDF = \text{angle } BDG$$

For since it has been shown that both angle CFD and angle CGD are right angles (III 45) the circle described about CD as a diameter will pass through points F and G therefore


$$\text{angle } DCG = \text{angle } DFG$$

for they are on the same segment of the circle And it was shown angle $DFG = \text{angle } ACF$ (III 45), and so

$$\text{angle } DCG = \text{angle } ACF$$

And likewise also

$$\text{angle } CDF = \text{angle } BDG$$



PROPOSITION 47


With the same things being so the straight line drawn from the point of meeting of the joined straight lines to the point of contact will be perpendicular to the tangent

For let the same things as before be supposed and let CG and FD meet each other at H and let CD and BA produced meet at K , and let EH be joined

I say that EH is perpendicular to CD

For if not let HL be drawn from H perpendicular to CD Since then
 $\text{angle } CDF = \text{angle } GBD$ (III 46),
 and also

$$\text{rt angle } DBC = \text{rt angle } DLH,$$



therefore triangle DGB is similar to triangle LHD Therefore

$$GD : DH :: BD : DL$$

But

$$GD : DH :: FC : CH$$

because the angles at F and G are right angles (III 45) and the angles at H are equal but

$$\frac{FC}{BD} = \frac{CH}{DL} = \frac{AC}{AC} = \frac{CL}{CL}$$

because of the similarity of the triangles AFC and LCH (III 46), therefore also

Alternately

$$BD : AC :: DL : CL$$

But

$$BD : AC :: BA : AA$$

therefore also

$$DL : CL :: BA : AA$$

Let EM be drawn from E parallel to AC , therefore it will have been dropped ordinate to AB (II 7), and

$$BA : KA = BM : MA$$

And

$$BM : MA = DE : EC$$


therefore also

$$DL : CL = DE : EC,$$


and this is absurd. Therefore HL is not perpendicular nor is any other straight line except HE *

PROPOSITION 48

With the same things being so, it must be shown that the straight lines drawn from the point of contact to the points produced by the application make equal angles with the tangent



*There is the analogous theorem for the parabola FD becomes a straight line parallel to



CE and CG a straight line parallel to AB . Again HE is perpendicular to CE and this can be proved rigorously as well as understood by analogy

For let the same things be supposed, and let EF and EG be joined
I say that

$$\text{angle } CEF = \text{angle } GED$$

For since angles DGH and DEH are right angles (in 45 47), the circle described about DH as a diameter will pass through the points E and G (Eucl in 31), and so

$$\text{angle } DHG = \text{angle } DLG \text{ (Eucl in 21),}$$

for they are in the same segment Likewise then also

$$\text{angle } CEF = \text{angle } CHF$$

But


$$\text{angle } CHF = \text{angle } DHG,$$

for they are vertical angles, therefore also


$$\text{angle } CEF = \text{angle } DLG *$$

PROPOSITION 49

With the same things being so if from one of the points (of application) a perpendicular is drawn to the tangent then the straight lines from that point to the ends of the axis make a right angle



Here there is another and important analogous theorem for the parabola EG becomes parallel to AB and



$$\text{angle } DEG = \text{angle } CEF$$

For let the same things be supposed, and let the perpendicular GH be drawn from G to CD , and let AH and BH be joined

I say that angle AHB is a right angle

For since angle DBG is a right angle and also angle DHG the circle described about DG as a diameter will pass through H and B , and
 angle $BHG = \text{angle } BDG$

But it was shown

$$\text{angle } AGC = \text{angle } BDG \text{ (III 46),}$$

therefore also

$$\text{angle } BHG = \text{angle } AGC = \text{angle } AHC \text{ (Eucl III 21) }$$

And so also

$$\text{angle } CHG = \text{angle } AHB$$

But angle CHG is a right angle, therefore also angle AHB is a right angle


PROPOSITION 50

With the same things being so if from the center of the section there falls to the tangent a straight line parallel to the straight line drawn through the point of contact and one of the points (of application), then it will be equal to one half the axis

For let there be the same things as before and let H be the center, and let EF be joined and let DC and BA meet at K and through H let HL be drawn parallel to EF

I say that

$$HL = HB$$



For let EG , AL , LG be joined and through G let GM be drawn parallel to EF . Since then

$$\text{rect } AF \cdot FB = \text{rect } AG \cdot GB \text{ (See III 45),}$$

therefore

$$AF = GB$$

But also

$$AH = HB,$$

therefore also

$$FH = HG$$

And so also

$$EL = LM$$

And since it was shown (III 48)

$$\text{angle } CEF = \text{angle } DEG,$$

and

$$\text{angle } CEF = \text{angle } EMG,$$

therefore also

$$\text{angle } EMG = \text{angle } DEG$$

And therefore

$$EG = GM$$

But it was also shown

$$EL = LM$$

therefore GL is perpendicular to EM And so through what was shown before (III 49) angle ALB is a right angle and the circle described about AB as a diameter will pass through L And

$$HA = HB$$

therefore also since HL is a radius of the semicircle,

$$HL = HB$$


PROPOSITION 51

If a rectangle equal to the fourth part of the figure is applied from both sides to the axis of an hyperbola or opposite sections and exceeding by a square figure and straight lines are deflected from the resulting points of application to either one of the sections, then the greater of the two straight lines exceeds the less by exactly as much as the axis

For let there be an hyperbola or opposite sections whose axis is AB and center C and let each of the rectangles $AD \cdot DB$ and $AE \cdot EB$ be equal to the fourth part of the figure and from points E and D let the straight lines EF and FD be deflected to the line of the section

I say that

$$EF = FD + AB$$



Let FKH be drawn tangent through F , and GCH through C parallel to FD therefore

$$\text{angle } KHG = \text{angle } KFD,$$

for they are alternate And

$$\text{angle } KFD = \text{angle } GFH \text{ (III 48),}$$

therefore

$$GF = GH$$

But

$$GF = GE,$$

since also

$$AE = BD$$

and

$$AC = CB$$

and

$$EC = CD,$$

and therefore

$$GH = EG$$

And so

$$FE = 2GH$$

And since it has been shown (III 50)

$$CH = CB,*$$

therefore

$$FE = 2(GC + CB)$$

But

$$FD = 2GC,$$

and

$$AB = 2CB,$$

therefore

$$FE = FD + AB$$

And so EF is greater than FD by AB

PROPOSITION 52

If in an ellipse a rectangle equal to the fourth part of the figure is applied from both sides to the major axis and deficient by a square figure, and from the points resulting from the application straight lines are deflected to the line of the section then they will be equal to the axis

Let there be an ellipse whose major axis is AB and let each of the rectangles $AC \cdot CB$ and $AD \cdot DB$ be equal to the fourth of the figure and from C and D let the straight lines CE and ED have been deflected to the line of the section

I say that

$$CE + ED = AB$$

*For

$$GF = GH$$

and by III 50 a line $C(\lambda)$ drawn parallel to GF is equal to CB But also

$$C(\lambda) = CH$$

Hence


$$CH = CB$$

Let FEH be drawn tangent and G be center and through it let GKH be drawn parallel to CE Since then

$$\text{Angle } CEF = \text{angle } HFA \text{ (III 48),}$$

and

$$\text{angle } CEF = \text{angle } EHA,$$



therefore also

$$\text{angle } EHA = \text{angle } HEK$$

Therefore also

$$HA = AE$$

And since

$$AG = GB$$

and

$$AC = DB,$$

therefore also

$$CG = GD,$$

and so also

$$FA = AD$$

And for this reason

$$ED = 2HA,$$

and

$$EC = 2AG$$

and

$$ED + EC = 2GH$$

But also

$$AB = 2GH \text{ (III 50),}$$

therefore

$$AB = FD + EC$$

PROPOSITION 53

If in an hyperbola or ellipse or circumference of a circle or opposite sections straight lines are drawn from the vertex of a diameter parallel to an ordinate, and straight lines drawn from the same ends to the same point on the line of the section cut the parallels then the rectangle contained by the straight lines cut off is equal to the figure on that same diameter

Let there be one of the aforesaid sections ABC whose diameter is AC , and let AD and CF be drawn parallel to an ordinate and let ABE and CBD be drawn across


I say that

$$\text{rect } AD \cdot FC = \text{figure on } AC$$

For let BF be drawn from B parallel to an ordinate

Therefore

rect AI', FC sq FB transverse side upright side
sq AC the figure (1 21)



But

rect $AF FC$ sq FB comp $AF FB FC FB$,

therefore

figure sq AC comp $FB AF, FB FC$

But

$AF FB AC CE$

and

$FC FB AC AD$

therefore

figure sq AC comp $CE AC AD AC$

And also

rect $AD CE$ sq AC comp $CE AC, AD AC$,

therefore

figure sq AC rect $AD CE$ sq AC

Therefore

rect $AD CE$ = figure on AC

PROPOSITION 54

If two tangents to a section of a cone or to a circumference of a circle meet and through the points of contact parallels to the tangents are drawn and from the points of contact to the same point of the line of the section straight lines are drawn across cutting the parallels then the rectangle contained by the straight lines cut

off to the square on the straight line joining the points of contact has a ratio compounded of the ratio which the inside segment line joining the point of meeting of the tangents and the midpoint of the straight line joining the points of contact has in square to the remainder and of the ratio which the rectangle contained by the tangents has to the fourth part of the square on the straight line joining the points of contact

Let there be a section of a cone or circumference of a circle ABC and tangents AD and CD , and let AC be joined and bisected at E , and let DBE be joined, and let AF be drawn from A parallel to CD , and CG from C parallel to AD and let some point H on the section be taken and let the straight lines AH and CH be joined and produced to G and F

I say that

$$\text{rect } AF, CG \text{ sq } AC \text{ comp sq } EB \text{ sq } BD, \text{ rect } AD, DC \\ \text{fourth sq } AC \text{ or rect } AE, EC$$

For let $KHO\lambda L$ be drawn from H parallel to AC and from B , MBN parallel to AC , then it is evident that MN is tangent (II 29 5, 6) Since then

$$AE = EC,$$

also


$$MB = BN$$

and

$$KO = OL$$

and


$$HO = O\lambda \text{ (II 7)}$$



and

$$KH = \sqrt{L}$$

Since then MB and MA are tangents and KHL has been drawn parallel to MB ,



$$\text{sq } AM \text{ sq } MB \text{ sq } AA \text{ rect } XA \text{ } KH \text{ (III 16)}$$

or

$$\text{sq } AM \text{ rect } MB, BV \text{ sq } AA \text{ rect } LH, HA$$

And

$$\text{rect } NC, AM \text{ sq } AM \text{ rect } LC, AK \text{ sq } AK \text{ (Eucl vi 2, v, 18),}$$

therefore *ex aequali*

$$\text{rect } NC, AM \text{ rect } MB, BN$$

$$\text{rect } LC, AK \text{ rect } LH, HK$$

But


$$\text{rect } LC, AK \text{ rect } LH, HK \text{ comp } LC \text{ } LH, AK \text{ } HK$$

or

$$\text{rect } LC, AK \text{ rect } LH, HK \text{ comp } FA \text{ } AC, GC \text{ } CA$$

which is the same as

$$\text{rect } GC \text{ } FA \text{ sq } CA$$



Therefore

$$\text{rect } NC, AM \quad \text{rect } MB, BN \quad \text{rect } GC, FA \quad \text{sq } CA$$

But with the rectangle ND, DM taken as a mean

$$\text{rect } NC, AM \quad \text{rect } MB, BN \quad \text{comp}$$

therefore $\text{rect } NC, AM \quad \text{rect } ND, DM \quad \text{rect } ND, DM \quad \text{rect } MB, BN,$

$$\text{rect } GC, FA \quad \text{sq } CA \quad \text{comp}$$

$$\text{rect } NC, AM \quad \text{rect } ND, DM \quad \text{rect } ND, DM \quad \text{rect } MB, BN$$

But

$$\text{rect } NC, AM \quad \text{rect } ND, DM \quad \text{sq } EB \quad \text{sq } BD,$$

and

$$\text{rect } ND, DM \quad \text{rect } NB, BM \quad \text{rect } CD, DA \quad \text{rect } CE, EA,$$

therefore

$$\text{rect } GC, FA \quad \text{sq } CA \quad \text{comp} \quad \text{sq } BE \quad \text{sq } BD, \text{rect } CD, DA \quad \text{rect } CE, EA$$

PROPOSITION 55


If two straight lines touching opposite sections meet and through the point of meeting a straight line is drawn parallel to the straight line joining the points of contact and from the points of contact parallels to the tangents are drawn across, and straight lines are produced from the points of contact to the same point of one of the sections cutting the parallels then the rectangle contained by the straight lines cut off will have to the square on the straight line joining the points of contact the ratio which the rectangle contained by the tangents has to the square on the straight line drawn through the point of meeting parallel to the straight line joining the points of contact as far as the section.

Let there be the opposite sections ABC and DEF , and tangents to them AG and GD and let AD be joined and from G let CGE be drawn parallel to AD , and from A AM parallel to DG and from D DM parallel to AG and let some point I be taken on the section DF , and let ANF and FDH be joined

I say that

$$\text{sq } CG \quad \text{rect } AG, GD \quad \text{sq } AD \quad \text{rect } HA, DN$$

For let ILK be drawn through I parallel to AD



Since then it has been shown that

$$\text{sq } EG \text{ sq } GD \text{ rect } BL \text{ } LF \text{ sq } DL \text{ (III 20),}$$

and

$$CG = EG \text{ (II 38),}$$

and

$$BK = LF \text{ (II 38),}$$

therefore

$$\text{sq } CG \text{ sq } GD \text{ rect } KF, FL \text{ sq } DL$$

And also

$$\text{sq } GD \text{ rect } AG \text{ } GD \text{ sq } DL \text{ rect } DL \text{ } AK \text{ (Eucl VI 2, 1),}$$

therefore *ex aequali*

$$\text{sq } GC \text{ rect } AG, GD \text{ rect } KF, FL \text{ rect } DL, AK$$

But

$$\text{rect } KF, FL \text{ rect } DL, AK \text{ comp } KF \text{ } AK, FL \text{ } DL$$

But

$$KF \text{ } AK \text{ } AD \text{ } DN,$$

and

$$FL \text{ } DL \text{ } AD \text{ } HA,$$

therefore

$$\text{sq } CG \text{ rect } AG, GD \text{ comp } AD \text{ } DN, AD \text{ } HA$$

And also

$$\text{sq } AD \text{ rect } HA, DN \text{ comp } AD \text{ } DN, AD \text{ } HA,$$

therefore

$$\text{sq } CG \text{ rect } AG \text{ } GD \text{ sq } AD \text{ rect } HA, DN$$

PROPOSITION 56

If two straight lines touching one of the opposite sections meet and parallels to the tangents are drawn through the points of contact and straight lines cutting the parallels are drawn from the points of contact to the same point of the other section, then the rectangle contained by the straight lines cut off will have to the square on the straight line joining the points of contact the ratio compounded of the ratio which of the straight line joining the point of meeting and the midpoint that part between the midpoint and the other section has in square to that part between the same section and the point of meeting and of the ratio which the rectangle contained by the tangents has to the fourth part of the square on the straight line joining the points of contact

Let there be the opposite sections AB and CD whose center is O and tangents $A'EFG$ and $BEHK$ and let AB be joined and bisected at L and let LE be joined and drawn across to D , and let AM be drawn from A parallel to BE and BN from B parallel to AE and let some point C be taken on the section CD and let CBM and CAN be joined

I say that

$$\text{rect } MA \text{ } BN \text{ sq } AB \text{ comp sq } LD \text{ sq } DE, \text{ rect } AE \text{ } EB \\ \text{fourth sq } AB \text{ or rect } AL \text{ } LB$$

For let GCK and HDF be drawn from C and D parallel to AB then it is evident that

$$HD = DF$$

and

$$AX = XG$$

TRANSLATOR'S APPENDIX ON THREE and
FOUR LINE LOCI

The three-line locus property of conics is easily deduced for the ellipse hyperbola parabola and circle from III 54, and for the opposite sections from III 55 and 56. The three-line locus property of conics can be stated thus: Any conic section or circle or pair of opposite sections can be considered as the locus of points whose distances from three given fixed straight lines (the distances being either perpendicular or at a given constant angle to each of the given straight lines) are such that the square of one of the distances is always in a constant ratio to the rectangle contained by the other two distances.

It is shown in III 54 that in the case of conic sections and circles $\text{rect } AF \cdot CG \text{ sq } AC \text{ comp sq } EB \text{ sq } BD, \text{ rect } AD, DC \text{ fourth sq } AC$. Now if we consider the straight lines AD, DC and AC as fixed and given and therefore straight line DE fixed and given as bisecting AC , then it is evident that the straight lines AC, EB, BD, AD, DC , and therefore the squares on them and the rectangles contained by them are also fixed and given. Then although as the point H is taken at different points along the conic the straight lines AF and CG change in magnitude nevertheless the magnitude of the rectangle $AF \cdot CG$ because of the above proportion remains constant.

For let HX be drawn parallel to BE and HY to AD and HZ to DC . Then HX is the distance from H to AC at a given angle and AY because of parallels represents the distance from H to AD at another given angle and ZC represents the distance from H to DC at another given angle. Then by similar triangles

$$\frac{CZ}{AY} = \frac{ZH}{YH} = \frac{AC}{AC} = \frac{AF}{CG},$$

therefore compounding
 $\text{rect } CZ, AY \text{ rect } ZH YH \text{ sq } AC \text{ rect } AF, CG$

Now we have seen that the rectangle $AF \cdot CG$ is a constant magnitude as the point H changes and the square on AC is constant therefore their ratio is constant. Therefore


$\text{rect } CZ, AY \text{ rect } ZH YH$ is a constant ratio (1)

Again by similar triangles

$$\frac{ZH}{YH} = \frac{HX}{HX} = \frac{CD}{AD} = \frac{DE}{DE},$$

therefore compounding
 $\text{rect } ZH YH \text{ sq } HX \text{ rect } CD, AD \text{ sq } DE$

But rectangle CD, AD and the square on DE are constant magni-



inverse of (α) is compounded with (β), and the inverse of (γ) with (δ) we would have two constant ratios

$$\text{pllpd } HY, HP \text{ } HP \text{ } \text{pllpd } HV, HR, HR, \quad (\epsilon)$$

$$\text{pllpd } HV, HS \text{ } HS \text{ } \text{pllpd } HY, HQ, HQ \quad (\zeta)$$

Again compounding the first of these with the second we would have finally
rect HP, HS rect HQ, HR , a constant ratio

And this is the property of the four line locus namely the locus of points H such that the rectangle contained by the distances from points H to any two given fixed straight lines FG and ID has to the rectangle contained by the distances from H to two other fixed straight lines IG, FD , a constant ratio

The rigorous method of effecting these compoundings is as follows For inverting (α), by Eucl xi 32 we have the constant ratios

$$\text{sq } HP \text{ rect } H\lambda, HV \text{ } \text{pllpd } HP \text{ } HP, HY \text{ } \text{pllpd } HX, HV, HY,$$

$$\text{rect } H\lambda \text{ } HY \text{ sq } HR \text{ } \text{pllpd } HX \text{ } HY, HV \text{ } \text{pllpd } HR, HR \text{ } HV$$

Hence, by definition the ratio (a constant one) compounded of these two is
 $\text{pllpd } H\lambda \text{ } HP, HP \text{ } \text{pllpd } HV, HR \text{ } HR$

And in the same way we find the constant ratio compounded of the inverse of (γ) and (δ) Now

$$\text{pllpd } HY \text{ } HP, HP \text{ } \text{pllpd } HV, HR \text{ } HR \text{ comp } HY \text{ } HV, \text{sq } HP \text{ } \text{sq } HR$$

$$\text{pllpd } HV, HS \text{ } HS \text{ } \text{pllpd } HY, HQ \text{ } HQ \text{ comp } HV \text{ } HY \text{ sq } HS \text{ } \text{sq } HQ$$

If then we take two lines M and N such that

$$\frac{HP}{HS} = \frac{HR}{HQ} = \frac{HR}{HQ} = \frac{M}{N}, \quad (\eta)$$

$$\frac{HS}{HS} = \frac{HQ}{HQ} = \frac{HQ}{HQ} = \frac{N}{N}, \quad (\theta)$$

then

$$\text{sq } HP \text{ } \text{sq } HR \text{ } \frac{HP}{HS} = \frac{M}{N},$$

$$\text{sq } HS \text{ } \text{sq } HQ \text{ } \frac{HS}{HS} = \frac{N}{N}$$

Hence

$$\text{ratio comp } H\lambda \text{ } HV \text{ sq } HP \text{ } \text{sq } HR \text{ ratio comp } H\lambda \text{ } HV \text{ } HP \text{ } M$$

$$\text{ratio comp } HV \text{ } HY, \text{sq } HS \text{ } \text{sq } HQ \text{ ratio comp } HV \text{ } HY, HS \text{ } N$$

But

$$\text{rect } HY \text{ } HP \text{ rect } HV \text{ } M \text{ comp } HY \text{ } HV, HP \text{ } M,$$

$$\text{rect } HV, HS \text{ rect } HY \text{ } N \text{ comp } HV \text{ } HY, HS \text{ } N$$

and

$$\text{pllpd } H\lambda \text{ } HP \text{ } HS \text{ } \text{pllpd } HV \text{ } M, HS \text{ rect } HY \text{ } HP \text{ rect } H\lambda \text{ } M,$$

$$\text{pllpd } HV \text{ } HS, M \text{ } \text{pllpd } HY \text{ } N \text{ } M \text{ rect } HV, HS \text{ rect } HY \text{ } N,$$

and these are constant ratios Hence compounding we get the constant ratio

$$\text{pllpd } HY, HP, HS \text{ } \text{pllpd } HY, N, M,$$

which is the same as the constant ratio

$$\text{rect } HP, HS \text{ rect } N, M$$

Now taking L and O as some constants

$$\text{rect } HP \text{ } HS \text{ rect } N \text{ } M \text{ } L \text{ } O$$

and

$$\text{rect } HP \text{ } HS \text{ rect } HR \text{ } HQ \text{ rect } HR \text{ } HQ \text{ rect } M \text{ } N$$

by compounding (η) and (θ) But equal ratios have equal duplicate ratios (Heath's note to Euclid vi 22) and hence

$$\text{rect } HP \text{ } HS \text{ rect } HR \text{ } HQ \text{ is constant}$$

In the case of opposite sections it is shown in III 56

$$\text{rect } MA \text{ } BV \text{ sq } AB \text{ comp sq } LD \text{ sq } DE \text{ rect } AE \text{ } FB \text{ fourth sq } AB$$

Then it is evident for the same reasons as before that for different points C the

magnitudes MA and BN may change, but the rectangle $MA BN$ is a constant magnitude

For as before, let CY be drawn parallel to DE , CY to EA , CZ to EB By similar triangles

$$\frac{AY}{BZ} = \frac{YC}{ZC} = \frac{AB}{AB} = \frac{BN}{MA},$$

therefore compounding


$$\text{rect } AY, BZ = \text{rect } YC, ZC \quad \text{sq } AB = \text{rect } MA, BN$$

Since rectangle MA, BN is constant as C changes, and also the square on AB is constant therefore

$$\text{rect } AY, BZ = \text{rect } YC, ZC \text{ is a constant ratio} \quad (1)$$

Again by similar triangles

$$\frac{ZC}{CY} = \frac{EB}{EL},$$



$$\frac{YC}{CY} = \frac{EA}{EL}$$

therefore compounding

$$\text{rect } YC, ZC = \text{sq } CY = \text{rect } EB, EA = \text{sq } EL$$

Hence

$$\text{rect } YC, ZC = \text{sq } CY \text{ is a constant ratio} \quad (2)$$

Compounding (1) and (2) we have a constant ratio

$$\text{rect } AY, BZ = \text{sq } CY$$


But AY and BZ are equal to CA and CB the distances from C . This is the property of the three line locus of section C with respect to the straight lines EA and EB tangents to the other section and LB the straight line joining their points of contact. And so one opposite section is a three line-locus to the tangents to the other of the opposite sections. That it is also a four line locus could be shown in the same way as before.

Again from III 55 we can conclude that both of the opposite sections together are a three line locus to the triangle formed by a tangent to each of the sections and the straight line joining their points of contact. For by III 55

$$\text{rect } HA, DV = \text{sq } AD = \text{rect } AG, GD = \text{sq } CG$$

Now since the three last terms of this proportion are evidently constants as the point F changes therefore also although HA and DA change with F yet rectangle HA, DV remains constant in magnitude. Then reproducing the figure of

in 55 we drop YF parallel to DL and FZ to KA , and FV to GE , where E is the midpoint of AD . Then by similar triangles



YD	FY	AD	HA,
AZ	FZ	AD	DN

therefore compounding

rect YD AZ rect FY, FZ sq AD rect HA, DN

But the last two terms are constant, therefore

rect $YD \cdot AZ$ rect FY, FZ is a constant ratio (1)

Again by similar triangles

$$\begin{array}{cc} \Gamma Y & F Y \\ F Z & F Y \end{array} \quad \begin{array}{cc} D G & L G \\ A G & E G. \end{array}$$

therefore compounding

$$\text{rect } FY, FZ \quad \text{sq } F\mathbf{X} \quad \text{rect } DG, AG \quad \text{rect } ED, EG$$

But the last two terms are constant therefore

rect $FY FZ$ sq FY is a constant ratio (2)

Compounding (1) and (2), we see that

rect $YD \cdot AZ$ sq FY is a constant ratio

But this is the definition of a three line locus that the rectangle contained by the distances from any point on the locus to two fixed straight lines have to the square on the distance to a third fixed straight line a constant ratio. But

$$DY = LF$$

$$AZ = \mathbf{1}F$$

and FY is the distance from F to AD . And so the ratio fulfills the definition

Furthermore if we consider B the point of intersection of the straight line AF drawn parallel to AD with the other opposite section and draw BS parallel to FY , BR to FV , and BT to FZ since they are parallels between parallels,

$$BR = FV$$

$$KF = AZ$$

$$TA = BK$$

magnitudes MA and BN may change, but the rectangle MA, BN is a constant magnitude

For as before let CX be drawn parallel to DE , CY to EA , CZ to EB By similar triangles

$$\frac{AY}{BZ} = \frac{YC}{ZC} = \frac{AB}{AB} = \frac{BN}{MA},$$

therefore compounding


$$\text{rect } AY, BZ = \text{rect } YC, ZC = \text{sq } AB = \text{rect } MA, BN$$

Since rectangle MA, BN is constant as C changes and also the square on AB is constant therefore

$$\text{rect } AY, BZ = \text{rect } YC, ZC \text{ is a constant ratio} \quad (1)$$

Again by similar triangles

$$\frac{ZC}{CX} = \frac{EB}{EL},$$



$$\frac{YC}{CX} = \frac{EA}{EL}$$

therefore compounding

$$\text{rect } YC, ZC = \text{sq } CX = \text{rect } EB, EA = \text{sq } EL$$

Hence

$$\text{rect } YC, ZC = \text{sq } CX \text{ is a constant ratio} \quad (2)$$

Compounding (1) and (2) we have a constant ratio

$$\text{rect } AY, BZ = \text{sq } CX$$

But AY and BZ are equal to CA' and CB the distances from C This is the property of the three line locus of section C with respect to the straight lines EA and EB tangents to the other section and EB the straight line joining their points of contact And so one opposite section is a three line-locus to the tangents to the other of the opposite sections That it is also a four line locus could be shown in the same way as before

Again from III 55 we can conclude that both of the opposite sections together are a three line locus to the triangle formed by a tangent to each of the sections and the straight line joining their points of contact For by III 55

$$\text{rect } HA, DV = \text{sq } AD = \text{rect } AG, GD = \text{sq } CG$$

Now since the three last terms of this proportion are evidently constants as the point F changes therefore also although HA and DA change with F , yet rectangle HA, DV remains constant in magnitude Then reproducing the figure of

INTRODUCTION TO ARITHMETIC

But it was shown in the course of III 55 that

$$BK = LF,$$

$$BL = AF$$

Hence

$$TA = BK = LD = LF,$$


$$AZ = AF = BL$$

Therefore

$$\text{rect } LF, AF \text{ sq } FV \quad \text{rect } BK, BL \text{ sq } BR$$

Hence any point B on one opposite section fulfill. the same constant ratio with respect to its distances from the three fixed lines AD , GD , and AG as any point F on the other opposite section

It can be similarly deduced that the opposite sections are together a four line locus with respect to any four fixed straight lines joining points, two on each section (the points being four points of contact of four tangents and the straight lines the straight lines joining them)



To sum up a parabola ellipse circle and hyperbola are three-line loci with respect to any two tangents to them and a straight line joining the points of contact. One opposite section is also a three line locus with respect to any two tangents to the other section together with the straight line joining their points of contact. The two opposite sections together are a three line-locus with respect to two tangents each to one of the sections together with the straight line joining their points of contact.

The parabola ellipse circle and hyperbola are four line loci with respect to any inscribed quadrilateral. One opposite section is also a four line locus with respect to any quadrilateral inscribed in the other section. The two opposite sections together are a four line locus to any four straight lines joining four points, two lying on each opposite section.

INTRODUCTION TO ARITHMETIC

BIOGRAPHICAL NOTE

NICOMACHUS, *fl c A D 100*

NICOMACHUS OF GERASA flourished around the end of the first century of our era. In one of his surviving books the *Introduction to Harmonics*, he mentions a certain Thrasyllus presumably Thrasyllus of Mendes, a writer on music who lived in the reign of Tiberius. Another book by Nicomachus, the *Introduction to Arithmetic* was translated into Latin by Apuleius under the Antonines. This places the life of Nicomachus somewhere between the middle of the first century and the middle of the second century. Perhaps the fact that Ptolemy, whose recorded astronomical observations were made between A D 127 and 151, is not mentioned in the *Introduction to Harmonics* makes it probable that he was not yet famous at the time Nicomachus was writing.

The manuscripts of Nicomachus' books and the scholia call him "of Gerasa." The best known city of that name was in Palestine and was primarily Greek. However, it can hardly be supposed that Nicomachus received all of his philosophical and mathematical education at Gerasa. He probably studied at Alexandria at this time the center of mathematical studies and of Neo-Pythagoreanism. Jamblichus says of Nicomachus: "The man is great in mathematics, and had as instructors those that were most skilled in the subject."

Nothing is known of the personal life of Nicomachus except what is said or implied in the dedication of the *Introduction to Harmonics* to an unknown lady: "But I must spur on all my zeal, most noble and august lady, since it is you that bid me. And if the gods are willing, just as soon as I shall have leisure and a rest from my journeyings, I will compile for you a better and more detailed *Introduction* dealing with this very subject, and so that you may the more easily follow the argument, I will take my beginning, say, from the same point as that at which I began your instruction when I was expounding the subject to you."

Nicomachus appears to have been an important member of the Neo-Pythagorean group, though his extant writings would seem to indicate that he was a popularizer and a compiler of manuals and not the head of a school. Besides the *Introduction to Arithmetic* and the *Introduction to Harmonics*, he also wrote a book on the mystical doctrine of number called *Theologoumena Arithmeticae*, which is one of the best sources on Neo-Pythagoreanism; extracts and paraphrases of this work survive in a later anonymous work of the same name and in the *Bibliotheca*, a collection of extracts from ancient works made in the ninth century by Photius, patriarch of Constantinople. Nicomachus also wrote an *Introduction to Geometry* and a *Life of Pythagoras*, which have not survived, and a larger work on music, possibly that promised in the dedication to the *Introduction to Harmonics*, of which we have only fragments. He may have written a book on the interpretation of Plato, though the evidence for it is slight, and also an *Introduction to Astronomy*, thereby completing the quadrivium series.

The success of the *Introduction to Arithmetic* must have been immediate. It

was used as a text book throughout later antiquity and in the Latin paraphrase of Boethius throughout the Middle Ages. It had a host of commentators. In the *Philopatrias* attributed to Lucian, a character says "You reckon like Nicomachus." This remark lends itself to more than one interpretation, but in any case it is evidence of his fame. Nicomachus also appears to have been considered one of the 'golden chain' or succession of true philosophers, for Proclus the fifth century Neo-Platonist who belonged to that 'chain,' claimed, on the basis of a dream, that he had within him the soul of Nicomachus.

CONTENTS

BIOGRAPHICAL NOTE	p 807
BOOK I,	p 811
BOOK II,	p 829

BOOK ONE

CHAPTER I

[1] The ancients who under the leadership of Pythagoras first made science systematic defined philosophy as the love of wisdom. In deed the name itself means this and before Pythagoras all who had knowledge were called wise indiscriminately—a carpenter for example a cobbler a helmsman and in a word anyone who was versed in any art or handicraft Pythagoras however restricting the title so as to apply to the knowledge and comprehension of reality and calling the knowledge of the truth in this the only wisdom naturally designated the desire and pursuit of this knowledge a philosophy as being desire for wisdom.

[2] He is more worthy of credence than those who have given other definitions since he makes clear the sense of the term and the thing defined. This wisdom he defined as the knowledge or science of the truth in real things conceiving science to be a steadfast and firm apprehension of the underlying substance and real things to be those which continue uniformly and the same in the universe and never depart even briefly from their existence these real things would be things immaterial by sharing in the substance of which everything else that exists under the same name and is so called is said to be this particular thing and exists.

[3] For bodily material things are to be sure forever involved in continuous flow and change—in imitation of the nature and peculiar quality of that eternal matter and substance which has been from the beginning and which was all changeable and variable throughout. The bodiless things however of which we conceive in connection with or together with matter such as qualities quantities configurations largeness smallness equality relations actualities dispositions places times all those things in a word whereby the qualities found in each body are comprehended—all these are of themselves immovable and unchangeable but accidentally they share in and partake of the affections of the body to which they belong.

[4] Now it is with such things that 'wisdom' is particularly concerned but accidentally also with things that share in them that is bodies.

CHAPTER II

[1] Those things however are immaterial eternal without end and it is their nature to persist ever the same and unchanging abiding by their own essential being and each one of them is called real in the proper sense. But what are involved in birth and destruction growth and diminution all kinds of change and participation are seen to vary continually, and while they are called real things by the same term as the former so far as they partake of them they are not actually real by their own nature for they do not abide for even the shortest moment in the same condition but are always passing over in all sorts of changes. [2] To quote the words of Timaeus in Plato¹

What is that which always is and has no birth and what is that which is always becoming but never is? The one is apprehended by the mental processes with reasoning and is ever the same the other can be guessed at by opinion in company with unreasoning sense a thing which becomes and passes away but never really is.

[3] Therefore if we crave for the goal that is worthy and fitting for man namely happiness of life²—and this is accomplished by philosophy alone and by nothing else and philosophy as I said means for us desire for wisdom and wisdom the science of the truth in things and of things some are properly so called others merely share the name—it is reasonable and most necessary to distinguish and systematize the accidental qualities of things.

[4] Things then both those properly so called and those that simply have the name are some of them unified and continuous for example an animal the universe a tree and

Timaeus 27

¹The word used by Nicomachus *ἐξωταρισ* once employed by Aristotle in the *Ethics* 1038^a 20 ff

the like which are properly and peculiarly called magnitudes others are discontinuous in a side-by side arrangement and, as it were in heaps which are called multitudes a flock for instance a people a heap a chorus and the like

[5] Wisdom then must be considered to be the knowledge of these two forms. Since however all multitude and magnitude are by their own nature of necessity infinite—for multitude starts from a definite root and never ceases in increasing and magnitude when division beginning with a limited whole is carried on cannot bring the dividing process to an end but proceeds therefore to infinity—and since sciences are always sciences of limited things and never of infinites it is accordingly evident that a science dealing either with magnitude *per se* or with multitude *per se* could never be formulated for each of them is limitless in itself multitude in the direction of the more and magnitude in the direction of the less. A science however would arise to deal with something separated from each of them with quantity set off from multitude and size set off from magnitude.

CHAPTER III

[1] Again to start afresh since of quantity one kind is viewed by itself having no relation to anything else as even odd perfect and the like and the other is relative to something else and is conceived of together with its relationship to another thing like double greater smaller half one and one-half time one and one-third times and so forth it is clear that two scientific methods will lay hold of and deal with the whole investigation of quantity arithmetic absolute quantity and music relative quantity.¹

[2] And once more inasmuch as part of size is in a state of rest and stability and another part in motion and revolution two other sciences in the same way will accurately treat of size geometry the part that abides and is at rest astronomy that which moves and revolves.

[3] Without the aid of these then it is not possible to deal accurately with the forms of being nor to discover the truth in things knowledge of which is wisdom and evidently not even to philosophize properly for just as painting contributes to the manual arts and correctness of theory so in truth lines numbers harmonic intervals and the revolutions of circles

bear aid to the learning of the doctrines of wisdom says the Pythagorean Androcydes [4] Likewise Archytas of Tarentum at the beginning of his treatise *On Harmony* says the same thing in about the same words. It seems to me that they do well to study mathematics and it is not at all strange that they have correct knowledge about each thing what it is. For if they knew rightly the nature of the whole, they were also likely to see well what is the nature of the parts. About geometry indeed, and arithmetic and astronomy they have handed down to us a clear understanding and not least also about music. For these seem to be sister sciences for they deal with sister subjects the first two forms of being.

[5] Plato too at the end of the thirteenth book of the *Laws* to which some give the title *The Philosopher* because he investigates and defines in it what sort of man the real philosopher should be in the course of his summary of what had previously been fully set forth and established adds. Every diagram system of numbers every scheme of harmony and every law of the movement of the stars ought to appear one to him who studies rightly and what we say will properly appear if one studies all things looking to one principle for there will be seen to be one bond for all the things and if any one attempts philosophy in any other way he must call on Fortune to assist him. For there is never a path without these this is the way these the studies be they hard or easy by this course must one go and not neglect it. The one who has attained all these things in the way I describe him I for my part call wisest and this I maintain through thick and thin. [6] For it is clear that the studies are like ladders and bridges that carry our minds from things apprehended by sense and opinion to those comprehended by the mind and understanding and from those material physical things our foster brethren known to us from childhood to the things with which we are unacquainted foreign to our senses but in their immateriality and eternity more akin to our souls and above all to the reason which is in our souls.

[7] And likewise in Plato's *Republic* when the interlocutor of Socrates appears to bring certain plausible reasons to bear upon the mathematical sciences to show that they are useful to human life arithmetic for reckoning distributions contributions exchanges and partnerships geometry for sieges the founding of cities and sanctuaries and the partition

¹Cf. Aristotle *Meta* IV 15 1070^b 26

of land music for festivals entertainment and the worship of the gods and the doctrine of the spheres or astronomy, for farming navigation and other undertakings, revealing beforehand the proper procedure and suitable season Socrates, reproaching him, says You amuse me because you seem to fear that these are useless studies that I recommend but that is very difficult nay impossible For the eye of the soul blinded and buried by other pursuits is rekindled and aroused again by these and these alone and it is better that this be saved than thousands of bodily eyes for by it alone is the truth of the universe beheld¹

CHAPTER IV

[1] Which then of these four methods must we first learn? Evidently the one which naturally exists before them all is superior and takes the place of origin and root and as it were of mother to the others [2] And thus is arithmetic² not solely because we said that it existed before all the others in the mind of the creating God like some universal and exemplary plan relying upon which as a design and archetypal example the creator of the universe sets in order his material creations and makes them attain to their proper ends but also because it is naturally prior in birth inasmuch as it abolishes other sciences with itself³ but is not abolished together with them For example animal is naturally antecedent to man for abolish an animal and man is abolished but if man be abolished it no longer follows that animal is abolished at the same time And again man is antecedent to schoolteacher for if man does not exist neither does 'schoolteacher' but if schoolteacher is nonexistent it is still possible for man to be Thus since it has the property of abolishing the other ideas with it self it is likewise the older

[3] Conversely that is called younger and posterior which implies the other thing with itself⁴ but is not implied by it like musician for this always implies man Again take horse animal is always implied along with horse but not the reverse for if animal exists it is not necessary that horse should exist nor if man exists must musician also be implied

¹Republic 527 ff

²Plato Rep 522

³Cf below II 22 3 Cf Aristotle, *Met* 1019^a 1 ff

⁴Cf Aristotle e.g. *Top* VI 6 144^b 17 also *Top* II 4 111 25 ff

[4] So it is with the foregoing sciences if geometry exists arithmetic must also needs be implied for it is with the help of this latter that we can speak of triangle quadrilateral octahedron, icosahedron double eightfold or one and one-half times⁵ or anything else of the sort which is used as a term by geometry, and such things cannot be conceived of without the numbers that are implied with each one For how can 'triple' exist, or be spoken of unless the number 3 exists beforehand or eightfold without 8? But on the contrary 3 4 and the rest might be without the figures existing to which they give names [5] Hence arithmetic abolishes geometry along with itself but is not abolished by it and while it is implied by geometry it does not itself imply geometry

CHAPTER V

[1] And once more is this true in the case of music not only because the absolute is prior to the relative as great to greater and rich to richer and man to 'father' but also because the musical harmonies diatessaron diapente and diapason are named for numbers similarly all of their harmonic ratios are arithmetical ones for the diatessaron is the ratio of 4 3 the diapente that of 3 2 and the diapason the double ratio and the most perfect the didiapason is the quadruple ratio

[2] More evidently still astronomy attains through arithmetic the investigations that pertain to it not alone because it is later than geometry⁶ in origin—for motion naturally comes after rest—nor because the motions of the stars have a perfectly melodious harmony but also because risings settings progressions retrogressions increase⁷ and all sorts of phases are governed by numerical cycles and quantities

[3] So then we have rightly undertaken first the systematic treatment of this as the science naturally prior more honorable and more venerable and as it were mother and nurse of the rest and here we will take our start for the sake of clearness

CHAPTER VI

[1] All that has by nature with systematic method been arranged in the universe seems both in part and as a whole to have been determined and ordered in accordance with number by the forethought and the mind of him that

⁵Cf Plato *Rep* 528

created all things for the pattern was fixed like a preliminary sketch by the domination of number preexistent in the mind of the world creating God number conceptual only and immaterial in every way, but at the same time the true and the eternal essence so that with reference to it as to an artistic plan should be created all these things time motion the heavens the stars all sorts of revolutions

[2] It must needs be then that scientific number being set over such things as these should be harmoniously constituted in accordance with itself not by any other but by itself

[3] Everything that is harmoniously constituted is knit together out of opposites and of course out of real things for neither can non-existent things be set in harmony nor can things that exist but are like one another nor yet things that are different but have no relation one to another It remains accordingly that those things out of which a harmony is made are both real different and things with some relation to one another

[4] Of such things therefore scientific number consists for the most fundamental species in it are two embracing the essence of quantity different from one another and not of a wholly different genus odd and even and they are reciprocally woven into harmony with each other inseparably and uniformly, by a wonderful and divine Nature as straightway we shall see

CHAPTER VII

[1] Number is limited multitude or a combination of units or a flow of quantity made up of units and the first division of number is even and odd

[2] The even¹ is that which can be divided into two equal parts without a unit intervening in the middle and the odd is that which cannot be divided into two equal parts because of the aforesaid intervention of a unit

[3] Now this is the definition after the ordinary conception by the Pythagorean doctrine however the even number is that which admits of division into the greatest and the smallest parts at the same operation greatest in size and smallest in quantity in accordance with the natural contrariety of these two genera and the odd is that which does not allow this to be done to it but is divided into two unequal parts

[4] In still another way by the ancient defini-

tion the even is that which can be divided alike into two equal and two unequal parts except that the dyad which is its elementary form admits but one division that into equal parts and in any division whatsoever it brings to light only one species of number, however it may be divided, independent of the other The odd is a number which in any division whatsoever, which necessarily is a division into unequal parts shows both the two species of number together never without intermixture one with another but always in one another's company

[5] By the definition in terms of each other, the odd is that which differs by a unit from the even in either direction that is toward the greater or the less and the even is that which differs by a unit in either direction from the odd that is, is greater by a unit or less by a unit

CHAPTER VIII

[1] Every number is at once half the sum of the two on either side of itself and similarly half the sum of those next but one in either direction and of those next beyond them and so on as far as it is possible to go [2] Unity alone because it does not have two numbers on either side of it is half merely of the adjoining number hence unity is the natural starting point of all number

[3] By subdivision of the even there are the even times even the odd times even and the even times odd The even times even and the even times odd are opposite to one another like extremes and the odd times even is common to them both like a mean term

[4] Now the even times even² is a number which is itself capable of being divided into two equal parts in accordance with the properties of its genus and with each of its parts similarly capable of division and again in the same way each of their parts divisible into two equals until the division of the successive subdivisions reaches the naturally indivisible unit [5] Take for example 64 one half of this is 32 and of this 16 and of this the half is 8 and of this 4 and of this 2 and then finally unity is half of the latter and this is naturally indivisible and will not admit of a half

[6] It is a property of the even times even that whatever part of it be taken it is always

¹ Euclid's definition is The even times even number is that which is measured by an even number an even number of times *Elements VII Def 8*

even times even in designation and at the same time by the quantity of the units in it even times even in value and that neither of these two things will ever share in the other class [7] Doubtless it is because of this that it is called even times even because it is itself even and always has its parts and the parts of its parts down to unity even both in name and in value in other words every part that it has is even times even in name and even times even in value

[8] There is a method of producing the even times even so that none will escape but all successively fall under it if you do as follows [9] As you proceed from unity as from a root by the double ratio to infinity as many terms as there are will all be even times even and it is impossible to find others besides these for instance 1 2 4 8 16 32 64 128 256 512

[10] Now each of the numbers set forth was produced by the double ratio, beginning with unity and is in every respect even times even and every part that it may be found to have is always named from some one of the numbers before it in the series and the sum of units in this part is the same as one of the numbers before it, by a system of mutual correspondence indeed and interchange If there is an even number of terms of the double ratio from unity, not one mean term can be found but always two from which the correspondence and interchange of factors and values values and factors will proceed in order going first to the two on either side of the means then to the next on either side until it comes to the extreme terms so that the whole will correspond in value to unity and unity to the whole For example if we set down 128 as the largest term the number of terms will be even for there are eight in all up to this number and they will not have one mean term for this is impossible with an even number but of necessity two 8 and 16 These will correspond to each other as factors for of the whole 128 16 is one eighth and conversely 8 is one sixteenth Thence proceeding in either direction we find that 32 is one fourth and 4 one thirty second and again 64 is one half and 2 one sixty fourth and finally at the extremes unity is one one hundred twenty eighth and conversely 128 is the whole to correspond with unity

[11] If however the series consists of an odd number of terms seven for example and we deal with 64 there will be of necessity one mean term in accordance with the nature of the odd the mean term will correspond to it-

self because it has no partner and those on either side of it in turn will correspond to one another until this correspondence ends in the extremes Unity for example will be one sixty fourth and 64 the whole corresponding to unity 32 is one half and 2 one thirty-second 16 is one fourth and 4 one sixteenth and 8 the eighth part with nothing else to correspond to it

[12] It is the property of all these terms when they are added together successively to be equal to the next in the series lacking one unit so that of necessity their summation in any way whatsoever will be an odd number, for that which fails by a unit of being equal to an even number is odd [13] This observation will be of use to us very shortly in the construction of perfect numbers¹ But to take an example the terms from unity preceding 256 in the series when added together, are within 1 of equaling 256 and all the terms before 128 the term immediately preceding are similarly equal to 128 save for one unit and to the next terms the sums of those below them are similarly related Thus unity itself² is within one unit of equaling the next term which is 2 and these two together fail by 1 of equaling the next and the three together are within 1 of the next in order and you will find that this goes on with out interruption to infinity

[14] This too it is very needful to recall If the number of terms of the even times even series dealt with is even the product of the extremes will always be equal to the product of the means if there is an odd number of terms the product of the extremes will be equal to the square of the mean For in the case of an even number of terms 1 times 128 is equal to 8 times 16 and further to 2 times 64 and again to 4 times 32 and this is so in every case and with an odd number of terms 1 times 64 equals 2 times 32 and this equals 4 times 16 and this again equals 8 times 8 the mean term alone multiplied by itself

CHAPTER IX

[1] The even times odd number is one which is by its genus itself even but is specifically opposed to the aforesaid even times even It is a number of which though it admits of the division into two equal halves after the fashion the genus common to it and the even the halves are not

¹See Chapter 16

²Cf on I 8 7

into two equals for example 6 10 14 18 22 26 and the like for after these have been divided their halves are found to be indivisible

[2] It is the property of the even times odd that whatever factor it may be discovered to have is opposite in name to its value and that the quantity of every part is opposite in value to its name and that the numerical value of its part never by any means ¹ of the same genus as its name To take a single example the number 18 its half with an even name is 9 odd in value its third part again with an odd designation is 6 even in value conversely, the sixth part is 3 and the ninth part 2 and in other numbers the same peculiarity will be found

[3] It is possibly for this reason that it received such a name that is because although it is even its halves are at once odd

[4] This number is produced from the series beginning with unity with a difference of 2, namely the odd numbers set forth in proper order as far as you like and then multiplied by 2 The numbers produced would be in order the 6 10 14 18 22 26 30 and so on as far as you care to proceed The greater terms all ways differ by 4 from the next smaller ones the reason for which is that their original basic forms the odd numbers exceed one another by 2 and were multiplied by 2 to make this series and 2 times 2 makes 4

[5] Accordingly in the natural series of numbers the even times odd numbers will be found fifth from one another exceeding one another by a difference of 4 passing over three terms and produced by the multiplication of the odd numbers by 2

[6] They are said to be opposite in properties to the even times even because of these the greatest extreme term alone is divisible while of these former the smallest only proved to be indivisible and in particular because in the former case the reciprocal arrangement of parts¹ from extremes to mean term or terms makes the product of the former equal to the square or product of the latter but in this case by the same correspondence and comparison the mean term is one half the sum of the extremes or if there should be two means their sum equals that of the two extremes

CHAPTER X

[1] The odd times even number is the one which displays the third form of the even belonging in common to both the previously men-

tioned species like a single mean between two extremes for in one respect it resembles the even times even and in another the even times odd and that property wherein it varies from the one it shares with the other, and by that property which it shares with the one it differs from the other

[2] The odd times even number is an even number which can be divided into two equal parts whose parts also can so be divided and sometimes even the parts of its parts but it cannot carry the division of its parts as far as unity Such numbers are 24 28 40 for each of these has its own half and indeed the half of its half and sometimes one is found among them that will allow the halving to be carried even farther among its parts There is none however that will have its parts divisible into halves as far as the naturally indivisible unit

[3] Now in admitting more than one division the odd times even is like the even times even and unlike the even times odd but in that its subdivision never ends with unity it is like the even times odd and unlike the even times even

[4] It alone has at once the proper qualities of each of the former two² and then again properties which belong to neither of them for of them one had only the highest term divisible, and the other only the smallest indivisible but this neither for it is observed to have more divisions than one in the greater term, and more than one indivisible in the lesser

[5] Furthermore there are in it certain parts whose names are not opposed to their values nor of the opposite genus³ after the fashion of the even times even and there are also always other parts of a name opposite and contrary in kind to their values after the fashion of the even times odd For example in 24 there are parts not opposed in name to their values the fourth part 6 the half 12 the sixth 4 and the twelfth 2 but the third part 8 the eighth 3 and the twenty fourth 1 are opposed and so it is with the rest

[6] This number is produced by a somewhat complicated method and shows after a fashion even in its manner of production that it is a mixture of both other kinds ¹ or whereas the even times even is made from even numbers the doubles from unity to infinity and the even times odd from the odd numbers from 3 progressing to infinity this must be woven

¹Cf I 9 6

²Cf I 8 7, 9 2

³Cf I 8 10

together out of both classes as being common to both [] Let us then set forth the odd numbers from 3 by themselves in due order in one series

3 5 7 9 11 13 15 17 19

and the even times even, beginning with 4 again one after another in a second series after their own order

4 8 16 32 64 128 256

as far as you please [8] Now multiply by the first number of either series—it makes no difference which—from the beginning and in order all those in the remaining series and note down the resulting numbers then again multiply by the second number of the same series the same numbers once more as far as you can and write down the results then with the third number again multiply the same terms anew and how ever far you go you will get nothing but the odd times even numbers

[9] For the sake of illustration let us use the first term of the series of odd numbers and multiply by it all the terms in the second series in order thus 3×4 3×8 3×16 3×32 and so on to infinity The results will be 12 24 48 96 which we must note down in one line Then taking a new start do the same thing with the second number 5×4 5×8 5×16 5×32 The results will be 20 40 80 160 Then do the same thing once more with 7 the third number 7×4 7×8 7×16 7×32 The results are 28 56

term or their product should there be two Thus this one species has the peculiar properties of them both because it is a natural mixture of them both

CHAPTER VI

[1] Again while the odd is distinguished over against the even in classification and has nothing in common with it since the latter is divisible into equal halves and the former is not thus divisible nevertheless there are found three species of the odd¹ differing from one another of which the first is called the prime and incomposite² that which is opposed to it the secondary and composite and that which is midway between both of these and is viewed as a mean among extremes namely the variety which in itself is secondary and composite, but relatively is prime and incomposite

[2] Now the first species the prime and incomposite is found whenever an odd number admits of no other factor save the one with the number itself as denominator which is always unity for example 3, 5, 7, 11, 13, 17, 19, 23, 29, 31 None of these numbers will by any chance be found to have a fractional part with a denominator different from the number itself but only the one with this as denominator and this part will be unity in each case for 3 has only a third part, which has the same denominator as the number and is of course uni-

Odd numbers	3	5	7	9	11	13	15
Even times even	4	8	16	32	64	128	256
Odd times even numbers	12	24	48	96	192	384	768
	20	40	80	160	320	640	1280
	28	56	112	224	448	896	1792
	36	72	144	288	576	1152	2304
	44	88	176	352	704	1408	2816

Length

112 224 and in the same way as far as you care to go you will get similar results

[10] Now when you arrange the products of multiplication by each term in its proper line making the lines parallel in marvelous fashion there will appear along the breadth of the table the peculiar property of the even times odd that the mean term is always half the sum of the extremes if there should be one mean and the sum of the means equals the sum of the extremes if two But along the length of the table the property of the even times even will appear for the product of the extremes is equal to the square of the mean should there be one mean

ty 5 a fifth 7 a seventh and 11 only an eleventh part and in all of them these parts are unity

[3] It has received this name because it can be measured only by the number which is first and common to all unity and by no other moreover because it is produced by no other number combined with itself save unity alone for 5 is 5×1 and 7 is 7×1 and the others in accordance with their own quantity To be sure when they are combined with themselves other

¹Cf Euclid Elem VII D ff 11 14

²Cf Euclid Elem VII Def 11 Aristotle VIII 2 157 39

numbers might be produced originating from them as from a fountain and a root wherefore they are called 'prime' because they exist beforehand as the beginnings of the others. For every origin is elementary and incomposite in to which every thing is resolved and out of which everything is made but the origin itself cannot be resolved into anything or constituted out of anything

CHAPTER XII

[1] The secondary composite number¹ is an odd number indeed because it is distinguished as a member of this same class but it has no elementary quality, for it gets its origin by the combination of something else. For this reason it is characteristic of the secondary number to have in addition to the fractional part with the number itself as denominator yet another part or parts with different denominators the former always as in all cases unity the latter never unity but always either that number or those numbers by the combination of which it was produced. For example 9 15 21 25 27 33 39 each one of these is measured by unity as other numbers are and like them has a fractional part with the same denominator as the number itself by the nature of the class common to them all but by exception and more peculiarly they also employ a part or parts with a different denominator 9 in addition to the ninth part has a third part besides 15 a third and a fifth besides a fifteenth 21 a seventh and a third besides a twenty first and 25 in addition to the twenty fifth which has as a denominator 25 itself also a fifth with a different denominator

[2] It is called secondary then because it can employ yet another measure along with unity and because it is not elementary but is produced by some other number combined with itself or with something else in the case of 9 3 in the case of 15 5 or by Zeus 3 and those following in the same fashion. And it is called composite for this or some such reason that it may be resolved into those numbers out of which it was made since it can also be measured by them. For nothing that can be broken down is incomposite but by all means composite

CHAPTER XIII

[1] Now while these two species of the odd are opposed to each other a third one² is con-

ceived of between them deriving as it were its specific form from them both namely the number which is in itself secondary and composite but relatively to another number is prime and incomposite. This exists when a number, in addition to the common measure unity is measured by some other number and is therefore able to admit of a fractional part or parts with denominator other than the number itself as well as the one with itself as denominator. When this is compared with another number of similar properties it is found that it cannot be measured by a measure common to the other nor does it have a fractional part with the same denominator as those in the other. As an illustration let 9 be compared with 25. Each in itself is secondary and composite but relatively to each other they have only unity as a common measure and no factors in them have the same denominator for the third part in the former does not exist in the latter nor is the fifth part in the latter found in the former

[2] The production of these numbers is called by Eratosthenes the sieve because we take the odd numbers mingled together and indiscriminate and out of them by this method of production separate as by a kind of instrument or sieve the prime and incomposite by themselves and the secondary and composite by themselves and find the mixed class by themselves

[3] The method of the sieve is as follows. I set forth all the odd numbers in order beginning with 3 in as long a series as possible and then starting with the first I observe what ones it can measure and I find that it can measure the terms two places apart as far as we care to proceed. And I find that it measures not as it chances and at random but that it will measure the first one that is the one two places removed by the quantity of the one that stands first in the series that is by its own quantity for it measures it 3 times and the one two places from this by the quantity of the second in order for this it will measure 5 times and again the one two places further on by the quantity of the third in order or 7 times and the one two places still farther on by the quantity of the fourth in order or 9 times and so *ad infinitum* in the same way

[4] Then taking a fresh start I come to the second number and observe what it can measure and find that it measures all the terms four places apart the first by the quantity of the first in order or 3 times the second by that of the second or 5 times the third by that of

¹Cf Euclid *Elements* VII Def 14

²Cf Euclid *Elements* VII Def 13

the third or 7 times and in this order *ad infinitum*

[5] Again as before the third term 7 taking over the measuring function will measure terms six places apart and the first by the quantity of 3 the first of the series the second by that of 5 for this is the second number and the third by that of 7, for this has the third position in the series

[6] And analogously throughout this process will go on without interruption so that the numbers will succeed to the measuring function in accordance with their fixed position in the series the interval separating terms measured is determined by the orderly progress of the even numbers from 2 to infinity or by the doubling of the position in the series occupied by the measuring term and the number of times a term is measured is fixed by the orderly advance of the odd numbers in series from 3

[7] Now if you mark the numbers with certain signs you will find that the terms which succeed one another in the measuring function neither measure all the same number—and sometimes not even two will measure the same one—nor do absolutely all of the numbers set forth submit themselves to a measure but some entirely avoid being measured by any number whatsoever some are measured by one only and some by two or even more [8] Now these that are not measured at all but avoid it are primes and incomposite sifted out as it were by a sieve those measured by only one measure in accordance with its own quantity will have but one fractional part with denominator different from the number itself in addition to the part with the same denominator and those which are measured by one measure only but in accordance with the quantity of some other number than the measure and not its own, or are measured by two measures at the same time will have several fractional parts with other denominators besides the one with the same as the number itself these will be secondary and composite

[9] The third division the one common to both the former which is in itself secondary and composite but primary and incomposite in relation to another will consist of the numbers produced when some prime and incomposite number measures them in accordance with its own quantity if one thus produced be compared to another of similar origin For example if 9 which was produced by 3 measuring by its own quantity for it is 3 times 3 be compared with 25 which was produced from 5 measuring by

its own quantity for it is 5 times 5 these numbers have no common measure except unity

[10] We shall now investigate how we may have a method of discerning whether numbers are prime and incomposite or secondary and composite, relatively to each other since of the former unity is the common measure but of the latter some other number also besides unity and what this number is

[11] Suppose there be given us two odd numbers and some one sets the problem and directs us to determine whether they are prime and incomposite relatively to each other or secondary and composite and if they are secondary and composite what number is their common measure We must compare the given numbers and subtract the smaller from the larger as many times as possible then after this subtraction subtract in turn from the other as many times as possible for this changing about and subtraction from one and the other in turn will necessarily end either in unity or in some one and the same number which will necessarily be odd [12] Now when the subtractions terminate in unity they show that the numbers are prime and incomposite relatively to each other and when they end in some other number odd in quantity and twice produced then say that they are secondary and composite relatively to each other and that their common measure is that very number which twice appears

For example if the given numbers were 23 and 45 subtract 23 from 45 and 22 will be the remainder subtracting this from 23 the remainder is 1 subtracting this from 22 as many times as possible you will end with unity Hence they are prime and incomposite to one another and unity which is the remainder is their common measure

[13] But if one should propose other numbers 21 and 49 I subtract the smaller from the larger and 28 is the remainder Then again I subtract the same 21 from this for it can be done and the remainder is 7 Thus I subtract in turn from 21 and 14 remains from which I subtract 7 again for it is possible and 7 will remain But it is not possible to subtract 7 from 7 hence the termination of the process with a repeated 7 has been brought about and you may declare the original numbers 21 and 49 secondary and composite relatively to each

This mode of determining common factors is found in Euclid (VII 1 & 2) and is commonly termed the Euclidean method of finding the greatest common divisor of numbers

other and 7 their common measure in addition to the universal unit

CHAPTER XIV

[1] To make again a fresh start of the simple even numbers some are superabundant some deficient like extremes set over against each other, and some are intermediary between them and are called perfect [2] Those which are said to be opposites to one another the superabundant and deficient are distinguished from one another in the relation of inequality¹ in the directions of the greater and the less for apart from these no other form of inequality could be conceived nor could evil² disease disproportion unseemliness nor any such thing save in terms of excess or deficiency I or in the realm of the greater³ there arise excesses overreaching and superabundance and in the less need deficiency privation and lack but in that which lies between the greater and the less, namely the equal are virtues wealth moderation propriety beauty and the like to which the aforesaid form of number the perfect is most akin

[3] Now the superabundant number is one which has over and above the factors which belong to it and fall to its share others in addition just as if an animal should be created with too many parts or limbs with ten tongues as the poet says⁴ and ten mouths or with nine lips or three rows of teeth or a hundred hands or too many fingers on one hand Similarly if when all the factors in a number are examined and added together in one sum it proves upon investigation that the number's own factors exceed the number itself this is called a superabundant number for it oversteps the symmetry which exists between the perfect and its own parts Such are 12 24 and certain others for 12 has a half 6 a third 4 a fourth 3 a sixth 2 and a twelfth 1 which added together make 16 which is more than the original 12 its parts therefore are greater than the whole itself [4] And 24 has a half a third fourth sixth eighth twelfth and twenty fourth, which are 12 8 6 4 3 2 1 Added together they make 36 which compared to the original number 24 is found to be greater than it, although made up solely of its factors Hence in this case also the parts are in excess of the whole

¹Cf I 17 2 4 6 also I 23 4

²Cf Arist *Ethics* II 6 1100^b 33

³Cf Arist *Ethics* II 6 1100^b 24 33

⁴Homer *Odyssey* XII 85 ff

CHAPTER XV

[1] The deficient number is one which has qualities the opposite of those pointed out and whose factors added together are less in comparison than the number itself It is as if some animal should fall short of the natural number of limbs or parts or as if a man should have but one eye as in the poem "And one round orb was fixed in his brow", or as though one should be one-handed or have fewer than five fingers on one hand or lack a tongue or some such member Such a one would be called deficient and so to speak maimed after the peculiar fashion of the number whose factors are less than itself such as 8 or 14 For 8 has the factors half fourth and eighth which are 4, 2, and 1 and added together they make 7 and less than the original number The parts therefore fall short of making up the whole [2] Again 14 has a half a seventh a fourteenth, 7 2 and 1 respectively and all together they make 10, less than the original number So this number also is deficient in its parts with respect to making up the whole out of them

CHAPTER XVI

[1] While these two varieties are opposed after the manner of extremes the so-called perfect number⁵ appears as a mean, which is discovered to be in the realm of equality and neither makes its parts greater than itself, added together nor shows itself greater than its parts but is always equal to its own parts For the equal is always conceived of as in the mid-ground between greater and less and is as it were moderation between excess and deficiency and that which is in tune between pitches too high and too low

[2] Now when a number comparing with itself the sum and combination of all the factors whose presence it will admit neither exceeds them in multitude nor is exceeded by them then such a number is properly said to be perfect as one which is equal to its own parts Such numbers are 6 and 28 for 6 has the factors half third and sixth 3 2 and 1, respectively and these added together make 6 and are equal to the original number and neither more nor less Twenty-eight has the factors half fourth seventh fourteenth and twenty-eighth which are 14 7 4 2 and 1 these added together make 28 and so neither are the

⁵Fuelid's definition *Flem* V II 22 is "A perfect number is one that is equal to its own parts."

parts greater than the whole nor the whole greater than the parts but their comparison is in equality which is the peculiar quality of the perfect number

[5] It comes about that even as fair and excellent things are few and easily enumerated while ugly and evil ones are widespread so also the superabundant and deficient numbers are found in great multitude and irregularly placed—for the method of their discovery is irregular—but the perfect numbers are easily enumerated and arranged with suitable order for only one is found among the units 6 only one other among the tens 28 and a third in the rank of the hundreds 496 alone and a fourth within the limits of the thousands that is below ten thousand 8128 And it is their accompanying characteristic to end alternately in 6 or 8 and always to be even

[4] There is a method of producing them neat and unailing which neither passes by any of the perfect numbers nor fails to differentiate any of those that are not such which is carried out in the following way

You must set forth the even times even numbers from unity advancing in order in one line as far as you please 1 2 4 8 16 32 64 128 2 6 512 1024 2048 4096 Then you must add them together one at a time and each time you make a summation observe the result to see what it is If you find that it is a prime incomposite number multiply it by the quantity of the last number added and the result will always be a perfect number If however the result is secondary and composite do not multiply but add the next and observe again what the resulting number is if it is secondary and composite again pass it by and do not multiply but add the next but if it is prime and incomposite multiply it by the last term added and the result will be a perfect number and so on to infinity In similar fashion you will produce all the perfect numbers in succession overlooking none

For example to 1 I add 2 and observe the sum and find that it is 3 a prime and incomposite number in accordance with our previous demonstrations for it has no factor with denominator different from the number itself but only that with denominator agreeing Therefore I multiply it by the last number to be taken into the sum that is 2 I get 6 and thus I declare to be the first perfect number in actuality and to have those parts which are beheld

in the numbers of which it is composed For it will have unity as the factor with denominator the same as itself, that is its sixth part and 3 as the half which is seen in 2 and conversely 2 as its third part

[6] Twenty-eight likewise is produced by the same method when another number 4 is added to the previous ones For the sum of the three 1 2 and 4 is 7, and is found to be prime and incomposite for it admits only the factor with denominator like itself the seventh part Therefore I multiply it by the quantity of the term last taken into the summation and my result is 28, equal to its own parts and having its factors derived from the numbers already adduced a half corresponding to 2 a fourth to 7 a seventh to 4 a fourteenth to offset the half and a twenty-eighth in accordance with its own nomenclature which is 1 in all numbers

[6] When these have been discovered, 6 among the units and 28 in the tens you must do the same to fashion the next [7] Again add the next number 8 and the sum is 15 Observing this I find that we no longer have a prime and incomposite number but in addition to the factor with denominator like the number itself it has also a fifth and a third with unlike denominators Hence I do not multiply it by 8 but add the next number 16 and 31 results As this is a prime incomposite number of necessity it will be multiplied in accordance with the general rule of the process by the last number added 16 and the result is 496 in the hundred and then comes 8128 in the thousands and so on as far as it is convenient for one to follow

[8] Now unity is potentially a perfect number but not actually for taking it from the series as the very first I observe what sort it is according to the rule and find it prime and incomposite for it is so in very truth not by participation like the rest but it is the primary number of all and alone incomposite [9] I multiply it therefore by the last term taken into the summation that is by itself and my result is 1 for 1 times 1 equals 1 [10] Thus unity is perfect potentially for it is potentially equal to its own parts the others actually

CHAPTER XVII

[1] Now that we have given a preliminary systematic account of absolute quantity we come in turn to relative quantity

[2] Of relative quantity then the highest general divisions are two equality and inequality

[1] Cf. Euclid (IX 36)

ity for everything viewed in comparison with another thing is either equal or unequal, and there is no third thing besides these

[5] Now the equal is seen when of the things compared one neither exceeds nor falls short in comparison with the other for example 100 compared with 100 10 with 10 2 with 2, a mina with a mina a talent with a talent, a cubit with a cubit and the like, either in bulk length weight or any kind of quantity [4] And as a peculiar characteristic also this relation is of itself not to be divided or separated, as being most elementary, for it admits of no difference For there is no such thing as this kind of equality and that kind, but the equal exists in one and the same manner [6] And that which corresponds to an equal thing to be sure, does not have a different name from it but the same like friend neighbor comrade" so also equal for it is equal to an equal

[6] The unequal on the other hand is split up by subdivisions and one part of it is the greater the other the less which have opposite names and are antithetical to one another in their quantity and relation For the greater is greater than some other thing and the less again is less than another thing in comparison, and their names are not the same but they each have different ones for example father and son striker and struck, teacher and pupil and the like

[7] Moreover of the greater separated by a second subdivision into five species one kind is the multiple another the superparticular another the superpartient another the multiple superparticular and another the multiple superpartient [8] And of its opposite the less there arise similarly by subdivision five species opposed to the foregoing five varieties of the greater the submultiple subsuperparticular subsuperpartient submultiple superparticular and submultiple-superpartient for as whole answers to whole smaller to greater so also the varieties correspond each to each in the aforesaid order with the prefix sub-

CHAPTER XVIII

[1] Once more then the multiple is the species of the greater first and most original by nature as straightway we shall see and it is a number which when it is observed in comparison with another contains the whole of that number more than once For example compared with unity all the successive numbers beginning with 2 generate in their proper order the regular forms of the multiple for 2 in the

first place is and is called the double 3 triple, 4 quadruple and so on for 'more than once' means twice, or three times, and so on in succession as far as you like

[2] Answering to this is the submultiple which is itself primary in the smaller division of inequality It is the number which when it is compared with a larger is able to measure it completely more than once and more than once starts with twice and goes on to infinity [3] If then it measures the larger number that is being compared twice only, it is properly called the subdouble as 1 is of 2 if thrice subtriple, as 1 of 3 if four times subquadruple as 1 of 4 and so on in succession

[4] While each of these the multiple and the submultiple is generically infinite the varieties by subdivision and the species also are observed naturally to make an infinite series For the double beginning with 2 goes on through all the even numbers as we select alternate numbers out of the natural series and these will be called doubles in comparison with the even and odd numbers successively placed beginning with unity [5] All the numbers from the beginning two places apart and third in order are triples for example 3 6 9 12 15, 18 21 24 It is their property to be alternately odd and even and they themselves in the regular series from unity are triples of all the numbers in succession as far as one wishes to go on with the process

[6] The quadruples are those in the fourth places three apart for instance 4 8 12 16, 20, 24 28 32 and so on These are the quadruples of the regular series of numbers from unity going on as far as one finds it convenient to follow It belongs to them all to be even, for one needs only to take the alternate terms out of the even numbers already selected Thus necessarily it is true that the even numbers with no further designation are all doubles the alternate ones quadruples those two places apart sextuples and those three places apart octuples and this series will go on on this same analogy indefinitely

[7] The quintuples will be seen to be those four places apart placed fifth from one another and them elses the quintuples of the successive numbers beginning with unity Alternately they are odd and even like the triples

CHAPTER XIX

[1] The superparticular the second species of the greater both naturally and in order, is a

number that contains within itself the whole of the number compared with it and some one factor of it besides

[9] If this factor is a half the greater of the terms compared is called specifically *sesquialter* and the smaller *subsesquialter* if it is a third *sesquitercian* and *subsesquitercian* and as you go on throughout it will always thus agree so that these species also will progress to infinity even though they are species of an unlimited genus

For it comes about that the first species the *sesquialter* ratio has as its consequents the even numbers in succession from 2 and no other at all and as antecedents the triples in succession from 3 and no other [8] These must be joined together regularly first to first, second to second third to third—3 2 6 4 9 6 12 8—and the analogous numbers to the ones corresponding to them in position

[4] If we care to investigate the second species of the superparticular the *sesquitercian* (for the fraction naturally following after the half is the third) we shall have this definition of it—a number which contains the whole of the number compared and a third of it in addition to the whole We may have examples of it in the proper order in the successive quadruples beginning with 4 joined to the triples from 3 each term with the one in the corresponding position in the series for example, 4 3 8 6 12 9 and so on to infinity [5] It is plain that that which corresponds to the *sesquialter* but is called with the prefix *sub-* *subsesquialter* is the number the whole of which is contained and a third part in addition for example 3 4 6 8 9 12 and the similar pairs of numbers in the same position in the series

[6] And we must observe the never failing corollary of all this that the first forms in each series the so-called root numbers are next to one another in the natural series the next after the root forms show an interval of only one number the third two the fourth three the fifth four and so on as far as you like [7] Furthermore that the fraction after which each of the superparticulars is named is seen in the lesser of the root numbers never in the greater

[8] That by nature and by no disposition of ours the multiple is a more elementary and an older form than the superparticular we shall shortly learn through a somewhat intricate process And here for a simple demonstration we must prepare in regular and parallel lines the multiples specified above according to their

varieties first the double in one line then in a second the triple then the quadruple in a third and so on as far as the tenfold multiples so that we may detect their order and variety, their regulated progress and which of them is naturally prior and indeed other corollaries delightful in their exactness Let the diagram be as follows [9]

1	2	3	4	5	6	7	8	9	10
2	4	6	8	10	12	14	16	18	20
3	6	9	12	15	18	21	24	27	30
4	8	12	16	20	24	28	32	36	40
5	10	15	20	25	30	35	40	45	50
6	12	18	24	30	36	42	48	54	60
7	14	21	28	35	42	49	56	63	70
8	16	24	32	40	48	56	64	72	80
9	18	27	36	45	54	63	72	81	90
10	20	30	40	50	60	70	80	90	100

[10] Let there be set forth in the first row the natural series from unity and then in order those species of the multiple which we were bidden to insert

[11] Now then in comparison with the first rows beginning with unity if we read both across and up and down in the form of the letter gamma the next rows both ways themselves in the form of a gamma beginning with 4 are multiples according to the first form of the multiple for they are doubles The first differs by unity from the first the second from the second by 2 the third from the third by 3 the next by 4 those following by 5 and you will find that this follows throughout

The third rows in both directions from 9, their common origin will be the triples of the terms in that same first row according to the second form of the multiple the cross lines like the letter chi ending in the term 3 in either direction are to be taken into consideration [12] The difference for these numbers will progress after the series of the even numbers being 2 for the first 4 for the next 6 for the third and this difference Nature has of her own accord interpolated for us between these rows that are being examined as is evident in the diagram

[13] The fourth row whose common origin in both directions is 16 and whose cross-lines end with the terms 4 exhibits the third species of multiple the quadruple when it is compared with that same first row according to corresponding positions first term with first second with second third with third and so on Again the differences of these numbers are 3 6 then 9 then 12 and the quantities that progress by steps of 3 These numbers are detected in the

structure of the diagram in places just above the quadruples and in the subsequent forms of the multiple the analogy will hold throughout.

[14] In comparison with the second line reading either way which begins with the common origin 4 and runs over in cross lines to the term 2 in each row, the lines which are next in order beneath display the first species of the superparticular that is, the sesquialter between terms occupying corresponding places. Thus by divine nature not by our convention or agreement the superparticulars are of later origin than the multiples. For illustration 3 is the sesquialter of 2 6 of 4 9 of 6 12 of 8 15 of 10 and throughout thus. They have as a difference the successive numbers from unity like those before them.

[15] The sesquiterterians the second species of superparticular proceed with a regular even advance from 4 3 8 6 12 9 16 12 and so on having also a regular increase of their differences. [16] And in the other multiple and superparticular relations you will see that the results are in harmony and not by any means inconsistent as you go on to infinity.

[17] The following feature of the diagram moreover is of no less exactness. The terms at the corners are units the one at the beginning a simple unit that at the end the unit of the third course and the other two units of the second course appearing twice so that the product (of the first two) is equal to the square (of the last). [18] Furthermore in reading either way there is an even progress from unity to the tens and again on the opposite sides two other progressions from 10 to 100.

[19] The terms on the diagonal from 1 to 100 are all square numbers the products of equals by equals and those flanking them on either side are all heteromecic unequal and the products of sides of which one is greater than the other by unity and so the sum of two successive squares and twice the heteromecic numbers between them is always a square and conversely a square is always produced from the two heteromecic numbers on the sides and twice the square between them.

[20] An ambitious person might find many other pleasing things displayed in this diagram upon which it is not now the time to dwell for we have not yet gained recognition of them from our Introduction and so we must turn to the next subject. For after these two generic relations of the multiple and the superparticular and the other two opposite to them with the prefix sub- the submultiple and the sub-

superparticular there are in the greater division of inequality the superpartient and in the less its opposite the sub-superpartient.

CHAPTER XX

[1] It is the superpartient relation when a number contains within itself the whole of the number compared and in addition more than one part of it and "more than one" starts with 2 and goes on to all the numbers in succession. Thus the root-form of the superpartient is naturally the one which has in addition to the whole two parts of the number compared and as a species will be called superbipartient after this the one with three parts besides the whole will be called supertripartient as a species then comes the superquadrupartient the superquintupartient, and so forth.

[2] The parts have their root and origin with the third for it is impossible in this case to begin with the half. For if we assume that any number contains two halves of the compared number besides the whole of it we shall inadvertently be setting up a multiple instead of a superpartient because each whole plus two halves of it, added together makes double the original number. Thus it is most necessary to start with two thirds then two fifths two sevenths and after these two ninths, following the advance of the odd numbers for two quarters, for example again are a half, two sixths a third and thus again superparticulars will be produced instead of superpartients which is not the problem laid before us nor in accord with the systematic construction of our science. [3] After the superpartient the subsuperpartient immediately is produced whenever a number is completely contained in the one compared with it, and in addition several parts of it 2 3 4 or 5 and so on.

CHAPTER XXI

[1] The regular arrangement and orderly production of both species are discovered when we set forth the successive even and odd numbers beginning with 3 and compare with them simple series of odd numbers only from 5 in succession first to first—that is 5 to 3—second to second—that is 7 to 4—third to third—that is 9 to 5—fourth to fourth—that is 11 to 6—and so on in the same order as far as you like. In this way the forms of the superpartient and the subsuperpartient in due order will be disclosed through the root forms of each species the superbipartient first then the supertripartient superquadrupartient and super-

quartupartient and further in succession in similar manner, for after the root forms of each species the ones which follow them will be produced by doubling or tripling both the terms and in general by multiplying after the regular forms of the multiple

nerically, it will have in its subdivisions according to species a sort of diversification and change of names proper both to the first part of the name and to the second. For instance in the first part that is the multiple it will have double triple, quadruple quintuple, and

TABLE OF THE SUPERPARTIENTS

Root-forms	5	3	7	4	9	5	11	6	13	7
	10	6	14	8	18	10	22	12	26	14
	15	9	21	12	27	15	33	18	39	21
	20	12	28	16	36	20	44	24	52	28
	25	15	35	20	45	25	55	30	65	35
	30	18	42	24	54	30	66	36	78	42
	35	21	49	28	63	35	77	42	91	49
	40	24	56	32	72	40	88	48	104	56
	45	27	63	36	81	45	99	54	117	63

[2] It must be observed that from the two parts in addition to the whole which are contained in the greater term we are to understand third in the case of three parts, fourth with four parts fifth with five sixth and so on so that the order of nomenclature is something like this superbi-partient supertripartient superquadrupartient then u-perquintupartient and similarly with the rest

[3] Now the simple uncompounded relations of relative quantity are these which have been enumerated. Those which are compounded of them and as it were woven out of two into one are the following of which the antecedents are the multiple superparticular and multiple superpartient and the consequents the ones that immediately arise in connection with each of the former named with the prefix sub together with the multiple superparticular the submultiple superparticular and with the multiple superpartient the submultiple superpartient. In the subdivision of the genera the species of the one will correspond to those of the other these also having names with the prefix sub

CHAPTER XXII

[1] Now the multiple superparticular is a relation in which the greater of the compared terms contains within it self the lesser term more than once and in addition some one part of it whatever this may be

[2] As a compound such a number is doubly diversified after the peculiarities of nomenclature of its components on either side for inasmuch as the multiple superparticular is compounded of the multiple and superparticular ge-

nerically, it will have in its subdivisions according to species a sort of diversification and change of names proper both to the first part of the name and to the second. For instance in the first part that is the multiple it will have double triple, quadruple quintuple, and

so forth and in the second part generically from the superparticular, its specific forms in due order the sesquialter sesquitercian sesquiquartan sesquiquintan and so on so that the combination will proceed in somewhat this order

Double sesquialter double sesquitercian, double sesquiquartan double sesquiquintan, double sesquiseptan and analogously

Beginning once more triple sesquialter triple sesquitercian triple sesquiquartan triple sesquiquintan

Again quadruple sesquialter quadruple sesquitercian quadruple sesquiquartan quadruple sesquiquintan and the forms analogous to these *ad infinitum*. Whatever number of times the greater contains the whole of the smaller by this quantity the first part of the ratio of the terms joined together in the multiple superparticular is named and whatever may be the factor in addition to the whole several times contained that is in the greater term from this is named the second kind of ratio of which the multiple superparticular is compounded

[3] Examples of it are these 5 is the double sesquialter of 2 7 the double sesquitercian of 3 9 the double sesquiquartan of 4 11 the double sesquiquintan of 5 You will furthermore always produce them in regular order in this fashion by comparing with the successive even and odd numbers from 2 the odd numbers, exclusively from 5 first with first second third with third and the with the one in the same position

The successive terms beginning with 5 and differing by 5 will be without exception double sesquialters of all the successive even numbers from 2 on, when terms in the same position in the series are compared and beginning with 3, if all those with a difference of 3 be set forth as 3 6 9 12 15 18 21 and in another series there be set forth those that differ by 7 to infinity as 7 14 21 28 35 42 49 and the greater be compared with the smaller first to first, second to second third to third fourth to fourth and so on the second species will appear the double sesquitercian disposed in its proper order

[4] Then again, to take a fresh start if the simple series of quadruples be set forth, 4 8 12 16 20 24 28 32 and then there be placed beside it in another series the successive numbers beginning with 9 and increasing by 9 as 9 18 27 36 45 54 we shall have revealed once more the multiple superparticular in a specific form that is the double sesquiquartan in its proper order and any one who desires can contrive this to an unlimited extent

[5] The second kind begins with the triple sesquialter such as 7 2 14 4 and in general the numbers that advance by steps of 7 compared with the even numbers in order from 2

[6] Then once more 10 3 is the first triplessesquitercian 20 6 the second and in a word the multiples of 10 in succession compared with the successive triples. This indeed we can observe with greater exactitude and clearness in the table studied above for in comparison with the first row the succeeding rows in order¹ compared as whole rows display the forms of the multiple in regular order up to infinity when they are all compared in each case to the same first row and when each row is compared to all those above it in succession the second row being taken as our starting point all the forms of the superparticular are produced in their proper order and if we start with the third row all of those beginning with the fifth that are odd in the series when they are compared with this same third row and those following it will show all the forms of the superpartient in proper order. In the case of the multiple superparticular the comparisons will have a natural order of their own if we start with the second row and compare the terms from the fifth first to first, second to second third to third and so on and then the terms of the seventh row to the third, those of the ninth to the

fourth and follow the corresponding order as far as we are able to go [7] It is plain that here too the smaller terms have names corresponding to the larger ones with the prefix sub- according to the nomenclature given them all

CHAPTER XXIII

[1] The multiple superpartient is the remaining relation of number. This and the relation called by a corresponding name with the prefix sub-, exist when a number contains the whole of the number compared more than once (that is twice, thrice or any number of times) and certain parts of it more than one either two three or four and so on, besides.^[2] These parts² are not halves for the reasons mentioned above but either thirds fourths or fifths and so on

[3] From what has already been said it is not hard to conceive of the varieties of this relation for they are differentiated in the same way as and consistently with those that precede double superbi-partient, double supertripartient double superquadrupartient and so on. For example 8 is the double superbi-partient of 3 16 of 6 and in general the numbers beginning with 8 and differing by 8 are double superbi-partients of those beginning with 3 and differing by 3 when those in corresponding places in the series are compared and in the case of the other varieties one could ascertain their proper sequence by following out what has already been said. In this case too we must conceive that the nomenclature of the number compared goes along and suffers corresponding changes with the addition of the prefix sub-

[4] Thus we come to the end of our speculation upon the ten arithmetical relations for a first Introduction. There is however a method very exact and necessary for all discussion of the nature of the universe which very clearly and indisputably presents to us the fact that that which is fair and limited and which subjects itself to knowledge³ is naturally prior to the unlimited incomprehensible and ugly and furthermore that the parts and varieties of the infinite and unlimited are given shape and boundaries by the former and through it attain to their fitting order and sequence and like objects brought beneath some real or measure all gain a share of likeness to it and similarity of name when they fall under its influence. For thus it is reasonable that the rational

¹Referring to the table in Chapter 19

²See 70 2 above

³Cf 1 2 5

part of the soul will be the agent which puts in order the irrational part and passion and appetite which find their places in the two forms of inequality will be regulated by the reasoning faculty as though by a kind of equality and sameness [5] And from this equalizing process there will properly result for us the so-called ethical virtues sobriety,¹ courage gentleness self control fortitude and the like

[6] Let us then consider the nature of the principle that pertains to these universal matters It is capable of proving that all the complex species of inequality and the varieties of these species are produced out of equality, first and alone as from a mother and root

[7] Let there be given us equal numbers in three terms first units then two's in another group of three then three's next four's five's and so on as far as you like For them as the setting forth of these terms has come about by a divine and not human contrivance may by Nature herself, multiples will first be produced and among these the double will lead the way the triple after the double the quadruple next and then the quintuple and following the order we have previously recognized *ad infinitum* second the superparticular and here again the first form, the sesquialter will lead and the next after it the sesquitercian will follow and after them the next in order the sesquiquartan the sesquiquintan the sesquiseptan and so on *ad infinitum* third the superpartient which once more the superbipartient will lead the supertripartient will follow immediately upon it and then will come the superquadripartient the superquintpartient and according to the foregoing as far as one may proceed

[8] Now you must have certain rules like invariable and inviolable natural laws following which the whole aforesaid advance and progress from equality may go on without failure These are the directions Make the first equal to the first the second equal to the sum of the first and second and the third to the sum of the first twice the second and the third For if you fashion according to these rules you would get first all the forms of the multiple in order out of the three given terms of the equality as it were sprouting and growing without your paying any heed or offering any aid From equality you will first get the double from the double the triple from the triple successively

the quadruple and from this the quintuple in due order and so on [9] From the same multiples in their regular order reversed there are immediately produced by a sort of natural necessity through the agency of the same three rules the superparticulars and these not as it chances and irregularly but in their proper sequence for from the first the double reversed comes the first the sesquialter and from the second the triple the second in this class the sesquitercian then the sesquiquartan from the quadruple and in general each one from the one of similar name [10] And with a fresh start if the superparticulars are set forth in the order of their production but with terms reversed the superpartients which naturally follow them are brought to light the superbipartient from the sesquialter, the supertripartient from the sesquitercian the superquadripartient from the sesquiquartan and so on *ad infinitum* [11] If, however the superparticulars are set forth with terms not in reverse but in direct order there are produced through the three rules the multiple superparticulars, the double sesquialter out of the first the sesquialter the double sesquitercian from the second the sesquitercian, the double sesquiquartan from the third the sesquiquartan and so on [12] From those produced by the reversal of the superparticular that is the superpartients and from those produced without such reversal the multiple superparticulars there are once more produced in the same way and by the same rules both when the terms are in direct or reverse order the numbers that show the remaining numerical relations

[13] The following must suffice as illustrations of all that has been said hitherto the production of these numbers and their sequence and the use of direct and of reverse order [14] From the relation and proportion in terms of the sesquialter reversed so as to begin with the largest term there arises a relation in superpartient ratios the superbipartient and from it in direct order beginning with the smallest term a multiple superparticular relation the double sesquialter For example from 9 6 4 we get either 9 15 25 or 4 10 25 From the relation in terms of sesquitercians beginning with the greatest term is derived a superpartient the supertripartient beginning with the smallest term a double sesquitercian For example from 16 12 9 comes either 16 28 49 or 9 21 49 And from the relation in terms sesquiquartans when it is arranged to with the largest term is derived a s

¹Cf Aristotle *Ethics* 1107^b 4 ff See *ibid* 1108 4 ff 1145^b 8 ff

tient the superquadrupartient when it starts with the smallest term a multiple superparticular, the double sesquiquintan for instance from 25 20, 16 comes either 25, 45, 81 or 16 36 81

[16] In the case of all the e relations that are thus differentiated and of the one from which both of the differentiated ones are derived the last term is always the same and a square the first term becomes the smallest and invariably the extremes are squares

[16] Moreover the multiple superpartients and superpartients of other kinds are made to appear in yet another way out of the superpartients for example from the superbipartient relation arranged so as to begin with the small

est term comes the double superbipartient but arranged so as to start with the greater, the superpartient ratio of 8 5 Thus from 9 15 25 comes either 9 24 64 or 25 40, 64 From the supertripartient beginning with the smallest term we have the double supertripartient, and beginning with the largest the ratio of 11 7 Thus from 16 28 49 comes either 16, 44, 121 or 49 77, 121 [17] Again from the superquintipartient as for example, 25, 45 81 beginning with the lesser term we derive the double superquintipartient in the terms 25 70 196 but beginning with the greater a superpartient again, the ratio of 14 9 in the terms 81 126 196 And you will find the results analogous and in agreement with the foregoing in all successive cases to infinity

BOOK TWO

CHAPTER I

[1] An element is said to be and is the smallest thing which enters into the composition of an object and the least thing into which it can be analyzed. Letters for example are called the elements of literate speech for out of them all articulate speech is composed and into them finally it is resolved. Sounds are the elements of all melody for they are the beginning of its composition and into them it is resolved. The so-called four elements of the universe in general are simple bodies fire water air and earth for out of them in the first instance we account for the constitution of the universe and into them finally we conceive of it as being resolved.

We wish also to prove that equality is the elementary principle¹ of relative number for of absolute number number per se unity and the dyad are the most primitive elements the least things out of which it is constructed even to infinity by which it has its growth and with which its analysis into smaller terms comes to an end. [2] We have however demonstrated that in the realm of inequality advance and increase have their origin in equality and go on to absolutely all the relations with a certain regularity through the operation of the three rules.² It remains then in order to make it an element in very truth to prove that analyses also finally come to an end in equality. Let this then be considered our procedure.

CHAPTER II

[1] Suppose then you are given three terms in any relation whatsoever and in any ratio whether multiple superparticular superpartient or a compound of these multiple superparticular or multiple superpartient provided only that the mean term is seen to be in the same ratio to the lesser as the greater to the mean and vice versa. Subtract always from

the mean the lesser term whether it be first or last in order, and set down the lesser term itself as the first term of your new series then put as your second term what remains from the second after the subtraction then after having subtracted the sum of the new first term and twice the new second term from the remaining number—that is the greater of the numbers originally given you—make the remainder your third term and the resulting numbers will be in some other ratio naturally more primitive. [2] And if again in the same way you subtract the remainder from these same terms it will be found that your three terms have passed back into three others more primitive and you will find that this always takes place as a consequence until they are reduced to equality whence by every necessity it appears evident that equality is the elementary principle of relative quantity.

[3] There follows upon this speculation a most elegant principle extremely useful in its application to the Platonic psychogony³ and the problem of all harmonic intervals for in the Platonic passage we are frequently bidden, for the sake of the argument to set up series of intervals of two three four five or an infinite number of sesquialter ratios or two sesquiter tians sesquiquartans sesquioctaves or super particulars of any kind whatsoever, and in each case three four or five of them or as many as may be directed. [4] It is reasonable that we should do this not in an unscientific unintelligent fashion it may be even blunderingly but artistically surely and quickly by the following procedure.

CHAPTER III

[1] Every multiple will stand at the head⁴ of as many superparticular ratios corresponding in name with itself as it itself chances to be removed from unity and no more nor less under any circumstances.

¹See on I 23 4

²See on I 23 8.

³See Plato *Timaeus* 35 ff

⁴Refers to the table in '.

[2] The doubles then, will produce¹ sesqui alters the first one the second two the third three the fourth four, the fifth five, the sixth six and neither more nor less, but by every necessity when the superparticulars that are generated attain the proper number, that is, when their number agrees with the multiples that have generated them at that point by a divine device as it were there is found the

Sixteen the fourth double will stand at the head of four sesquialters 24, 36, 54 and finally 81, so that they may of necessity be equal in number to what generated them for 81 by its nature is not divisible by 2. And this as you go on you will find holds true in similar fashion to infinity.

[4] For the sake of illustration let there be set down the table of the doubles thus

The double ratio in the breadth of the table

	1	2	4	8	16	32	64	
		3	6	12	24	48	96	
The triple ratio along the hypotenuse			9	18	36	72	144	The sesquialter ratio in the depth
				27	54	108	216	
					81	162	324	
						243	486	
							729	

number which terminates them all because it naturally is not divisible by that factor where-by the progression of the superparticular ratios went on

CHAPTER IV

[1] We must make a similar table in illustration of the triple

The triple ratio in the breadth

	1	3	9	27	81	243	729	
		4	12	36	108	324	972	
The quadruple ratio on the hypotenuse			16	48	144	432	1296	The sesquitercian ratio in the depth
				64	192	576	1728	
					256	768	2304	
						1024	3072	
							4096	

From the triples all the sesquitercians will proceed likewise equal in number to the number of the generating terms and coming to an end after the independence of their advance is lost in numbers not divisible by 3. Similarly the sesquiquartans come from the quadruples reaching a culmination after their independent progression in a number that is not divisible by 4.

[3] As an example since doubles generate sesquialters corresponding to them in number the first row of multiples will be 1 2 4 8 16 32 64. Now since 2 is the first after unity this will be the origin of one sesquialter only 3 which number is not divisible by 2 so that another sesquialter might arise out of it. The first double therefore is productive of but one sesquialter, and the second 4 of two. For it produces its own sesquialter 6 and that of 6 9 but there is none for 9 because it has no half. Eight which is the third double is father to three sesquialters one its own 12 the second 18 the sesquialter of 12 and third 27 that of 18 there is no fourth one however because of the general rule for 27 is not divisible by 2.

¹See I 19 2

In the foregoing table we shall observe that in the same way the first triple 3 stands at the head of but one sesquitercian ratio 4 its own sesquitercian which immediately shuts off the development of another like it for 4 is not divisible by 3, and hence will not have a sesquitercian. The second triple is 9 and hence will begin a series of only two sesquitercian ratios 12 its own and 16 that of 12 but 16 cuts off further progress for it is not divisible by 3 and hence will not have a sesquitercian. [2] Next in order is the triple 27 three times removed from 1 for the triples progress thus 1 3, 9 27. Therefore this number will stand at the head of three sesquitercian ratios and no more. The first is its own 36 the second the sesquitercian of 36 48 the third that of the last 64 and this no longer has a third part and therefore will not admit of a sesquitercian. The fourth leads a series of four sesquitercians and the fifth of course five.

[3] Such then is the illustration and for the other multiples let the manner of your tables be the same. Observe that likewise here as we found to be true in our previous discus-

mon Nature shows us that the doubles are more nearly original than the triples the triples than the quadruples the quadruples than the quintuples and so on throughout For the highest rows of figures across the breadth of the tables if they are doubles will have doubles lying parallel to them and the numbers lying diagonally on the hypotenuse will be of the next succeeding variety greater by 1 than is triples seen also in a series of parallel lines If however there are triples across the breadth the diagonals will by all means be quadruples if the former are quadruples then the latter are quintuples and so forth

CHAPTER V

[1] It remains after we have explained what other ratios are produced by combination of ratios to pass on to the succeeding topics of the *Introduction*

[2] Now the first two ratios of the superparticular combined produce the first ratio of the multiple namely the double for every double is a combination of sesquialter and sesquitercian and every sesquialter and sesquitercian combined will invariably produce a double

For example since 3 is the sesquialter of 2 and 4 the sesquitercian of 3 4 will be the double of 2 and is a combination of sesquialter and sesquitercian Again as 6 is the double of 3 we shall find between them some number that will of necessity preserve the sesquitercian ratio to the one and the sesquialter to the other and indeed 4 lying between 6 and 3 gives the sesquitercian ratio to 3 and the sesquialter to 6

[3] It was rightly said then that the double when resolved is resolved into the sesquialter and the sesquitercian and that when sesquialter and sesquitercian are combined there arises the double and that the first two forms of the superparticular combined make the first form of the multiple

[4] But again to take another start this first form of the multiple which has thus been produced together with the first form of the superparticular will produce the next form of the same class that is the second multiple the triple for from every multiple and sesquialter combined a triple of necessity arises For example as the double of 6 is 12 and the sesquialter of this is 18 then immediately 18 is the triple of 6 and to take another method if I do not care to make 12 the mean term but rather 9 the sesquialter of 6 the same result will come about without deviation and harmoniously

for while 18 is the double of 9 it will preserve the triple ratio to 6 Hence from the sesquialter and the double, the first forms of the superparticular and the multiple there arises by combination the second form of the multiple, the triple and into them it is always resolved [6] For look you 6 which is the triple of 2 will have a mean term 3 which will exhibit two ratios the sesquialter with regard to 2 and the double ratio of 6 to itself

But if this triple ratio likewise the second form of the multiple is combined with the sesquitercian which is the second form of the superparticular there would be produced from them the next form of the multiple namely the quadruple and this also will of necessity be resolved into them after the same fashion as the cases previously set forth and the quadruple taking into combination the sesquiquartan will make the quintuple and once more the latter with the sesquiquintan will make the sextuple and so on to the end Thus the multiples in regular order from the beginning with the superparticulars in regular order from the beginning will be found to produce the next larger multiples For the double with the sesquialter makes the triple the triple with the sesquitercian the quadruple the quadruple with the sesquiquartan the quintuple and as far as you wish to proceed no contrary result will appear

CHAPTER VI

[1] Up to this point then we have sufficiently discussed relative number by a process of selection measuring out what is easily comprehended and appropriate to the nature of the matters thus far introduced Whatever remains to be said on this topic will be filled in after we have put it aside and have first discussed certain subjects which involve a more serviceable inquiry having to do with the properties of absolute number not relative For mathematical speculations are always to be interlocked and to be explained one by means of another The subjects which we must first survey and observe are concerned with linear plane and solid numbers cubical and spherical equilateral and scalene bricks beams wedges and the like the tradition concerning which to be sure since they are more closely related to magnitude is properly given in the *Geometrical Introduction* Yet the germs of these ideas are taken over into arithmetic, as the science which is the mother of geometry and more elementary than it For we recall that a short time ago we saw that arithmetic abolishes the other science

with itself but is not abolished by them, and conversely is of necessity implied by them but does not itself imply them

[2] First however, we must recognize that each letter by which we indicate a number, such as *iota* the sign for 10 *kappa* for 20 and *omega* for 800 designates that number by man's convention and agreement not by nature. On the other hand the natural unartificial and therefore simplest indication of numbers would be the setting forth one beside the other of the units contained in each. For example the writing of one unit by means of one *alpha* will be the sign for 1 two units side by side that is, a series of two *alphas* will be the sign for 2 when three are put in a line it will be the character for 3 four in a line for 4 five for 5 and so on. For by means of such a notation and indication alone could the schematic arrangement of the plane and solid numbers mentioned be made clear and evident thus

The number 1	α
The number 2	$\alpha \alpha$
The number 3	$\alpha \alpha \alpha$
The number 4	$\alpha \alpha \alpha \alpha$
The number 5	$\alpha \alpha \alpha \alpha \alpha$

and further in similar fashion

[3] Unity then occupying the place and character of a point will be the beginning of intervals and of numbers but not itself an interval or a number, just as the point is the beginning of a line or an interval but is not itself line or interval. Indeed when a point is added to a point it makes no increase for when a non-dimensional thing is added to another non-dimensional thing it will not thereby have dimension just as if one should examine the sum of nothing added to nothing which makes nothing. We saw¹ a similar thing also in the case of equality among the relatives for a proportion is preserved—as the first is to the second so the second is to the third—but no interval is generated in the relation of the extremes to each other as there is in all the other relations with the exception of equality. In exactly the same way unity alone out of all number when it multiplies itself produces nothing greater than itself

Unity, therefore, is non-dimensional and elementary and dimension first is found and seen in 2 then in 3 then in 4 and in succession in the following numbers for dimension is that

which is conceived of as between two limits

[4] The first dimension is called 'line' for 'line' is that which is extended in one direction. Two dimensions are called 'surface' for a surface is that which is extended in two directions. Three dimensions are called 'solid' for a solid is that which is extended in three directions and it is by no means possible to conceive of a solid which has more than three dimensions depth breadth and length. By these are defined the six directions which are said to exist in connection with every body and by which motions in space are distinguished forward backward² up down right and left for of necessity two directions opposite to each other follow upon each dimension up and down upon one, forward and backward upon the second and right and left upon the third.

[5] The statement also as it happens can be made conversely thus. If a thing is solid it has by all means three dimensions length depth and breadth and conversely if it has the three dimensions it is always a solid, and nothing else.

[6] That which has but two dimensions therefore will not be a solid but a surface for the latter admits of but two dimensions. Here too it is possible similarly to reverse the statement directly stated a surface is that which has two dimensions and conversely that which has two dimensions is always a surface.

[7] The surface then, is exceeded by the solid by one dimension and the line is exceeded by the surface by one for the line is that which is extended in but one direction and has only one dimension and it falls short of the solid by two dimensions. The point falls short of the latter by one dimension and hence it has already been stated that it is non-dimensional since it falls short of the solid by three dimensions of the surface by two and of the line by one.

CHAPTER VII

[1] The point then is the beginning of dimension but not itself a dimension and likewise the beginning of a line but not itself a line the line is the beginning of surface but not surface and the beginning of the two-dimensional but not itself extended in two directions. [2] Naturally too surface is the beginning of body, but not itself body and likewise the beginning of the three-dimensional but not itself extended in three directions.

[3] Exactly the same in numbers unity is the beginning of all number that advances unit by unit in one direction linear number is the beginning of plane number which spreads out like a plane in one more dimension and plane number is the beginning of solid number which possesses a depth¹ in the third dimension besides the original ones To illustrate and classify, linear numbers are all those which begin with 2 and advance by the addition of 1 in one and the same dimension and plane numbers are those² that begin with 3 as their most elementary root and proceed through the next succeeding numbers They receive their names also in the same order for there are first the triangles then the squares the pentagons after these then the hexagons the heptagons and so on indefinitely and as we said they are named after the successive numbers beginning with 3

[4] The triangle therefore is found to be the most original and elementary form of the plane number Thus we can see from the fact that, among plane figures³ graphically represented if lines are drawn from the angles to the centers each rectilinear figure will by all means be resolved into as many triangles as it has sides but the triangle itself if treated like the rest will not change into anything else but itself Hence the triangle is elementary among these figures for everything else is resolved in to it but it into nothing else From it the others likewise would be constituted but it from no other It is therefore the element of the others and has itself no element [5] Likewise as the argument proceeds in the realm of numerical forms it will confirm this statement

CHAPTER VIII

[1] Now a triangular number is one which when it is analyzed into units shapes into triangular form the equilateral placement of its parts in a plane 3 6 10 15 21 28 and so on are examples of it for their regular formations expressed graphically will be at once triangular and equilateral As you advance you will find that such a numerical series as far as you like takes the triangular form if you put as the most elementary form the one that arises from unity so that unity may appear to be potentially a triangle and 3 the first actually

[2] Their sides will increase by the succes-

sive numbers for the side of the one potentially first is unity that of the one actually first that is 3 is 2 that of 6 which is actually second 3 that of the third 4 the fourth, 5 the fifth 6, and so on

[3] The triangular number is produced from the natural series of number set forth in a line and by the continued addition of successive terms one by one, from the beginning for by the successive combinations and additions of another term to the sum the triangular numbers in regular order are completed For example from this natural series 1 2 3 4 5 6 7 8 9 10 11 12 13, 14 15 I take the first term and have the triangular number which is potentially first, 1,



then adding the next term I get the triangle actually first for 2 plus 1 equals 3 In its graphic representation it is thus made up Two units side by side are set beneath one unit and the number three is made a triangle



Then when next after these the following number 3 is added simplified into units and joined to the former it gives 6 the second triangle in actuality and furthermore it graphically represents this number



Again the number that naturally follows 4 added in and set down below the former reduced to units gives the one in order next after the aforesaid 10 and takes a triangular form




5 after this then 6 then 7 and all the numbers in order are added so that regularly the sides of each triangle will consist of as many

¹Cf Plato *Timaeus* 53

²But cf Euclid in *Elements* VII Def 17

³Cf Plato *Timaeus* 53 ff


numbers as have been added from the natural series to produce it



Side 5




Side 6



Side 7

CHAPTER IX

[1] The square is the next number after this which shows us no longer 3 like the former but 4 angles in its graphic representation but is none the less equilateral. Take for example 1 4 9 16 25 36 49 64 81, 100 for the representations of these numbers are equilateral, square figures as here shown, and it will be similar as far as you wish to go



1

4

9



16

25

[2] It is true of these numbers as it was also of the preceding that the advance in their sides progresses with the natural series. The side of the square potentially first 1 is 1 that of 4 the first in actuality 2 that of 9 actually the second 3 that of 16 the next actually the

third 4 that of the fourth 5 of the fifth, 6, and so on in general with all that follow

[3] This number also is produced if the natural series is extended in a line increasing by 1, and no longer the successive numbers are added to the numbers in order, as was shown before but rather all those in alternate places that is, the odd numbers. For the first 1 is potentially the first square the second 1 plus 3 is the first in actuality, the third 1 plus 3 plus 5 is the second in actuality the fourth 1 plus 3 plus 5 plus 7 is the third in actuality the next is produced by adding 9 to the former numbers the next by the addition of 11 and so on

[4] In these cases also it is a fact that the side of each consists of as many units as there are numbers taken into the sum to produce it

CHAPTER X

[1] The pentagonal number is one which likewise upon its resolution into units and depiction as a plane figure assumes the form of an equilateral pentagon 1, 5 12 22 35 51 70 and analogous numbers are examples [2] Each side of the first actual pentagon 5 is 2 for 1 is the side of the pentagon potentially first 1 3 is the side of 12 the second of those listed 4 that of the next 22 5 that of the next in order 35 and 6 of the succeeding one, 51 and so on. In general the side contains as many units as are the numbers that have been added together to produce the pentagon chosen out of the natural arithmetical series set forth in a row. For in a like and similar manner there are added together to produce the pentagonal numbers the terms beginning with 1 to any extent whatever that are two places apart that is those that have a difference of 3

Unity is the first pentagonal potentially and is thus depicted



5 made up of 1 plus 4 is the second similarly represented



12 the third is made up out of the two former numbers with 7 added to them so that it may have 3 as a side as three numbers have been

added to make it. Similarly the preceding pentagon 5 was the combination of two numbers and had 2 as its side. The graphic representation of 12 is this:



The other pentagonal numbers will be produced by adding together one after another in due order the terms after 7 that have the difference 3, as for example 10 13 16 19 22 25 and so on. The pentagons will be 22 35 51 70 92, 117 and so forth.

CHAPTER XI

[1] The hexagonal, heptagonal and succeeding numbers will be set forth in their series by following the same process: if from the natural series of number there be set forth series with their differences increasing by 1. For as the triangular number was produced by admitting into the summation the terms that differ by 1 and do not pass over any in the series, as the square was made by adding the terms that differ by 2 and are one place apart, and the pentagon similarly by adding terms with a difference of 3 and two places apart (and we have demonstrated these by setting forth examples both of them and of the polygonal numbers made from them), so likewise the hexagons will have as their root numbers those which differ by 4 and are three places apart in the series which added together in succession will produce the hexagons. For example 1 5 9 13 17 21 and so on, so that the hexagonal numbers produced will be 1 6 15 28 45 66 and so on as far as one wishes to go.

[2] The heptagonals which follow these have as their root numbers terms differing by 5 and four places apart in the series, like 1 6 11 16 21 26 31 36 and so on. The heptagons that thus arise are 1 7 18 34 55 81 112 148 and so forth.

[3] The octagonals increase after the same fashion with a difference of 6 in their root-

numbers and corresponding variation in their total constitution.

[4] In order that as you survey all cases you may have a rule generally applicable, note that the root-numbers of any polygonal differ by 2 less than the number of the angles shown by the name of the polygonal—that is, by 1 in the triangle, 2 in the square, 3 in the pentagon, 4 in the hexagon, 5 in the heptagon, and so on with similar increase.

CHAPTER XII

[1] Concerning the nature of plane polygon, as this is sufficient for a first *Introduction*. That however the doctrine of these numbers is to the highest degree in accord with their geometrical representation, and not out of harmony with it, would be evident not only from the graphic representation in each case, but also from the following. Every square figure diagonally divided is resolved into two triangles, and every square number is resolved into two consecutive triangular numbers, and hence is made up of two successive triangular numbers. For example 1 3 6 10 15 21 28 36 45 55 and so on are triangular numbers, and 1 4 9 16 25 36 49 64 81 100 squares. [2] If you add any two consecutive triangles that you please, you will always make a square, and hence whatever square you resolve, you will be able to make two triangles of it.

Again, any triangle joined to any square figure makes a pentagon; for example, the triangle 1 joined with the square 4 makes the pentagon 5, the next triangle 3 of course with 9 the next square makes the pentagon 12, the next 6 with the next square 16 gives the next pentagon 22, 10 and 25 give 35, and so on.

[3] Similarly, if the triangles are added to the pentagons following the same order, they will produce the hexagonals in due order, and again the same triangles with the latter will make the heptagonals in order, the octagonals after the heptagonals, and so on to infinity.

[4] To remind us, let us set forth rows of the polygonals written in parallel lines as follows. The first row triangles, the next squares, after them pentagonals, then hexagonals, then heptagonals, then if one wishes the succeeding polygonals.

Triangles	1	3	6	10	15	21	28	36	45	55
Squares	1	4	9	16	25	36	49	64	81	100
Pentagonals	1	5	12	22	35	51	70	92	117	145
Hexagonals	1	6	15	28	45	66	91	120	153	190
Heptagonals	1	7	18	34	55	81	112	148	189	"

You can also set forth the succeeding polygon also in similar parallel lines

[5] In general you will find that the squares are the sum of the triangles above those that occupy the same place in the series, plus the numbers of that same class in the next place back for example 4 equals 3 plus 1 9 equals 6 plus 3 16 equals 10 plus 6 25 equals 15 plus 10, 36 equals 21 plus 15 and so on

The pentagons are the sum of the squares above them in the same place in the series, plus the elementary triangles that are one place further back in the series for example 5 equals 4 plus 1 12 equals 9 plus 3 22 equals 16 plus 6, 35 equals 25 plus 10 and so on

[6] Again the hexagonals are similarly the sums of the pentagons above them in the same place in the series plus the triangles one place back for instance, 6 equals 5 plus 1 15 equals 12 plus 3, 28 equals 22 plus 6 45 equals 35 plus 10 and as far as you like

[7] The same applies to the heptagonals for 7 is the sum of 6 and 1 18 equals 15 plus 3 34 equals 28 plus 6 and so on Thus each polygonal number is the sum of the polygonal in the same place in the series with one less angle plus the triangle in the highest row one place back in the series

[8] Naturally then the triangle is the element of the polygon both in figures and in numbers and we say this because in the table reading either up and down or across the successive numbers in the rows are discovered to have as differences the triangles in regular order

CHAPTER XIII

[1] From this it is easy to see what the solid number is and how its series advances with equal sides for the number which in addition to the two dimensions contemplated in graphic representation in a plane length and breadth has a third dimension which some call depth others thickness and some height that number would be a solid number extended in three directions and having length depth and breadth

[2] This first makes its appearance in the so-called pyramids These are produced from rather wide bases narrowing to a sharp apex first after the triangular form from a triangular base second after the form of the square from a square base, and succeeding these after the pentagonal form from a pentagonal base then similarly from the hexagon heptagon octagon and so on indefinitely

[3] Exactly so among the geometrical solid

figures if one imagines three lines from the three angles of an equilateral triangle equal in length to the sides of the triangle converging in the dimension height to one and the same point a pyramid would be produced, bounded by four triangles equilateral and equal one to the other one the original triangle and the other three bounded by the aforesaid three line [4] And again if one conceives of four lines starting from a square equal in length to the sides of the square each to each and again converging in the dimension height to one and the same point a pyramid would be completed with a square base and diminishing in square form, bounded by four equilateral triangles and one square the original one [5] And starting from a pentagon hexagon heptagon and however far you care to go lines equal in number to the angles erected in the same fashion from the angles and converging to one and the same point will complete a pyramid named from its pentagonal hexagonal, or heptagonal base, or similarly

[6] So likewise among numbers each linear number increases from unity as from a point as for example 1, 2, 3, 4, 5 and successive numbers to infinity and from the same numbers which are linear and extended in one direction combined in no random manner the polygonal and plane numbers are fashioned—the triangles by the combination of root numbers immediately adjacent the square by adding every other term the pentagons every third term and so on [7] In exactly the same way, if the plane polygonal numbers are piled one upon the other and as it were built up the pyramids that are akin to each of them are produced the triangular pyramid from the triangles the square pyramid from the squares the pentagonal from the pentagons the hexagonal from the hexagons and so on throughout

[8] The pyramids with a triangular base then in their proper order are these 1, 4, 10, 20, 35, 56, 84 and so on and their origin is the piling up of the triangular numbers one upon the other first 1, then 1 3 then 1 3 6 then 10 in addition to these and next 15 together with the foregoing then 21 besides these next 28 and so on to infinity

[9] It is clear that the greatest number is conceived of as being lowest for it is discovered to be the base the next succeeding one is on top of it and the next on top of that until unity appears at the apex and so to speak tapers off the completed pyramid into a point

CHAPTER XIV

[1] The next pyramids in order are those with a square base which rise in this shape to one and the same point. These are formed in the same way as the triangular pyramids of which we have just spoken. For if I extend in series the square numbers in order beginning with unity thus 1 4, 9 16 25 36 49 64 81 100 and again set the successive terms as in a pile one upon the other in the dimension height when I put 1 on top of 4 the first actual pyramid with square base 5 is produced for here again unity is potentially the first [2] Once more I put this same pyramid entire composed of 5 units just as it is upon the square 9 and there is made up for me the pyramid 14 with square base and side 3—for the former pyramid had the side 2 and the one potentially first 1 as a side 1 or here too each side of any pyramid whatsoever must consist of as many units as there are polygonal numbers piled together to create it.

[3] Again I place the whole pyramid 14 with the square 9 as its base upon the square 16 and I have 30 the third actual pyramid of those that have a square base and by the same order and procedure from a pentagonal hexagonal or heptagonal base, and even going on farther we shall produce pyramids by piling upon one another the corresponding polygonal numbers starting with unity as the smallest and going on to infinity in each case.

[4] From this too it becomes evident that triangles are the most elementary for absolutely all of the pyramids that are exhibited and shown, with the various polygonal bases are bounded by triangles up to the apex.

[5] But lest we be heedless of truncated bi-truncated and tri-truncated pyramids the names of which we are sure to encounter in scientific writings you may know that if a pyramid with any sort of polygon as its base triangular square pentagon or any of the succeeding polygons of the kind when it increases by this process of piling up does not taper off into unity it is called simply truncated when it is left without the natural apex that belongs to all pyramids for it does not terminate in the potential polygon unity as in some one point but in another polygon and an actual one and unity is not its apex but its upper boundary becomes a plane figure with the same number of angles as the base. If however in addition to the failure to terminate in unity it does not even terminate in the polygon next to unity and

the first in actuality, such a pyramid is called bi-truncated and if, still further it does not have the second actual polygon at its upper limit but only the one next beneath it will be called tri-truncated yes even four times truncated if it does not have the next one as its limit, or five times truncated at the next step and so on as far as you care to carry the nomenclature.

CHAPTER XV

[1] While the origin advance increase and nature of the equilateral solid numbers of pyramidal appearance is the foregoing, with its seed and root in the polygonal numbers and the piling up of them in their regular order there is another series of solid numbers of a different kind consisting of the so-called cubes, beams, 'bricks' 'wedges' spheres and parallelepipeds which has the order of its progress somewhat as follows.

[2] The foregoing squares 1 4 9 16 25 36 49 64 and so on which are extended in two directions and in their graphic representation in a plane have only length and breadth will take on yet a third dimension and be solids and extended in three directions if each is multiplied by its own side 4 which is 2 times 2 is again multiplied by 2 to make 8 9 which is 3 times 3 is again increased by 3 in another dimension and gives 27 16 which is 4 times 4 is multiplied by its own side 4 and 64 results and so on with the succeeding squares throughout.

[3] Here too the sides will be composed of as many units as were in the sides of the squares from which they arose in each case the sides of 8 will be 2 like those of 4 those of 27 3, like those of 9 those of 64 4 like those of 16 and so on so that likewise the side of unity the potential cube will be 1 which is the side of the potential square 1.

In general each square is a single plane and has four angles and four sides while each several cube having increased out of some one square multiplied by its own side will have always six plane surfaces each equal to the original square and twelve edges each equal to and containing exactly the same number of units as each side of the original square and eight solid angles each of which is bounded by three edges like in each case to the sides of the original square.

CHAPTER XVI

[1] Now since the cube is a solid figure with equal sides in all dimensions in length depth,

and breadth and is equally extended in all the six so called directions¹ it follows that there is opposed to it that which has its dimensions in no case equal to one another but its depth unequal to its breadth and its length unequal to either of these for example 2 times 3 times 4 or 2 times 4 times 8, or 3 times 5 times 12 or a figure which follows some other scheme of inequality

[2] Such solid figures in which the dimensions are everywhere unequal one to another, are called scalene in general. Some however, using other names call them 'wedges' for carpenters house builders and blacksmiths wedges and those used in other crafts having unequal sides in every direction are fashioned so as to penetrate they begin with a sharp end and continually broaden out unequally in all the dimensions. Some also call them *sphekiskoi* wasps because wasps bodies also are very like them compressed in the middle and showing the resemblance mentioned. From this also the *sphekoma* point of the helmet must derive its name for where it is compressed it imitates the waist of the wasp. Others call the same numbers altars using their own metaphor for the altars of ancient style particularly the Ionic do not have the breadth equal to the depth nor either of these equal to the length nor the base equal to the top but are of varied dimensions everywhere

[3] Now whereas the two kinds of numbers cube and scalene are extremes the one equally extended in every dimension the other unequally the so-called parallelepipedons are solid numbers like means between them. The plane surfaces of these are heteromecic numbers² just as in the case of the cubes the faces were squares as has been shown

CHAPTER XVII

[1] Again then to take a fresh start a number is called heteromecic if its representation when graphically described in a plane is quadrilateral and quadrangular to be sure but the sides are not equal one to another nor is the length equal to the breadth but they differ by 1. Examples are 2 6 12 20 30 42 and so on for if one represents them graphically he will always construct them thus 1 times 2 equals 2 2 times 3 equals 6 3 times 4 equals 12 and the succeeding ones similarly 4 times 5 5 times 6, 6 times 7, 7 times 8 and thus indefinitely

provided only that one side is greater than the other by 1 and by no other number. If however the sides differ otherwise than by 1 for instance by 2 3 4 or succeeding numbers as in 2 times 4 3 times 6 4 times 8 or however else they may differ then no longer will such a number be properly called a heteromecic but an oblong number. For the ancients of the school of Pythagoras and his successors saw the other³ and otherness primarily in 2 and 'the same and sameness in 1 as the two beginnings of all things and these two are found to differ from each other only by 1. Thus the other⁴ is fundamentally other by 1, and by no other number, and for this reason customarily other is used among those who speak correctly of two things and not of more than two

[2] Moreover, it was shown that all odd number is given its specific form⁵ by unity and all even number by 2. Hence we shall naturally say that the odd partakes of the nature of the same and the even of that of the other for indeed there are produced by the successive additions of each of these—naturally and not by our decree—by the addition of the odd numbers from 1 to infinity the class of the squares and by the addition of the evens from 2 to infinity that of the heteromecic numbers⁶

[3] There is accordingly every reason to think that the square once more shares in the nature of the same for its sides display the same ratio alike unchanging and firmly fixed in equality to themselves while the heteromecic number partakes of the nature of the other for just as 1 is differentiated from 2 differing by 1 alone thus also the sides of every heteromecic number differ from one another, one differing from the other by 1 alone

To illustrate if I have set out before me the successive numbers in series beginning with 1 and select and arrange by themselves the odd numbers in the line and the even by themselves in another there are obtained these two series

1 3 5 7 9 11 13 15 17 19 21 23 25 27
2 4 6 8 10 12 14 16 18 20 22 24 26 28

[4] Now then the beginning of the odd series is unity which is of the same class as the series and possesses the nature of the same and so whether it multiplies itself in two dimensions

¹Cf Plato *Timaeus* 35 ff

²Cf 1 7 2

³Cf II 18 2 and 20 3

⁴Cf II 6 4

⁵See the following chapter

or in three it is not made different nor yet does it make any other number depart from what it was originally¹ but keeps it just as it was. Such a property it is impossible to find in any other number. [6] Of the other series the beginning is 2 which is similar in kind to this series and imitates otherness for whether it multiplies itself or another number it causes a change² for example 2 times 2 2 times 3

[6] But in cases like 8 times 8 times 2 or 8 times 8 times 3 such solid forms are called 'bricks,' the product of a number by itself and then by a smaller number if however a greater height is joined to the square as in 3 times 3 times 7, 3 times 3 times 8 or 3 times 3 times 9 or however many times the square be taken provided only it be a greater number of times than the square itself then the number is a beam the product of a number by itself and then by a larger number. The wedges to be sure were the products of three unequal numbers and cubes of three equal ones.

[7] Among the cubes some of them in addition to being the product of three equal numbers have the further property of ending at every multiplication in the same number as that from which they began these are called spherical and also recurrent. Such indeed are those with sides 5 or 6 for however many times I increase each one of these it will by all means end each time in the same figure the derivative of 6 in 6 and that of 5 in 5. For example the product of 5 times 5 will end in 5 and so will 5 times this product and if necessary 5 times this again and to infinity no other concluding term will be found except 5. From 6 too in the same fashion 6 and no other will be the concluding term and so 1 likewise is potentially spherical and recurrent for as is reasonable it has the same property as the spheres and circles. For each one of them circling and turning around ends where it begins. And so these numbers afore said are the only ones of the products of equal factors to return to the same starting point from which they began in the course of all their increases. If they increase in the manner of planes in two dimensions they are called circular like 1 25 and 36 derived from 1 times 1 5 times 5 and 6 times 6 but if they have three dimensions or are multiplied still further than this they are called spherical solid numbers for example 1 125 216 or again 1 625 1 296

CHAPTER XVIII

[1] Regarding the solid numbers this is for the present sufficient. The physical philosophers however and those that take their start with mathematics call the same and the other 'the principles of the universe and it has been shown that "the same inheres in unity and the odd numbers to which unity gives specific form and to an even greater degree in the squares made by the continued addition of odd numbers because in their sides they share in equality while "the other" inheres in 2 and the whole even series which is given specific form by 2 and particularly in the heteromecic numbers which are made by the continued addition of the even numbers because of the share of the original inequality' and otherness which they have in the difference between their sides. Therefore it is most necessary further to demonstrate how in these two as in origins and seeds there are potentially existent all the peculiar properties of number of its forms and subdivisions of all its relations of polygonals and the like.

[2] First however we must make the distinction whereby the oblong (promecic) number differs from the heteromecic. The heteromecic is as was stated above 'the product of a number multiplied by another larger than the first by 1 for example 6 which is 2 times 3 or 12 which is 3 times 4. But the oblong is similarly the product of two differing numbers differing however not by 1 but by some larger number as 2 times 4 3 times 6 4 times 8 and similar numbers which in a way exceed in length and overstep the difference of 1.

[3] Therefore since squares are produced from the multiplication of numbers by their own length and have their length the same as their breadth properly speaking they would be called idiomecic or tautomecic for example 2 times 2 3 times 3 4 times 4 and the rest. And if this is true they will admit in every way of sameness and equality and for this reason are limited and come to an end for the equal and the same are so in one definite way. But since the heteromecic numbers are produced by the multiplication of a number by not its own but another number's length they are therefore called heteromecic and admit of infinity and boundlessness.

[4] In this way then all numbers and the

¹Cf II 6 3

²Cf Aristotle, *On the Soul* 406^b 13

¹Cf II 17 2

²Cf II 17 1

objects in the universe which have been created with reference to them are divided and classified and are seen to be opposite one to another and well do the ancients at the very beginning of their account of Nature make the first subdivision in their cosmogony on this principle. Thus Plato¹ mentions the distinction between the natures of the same and "the other" and again that between the essence which is indivisible and always the same and the one which is divided. and Philolaus says that existent things must all be either limitless or limited or limited and limitless at the same time, by which it is generally agreed that he means that the universe is made up out of limited and limitless things at the same time obviously after the image of number for all number is composed of unity and the dyad even and odd and these in truth display equality and inequality sameness and otherness the bounded and the boundless the defined and the undefined

CHAPTER XIV

[1] That we may be clearly persuaded of what is being said namely that things are made up of warring and opposite elements² and have in all likelihood taken on harmony—and harmony always arises from opposites for harmony is the unification of the diverse and the reconciliation of the contrary minded—let us set forth in two parallel lines no longer as just previously the even numbers from 2 by themselves and the odd numbers from 1 but the numbers that are produced from these by adding them successively together the squares from the odd numbers and the heteromeic from the even. For if we give careful attention to their setting forth we shall admire their mutual friendship and their cooperation to produce and perfect the remaining forms to the end that we may with probability conceive that also in the nature of the universe from some such source as this a similar thing was brought about by universal providence

[2] Let the two series then be as follows That of the squares from unity 1 4 9 16 25 36 49 64 81 100 121 144 169 196 225 and that of the heteromeic numbers beginning with 2 and proceeding thus 2 6 12 20 30 42 56 72 90 110 132 156 182 210 240

[3] In the first place then the first square is the fundamental multiple of the first hetero-

meic number the second compared to the second is its sesquialter the third, sesquitercian of the third the fourth sesquiquartan of the fourth then sesquiquintan sesquisextan and so on similarly *ad infinitum*. Their differences too will increase according to the successive numbers from 1 the difference of the first terms is 1 of the second 2 of the third 3, and so on. Next if first the second term of the squares be compared with the first heteromeic number, the third with the second the fourth with the third and the rest similarly they will keep unchanged the same ratios as before but their differences will begin to progress no longer from 1 but from 2 remaining the same as before and according to the advance observed in the former comparison the first to the first will be the first or root form multiple the second to the second the second sesquialter from the root-form the third to the third the third sesquitercian from the root-form and the succeeding terms will go on in similar fashion

[4] Furthermore the squares among themselves will have only the odd numbers as differences the heteromeic even numbers. And if we put the first heteromeic number as a mean term between the first two squares the second between the next two the third between the two following and the fourth between the two next succeeding therein will be seen still more regularly the numerical relations in groups of three terms. For as 4 is to 2 so is 2 to 1 and as 9 is sesquialter to 6 so is 6 to 4 and as 16 to 12 so is 12 to 9 and so on with both numbers and ratios regularly advancing. As the greater is to the mean so will the mean be to the lesser and not in the same ratio but always a different one by an increase. In all the groupings too the product of the extremes is equal to the square of the mean and the extremes plus twice the mean by exchange will always give a square. What is neatest of all from the addition of both there comes about the production of the triangles in due order showing that the nature of these is more ancient³ than the origin of all things thus 1 plus 2 2 plus 4 4 plus 6 6 plus 9 9 plus 12 12 plus 16 16 plus 20 and by this process the triangles which give rise to the polygons come forth in order

CHAPTER XX

[1] Still further every square plus its own side becomes heteromeic or by Zeus if its side is subtracted from it. Thus the other⁴ is con-

¹Cf Plato *Timaeus* 30

²Plato *Timaeus* 30

³Cf II 17 3 18 1 and II 7 4 cf 12 8

ceived of as being both greater and smaller than 'the same' since it is produced both by addition and by subtraction in the same way that the two kinds of inequality¹ also the greater and the less have their origin from the application of addition or subtraction to equality [2] This also is sufficient evidence that the two forms partake of sameness and otherness of otherness in an indefinite fashion but of sameness definitely 1 and 2 generically but the odd of sameness after the manner of a subordinate species because it belongs to the same class as 1, and the even of otherness because it is homogeneous with 2

[3] There is also a still clearer reason why the square since it is the product of the addition of odd numbers is akin to sameness and the heteromeic numbers to otherness because it is made up by adding even numbers for as though they were friends of one another these two forms share in their two rows the same differences when they do not have the same ratios and conversely the same ratios when they do not have the same differences For the difference between 4 and 2 in the double ratio is found between 6 and 4 as a superparticular and again the difference between 9 and 6 as a sesquialter is found between 12 and 9 as a sesquitercian and so on What is the same in quality is different in quantity and just the opposite what is the same in quantity is different in quality [4] Again it is clear that in all their relations the same difference between two terms will necessarily be called fractions with names that differ by 1 and be the half of one and the third of the other or the third of one and the quarter of the other or the fourth of one and the fifth of the other and so on

[5] But what will most of all confirm the fact that the odd and never the even is pre-eminently the cause of sameness is to be demonstrated in every series beginning with 1 following some ratio for example the double ratio 1 2 4 8 16 32 64 128 256 or the triple 1 3 9 27 81 243 729 2187 and as far as you like You will find that of necessity all the terms in the odd places in the series are squares and no others by any device whatsoever and that no square is to be found in an even place

But all the products of a number multiplied twice into itself that is the cubes which are extended in three dimensions and seen to share in sameness to an even greater extent are the product of the odd numbers not the even 1

8 27 64 125 and 216 and those that go on analogously, in a simple unvaried progression as well For when the successive odd numbers are set forth indefinitely beginning with 1 observe this The first one makes the potential cube the next two added together the second the next three the third the four next following the fourth the succeeding five the fifth the next six the sixth and so on

CHAPTER XXI

[1] After this it would be the proper time to incorporate the nature of proportions a thing most essential for speculation about the nature of the universe and for the propositions of music astronomy, and geometry and not least for the study of the works of the ancients and thus to bring the *Introduction to Arithmetic* to the end that is at once suitable and fitting

[2] A proportion then is in the proper sense the combination of two or more ratios but by the more general definition the combination of two or more relations even if they are not brought under the same ratio but rather a difference or something else

[3] Now a ratio² is the relation of two terms to one another and the combination of such is a proportion so that three is the smallest number of terms of which the latter is composed although it can be a series of more subject to the same ratio or the same difference For example 1 2 is one ratio where there are two terms but 2 4 is another similar ratio hence 1 2 4 is a proportion for it is a combination of ratios or of three terms which are observed to be in the same ratio to one another [4] The same thing may be observed also in greater numbers and longer series of terms for let a fourth term 8 be joined to the former after 4 again in a similar relation the double and then 16 after 8 and so on

[5] Now if the same term one and unchanging is compared to those on either side of it to the greater as consequent and to the lesser as antecedent such a proportion is called continued for example 1 2 4 is a continued proportion as regards quality³ for 4 2 equals 2 1 and conversely 1 2 equals 2 4 In quantity 1 2 3 for example is a continued proportion for as 3 exceeds 2 so 2 exceeds 1 and conversely as 1 is less than 2 by so much 2 is less than 3

[6] If however one term answers to the lesser term and becomes its antecedent and a

¹Cf I 17 6

²Cf Euclid *Elements* I viii

³Cf II 22 2 23 4 below

greater term and another not the same takes the place of consequent and lesser term with reference to the greater such a mean and such a proportion is called *no longer continued* but *disjunct* for example as regards quality, 1 2 4 8 for 2 1 equals 8 4 and conversely 1 2 equals 4 8 and again 1 4 equals 2 8 or 4 1 equals 8 2 and in quantity, 1 2 3 4 for as 1 is exceeded by 2 by so much 3 is exceeded by 4 or as 4 exceeds 3 so 2 exceeds 1 and by interchange as 3 exceeds 1 so 4 exceeds 2 or as 1 is exceeded by 3 by so much 2 is exceeded by 4

CHAPTER XXII

[1] The first three proportions then which are acknowledged by all the ancients Pythagoras Plato and Aristotle are the arithmetic geometric and harmonic and there are three others *subcontrary* to them which do not have names of their own but are called in more general terms the fourth fifth and sixth forms of mean after which the moderns discover four others as well making up the number ten which according to the Pythagorean view is the most perfect possible It was in accordance with this number indeed that not long ago the ten relations were observed to take their proper number the so-called ten categories the divisions and forms of the extremities of our hands and feet and countless other things which we shall notice in the proper place

[2] Now however we must treat from the beginning first that *form of proportion* which by *quantity reconciles and binds together* the comparison of the terms which is a quantitative equality as regards the difference of the several terms to one another This would be the arithmetic proportion for it was previously reported that quantity is its peculiar belonging [3] What then is the reason that we shall treat of this first and not another? Is it not clear that Nature shows it forth before the rest? For in the natural series of simple numbers beginning with 1 with no term passed over or omitted the definition of this proportion alone is preserved moreover in our previous statement we demonstrated that the *Arithmetical Introduction* itself is antecedent to all the others because it abolishes them together with itself but is not abolished together with them and because it is implied by them but does not imply them Thus it is natural that the mean which shares the name of arithmetic will not

unreasonably take precedence of the means which are named for the other sciences the geometric and harmonic for it is plain that all the more will it take precedence over the *subcontraries* over which the first three hold the leadership [4] As the first and original therefore since it is most deserving of the honor let the arithmetic proportion have its discussion at our hands before the others

CHAPTER XXIII

[1] It is an arithmetic proportion then, whenever three or more terms are set forth in succession or are so conceived and the same quantitative difference is found to exist between the successive numbers but not the same ratio among the terms, one to another For example 1 2 3 4, 5 6 7 8 9 10 11, 12 13 for in this natural series of numbers examined consecutively and without any omissions every term whatsoever is discovered to be placed between two and to preserve the arithmetic proportion to them For its differences as compared with those ranged on either side of it are equal the same ratio however is not preserved among them

[2] And we understand that in such a series there comes about both a continued and a disjunct proportion for if the same middle term answers to those on either side as both antecedent and consequent it would be a continued proportion but if there is another mean along with it a disjunct proportion comes about

[3] Now if we separate out of this series any three consecutive terms whatsoever after the form of the continued proportion or four or more terms after the disjunct form and consider them the difference of them all would be 1 but their ratios would be different throughout If however again we select three or more terms not adjacent but separated separated nevertheless by a constant interval if one term was omitted in setting down each term the difference in every case will be 2 and once more with three terms it will be a continued proportion with more disjunct If two terms are omitted the difference will always be 3 in all of them continued or disjunct if three 4 if four 5 and so on

[4] Such a proportion therefore partakes in equal quantity in its differences but of unequal quality for this reason it is arithmetic If on the contrary it partook of similar quality but not quantity it would be geometric instead of arithmetic

[5] A thing is peculiar to this proportion

that does not belong to any other namely, the mean is either half of or equal to the sum of the extremes whether the proportion be viewed as continuous or disjunct or by alternation for either the mean term with itself or the mean terms with one another are equal to the sum of the extremes

[6] It has still another peculiarity, what ratio each term has to itself, thus the differences have to the differences that is they are equal

Again the thing which is most exact and which has escaped the notice of the majority the product of the extremes when compared to the square of the mean is found to be smaller than it by the product of the differences whether they be 1 2, 3 4 or any number whatever

In the fourth place a thing which all previous writers also have noted the ratios between the smaller terms are larger as compared to those between the greater terms It will be shown that in the harmonic proportion on the contrary the ratios between the greater terms are greater than those between the smaller for this reason the harmonic proportion is subcontrary to the arithmetic and the geometric is midway between them as it were between extremes for this proportion has the ratios between the greater terms and those between the smaller equal and we have seen that the equal is in the middle ground between the greater and the less So much then about the arithmetic proportion

CHAPTER XXIV

[1] The next proportion after this one the geometric is the only one in the strict sense of the word to be called a proportion because its terms are seen to be in the same ratio It exists whenever of three or more terms as the greatest is to the next greatest so the latter is to the one following and if there are more terms as this again is to the one following it but they do not however differ from one another by the same quantity but rather by the same quality of ratio the opposite of what was seen to be the case with the arithmetic proportion

[2] For an example set forth the numbers beginning with 1 that advance by the double ratio 1 2 4 8 16 32 64 and so on or by the triple ratio 1 3 9 27 81 243 and so on or by the quadruple or in some similar way In each one of these series three adjacent terms or four or any number whatever that may be taken

will give the geometric proportion to one another as the first is to the next smaller so is that to the next smaller and again that to the next smaller and so on as far as you care to go and also by alternation For instance, 2 4 8 the ratio which 8 bears to 4 that 4 bears to 2, and conversely they do not, however have the same quantitative difference Again 2, 4 8, 16 for not only does 16 have the same ratio to 8 as before though not the same difference but also by alternation it preserves a similar relation—as 16 is to 4 so 8 is to 2 and conversely as 2 is to 8 so 4 is to 16 and disjunctly as 2 is to 4 so 8 is to 16 and conversely and in disjunct form as 16 is to 8 so 4 is to 2 for it has the double ratio

[3] The geometric proportion has a peculiar property shared by none of the rest that the differences of the terms are in the same ratio to each other as the terms to those adjacent to them the greater to the less and vice versa Still another property is that the greater terms have as a difference with respect to the lesser the lesser terms themselves and similarly difference differs from difference by the smaller difference itself if the terms are set forth in the double ratio in the triple ratio both terms and differences will have as a difference twice the next smaller in the quadruple ratio thrice in the quintuple four times and so on

[4] Geometric proportions come about not only among the multiples but also among all the superparticular, superpartient and mixed forms and the peculiar property of this proportion in all cases is preserved that in the continued proportions the product of the extremes is equal to the square of the mean but in disjunct proportions or those with a greater number of terms even if they are not continued but with an even number of terms that the product of the extremes equals that of the means

[5] As an illustration of the fact that in all the relations all kinds of multiple superparticulars superpartients and mixed ratios the peculiar property of this proportion is preserved let that suffice and be sufficient for us wherein we fashioned beginning with equality by the three rules all the kinds of inequality out of one another when they were in both direct and reverse order for each act of fashioning and each series set forth is a geometric proportion with all the aforesaid properties as well as a fourth namely that they keep the

sponse for diagnostic purposes (1175) Dehydroisoandrosterone excretion may aid in the differential diagnosis of adrenal hyperfunction (370) Similarly increased urinary pregnanediol and pregnanetriol, in cases of adrenal hyperplasia has been reduced by cortisone therapy (100) and urinary estrogens may also be reduced (371-771) The therapeutic use of cortisone to depress adrenal secretion in nonmalignant cases has been encouraging

Adrenal tumors in the mouse have been discussed in a review by Woolley (1237), and the adrenal genital relationship reviewed by Zuckerman (1262)

B OVARIAN DISTURBANCES

The use of cortisone in various ovarian disturbances and its use in cases of sterility have been described (598) ACTH caused a two- to eight fold increase in urinary output of gonadotropins in eight of nine patients treated (1042) From the studies on adrenalectomized patients it appears that the adrenals normally produce estrogens (203) In rats, cortisone caused a slight augmentation of the increase in ovarian weight after administration of chorionic gonadotrophin (330) Cortisone alone did not exert any sex hormone or gonadotrophic activity (787) Szego found that the acute stimulation of the uterus by estrogens could be antagonized by cortisone or hydrocortisone but not by DCA Such action was independent of pituitary function (1128) Others observed a similar action using ovariectomized rats (1132) In castrated male guinea pigs cortisone in small doses exerted a stimulating estrogen like action on nipple growth Larger doses however inhibited nipple growth (1202) The correlation between ovarian and adrenal cyclic function in guinea pigs has been suggested as related to the estrogen stimulation of the anterior pituitary to release ACTH (1259) In the dog cortisone did not exert androgenic effects nor did it antagonize the action of testosterone propionate on the prostate (1132) A recent detailed review containing extensive experimental data on the influence of the adrenals on the reproductive system has been published by Moore (787)

C MAINTENANCE OF LACTATION

The controversy over the effectiveness of DCA or cortisone and 11 dehydrocortisone in maintaining lactation in adrenalectomized rats has been reinvestigated by Cowie Whereas no agent alone was completely effective it was found that in the rat cortisone pellets (2 of 11 mg) and DCA (1 of 50 mg) allowed complete restoration of lactation (184) The relationship between the adrenal cortex and the mammary gland has been reviewed by Folley (333)

REFERENCES

1. Abernathy R. J. *Clin Invest* 30: 676 (1951)
2. Abernathy R. and Spink W. W. *J Clin Invest* 31: 947 (1952)

- 3 Abrams W B and Harris T A *Am Heart J* **42** 816 (1951)
- 4 Adams I and White A *Proc Soc Exptl Biol Med* **75** 570 (1950)
- 5 Adlersberg D Schaefer I L and Drachman S R *J Am Med Assoc* **144** 909 (1950)
- 6 Adlersberg D Schaefer I E and Drachman S R *J Clin Endocrinol* **11** 67 (1951)
- 7 Adlersberg D Schaefer I I and Ditch R *Proc Soc Exptl Biol Med* **74** 877 (1950)
- 8 Adlersberg D Schaefer I I and Ditch R *J Clin Endocrinol* **10** 814 (1950)
- 9 Adlersberg D Drachman S R Schaefer I I and Ditch R *J Clin Invest* **30** 676 (1951)
- 10 Agosin M Christen R Badinez O Gasic C Neghme A Iizarro O and Jarpa A *Proc Soc Exptl Biol Med* **80** 128 (1957)
- 11 Aikawa J K *Proc Soc Exptl Biol Med* **82** 105 (1953)
- 12 Anshe J D Francis T and Brown G C *J Lab Clin Med* **33** 344 (1951)
- 13 Aird R B and Gordan G S *J Am Med Assoc* **145** 715 (1951)
- 14 Albaum H G Harshfeld A I Tonhazy N L and Umbreit W W *Proc Soc Exptl Biol Med* **76** 546 (1951)
- 15 Albert A Tenney A and Ford C *Endocrinology* **50** 374 (1952)
- 16 Alrich E M Carter J P and Lehman E P *Ann Surg* **133** 783 (1951)
- 17 Alrich F M Carter J P and Lehman E P *Surgery* **32** 376 (1952)
- 18 Alpert M *Endocrinology* **45** 166 (1950)
- 19 American Trudeau Society Committee on Medical Research *Am Rev Tuberc* **65** 257 (1952)
- 20 American Trudeau Society Committee on Therapy *Am Rev Tuberc* **66** 251 (1957)
- 21 Anderson G E Wiesel L L Hillman R W and Stumpe W M *Proc Soc Exptl Biol Med* **76** 825 (1951)
- 22 Andrae W A and Browne J S L *Can Med Assoc J* **55** 423 (1946)
- 23 Antopol W *Proc Soc Exptl Biol Med* **73** 262 (1950)
- 24 Appel S B and Orr K D *U S Armed Forces Med J* **3** 201 (1957)
- 25 Aronson S M Teodoru C V Adler M and Schwartzman G *Proc Soc Exptl Biol Med* **85** 214 (1954)
- 26 Asboe Hansen G *Ann Rheumatic Diseases* **9** 149 (1950)
- 27 Asboe Hansen G *Scand J Clin & Lab Invest* **2** 271 (1950)
- 28 Asboe Hansen G *Acta Endocrinol* **9** 29 (1957)
- 29 Asboe Hansen G *Proc Soc Exptl Biol Med* **80** 617 (1952)
- 30a Asboe Hansen G *Cancer Research* **14** 94 (1954)
- 30 Ashton N and Cook C *Brit J Ophthalmol* **35** 708 (1951)
- 31 Aterman K *Lancet* **ii** 517 (1950)
- 32 Aterman K and Greenberg S M *Endocrinology* **52** 510 (1953)
- 33 Awapara J Marvin H N and Wells B B *Endocrinology* **44** 38 (1949)
- 34 Baccus H *Proc Soc Exptl Biol Med* **77** 167 (1951)
- 35 Baccus H *Endocrinology* **49** 789 (1951)
- 36 Baccus H *Endocrinology* **51** 516 (1952)
- 37 Baccus H and Altszuler N *Endocrinology* **51** 1 (1952)
- 38 Baccus H and Toompas C A *Science* **113** 260 367 (1951)
- 39 Baccus H Altszuler N and Hauffer M H *Proc Soc Exptl Biol Med* **80** 88 (1957)

sponse for diagnostic purposes (1175). Dehydroisoandrosterone excretion may aid in the differential diagnosis of adrenal hyperfunction (370). Similarly, increased urinary pregnanediol and pregnanetriol in cases of adrenal hyperplasia has been reduced by cortisone therapy (100) and urinary estrogens may also be reduced (371, 771). The therapeutic use of cortisone to depress adrenal secretion in nonmalignant cases has been encouraging.

Adrenal tumors in the mouse have been discussed in a review by Woolley (1237), and the adrenal genital relationship reviewed by Zuckerman (1262).

B. OVARIAN DISTURBANCES

The use of cortisone in various ovarian disturbances and its use in cases of sterility have been described (598). ACTH caused a two- to eight-fold increase in urinary output of gonadotropins in eight of nine patients treated (1042). From the studies on adrenalectomized patients it appears that the adrenals normally produce estrogens (203). In rats, cortisone caused a slight augmentation of the increase in ovarian weight after administration of chorionic gonadotrophin (330). Cortisone alone did not exert any sex hormone or gonadotrophic activity (787). Szego found that the acute stimulation of the uterus by estrogens could be antagonized by cortisone or hydrocortisone but not by DCA. Such action was independent of pituitary function (1128). Others observed a similar action using ovariectomized rats (1132). In castrated male guinea pigs cortisone in small doses exerted a stimulating estrogen-like action on nipple growth. Larger doses, however, inhibited nipple growth (1202). The correlation between ovarian and adrenal cyclic function in guinea pigs has been suggested as related to the estrogen stimulation of the anterior pituitary to release ACTH (1209). In the dog cortisone did not exert androgenic effects nor did it antagonize the action of testosterone propionate on the prostate (1132). A recent detailed review containing extensive experimental data on the influence of the adrenals on the reproductive system has been published by Moore (787).

C. MAINTENANCE OF LACTATION

The controversy over the effectiveness of DCA or cortisone and 11-dehydrocortisone in maintaining lactation in adrenalectomized rats has been reinvestigated by Cowie. Whereas no agent alone was completely effective it was found that in the rat cortisone pellets (2 of 11 mg) and DCA (1 of 50 mg) allowed complete restoration of lactation (184). The relationship between the adrenal cortex and the mammary gland has been reviewed by Folley (333).

REFERENCES

1. Abernathy R. *J. Clin. Invest.* 30: 626 (1951).
2. Abernathy R. and Spink W. W. *J. Clin. Invest.* 31: 947 (1952).

- 3 Abrams W B and Harris T N *Am Heart J* 42 56 (1951)
- 4 Adams F and White A *Proc Soc Exptl Biol Med* 75 190 (1950)
- 5 Allersberg D Schaefer I I and Drachman S R *J Clin Med* 11 99 (1950)
- 6 Allersberg D Schaefer I I and Drachman S R *J Clin Endocrinol* 11 61 (1951)
- 7 Allersberg D Schaefer I I and Ditch R *Proc Soc Exptl Biol Med* 71 5 (1950)
- 8 Allersberg D Schaefer I I and Ditch R *J Clin Endocrinol* 10 814 (1950)
- 9 Allersberg D Drachman S R Schaefer I I and Ditch R *J Clin Invest* 30 676 (1951)
- 10 Aguirre M Christen R Badinez O Case C Noguera A Lizarri O and Jarpa A *Proc Soc Exptl Biol Med* 80 125 (1953)
- 11 Aikawa J K *Proc Soc Exptl Biol Med* 82 105 (1953)
- 12 Anshel J D Francis T and Brown C C *J Lab Clin Med* 38 314 (1951)
- 13 Ard R B and Cordan G S *J Am Med Assoc* 145 15 (1951)
- 14 Albano H G Hirschfeld A I Tonhazy N I and Liment W W *Proc Soc Exptl Biol Med* 75 516 (1951)
- 15 Albert A Tenney A and Lord I *Endocrinology* 50 34 (1952)
- 16 Alrich F M Carter J I and Lehman I I *Ann Surg* 133 783 (1951)
- 17 Alrich F M Carter J I and Lehman I I *Surgery* 32 356 (1952)
- 18 Apert M *Endocrinology* 46 166 (1950)
- 19 American Trudeau Society Committee on Medical Research *Am J Med* 11 66 25 (1952)
- 20 American Trudeau Society Committee on Therapy *Am Rev Tuberc* 66 254 (1952)
- 21 Anderson C F Wiesel I I Hillman R W and Stumpe W M *Proc Soc Exptl Biol Med* 78 825 (1951)
- 22 Andress W A and Browne J S L *Can Med Assoc J* 55 423 (1946)
- 23 Antopol W *Proc Soc Exptl Biol Med* 73 762 (1950)
- 24 Appel S B and Orr H D *US Armed Forces Med J* 3 207 (1952)
- 25 Aronson S M Teodoru C V Adler M and Schwartzman C *Proc Soc Exptl Biol Med* 85 214 (1954)
- 26 Ashoe Hansen G *Ann Rheumatic Diseases* 9 141 (1950)
- 27 Ashoe Hansen G *Scand J Clin Lab Invest* 2 271 (1950)
- 28 Ashoe Hansen G *Acta Endocrinol* 9 29 (1952)
- 29 Ashoe Hansen G *Proc Soc Exptl Biol Med* 80 677 (1952)
- 30 Ashoe Hansen G *Cancer Research* 14 91 (1954)
- 31 Ashton N and Cook C *Brit J Ophthalmol* 35 708 (1951)
- 32 Aterman H *Lancet* II 517 (1950)
- 33 Aterman H and Greenberg S M *Endocrinology* 52 510 (1953)
- 34 Awapara J Marvin H N and Wells B B *Endocrinology* 44 38 (1949)
- 35 Baccus H *Proc Soc Exptl Biol Med* 77 167 (1951)
- 36 Baccus H *Endocrinology* 49 789 (1951)
- 37 Baccus H *Endocrinology* 51 516 (1952)
- 38 Baccus H and Altszuler N *Endocrinology* 51 1 (1952)
- 39 Baccus H and Toompas C A *Science* 113 269 367 (1951)
- 40 Baccus H Altszuler N and Heffer M H *Proc Soc Exptl Biol Med* 80 83 (1952)

- 40 Bacehus H, Heffer M H and Mitzuler N *Proc Soc Exptl Biol Med* 79 619 (1952)
- 41 Badenoch J *Brit Med J* 1 306 (1952)
- 42 Bagnall A W *Can Med Assoc J* 65 125 (1951)
- 43 Baker B I *Ann N Y Acad Sci* 63 690 (1951)
- 44 Baker B I *Ann N Y Acad Sci* 66 684 (1953)
- 45 Baker B I and Scharrer M A *Proc Soc Exptl Biol Med* 82 235 (1953)
- 46 Baker B I and Whitaker W I *Am J Physiol* 169 118 (1945)
- 47 Baker B I and Whitaker W I *Endocrinology* 46 544 (1950)
- 48 Baker B I, Ingle D J, In C H and Evans H M *Anat Record* 102 313 (1948)
- 49 Baldrige G D, Kligman A M, Lipnik M J and Pillsbury D M *Arch Pathol* 61 593 (1951)
- 50 Balmain J H and Folley S J *Arch Biochem* 39 188 (1952)
- 51 Balmain J H, Folley S J and Glascock R F *Nature* 169 447 (1952)
- 52 Banerjee S and Deb C *J Biol Chem* 190 177 (1951)
- 53 Banerjee S and Deb C *Endocrinology* 51 572 (1952)
- 54 Banerjee S and Deb C *J Biol Chem* 194, 515 (1952)
- 55 Bangham A D *Brit J Exptl Pathol* 32 77 (1951)
- 56 Barber A and Nothacker W C *Arch Pathol* 64 334 (1952)
- 57 Barberio J R, Patc J W, Sawyer I N and Hufnagel C A *Surgery* 33 827 (1953)
- 58 Barton A D *Federation Proc* 10 160 (1951)
- 59 Bartter I C, Albright F, Forbes A I, Leaf A, Dempsey I and Carroll I *J Clin Invest* 30 237 (1951)
- 60 Baschieri I and Rossi A *Folia Endocrinol* 3 57 (1950)
- 60a Baserga R and Shubik P *Cancer Research* 14 12 (1954)
- 61 Baumann I J and Marine D *Endocrinology* 36 400 (1945)
- 62 Beck L V *Cancer Research* 11 235 (1951)
- 63 Beck I V and Voloshin T *Am J Physiol* 163 696 (1950)
- 64 Beierwaltes W H, Wolfson W Q, Jones J R, Knorpp C T and Siemieniski J S *J Lab Clin Med* 36 799 (1950)
- 65 Bell C I, Bell R I and Wilson D R *Can Med Assoc J* 63 63 (1950)
- 66 Benditt L I, Schuller S, Wong H and Dorfman A *Proc Soc Exptl Biol Med* 75 782 (1950)
- 67 Benedict W H and Hollenhorst R W *Am J Ophthalmol* 36 247 (1953)
- 68 Benedict J D, Forsham P H, Roche M, Soloway S and Stetten D Jr *J Clin Invest* 29 1104 (1950)
- 69 Benjamin F B, Cornbleet T and Grossman M I *J Invest Dermatol* 19 109 (1952)
- 70 Bennett H S *Proc Soc Exptl Biol Med* 42 786 (1939)
- 71 Bennett H S *Am J Anat* 67 151 (1940)
- 72 Bennett I I, Garcia J I and In C H *Proc Soc Exptl Biol Med* 69 52 (1948)
- 73 Bennett I I, Liddle C W and Bentinck R C *J Clin Endocrinol and Metabolism* 13 392 (1953)
- 74 Bennett I I, Slessor A and Thorn G W *J Clin Endocrinol and Metabolism* 9 675 (1949)
- 75 Bennett L L, Kreiss R F, Li C H and Evans H M *Am J Physiol* 152 210 (1948)

- 6 Bennett W A 1m *J Pathol* 27 704 (1951)
- 7 Bennett W A and Kilby R A 1m *J Pathol* 29 586 (1953)
- 8 Bentinck R C Cordan G S Adams J F Arnstein I H and Leake T B
J Clin Invest 30 200 (1951)
- 9 Berlin B S Johnson C Hawk W D and Lawrence A C *J Lab Clin Med* 40 82 (1952)
- 80 Bernstein D E *Proc Soc Exptl Biol Med* 73 175 (1950)
- 81 Bernstein D E *Proc Soc Exptl Biol Med* 75 847 (1950)
- 82 Berson S A and Yalow I S *J Clin Endocrinol and Metabolism* 12 407 (1952)
- 83 Best W R and Samter M *Blood* 6 61 (1951)
- 84 Best W R Muchrecke R C and Kark R H *J Clin Invest* 31 733 (1952)
- 85 Bibile S W *J Endocrinol* 9 357 (1953)
- 85a Bien F J and Liff M *Proc Soc Exptl Biol Med* 78 377 (1951)
- 86 Billingham R E Krohn P L and Medawar P B *Brit Med J* II 1049 (1951)
- 87 Bisio A *J Clin Endocrinol and Metabolism* 13 1226 (1953)
- 88 Björneboe M Fischel F E and Stoerk H C *J Exptl Med* 93 37 (1951)
- 89 Blalock A and Levy S F *Ann Surg* 106 876 (1937)
- 90 Bloodworth J M B *Endocrinology* 50 174 (1952)
- 91 Bloom F *Proc Soc Exptl Biol Med* 79 651 (1952)
- 92 Bloom B and Pierce F T *Metabolism Clin and Exptl* 1 155 (1952)
- 93 Blunt J W Plotz C M Lattes R Howes F I Meyer K and Ragan C *Proc Soc Exptl Biol Med* 73 678 (1950)
- 94 Boland F W and Headley N T *J Am Med Assoc* 141 301 (1949)
- 95 Boland E W and Headley N T *J Am Med Assoc* 148 981 (1952)
- 96 Bondy P K *Endocrinology* 45 605 (1949)
- 97 Bondy P K Engel F I and Farrar B *Endocrinology* 44 46 (1949)
- 98 Bondy P K and Hagewood M A *Proc Soc Exptl Biol Med* 81 38 (1952)
- 99 Bondy P K and Wilhelm A E *J Biol Chem* 186 245 (1950)
- 100 Bongiovanni A M *Bull Johns Hopkins Hosp* 92 244 (1953)
- 101 Bonner C D Fishman W H and Homburger F *Geriatrics* 5 203 (1950)
- 102 Booker W M Dent F M Hays R L Harris W and Green S *Proc Soc Exptl Biol Med* 78 110 (1951)
- 103 Bookman J J Drachman S R Schaefer I E and Adlersberg D *Diabetes* 2 100 (1953)
- 104 Borden A L Brodie F C Wallraff E B Holbrook W P Hill D F Stephens C A I Johnson R B and Kemmerer A R *J Clin Invest* 37 315 (1952)
- 105 Boss W R Birnie J H and Gaunt R *Endocrinology* 46 307 (1950)
- 106 Boss W R Osborn C M and Renzi A A *Endocrinology* 51 66 (1952)
- 107 Bourne G H *J Anat* 83 10 (1949)
- 108 Bowers J Z Nelson D H Bay R and Samuels L T *J Clin Endocrinol and Metabolism* 12 971 (1952)
- 109 Braun Menendez E and Covian M P *Per soc argentina biol* 24 31 (1948)
- 110 Bristol W R and Drill A A *Endocrinology* 50 671 (1952)
- 111 Brodie E C Wallraff E B Borden A L Holbrook W P Stephens C A I Hill D F Kent I J and Kemmerer A R *Proc Soc Exptl Biol Med* 75 285 (1950)

- 40 Beechus H Heffer M H and Mtszuler N *Proc Soc Exptl Biol Med* **79** 648 (1952)
- 41 Badenoch J *Brit Med J* **1**, 356 (1952)
- 42 Bagnall A W *Can Med Assoc J* **65** 125 (1951)
- 43 Baker B I *Ann N Y Acad Sci* **53** 640 (1951)
- 44 Baker B I *Ann N Y Acad Sci* **56** 684 (1953)
- 45 Baker B I and Scharrer M A *Proc Soc Exptl Biol Med* **82** 735 (1953)
- 46 Baker B I and Whitaker W I *Am J Physiol* **159** 118 (1949)
- 47 Baker B I and Whitaker W I *Endocrinology* **46** 544 (1950)
- 48 Baker B I Ingle D J Jr, C H and Evans H M *Anat Record* **102**, 313 (1948)
- 49 Baldrige G D Kligman A M Lipnik M J and Pillsbury D M *Arch Pathol* **51** 503 (1951)
- 50 Balmain J H and Folley S J *Arch Biochem* **39** 188 (1952)
- 51 Balmain J H Folley S J and Glascock R F *Nature* **169** 447 (1952)
- 52 Banerjee S and Deb C *J Biol Chem* **190** 177 (1951)
- 53 Banerjee S and Deb C *Endocrinology* **51** 52 (1952)
- 54 Banerjee S and Deb C *J Biol Chem* **194** 575 (1952)
- 55 Bingham A D *Brit J Exptl Pathol* **32** 77 (1951)
- 56 Barber A and Nothacker W C *Arch Pathol* **54** 334 (1952)
- 57 Barberio J R Late J W Sawyer L N, and Hufnagel C A *Surgery* **33** 877 (1953)
- 58 Barton A D *Federation Proc* **10** 160 (1951)
- 59 Bartter I C Albright F Forbes A I Leif A Dempsey F and Carroll L *J Clin Invest* **30** 237 (1951)
- 60 Baehre I and Rossi A *Folia Endocrinol* **3** 57 (1950)
- 60a Baserga R and Shubik I *Cancer Research* **14** 12 (1954)
- 61 Baumann E J and Marine D *Endocrinology* **36** 400 (1945)
- 62 Beck I V *Cancer Research* **11**, 235 (1951)
- 63 Beck I V and Voloshin T *Am J Physiol* **163** 696 (1950)
- 64 Beierwaltes W H Wolfson W Q Jones J R Knorpp C T and Siemieniski J S *J Lab Clin Med* **36** 799 (1950)
- 65 Bell G I Bell R I and Wilson D R *Can Med Assoc J* **63** 63 (1950)
- 66 Benditt I I Schiller S Wong H and Dorfman A *Proc Soc Exptl Biol Med* **75** 782 (1950)
- 67 Benedict W H and Hollenhorst R W *Am J Ophthalmol* **36** 247 (1953)
- 68 Benedict J D Forsham P H Roche M Soloway S and Stetten D Jr *J Clin Invest* **29** 1104 (1950)
- 69 Benjamin F B Cornbleet T and Grossman M I *J Invest Dermatol* **19** 109 (1952)
- 70 Bennett H S *Proc Soc Exptl Biol Med* **42** 786 (1939)
- 71 Bennett H S *Am J Anat* **67** 151 (1940)
- 72 Bennett L I Garcia J F and Ii C H *Proc Soc Exptl Biol Med* **69** 52 (1948)
- 73 Bennett L I Iiddle G W and Bentinck R C *J Clin Endocrinol and Metabolism* **13** 302 (1953)
- 74 Bennett L I Slessor A and Thorn C W *J Clin Endocrinol and Metabolism* **9** 615 (1949)
- 75 Bennett L L Kreiss R E Ii C H and Evans H M *Am J Physiol* **152** 210 (1948)

- 151 Clark I *Federation Proc* **9** 161 (1950)
- 152 Clark I *J Biol Chem* **200** 69 (1952)
- 153 Clark I D Quirton C C Cobb S and Bauer W *New Engl J Med* **249** 178 (1953)
- 154 Clayton B I and Lurmy I I C *Brit Med J* **41** 977 (1951)
- 155 Clayton B I and Lurmy I T C *J Endocrinol* **9** 340 (1952)
- 156 Cleghorn R A *J Clin Endocrinol and Metabolism* **13** 1701 (1953)
- 157 Cleghorn R A Fowler J I A Greenwood W F and Clarke A P W *Am J Physiol* **161** 21 (1950)
- 158 Cochrane C C Jahn J I Foreman N and Kinsell I W *J Clin Endocrinol and Metabolism* **13** 993 (1953)
- 159 Cohen R S *Am J Anat* **56** 143 (1952)
- 160 Cohn C Katz B Huddleston B Kolinsky M and Levine I *Am J Physiol* **170** 87 (1952)
- 161 Cole D F *J Endocrinol* **6** 215 (1950)
- 162 Cole D F *J Endocrinol* **6** 251 (1950)
- 163 Cole D F *Acta Endocrinol* **14** 245 (1953)
- 164 Cole J W and Holden W D *Proc Soc Exptl Biol Med* **77** 363 (1951)
- 165 Cole J W Orbison J I Holden W D Hancock T J and Lindsay J F *Surg Gynecol Obstet* **93** 321 (1951)
- 166 Collins D A and Wood L H *Am J Physiol* **123** 224 (1955)
- 167 Conn J W *Arch Intern Med* **83** 416 (1949)
- 168 Conn J W and Fajans S S *Ann Rev Physiol* **14** 453 (1952)
- 169 Conn J W and Louis I H *J Clin Endocrinol and Metabolism* **10** 12 (1950)
- 170 Conn J W Fajans S S Louis I H and Johnson B *J Lab Clin Med* **36** 813 (1950)
- 171 Conn J W Johnston M W and Louis I H *J Clin Invest* **25** 917 (1946)
- 172 Conn J W Louis L H and Fajans S S *Science* **113** 113 (1951)
- 173 Conn J W Louis I H and Johnston M W *Science* **109** 219 (1949)
- 174 Conn J W Louis I H and Johnston M W *J Lab Clin Med* **34** 255 (1949)
- 175 Conn J W Louis L H and Johnston M W *J Clin Invest* **28** 775 (1949)
- 176 Conn J W Louis L H and Wheeler C L *J Lab Clin Med* **33** 651 (1948)
- 177 Conn J W Louis L H Fajans S S Johnson B J Barott G W Hogg L W Blood J I and Weiss M *J Lab Clin Med* **38** 799 (1951)
- 178 Cook C and MacDonald R K *Brit J Ophthalmol* **35** 130 (1951)
- 179 Cook D I Ray R Davidson E Feldstein L M Calvin L D and Green D M *J Exptl Med* **96** 27 (1952)
- 180 Cosniff S W Diefenthal A F and Vogt W Jr *Am J Med* **9** 752 (1950)
- 181 Coste F Delbarre F and Bassett G *Rev rhumat* **19** 327 (1952)
- 182 Coste F Piquet B Gorniche P and Cayala J *Ann med* **52** 747 (1951)
- 183 Courier R and Collonge A *J Am Med Assoc* **146** 493 (1951)
- 184 Cowi A T *Endocrinology* **51** 217 (1950)
- 185 Cowie A T and Stewart J *J Endocrinol* **6** 191 (1949)
- 186 Crabb E D and Kelsall M A *J Natl Cancer Inst* **12** 91 (1951)
- 187 Crag J M *Am J Pathol* **28** 629 (1952)
- 188 Crandon J H Lund C C and Dill D B *New Engl J Med* **223** 353 (1940)

- 112 Brody S *Psychosom Med* 14 91 (1952)
- 113 Broth Kahn R H and Mirsky I A *Science* 106 148 (1941)
- 114 Brown J H U *Fraternation Proc* 12 21 (1953)
- 115 Brown H Hargreaves H P and Tyler F H *Arch Intern Med* 89 951 (1952)
- 116 Brownlee C *Lancet* 1 157 (1950)
- 117 Buchwald H W Hudson I and Bellanca J *Endocrinology* 47 228 (1950)
- 118 Bullough W S *J Endocrinol* 8 261 (1952)
- 119 Burchenal J H Stock C C and Rhoads C P *Cancer Research* 10 709 (1950)
- 120 Burchenal J H Webber I F and Johnston S I *Proc Soc Exptl Biol Med* 78 352 (1951)
- 121 Burkell C C *J Can Assoc Radiologists* 3 30 (1952)
- 122 Bush I F *J Physiol* 115 12 (1951)
- 123 Bush I L *J Endocrinol* 9 95 (1953)
- 124 Butcher I O *Endocrinology* 43 30 (1948)
- 125 Butler I C and Morgan A F *Proc Soc Exptl Biol Med* 83 655 (1953)
- 126 Cagin R N Gray J I and Jensen H *J Biol Chem* 183 11 (1950)
- 127 Cagin R N Klein R I and Loewe I *J Clin Endocrinol and Metabolism* 13 429 (1952)
- 128 Cameron A T and Carmichael J *Can J Research* 24 37 (1946)
- 129 Cameron C R In J M Yoffey *The Suprarenal Cortex* Academic Press New York 1953 p 155
- 130 Cannon J A and Longmire W P *Ann Surg* 135 60 (1952)
- 131 Cape R D T Thomas J W and Palmer R A *Can Med Assoc J* 66 441 (1952)
- 132 Carroll H K *Endocrinology* 48 101 (1951)
- 133 Carroll H K *J Biol Chem* 200 287 (1953)
- 134 Carroll H K and Noble R I *Endocrinology* 51 476 (1952)
- 135 Carroll H K McWipine H T and Noble R I *Can Med Assoc J* 65 363 (1951)
- 136 Carrier H M and Code C F *J Allergy* 21 258 310 (1950)
- 137 Castor C W and Baker B I *Endocrinology* 47 231 (1950)
- 138 Castor C W Baker B I Ingle D J and Ito C H *Proc Soc Exptl Biol Med* 76 353 (1951)
- 139 Cavallero C and Braccini C *Proc Soc Exptl Biol Med* 78 141 (1951)
- 140 Cavallero C and Silva C *Lancet* 1 175 (1951)
- 141 Cavallero C Di Marco A Lucio I and Silva G *Proc Soc Exptl Biol Med* 81 619 (1952)
- 142 Chalmers T M and Lewis A A G *Lancet* 1 1158 (1951)
- 143 Chapman D W Slaggs R H Thomas J R and Green J A *Am J Med Sci* 223 41 (1952)
- 144 Chase H B *Physiol Revs* 34 113 (1954)
- 145 Chen C and Wickel A *Endocrinology* 51 51 (1952)
- 146 Chen C I and Sayers C *Endocrinology* 44 400 (1949)
- 147 Chesterton F C *Lancet* 1 1953 (1953)
- 148 Chiu C Y *Biochem J* 46 120 (1950)
- 149 Chiu C Y and Neillham D M *Biochem J* 46 114 (1950)
- 150 Clark D F Nelson T S and Raiman R J *J Am Med Assoc* 151 551 (1953)

- 225 Dempsey E W Greep R O and Deane H W *Endocrinology* 44 88 (1949)
- 226 Derbes V J and Weiss T *Quart Rev Allergy and Appl Immunol* 6 153 (1951)
- 227 Derbes V J Dent J H Weaver V K and Vaughn D D *Proc Soc Exptl Biol Med* 75 423 (1950)
- 228 Derranis A Scifter J and Ehrlich W E *Federation Proc* 11 338 (1952)
- 229 Desaulles P Tripodi J and Schuler W *Schweiz med Hochschr* 83 1088 (1953)
- 230 Devitt J E Pirozynski W J and Samuels P B *Proc Soc Exptl Biol Med* 83 335 (1953)
- 231 DeVries J A *J Immunol* 65 1 (1950)
- 232 Dews P B and Code C F *Proc Soc Exptl Biol Med* 77 141 (1951)
- 233 Dews P B and Code C F *J Immunol* 70 199 (1953)
- 234 Dhyse F G Fisher C P Tullner W W and Hertz R *Endocrinology* 53 447 (1953)
- 235 Diamond M C *Am J Physiol* 171 719 (1952)
- 236 Dickel H *Munch med Hochschr* 93 1 (1951)
- 237 Diezfelussy E and Westman A *Lancet* II 41 (1950)
- 238 Dillaha C J and Rothman S *J Invest Dermatol* 18 5 (1952)
- 239 Dillaha C J and Rothman S *Am Practitioner* 3 616 (1952)
- 240 Dillaha C J and Rothman S *J Am Med Assoc* 150 546 (1952)
- 241 Diller I C Beck L V and Blanch B *Cancer Research* 8 581 (1948)
- 242 Doerner A A Naegle C F Regan F D Shanaphy J F and Edwards W B *J Am Med Assoc* 147 1099 (1951)
- 243 Donald T C and Higgins G M *Cancer Research* 11 907 (1951)
- 244 Done A H Ely R S Raile R B and Kelley V C *Proc Soc Exptl Biol Med* 81 667 (1952)
- 245 Dorfman A Ann A I *Acad Sci* 56 698 (1953)
- 246 Dorfman A Apter N S Smull K Bergenstal D M and Richter R B *J Am Med Assoc* 146 25 (1941)
- 247 Dorfman P I Ann A I *Acad Sci* 50 506 (1949)
- 248 Dorfman R I *Proc Soc Exptl Biol Med* 70 732 (1949)
- 249 Dorfman R I *Recent Progr Hormone Research* 8 87 (1952)
- 250 Dorfman R I *Physiol Revs* 34 138 (1954)
- 251 Dorfman R I Potts A M and Feil M S *Endocrinology* 41 464 (1947)
- 252 Dorfman R I Ross E and Shipley R A *Endocrinology* 38 178 (1946)
- 253 Dougherty T F *Am J Pathol* 27 714 (1951)
- 254 Dougherty T F *Recent Progr Hormone Research* 7 307 (1952)
- 255 Dougherty T F Ann A I *Acad Sci* 56 748 (1953)
- 256 Dougherty T F and Kumagai L F *Endocrinology* 48 691 (1951)
- 257 Dougherty T F and Schneebehl G L *Proc Soc Exptl Biol Med* 75 854 (1950)
- 258 Dougherty T F and White A *Endocrinology* 35 1 (1944)
- 259 Dougherty T F Chase J H and White A *Proc Soc Exptl Biol Med* 58 135 (1945)
- 260 Doyle L P Andrews F M and Hutchings L M *Proc Soc Exptl Biol Med* 74 373 (1950)
- 261 Draper H H and Johnson B C *Proc Soc Exptl Biol Med* 82 73 (1953)
- 262 Drury M I Hickey M D and Malone J P *Brit Med J* II 1487 (1951)

- 189 Crepea S B Mignin G I and Scastone C A *Proc Exptl Biol Med* 77 701 (1951)
- 190 Cronkite I I and Chapman W H *Proc Soc Exptl Biol Med* 74 33 (1950)
- 191 Cummings M M and Hudgins I C *Diseases of the Chest* 22 289 (1952)
- 192 Cummings M M and Hudgins P C *J Immunol* 69 331 (1952)
- 193 Cummings M M Hudgins P C Wharton M C and Sheldon W H *Am Rev Tuberc* 65 596 (1952)
- 194 Curtiss P H and Wilson P D *Surg Gynecol Obstet* 96 150 (1953)
- 195 Daft I S and Sebrell W H *Public Health Reps (U S)* 64 2 17 (1939)
- 196 Dalton A J Morris H P and Dubnik C S *J Natl Cancer Inst* 5 451 (1945)
- 197 Dameron J T *Proc Soc Exptl Biol Med* 73 343 (1950)
- 198 Danford I A and Danford H G *Am J Physiol* 164 690 (1951)
- 199 Danford, H G and Danford I A *Am J Physiol* 166 521 (1951)
- 200 D Angelo S A Cordon A S and Charipper H A *Proc Soc Exptl Biol Med* 68 527 (1948)
- 201 Danowski T S Tarant R Peters J H Weigand F A Mateer, F M and Greenman I *Proc Soc Exptl Biol Med* 81 445 (1952)
- 202 Danto J I and Maddin S *J Invest Dermatol* 18 351 (1952)
- 203 Dao T I Y *Science* 118 21 (1953)
- 204 Davenport H W and Chavre V J *Endocrinology* 47 193 (1950)
- 205 Davis A K Bass A C and Overman R R *Am J Physiol* 166 493 (1951)
- 206 Davis J O and Howell D S *Endocrinology* 62 245 (1953)
- 207 Davis M L Plotz I J and Plotz I *Endocrinology* 62 161 (1953)
- 208 Davis M L and Plotz I J *Endocrinology* 64 381 (1954)
- 209 De Andino A M Rivero Fontan J I and Paschalis K L *Proc Soc Exptl Biol Med* 77 700 (1951)
- 210 Deane H W and Greep R O *Endocrinology* 41 243 (1947)
- 211 Deane H W and McHibbin J M *Endocrinology* 38 380 (1946)
- 212 Deane H W and Seligman A M *Vitamins and Hormones* 11 1,3 (1953)
- 213 DeBodo R C and Sinkoff M W *Recent Progr Hormone Research* 8 511 (1953)
- 214 DeBodo R C Kurtz M Sinkoff M W and Hwang S P *Proc Soc Exptl Biol Med* 80 345 (1952)
- 215 DeBodo R C Sinkoff M W and Hwang S P *Proc Soc Exptl Biol Med* 80 350 (1952)
- 216 DeBodo R C Sinkoff M W Hwang S P and Den H *Proc Soc Exptl Biol Med* 81 470 (1952)
- 217 Debre R Mozziconacci P and Nekhorochell I *Presse med* 60 502 (1952)
- 218 DeCosta F J and Abelman M A *Am J Obstet Gynecol* 64 746 (1952)
- 219 DeCosta F J and Abelman M A *Obstet and Gynecol* 1 269 (1953)
- 220 DeIamater L D Saurino V R and Urbach F *Am J Syphilis Gonorrhea Venereal Diseases* 36 127 (1952)
- 221 Demartini F Grokoest A W and Ragan C *J Am Med Assoc* 149 750 (1952)
- 222 Deming Q B and Luetscher J A Jr *Proc Soc Exptl Biol Med* 73 171 (1950)
- 223 DeMowbray R R and Bishop P M F *Brit Med J* 1 17 (1953)
- 224 DeMoor P *Ann Endocrinol* 12 1073 (1952)

- 303 Engel G L and Romano J *Arch Neurol Psychiat* **51** 348 (1944)
- 303a Engellbreth Holm J and Asboe Hansen C *Acta Pathol Microbiol Scand* **32** 560 (1953)
- 304 Engstrom W W and Liebman A *Am J Med* **15** 180 (1953)
- 305 Ershoff B H *Proc Soc Exptl Biol Med* **78** 836 (1951)
- 306 Ershoff B H *Metabolism Clin and Exptl* **2** 9 (1953)
- 307 Essellier A F and Wagner K F *Acta Haematol* **8** 63 (1952)
- 308 Essellier A F Jenneret I Kopp F and Morandi I *Endocrinology* **54** 474 (1954)
- 309 Etheridge L M and Hoch Ingels C *Am J Pathol* **28** 315 (1952)
- 310 Evans V J *J Natl Cancer Inst* **14** 741 (1953)
- 311 Everall J and Fisher I *J Invest Dermatol* **19** 97 (1952)
- 312 Fahey J I *Proc Soc Exptl Biol Med* **77** 491 (1951)
- 313 Fajans S S Louis I H and Conn J W *J Clin Endocrinol and Metabolism* **11** 455 (1951)
- 314 Falk M S Allende M F and Bennett J H *J Invest Dermatol* **18** 304 (1952)
- 315 Farnsworth E B *J Clin Endocrinol and Metabolism* **13** 1169 (1953)
- 316 Feldman J D *Endocrinology* **46** 552 (1950)
- 317 Feldman J D Rachmiewitz M Stein O and Stein Y *Blood* **8** 342 (1953)
- 318 Fienberg R and Colpoys F I *Am J Pathol* **27** 925 (1951)
- 319 Finch S C Crockett C I Ross J F and Bayles T B *Blood* **6** 1034 (1951)
- 320 Findlay C W and Howes L I *New Engl J Med* **246** 597 (1952)
- 321 Finerty J C and Ianos T C *Proc Soc Exptl Biol Med* **76** 83 (1951)
- 322 Fingal E Olsen I J Harding E W Cockett A T and Goodman L S *J Pharmacol Exptl Therap* **105** 37 (1952)
- 323 Fischel L E *Bull N Y Acad Med* **26** 255 (1950)
- 324 Fischel L E LeMay M and Kabat L A *J Immunol* **61** 89 (1949)
- 325 Fischel E F Stoerk H C and Bjørneboe M *Proc Soc Exptl Biol Med* **77** 111 (1951)
- 326 Fischel L E Vaughan J H and Photopoulos C *Proc Soc Exptl Biol Med* **81** 344 (1952)
- 327 Flanagan J B Davis A K and Overman R R *Am J Physiol* **160** 89 (1950)
- 328 Fletcher A A Dauphinee J A and Ogryzlo M A *J Clin Invest* **31** 561 (1952)
- 329 Flink E B and Thorn G W *J Clin Endocrinol and Metab* **9** 660 (1949)
- 330 Fluhmann C F *Proc Soc Exptl Biol Med* **80** 504 (1951)
- 331 Foley I J *Proc Soc Exptl Biol Med* **80** 669 (1952)
- 332 Foley E J and Silverstein R *Proc Soc Exptl Biol Med* **77** 713 (1951)
- 333 Foley S J In J M Loffey *The Suprarenal Cortex* Academic Press New York 1953 p 85
- 334 Foley S J and Greenbaum A L *Biochem J* **60** 40 46 (1946)
- 335 Foley S J and Watson S C *Proc Soc Exptl Biol Med* **78** 443 (1951)
- 336 Foliss R H *Proc Soc Exptl Biol Med* **76** 722 (1951)
- 337 Foliss R H *Proc Soc Exptl Biol Med* **78** 123 (1951)
- 338 Foliss R H *Bull Johns Hopkins Hosp* **90** 337 (1952)

- 263 Duff I F Robinson W D and Smith I M *J Lab Clin Med* 38 803 (1951)
- 264 Duffy B J and Morgan H R *Proc Soc Exptl Biol Med* 78 687 (1951)
- 265 Dugil I I and Ichlune J *Acta can biol* 8 440 (1949)
- 266 Dugil I I and Thurnen M *Endocrinology* 44 470 (1949)
- 267 Dumm M F and Rall H I *Metabolism (lin and Exptl)* 2 153 (1953)
- 268 Dumm M I (erschberg H Beck I M and Rall F I *Proc Soc Exptl Biol Med* 82 659 (1953)
- 269 Durgin M I and Meyer R K *Endocrinology* 48 518 (1951)
- 270 Dury A *Am J Physiol* 160 73 (1950)
- 271 Dury A *Proc Soc Exptl Biol Med* 77 193 (1951)
- 272 Dury A and Johnston T N *Proc Soc Exptl Biol Med* 78 425 (1951)
- 273 Dury A and Moss I D *Proc Soc Exptl Biol Med* 80 199 (1952)
- 274 Dworetzky M Code C F and Higgins C M *Proc Soc Exptl Biol Med* 75 201 (1950)
- 275 Eastcott H H G Lawcett J K and Rob C G *Lancet* i 1068 (1953)
- 276 Ebert R H *J Clin Invest* 30 636 (1951)
- 277 Ebert R H *Am Rev Tuberc* 65 64 (1952)
- 278 Ebert R H and Barclay W R *Ann Internat Med* 37 506 (1952)
- 279 Edelmann A *Federation Proc* 12 37 (1953)
- 280 Eichman P I Havens W P Jr and Miller R W *J Clin Endocrinol and Metabolism* 13 648 (1953)
- 281 Eklves K Nelson D H and Samuels I T *J Clin Endocrinol and Metabolism* 13 1250 (1953)
- 282 Fischel A M and Kochakian C D *Proc Soc Exptl Biol Med* 85 339 (1954)
- 283 Eisen H V Mayer M M Moore D H Tarr R and Stoerk H C *Proc Soc Exptl Biol Med* 65 301 (1947)
- 284 Eisenberg E Gordan G S Elliott H W and Talbot J *Proc Soc Exptl Biol Med* 73 140 (1950)
- 285 Eisenstein A B and Boniface J *Federation Proc* 11 20 (1952)
- 286 Eisenstein A B and Shank R L *Proc Soc Exptl Biol Med* 78 619 (1951)
- 287 Eliel I P and Pearson O H *N J State J Med* 51 1839 (1951)
- 288 Eliel I P Pearson O H Katz B and Krautitz F W *Federation Proc* 9 168 (1950)
- 289 Ellinger F *Proc Soc Exptl Biol Med* 64 31 (1947)
- 290 Ellinger F *Proc Soc Exptl Biol Med* 80 215 (1952)
- 291 Ellinger F *Proc Soc Exptl Biol Med* 80 214 (1952)
- 292 Elliott K A C and Yarrarazaval S *Nature* 169 416 (1952)
- 293 Elmajian F and Pineus G *J Clin Endocrinol and Metabolism* 6 266 (1946)
- 294 Elster S K and Lowry C I *Proc Soc Exptl Biol Med* 73 49 (1950)
- 295 Engel F I *Endocrinology* 49 127 (1951)
- 296 Engel F I *Recent Progr Hormone Research* 6 277 (1951)
- 297 Engel F L *Endocrinology* 49 538 (1951)
- 298 Engel F L *Endocrinology* 50 462 (1952)
- 299 Engel F I and Schwartz T B *Proc Soc Exptl Biol Med* 77 615 (1951)
- 300 Engel F I and Scott J I Jr *Endocrinology* 48 56 (1951)
- 301 Engel F I Schuller S and Pentz F I *Endocrinology* 44 458 (1949)
- 302 Engel G L and Margolin S *Arch Internat Med* 70 236 (1952)

- 309 Gabrilove J L Dorrance W R and Soffer L J *Am J Physiol* **169** 565 (1957)
- 310 Gardner L I and Migeon C J *J Clin Endocrinol and Metabolism* **12** 1117 (1957)
- 311 Gastineau C F Logan G B Albert A and Mason H I *J Clin Endocrinol and Metabolism* **13** 724 (1953)
- 312 Gaudino N M *Rev soc argentina biol* **20** 470 (1941)
- 313 Gaudino N M and Levitt M F *J Clin Invest* **28** 1484 (1949)
- 314 Gaunt R *Recent Progr Hormone Research* **6** 247 (1951)
- 315 Gaunt R Birnie J H and Liversole W J *Physiol Revs* **29** 281 (1949)
- 316 Gaunt R Leatham J H Howell C and Antonchak N *Federation Proc* **11** 53 (1952)
- 317 Gaunt R Leatham J H Howell C and Antonchak N *Endocrinology* **50** 571 (1952)
- 318 Gaunt R Tuthill C H Antonchak N and Leatham J H *Endocrinology* **52** 407 (1953)
- 319 Gemzell C A and Samuels L T *Endocrinology* **47** 48 (1950)
- 320 Gemzell C A Van Dyke D C Tobias C A and Evans H M *Endocrinology* **49** 375 (1951)
- 321 Gemzell C A Burke G Hellstrom J Franksson C E and Plantin I O *Acta Endocrinol* **12** 1 (1953)
- 322 Germuth F G *Am J Pathol* **28** 565 (1952)
- 323 Germuth F G *J Exptl Med* **98** 1 (1953)
- 324 Germuth F G and Ottinger B *Proc Soc Exptl Biol Med* **74** 815 (1950)
- 325 Germuth F G Ottinger B and Oyama J *Bull Johns Hopkins Hosp* **91** 2 (1957)
- 326 Germuth F G Ottinger B and Oyama J *Proc Soc Exptl Biol Med* **80** 188 (1952)
- 327 Germuth F G Nedzel G A Ottinger B and Oyama J *Proc Soc Exptl Biol Med* **76** 177 (1951)
- 328 Giroud A *Presse med* **48** 841 (1940)
- 329 Giroud A and Leblond C P *Arch anat microscop* **31** 111 (1935)
- 330 Giroud A and Ratsimamanga A R *Presse med* **48** 449 (1940)
- 331 Giroud A and Ratsimamanga A R *Paris med* **30** 356 (1940)
- 332 Giroud A and Santa N *Compt rend soc biol* **131** 1176 (1939)
- 333 Giroud A Santa N and Martinet M *Compt rend soc biol* **134** 23 100 (1940)
- 334 Glaser G H *Psychosomat Med* **15** 280 (1953)
- 335 Glaser G H and Merritt H H *J Am Med Assoc* **148** 898 (1952)
- 336 Glaser R J Berry J W Loeb L H and Wood W B *J Clin Invest* **30** 640 (1951)
- 337 Glaser R J Berry J W Loeb L H Wood W B and Hamlin A J *Lab Clin Med* **38** 363 (1951)
- 338 Glaubach S *J Mt Sinai Hosp N Y* **19** 84 (1952)
- 339 Glaubach S Antopol W and Graff S *Bull N Y Acad Med* **27** 398 (1951)
- 340 Glick D and Biskind G R *J Biol Chem* **115** 551 (1936)
- 341 Glock G E *Nature* **168** 518 (1945)
- 342 Godlowski Z Z *Brit Med J* **1** 46 (1948)
- 343 Godlowski Z Z *J Endocrinol* **8** 102 (1952)
- 344 Goldblatt H *Ann Internal Med* **11** 69 (1937)

- 339 Forbes A I Criswold C C and Albright I J *J Clin Endocrinol and Metabolism* 10 230 (1950)
- 340 Forsham I H Thorn G W Prunty F T G and Hills A G *J Clin Endocrinol and Metabolism* 8 15 (1948)
- 341 Forsham I H Bennett I L Roche M Reiss R S Slesor A Flink F R and Thorn G W *J Clin Endocrinol and Metabolism* 9 660 (1949)
- 342 Forsyth B T *Endocrinology* 62, 65 (1953)
- 343 Fourman I Bartter I C Albright I Dempsey F Carroll I and Alexander J A *J Clin Invest* 29 1462 (1950)
- 344 Fraenkel Conrat H Simpson M F and Evans H M *J Biol Chem* 147 99 (1943)
- 345 Frawley T F and Forsham P H *J Clin Endocrinol and Metabolism* 11 772 (1951)
- 346 Fredrickson D S Forsham P H and Thorn G W *J Clin Endocrinol and Metabolism* 12 541 (1952)
- 347 Freed S C Roenman R H and Friedman M Inn A I *Acad Sci* 56 637 (1953)
- 348 Freedman H H and Gordon A S *Proc Soc Exptl Biol Med* 75 729 (1950)
- 349 Frenk S Wolfe S and Paschkis H L *Endocrinology* 47 386 (1950)
- 350 Freyberg R H Traeger C T Adams C H Kuseu T Wainerd H and Bonomo I *Science* 112 479 (1950)
- 350a Fried J and Sabo I F *J Am Chem Soc* 76 1455 (1954)
- 350b Fried J Thomas R W Perlman D Herz J F and Borman A *Recent Progr Hormone Research* 11 149 (1955)
- 351 Friedberg I *Euclides (Madrid)* 10 116 (1950)
- 352 Friedberg F and Greenberg D M *Arch Biochem* 17 193 (1948)
- 353 Friedgood C E Vars H M and Zerbe J W *Am J Physiol* 163 354 (1950)
- 354 Friedlaender S and Friedlaender A S *J Allergy* 21 303 (1950)
- 355 Friedlander W J and Rottger L *J Electroenceph and Clin Neurophysiol* 3 311 (1951)
- 356 Friedman M and Byers S O *Am J Physiol* 163 684 (1950)
- 357 Friedman S M and Friedman C I *Endocrinology* 49 318 (1951)
- 358 Friedman S M Friedman C L and Nakashima M *Am J Physiol* 163 319 (1950)
- 359 Friedman S M Friedman C L and Nakashima M *Endocrinology* 51 401 (1952)
- 360 Friedman S M Mackenzie K R and Friedman C I *Endocrinology* 43 123 (1948)
- 361 Friedman S M Polley J R and Friedman C L *J Exptl Med* 87 329 (1948)
- 362 Frost R C and Talmage R V *Endocrinology* 49 606 (1951)
- 363 Fruhman G J and Gordon A S *Federation Proc* 11 51 (1952)
- 364 Fruhman G J and Gordon A S *Proc Soc Exptl Biol Med* 82 162 (1953)
- 365 Fuenzalida F and Lipschutz A *J Clin Endocrinol and Metabolism* 13 1701 (1953)
- 366 Fulton G P and Maynard F L *Proc Soc Exptl Biol Med* 84 759 (1953)
- 367 Gabrielove J I and Soffer I J *Endocrinology* 47 461 (1950)
- 368 Gabrielove J L and Soffer L J *J Clin Invest* 29 814 (1950)

- 441 Gross F and Tschopp T *Experientia* 8 75 (1952)
- 442 Gross J Ann N Y Acad Sci 66 64 (1953)
- 443 Gross R and Siecke A Klin Wochschr 30 456 (1952)
- 444 Grundy H M Simpson S A and Tut J F Nature 169 190 (1952)
- 445 Hackel D B Portfolio A C and Kinney T D Proc Soc Exptl Biol Med 74 458 (1950)
- 446 Hahn L O Houser H B Rummelkamp C H Denny F W and Wannamaker L W J Clin Invest 30 274 (1951)
- 447 Halberg F and Ulstrom R A Proc Soc Exptl Biol Med 80 747 (1952)
- 448 Halberg F and Visscher M B Proc Soc Exptl Biol Med 75 846 (1950)
- 449 Halberg F and Visscher M B Endocrinology 51 370 (1952)
- 450 Halberg F Cohen S I and Flink F B J Lab Clin Med 38 817 (1951)
- 451 Hall C F and Hall O Proc Soc Exptl Biol Med 71 690 (1949)
- 452 Hall C F and Hall O Proc Soc Exptl Biol Med 79 536 (1952)
- 453 Hall C E and Hall O Endocrinology 52 15 (1953)
- 454 Hall M G Darling R C and Taylor F H I Ann Internal Med 13 415 (1939)
- 455 Hall T C McCracken B H and Thorn C W J Clin Endocrinol and Metabolism 13 243 (1953)
- 456 Hallett J W Leopold I H and Steinmetz C G Arch Ophthalmol 46 768 (1951)
- 457 Halmi N S Proc Soc Exptl Biol Med 80 175 (1952)
- 458 Halmi N S and Barker S B Endocrinology 51 177 (1952)
- 459 Halmi N S and Bogdanove F M Proc Soc Exptl Biol Med 77 518 (1951)
- 460 Halpern B N Benacerraff B and Briot M Proc Soc Exptl Biol Med 79 37 (1952)
- 461 Halpern B N Maurie G Holtzer A and Briot M J Allergy 23 303 (1952)
- 462 Hamburger C Acta Endocrinol 14 375 (1953)
- 463 Hamilton I D Cubler C J Ashenbrucker H Cartwright G F and Wintrobe M M Endocrinology 48 44 (1951)
- 464 Hanan R and Overman J R Proc Soc Exptl Biol Med 84 470 (1953)
- 465 Hanan R and Oyama J Proc Soc Exptl Biol Med 85 33 (1954)
- 466 Handler P and Bernheim F Federation Proc 10 104 (1951)
- 467 Hardy J D Riegel C and Erisman E P Am J Med Sci 219 582 (1950)
- 468 Harris L J Bland M U Hughes P E and Constable B J Lancet i 1071 (1953)
- 469 Harris L J and Ray S N Biochem J 27 303 (1933)
- 470 Harris S and Harris T N Proc Soc Exptl Biol Med 74 186 (1950)
- 471 Harrison C U King L H Dole J C and Siegel I Brit J Ind Med 9 165 (1952)
- 472 Harrison J H Thorn G W and Crisantiello M G J Urol 67 405 (1952)
- 473 Harrison J H Thorn G W and Jenkins D New Engl J Med 248 86 (1953)
- 474 Hart P D and Rees R J W Lancet ii 591 (1950)
- 475 Hartroft P M and Hartroft W S J Exptl Med 97 415 (1953)
- 476 Haskin D Iasher N and Rothman S J Invest Dermatol 20 207 (1953)
- 477 Havens W P Jr Shaffer J M and Hopke C J J Immunol 68 380 (1952)
- 478 Havano M and Dorfman R I Ann N Y Acad Sci 54 608 (1951)

- 405 Colden W R C and Long C N H *Endocrinology* 30 615 (1912)
- 406 Goldman I and Richfield D I *J Am Med Assoc* 147 941 (1951)
- 407 Goldman I Ireston R and Rockwell F *J Invest Dermatol* 18 89 (1952)
- 408 Goldman I Rockwell I and Richfield D I *Am J Trop Med Hyg* 1 511 (1952)
- 409 Goldstein M S Mendel B and Levine R *Am J Physiol* 163 714 (1950)
- 410 Goldzieher J W Epstein I R Rawls W B and Goldzieher M A *J Clin Endocrinol and Metabolism* 13 662 (1953)
- 411 Gordon A S and Katsch C F *Ann N Y Acad Sci* 52 1 (1949)
- 412 Gordon A S Ehlers S J and Lindau D *Endocrinology* 49 491 (1951)
- 413 Gordon G S and Elliott H W *Endocrinology* 41 517 (1941)
- 414 Gordon G S Bentuck R C and Eisenberg I *Ann N Y Acad Sci* 64 515 (1951)
- 415 Gottschalk R C and Crollman A *Cancer Research* 12 651 (1952)
- 416 Grabar I Benacerraff B and Biozzi G *Ann Inst Pasteur* 81 187 (1951)
- 417 Grad B J *J Clin Endocrinol and Metabolism* 12 708 (1952)
- 418 Graham A B Werder A A Syvertson J T and Friedman J *Federation Proc* 11 470 (1952)
- 419 Graham J B Graham R M and Graffeo A J *Endocrinology* 46 431 (1950)
- 420 Gray S J Benson J A and Reifenstein R W *Proc Soc Exptl Biol Med* 78 338 (1951)
- 421 Gray S J Benson J A Reifenstein R W and Ramsey C *J Clin Invest* 31 633 (1952)
- 422 Gray S J Benson J A Reifenstein R W and Spiro H M *J Am Med Assoc* 147 1529 (1951)
- 423 Gray W D and Munson I I *Endocrinology* 48 471 (1951)
- 424 Gray W D Pedrick I and Winne R *Proc Soc Exptl Biol Med* 78 679 (1951)
- 425 Green D M Farah A and Klempner W W *Endocrinology* 47 281 (1950)
- 426 Green D M Johnson A D Bridges W E Lehmann J H and Gray F *Endocrinology* 47 102 (1950)
- 427 Green D M Saunders F J Wahlgren N and Craig R I *Am J Physiol* 170 94 (1950)
- 428 Green H N *Brit Med J* i 1165 (1950)
- 429 Greenspan F S Clifford H and Deming Q B *Endocrinology* 52 638 (1953)
- 430 Greep R O and Deane H W *Ann N Y Acad Sci* 50 596 (1949)
- 431 Greep R O Knobil F Hofmann I G and Jones T I *Endocrinology* 60 664 (1952)
- 432 Grenell R G and McCrawley T L *Federation Proc* 6 116 (1947)
- 433 Grob D and Harvey A M *Bull Johns Hopkins Hosp* 91 124 (1952)
- 434 Groen H Ieiser H Frenkel M Kaminga C L and Willebrands A F *New Engl J Med* 244 471 (1951)
- 435 Groen H *J Clin Invest* 31 87 (1952)
- 436 Grokoest A W Vaillancourt deG Gattisgen R and Ragan C *J Clin Invest* 30 614 (1951)
- 437 Grollman A and Konnerth A *Endocrinology* 48 213 (1951)
- 438 Grolnick M *J Allergy* 23 416 (1952)
- 439 Gross F *Schweiz med Wochschr* 80 697 (1950)
- 440 Gross F and Gysel H *Acta Endocrinol* 15 199 (1954)

- 19 Haggins G M *Am J Clin Pathol* 23 770 (1953)
- 511 Haggins G M Woods R A and Bennett W A *Cancer Research* 10 203 (1950)
- 51 Haggins G M Woods R A and Kendall L C *Endocrinology* 48 170 (1951)
- 513 Hall S R Reiss R S Forsham P H and Thorn G W *J Clin Endocrinol and Metabolism* 10 1370 (1950)
- 514 Hills A G Forsham P H and Finch C A *Blood* 3 755 (1948)
- 515 Hoagland H *Psychosomat Med* 12 142 (1950)
- 516 Hoagland H *Intern Record of Med* 166 183 (1953)
- 517 Hoagland H *In J M Yoffey The Suprarenal Cortex* Academic Press New York 1953 p 125
- 518 Hoagland H Bergen J R Blocombe A C and Hunt C *Ann N Y Acad Sci* 56 659 (1953)
- 519 Hoberman H D *Yale J Biol and Med* 22 311 (1950)
- 520 Hoch Liggett C and Hau Y T *Am J Pathol* 29 614 (1953)
- 521 Hoefler I F A and Glaser G H *J Am Med Assoc* 143 620 (1950)
- 522 Hoffman W C Lewis R A and Thorn G W *Bull Johns Hopkins Hosp* 70 335 (1947)
- 523 Hofmann F G *Endocrinology* 50 706 (1952)
- 524 Hogness J R Lee A D and Williams R H *Endocrinology* 52 378 (1953)
- 525 Hogness J R Williams R H and Lance M *Proc Soc Exptl Biol Med* 79 43 (1952)
- 526 Holden M Seegal B C and Rybo I *Am J Pathol* 27 743 (1951)
- 527 Hollander J I Brown L M Jessar R A and Brown C Y *J Clin Invest* 30 650 (1951)
- 528 Hollander J I Brown L M Jessar R A and Brown C Y *J Am Med Assoc* 147 1679 (1951)
- 529 Holley H I and Mc Lester J S *Arch Internal Med* 88 760 (1951)
- 530 Holman H R White A and Fruton J S *Proc Soc Exptl Biol Med* 65 106 (1944)
- 531 Holmes J H and Percefull S C *Ann Internal Med* 33 608 (1951)
- 532 Holtkamp D E Mansor L F and Heming A L *Federation Proc* 11 338 (1952)
- 533 Homburger F and Bonner C D *Gastroenterics* 8 385 (1953)
- 534 Homburger F Dart R M Bonner C D Branche G Hasdon S C and Fishman W H *J Clin Endocrinol and Metabolism* 13 64 (1953)
- 535 Housay A B *Rev assoc méd argentina* 66 132 (1952)
- 536 Howes E L *Yale J Biol and Med* 23 451 (1951)
- 537 Howes E L Plotz C M Blunt J W and Ragan C *Surgery* 28 177 (1950)
- 538 Hubay C A Weckesser E C and Holden W D *Surg Gynecol Obstet* 96 65 (1953)
- 539 Hubble D *Lancet II* 464 (1952)
- 540 Hudson G Herdan G and Yoffey J M *Brit Med J* 1 999 (1952)
- 541 Hudson P B Mittleman A and Mann I *J Clin Endocrinol and Metabolism* 13 1064 (1953)
- 542 Huggins C and Bergenstal D M *J Am Med Assoc* 147 191 (1951)
- 543 Huggins C and Bergenstal D M *Proc Natl Acad Sci U S* 38 73 (1952)
- 544 Huggins C and Dao T I Y *J Am Med Assoc* 161 1388 (1953)

- 479 Hayano M Dorfman R I and Yamada F Y *J Biol Chem* 186 603 (1950)
- 480 Hayano M Schiller S and Dorfman R I *Endocrinology* 46 357 (1950)
- 481 Hayes M A *Endocrinology* 52 646 (1953)
- 482 Hayes M A and Baker B I *Endocrinology* 49 349 (1951)
- 483 Hayes M A Reed T C and Baker B I *Proc Soc Exptl Biol Med* 75 357 (1950)
- 484 Hechter O and Johnson S *Endocrinology* 45 351 (1949)
- 485 Heidelberger M and Kendall F I *J Exptl Med* 62 697 (1935)
- 486 Heilmann D N *Proc Staff Meetings Mayo Clinic* 20 310 (1945)
- 487 Heilmann F R and Kendall F I *Endocrinology* 34 416 (1944)
- 488 Heller H *In J M Yoffey The Suprarenal Cortex* Academic Press New York 1953 p 187
- 489 Heller J H *Federation Proc* 12 65 (1953)
- 490 Heller J H *Federation Proc* 13 69 (1954)
- 491 Hellman I *Science* 109 280 (1949)
- 492 Heming A L Holtkamp D F and Mansor I F *J Clin Endocrinol and Metabolism* 12 961 (1952)
- 493 Heming A L Holtkamp D I Kerwin J F Mansor L F and Dacanay, J G *Proc Soc Exptl Biol Med* 80 151 (1952)
- 494 Hemphill R E *In J M Yoffey The Suprarenal Cortex* Academic Press New York 1953 p 135
- 495 Hench P S Kendall E C Slocumb C H and Polley H F *Proc Staff Meetings Mayo Clinic* 24 181 (1949)
- 496 Hench P S Kendall E C Slocumb C H and Polley H F *Arch Intern Med* 85 545 (1950)
- 497 Hench P S Slocumb C H Barnes A R Smith H L Polley H F and Kendall E C *Proc Staff Meetings Mayo Clinic* 24 277 (1949)
- 498 Henderson J W and Hollenhorst R W *Proc Staff Meetings Mayo Clinic* 25 459 (1950)
- 499 Henriques O B Henriques S B and Selye H *Proc Soc Exptl Biol Med* 73 611 (1950)
- 500 Henriques S B Henriques O B and Mattos I *Endocrinology* 47 457 (1950)
- 501 Henry W I Oliver L and Ramey E R *Federation Proc* 12 66 (1953)
- 502 Hensler N M Wurl O A and Gillespie J O *U S Armed Forces Med J* 3 199 (1952)
- 503 Hentstell H H and Freedman R I *Proc Soc Exptl Biol Med* 76 238 (1951)
- 504 Herrick E H Mead F R Egerton B W and Hughes J S *Endocrinology* 50 259 (1952)
- 505 Hertz R and Tullner W W *Proc Soc Exptl Biol Med* 78 737 (1951)
- 506 Hertz R Allen M J and Tullner W W *Proc Soc Exptl Biol Med* 75 627 (1950)
- 507 Hertz R Tullner W W and Allen M J *Proc Soc Exptl Biol Med* 77 480 (1951)
- 508 Hess W C Kyle L H and Doolan P D *Proc Soc Exptl Biol Med* 76 418 (1951)
- 509 Hidalgo J McClure C D Henderson J B Whitehead R W and Smith C J *Proc Soc Exptl Biol Med* 80 97 (1952)

- 510 Higgins G M *Am J Clin Pathol* **23** 775 (1953)
- 511 Higgins G M Woods K A and Bennett W A *Cancer Research* **10** 203 (1950)
- 512 Higgins C M Woods K A and Kendall L C *Endocrinology* **48** 175 (1951)
- 513 Hill S R Reiss R S Forsham P H and Thorn G W *J Clin Endocrinol and Metabolism* **10** 1345 (1950)
- 514 Hills A G Forsham P H and Finch C A *Blood* **13** 755 (1948)
- 515 Hoagland H *Psychosomat Med* **12** 142 (1950)
- 516 Hoagland H *Intern Record of Med* **166** 183 (1953)
- 517 Hoagland H *In* J M Yoffey *The Suprarenal Cortex* Academic Press New York 1953 p 125
- 518 Hoagland H Bergen J R Slocombe A G and Hunt C *Ann N Y Acad Sci* **56** 659 (1953)
- 519 Holberman H D *Yale J Biol and Med* **22** 341 (1950)
- 520 Hoch Ligeti C and Hsu Y T *Am J Pathol* **29** 614 (1953)
- 521 Hoefler P F A and Glaser G H *J Am Med Assoc* **143** 620 (1950)
- 522 Hoffman W C Lewis R A and Thorn G W *Bull Johns Hopkins Hosp* **70** 335 (1942)
- 523 Hofmann F G *Endocrinology* **60** 206 (1952)
- 524 Hogness J R Lee N D and Williams R H *Endocrinology* **62** 378 (1953)
- 525 Hogness J R Williams R H and Lance M *Proc Soc Exptl Biol Med* **79** 43 (1952)
- 526 Holden M Seegal B C and Ryby I *Am J Pathol* **27** 748 (1951)
- 527 Hollander J L Brown I M Jessar R A and Brown C Y *J Clin Invest* **30** 650 (1951)
- 528 Hollander J L Brown E M Jessar R A and Brown C Y *J Am Med Assoc* **147** 1629 (1951)
- 529 Holley H L and McLester J S *Arch Internal Med* **88** 760 (1951)
- 530 Holman H R White A and Fruton J S *Proc Soc Exptl Biol Med* **65** 196 (1947)
- 531 Holmes J H and Percetull S C *Ann Internal Med* **35** 608 (1951)
- 532 Holtkamp D E Mansoor L F and Heming A I *Federation Proc* **11** 358 (1952)
- 533 Homburger F and Bonner C D *Geriatrics* **8** 385 (1953)
- 534 Homburger F Dart R M Bonner C D Branche G Kasdon S C and Fishman W H *J Clin Endocrinol and Metabolism* **13** 704 (1953)
- 535 Houssay A B *Rev assoc méd argentina* **66** 132 (1952)
- 536 Howes E L *Yale J Biol and Med* **23** 44 (1951)
- 537 Howes E I Plotz C M Blunt J W and Ragan C *Surgery* **28** 177 (1950)
- 538 Hubay C A Weckesser E C and Holden W D *Surg Gynecol Obstet* **96** 65 (1953)
- 539 Hubble D *Lancet* **ii** 464 (1952)
- 540 Hudson G Herdan G and Yoffey J M *Brit Med J* **i** 999 (1952)
- 541 Hudson P B Mittleman A and Mann P *J Clin Endocrinol and Metabolism* **13** 1064 (1953)
- 542 Huggins C and Bergenstal D M *J Am Med Assoc* **147** 191 (1951)
- 543 Huggins C and Bergenstal D M *Proc Natl Acad Sci U S* **38** 73 (1952)
- 544 Huggins C and Dao T L Y *J Am Med Assoc* **151** 1388 (1953)

- 545 Huggins C, Bergen tal D M and Cleveland A S *Recent Progr Hormone Research* 8 273 (1953)
- 546 Human T H J *Acta Endocrinol* 13 55 (1953)
- 547 Hull R B and Loush C C *J Lab Clin Med* 37 603 (1951)
- 548 Humphrey J H *Brit J Exptl Pathol* 32 274 (1951)
- 549 Humphreys R J and Rabb W *Proc Soc Exptl Biol Med* 74 302 (1950)
- 550 Hundley I M and Ing K B *Endocrinology* 48 482 (1951)
- 551 Hungerford C F Reinhardt W O and Ii C H *Blood* 7 1125 (1952)
- 552 Hungerford C F Reinhardt W O and Ii C H *Proc Soc Exptl Biol Med* 81 320 (1952)
- 553 Hunter O B and Ross J B *Southern Med J* 45 732 (1952)
- 554 Hunter S F *Proc Soc Exptl Biol Med* 82 14 (1953)
- 555 Hyman G A Rakin C and Turner J C *Proc Soc Exptl Biol Med* 75 410 (1950)
- 556 Hyman G A Rakin C and Turner J C *Trans N Y Acad Sci* 13 161 (1951)
- 557 Ingbar S H Kass I H Burnett C H Reiman A S Burrows B A and Sisson J H *J Lab Clin Med* 38 533 (1951)
- 558 Ingbar S H Otto J I and Kass L H *Proc Soc Exptl Biol Med* 77 20 (1951)
- 559 Ingle D J *Ann N Y Acad Sci* 50 576 (1950)
- 560 Ingle D J *J Clin Endocrinol and Metabolism* 10 1312 (1950)
- 561 Ingle D J *Recent Progr Hormone Research* 7 315 (1952)
- 562 Ingle D J *Proc Soc Exptl Biol Med* 79 184 (1952)
- 563 Ingle D J and Baker B I *Endocrinology* 48 313 (1951)
- 564 Ingle D J and Baker B I *Endocrinology* 48 764 (1951)
- 565 Ingle D J and Kendall I C *Am J Physiol* 122 585 (1953)
- 566 Ingle D J and Meeks R C *Am J Physiol* 170 77 (1957)
- 567 Ingle D J and Nezamis J I *Am J Physiol* 156 365 (1949)
- 568 Ingle D J and Nezamis J I *Endocrinology* 48 484 (1951)
- 569 Ingle D J and Nezamis J I *Endocrinology* 52 361 (1953)
- 570 Ingle D J and Prestrud M C *Endocrinology* 46 143 (1949)
- 571 Ingle D J and Thorn C W *Am J Physiol* 132 610 (1941)
- 572 Ingle D J Meeks R C and Beary D F *Proc Soc Exptl Biol Med* 84 334 (1953)
- 573 Ingle D J Morley I H and Nezamis J F *Proc Soc Exptl Biol Med* 78 220 (1951)
- 574 Ingle D J Morley I H and Nezamis J E *Endocrinology* 51 487 (1952)
- 575 Ingle D J Meeks R C and Thomas K E *Endocrinology* 49 103 (1951)
- 576 Ingle D J Nezamis J I and Jeffries J W *Am J Physiol* 167 99 (1949)
- 577 Ingle D J Nezamis J I and Morley E H *Proc Soc Exptl Biol Med* 78 79 (1951)
- 578 Ingle D J Nezamis J E and Morley E H *Endocrinology* 50 1 (1957)
- 579 Ingle D J Nezamis J E and Morley E H *Am J Physiol* 171 378 (1957)
- 580 Ingle D J Prestrud M C and Ii C H *Am J Physiol* 166 165 (1951)
- 581 Ingle D J Prestrud M C and Nezamis J F *Proc Soc Exptl Biol Med* 75 801 (1950)
- 582 Ingle D J Prestrud M C and Nezamis J E *Am J Physiol* 166 111 (1951)
- 583 Ingle D J Prestrud M C and Rice K L *Endocrinology* 46 510 (1950)

- 584 Ingle D J Sheppard R Oberle F A and Kuizenga M H *Endocrinology* 39 52 (1946)
- 585 Ivon Sillmann I Jones I S Wiggins R I and Locke B D *Klin Wochschr* 63 684 (1951)
- 586 Jackson C M *Am J Anat* 21 321 (1917)
- 587 Jacob J and Szerb J *Arch intern pharmacodynamie* 90 301 (1952)
- 588 Jailer J W and Bois V F *Endocrinology* 46 314 (1950)
- 589 Jailer J W Touchart J and Cahill G F *J Am Med Assoc* 160 575 (1952)
- 590 Jailer J W Touchart J and Cahill G F *J Clin Invest* 31 883 (1955)
- 591 Jailer J W Wong A S H and Ingle E T *J Clin Endocrinol and Metabolism* 11 186 (1951)
- 592 Jarvinen K A J *Brit Med J* 11 1377 (1951)
- 593 Jensen H and Gray J I *Ann N Y Acad Sci* 54 619 (1951)
- 594 Johnson A S Scheinberg S R Gerisch R and Saltzstein H C *Harper Hosp Bull* 9 187 (1951)
- 595 Johnson A S Scheinberg S R Gerisch R and Saltzstein H C *Circulation* 7 224 (1953)
- 596 Johnson B B *Endocrinology* 54 196 (1954)
- 597 Joiner C I *Brit Med J* 11 642 (1952)
- 598 Jones G E S Howard J F and Langford H *Fertility and Sterility* 4 49 (1953)
- 599 Jones I S and Meyer K *Proc Soc Exptl Biol Med* 74 102 (1950)
- 600 Josey A I and Lawrence J S *Folia Haematol* 48 303 (1952)
- 601 Jost A *Recent Progr Hormone Research* 8 349 (1953)
- 602 Kabat E A and Heidelberger M *J Exptl Med* 66 229 (1937)
- 603 Kabat E A Wolf A and Bezer A F *J Immunol* 68 265 (1952)
- 604 Kagawa C M Shipley E G and Meyer R K *Proc Soc Exptl Biol Med* 80 281 (1952)
- 605 Kaplan H S Brown M B and Varder S N *Cancer Research* 11 262 (1951)
- 606 Kaplan N O and Lipmann F *J Biol Chem* 174 37 (1948)
- 607 Kark R M and Muehrcke R C *Lancet* 1 1189 (1952)
- 608 Kark R M Chapman R E Corsolazio C F and Nesby C *J Lab Clin Med* 40 817 (1952)
- 609 Karlson A C and Gainer J H *Proc Staff Meetings Mayo Clinic* 27 465 (1952)
- 610 Karnofsky D A Ridgway L P and Patterson P A *Endocrinology* 48 596 (1951)
- 611 Karnofsky D A Ridgway L P and Stock C C *Federation Proc* 10 204 (1951)
- 612 Kass E H and Finland M *New Engl J Med* 243 693 (1950)
- 613 Kass E H Geiman Q M and Finland M *New Engl J Med* 245 1006 (1951)
- 614 Kass E H Lundgren M M and Finland M *Ann N Y Acad Sci* 66 765 (1953)
- 615 Kass E H Ingbar S H and Finland M *Ann Internal Med* 33 1081 (1950)
- 616 Kass E H Ingbar S H and Finland M *Proc Soc Exptl Biol Med* 73 669 (1950)

- 617 Kass E H Hendrick, M I and Finland M *Ann N Y Acad Sci* 66 737 (1953)
- 618 Kass E H Ingbar S H Lundgren M M and Finland M *J Lab Clin Med* 37 780 (1951)
- 619 Kaufman N Mason L J and Kinney T D *Am J Pathol* 29 761 (1953)
- 620 Kay J H *Surgery* 32 686 (1952)
- 621 Kayahan S *J Endocrinol* 8 211 (1952)
- 622 Kellgren J H *Brit Med J* 1 1093 (1952)
- 623 Kelsall M A and Crabb F D *Inat Record* 115 331 (1953)
- 624 Kelley V C Fly R S Raile R B and Bray P F *Proc Soc Exptl Biol Med* 81 611 (1952)
- 625 Kendall E C *Science* 111 457 (1950)
- 626 Kennedy T H and Purves H D *Brit J Exptl Pathol* 22 241 (1941)
- 627 Kerppola W *Endocrinology* 51 192 (1952)
- 628 Key J A Odell R T and Taylor I W *J Bone and Joint Surgery* 34 665 (1952)
- 629 Keyes P H *Endocrinology* 44 274 (1949)
- 630 Kilbourne E D and Horsfall F I *Proc Soc Exptl Biol Med* 77 135 (1951)
- 631 Kilbourne E D and Horsfall F I *Proc Soc Exptl Biol Med* 76 116 (1951)
- 632 Kilbourne E D and Tateno I *Proc Soc Exptl Biol Med* 82 274 (1953)
- 633 Kimura S J Thygeson P and Geller H O *Am J Ophthalmol* 36 116 (1953)
- 634 Kinsell I W Partridge J W and Foreman N *Ann Internal Med* 38 913 (1953)
- 635 Kinsell I W Partridge J W Boling L and Margen S *Ann Internal Med* 37 921 (1952)
- 636 Kinsell L W Michaels C D Margen S Boling I and Partridge J W *Am J Med* 13 96 (1952)
- 637 Kinsell L W Margen S Michaels G D Russ R Frantz R and Carbone J *J Clin Invest* 30 1491 (1951)
- 638 Mitchell R L and Wells L J *Endocrinology* 50 83 (1952)
- 639 Kit S and Guzman Barron F S *Endocrinology* 62 1 (1953)
- 640 Klein R *J Clin Invest* 30 318 (1951)
- 641 Klein R and Hanson J *Pediat* 6 192 (1950)
- 642 Klein R and Livingstone S *J Pediat* 37 733 (1950)
- 643 Kleigman A M Baldrige G D Rebell G and Pillsbury D M *J Lab Clin Med* 37 615 (1951)
- 644 Kline D I *Endocrinology* 45 596 (1949)
- 645 Knauff R E Nielson E D and Haines A J *J Am Chem Soc* 75 4809 (1953)
- 645a Knobil E Hofmann F G and Greep O *Endocrinology* 53 242 (1953)
- 646 Knowlton A I Loeb F N Stoerk H C and Seegal B C *Proc Soc Exptl Biol Med* 72 722 (1949)
- 647 Knowlton A I Loeb E N Seegal B C Stoerk H C and Berg J L *Proc Soc Exptl Biol Med* 74 661 (1950)
- 648 Kobernick C D and More R H *Proc Soc Exptl Biol Med* 74 602 (1950)
- 649 Kochakian C D *Ann N Y Acad Sci* 54 534 (1951)
- 650 Kochakian C D and Robertson F *J Biol Chem* 190 481 (1951)

- 651 Kochakian C D and Robertson L *J Biol Chem* 190 495 (1951)
- 652 Kochakian C D and Vail V N *J Biol Chem* 169 1 (1951)
- 653 Kountz W B Ackermann I G and Kheim T *J Clin Endocrinol and Metabolism* 13 534 (1953)
- 654 Kowalewski K and Bastenie I A *Acta Endocrinol* 11 231 (1950)
- 655 Krali M E *Ann N Y Acad Sci* 54 649 (1951)
- 656 Kramar J and Simay Kramar M *Endocrinology* 52 453 (1953)
- 657 Krohn I L In J M Yoffey *The Suprarenal Cortex* Academic Press New York 1953 p 167
- 658 Kuhl W J and Ziff M *J Clin Endocrinol and Metabolism* 12 554 (1952)
- 659 Kumagai I F and Dougherty T F *Am J Physiol* 167 804 (1951)
- 660 Kurland C S and Freedberg A S *Proc Soc Exptl Biol Med* 78 28 (1951)
- 661 Kuusisto A N and Telkka A *Acta Endocrinol* 13 61 (1953)
- 662 Kuzell W C and Mankle E A *Proc Soc Exptl Biol Med* 74 677 (1950)
- 663 Lachance J P and Page E *Endocrinology* 52 57 (1953)
- 664 Landau W W Nelson W A and Gay L N *Bull Johns Hopkins Hosp* 88 395 (1951)
- 665 Landauer W *Endocrinology* 41 489 (1947)
- 666 Lanman J T *Pediatrics* 12 62 (1953)
- 667 Iaqueur G I *Am J Pathol* 28 591 (1952)
- 668 Iast J H Jordan P H Pitesky I and Siegel B M *Proc Soc Exptl Biol Med* 74 96 (1950)
- 669 Lattes R Blunt J W Rose H M Jessar R A Vaillancourt de G and Ragan C *Am J Pathol* 29 1 (1953)
- 670 Lauridsen J Belko J S and Warren R *Proc Soc Exptl Biol Med* 79 709 (1952)
- 671 Law L W and Speirs R S *Proc Soc Exptl Biol Med* 66 226 (1947)
- 672 Layton I L *Proc Soc Exptl Biol Med* 76 598 (1951)
- 673 Lazarow A *Proc Soc Exptl Biol Med* 74 707 (1950)
- 674 Lazarow A and Berman J *Anat Record* 106 215 (1950)
- 675 Leaf A Mamby A R Rasmussen H and Marasco J P *J Clin Invest* 31 914 (1952)
- 676 Leatham J H *Proc Soc Exptl Biol Med* 74 855 (1950)
- 677 Leblond C P and Hoff H E *Endocrinology* 35 223 (1944)
- 678 Leblond C P and Segal C *Am J Roentgenol Radium Therapy* 47 302 (1942)
- 679 Lee N D and Williams R H *Endocrinology* 51 454 (1952)
- 680 Lee N D and Williams R H *Proc Soc Exptl Biol Med* 79 669 (1952)
- 681 Lee R F and Pfeiffer C C *Proc Soc Exptl Biol Med* 77 752 (1951)
- 682 Leger J Leith W and Rose B *Proc Soc Exptl Biol Med* 69 465 (1948)
- 683 LeMaistre C and Tompsett R *Am Rev Tuberc* 64 235 (1951)
- 684 LeMaistre C and Tompsett R *J Exptl Med* 95 393 (1952)
- 685 LeMaistre C Tompsett R and McDermott W *Ann N Y Acad Sci* 56 772 (1953)
- 686 Lemonde P Panisset M Dobija M and Selye H *J Clin Endocrinol and Metabolism* 12 93 (1952)
- 687 Leopold I H and Maylath F R *Am J Ophthalmol* 35 1125 (1952)
- 688 Leopold I H and Maylath F R *Arch Ophthalmol* 48 110 (1952)
- 689 Leopold I H Purnell J E Cannon E J Steinmetz C G and McDonald P R *Am J Ophthalmol* 34 361 (1951)

- 690 Lever J D *J Endocrinol* 10 133 (1951)
- 691 Levin I *J Clin Endocrinol and Metabolism* 9 617 (1919)
- 692 Levin L and Farber R K *Proc Soc Exptl Biol Med* 74 758 (1950)
- 693 Levitt M F and Bader M F *J Clin Invest* 30 655 (1951)
- 694 Lewin L and Wassén I *Lancet* II 993 (1919)
- 695 Lewis L A and Page I H *Endocrinology* 43 415 (1948)
- 696 Lewis L A and Page I H *Ann N Y Acad Sci* 50 547 (1919)
- 697 Lewis R A Rosenberg F and Wilkins I *Endocrinology* 47 414 (1950)
- 698 Lewis R B and Freytag L *Proc Soc Exptl Biol Med* 77 816 (1951)
- 699 Jippman R W and Marti H U *Proc Natl Acad Sci U S* 37 447 (1951)
- 700 Lipsett M N and Moore F J *J Biol Chem* 192 743 (1951)
- 701 Lipsett M N and Moore F J *J Biol Chem* 197, 303 (1952)
- 702 Lister A and Greaves D I *Brit J Ophthalmol* 35 725 (1951)
- 703 Locke W Talbot N B Jones H S and Worcester J *J Clin Invest* 30 325 (1951)
- 704 Lockett M F *J Physiol* 109 250 (1949)
- 705 Long C N H *Federation Proc* 6 461 (1917)
- 706 Long C N H *Science* 111 458 (1950)
- 706a Long D A *Lancet* I 579 (1951)
- 707 Long J B and Favour C B *Bull Johns Hopkins Hosp* 87 186 (1950)
- 708 Loosli C G Hull R B Berlin B S and Alexander, L R *J Lab Clin Med* 37 464 (1951)
- 709 Lotspeich W D *Endocrinology* 44 314 (1949)
- 710 Love W D *Proc Soc Exptl Biol Med* 75 639 (1950)
- 711 Lowe C U Williams W I and Thomas L *Proc Soc Exptl Biol Med* 78 818 (1951)
- 712 Lowe J *Brit Med J* II 746 (1952)
- 713 Luft R and Sjörgen B *Acta Endocrinol* 7 211 (1951)
- 714 Luft R and Sjörgen B *Acta Endocrinol* 10 49 (1952)
- 715 Luongo M A Reid D H and Weiss W W *New Engl J Med* 245 757 (1951)
- 716 Lurie M B Zappasodi P Dannenburg A M Jr and Cardona Lynch E *Ann N Y Acad Sci* 66 779 (1953)
- 717 Lurie M B Zappasodi P Dannenburg A M Jr and Swartz I B *Science* 113 234 (1951)
- 718 Madden R J and Ramsburg H H *Endocrinology* 49 82 (1951)
- 719 Madison L L *J Clin Invest* 29 789 (1950)
- 720 Magarey F R and Gough J *Brit J Exptl Pathol* 33 510 (1952)
- 721 Maqsood M *J Endocrinol* 7 82 (1950)
- 722 Malkiel S *Proc Soc Exptl Biol Med* 77 333 (1951)
- 723 Malkiel S and Hargis B J *J Immunol* 69 217 (1957)
- 724 Mancini R E and de Lustig E S *J Natl Cancer Inst* 13 247 (1952)
- 725 Mandel W Singer M J Gudmundson H R Meister L and Modern F W S *J Am Med Assoc* 146 516 (1951)
- 726 Marcus S and Donaldson D M *J Immunol* 69 101 (1952)
- 727 Marcus F Romanoff I P and Pincus G *Endocrinology* 50 286 (1952)
- 728 Margen S Kissell L W Flanagan E K Sutter L E and Rapaport, E *J Clin Endocrinol and Metabolism* 9 662 (1949)
- 729 Marshall I M and Friedberg F *Endocrinology* 48 113 (1951)
- 730 Martin S P and Kerby G P *Proc Soc Exptl Biol Med* 81 73 (1952)

- 31 Martinez C, Visscher M B, King J T and Bittner J J *Proc Soc Exptl Biol Med* 80 81 (1952)
- 32 Mason A S and Morris C J O R *Lancet* 1 116 (1953)
- 33 Masson G M C *Am J Pathol* 27 717 (1951)
- 34 Masson G M C, Corcoran A C and Page I H *J Lab Clin Med* 34 1416 (1949)
- 35 Masson G M C, Lewis I A, Corcoran A C and Page I H *J Clin Endocrinol and Metabolism* 13 300 (1953)
- 36 Masson G M C, Page I H and Corcoran A C *Proc Soc Exptl Biol Med* 73 434 (1950)
- 37 Mattox V R, Mason H I, Albert A and Code C F *J Am Chem Soc* 75 4869 (1953)
- 38 Maumence A I *Am J Ophthalmol* 34 142 (1951)
- 39 McCracken B H and Hall T C *J Clin Endocrinol and Metabolism* 12 923 (1952)
- 40 McDermott W, Fry I G, Brobeck J R and Long C N H *Proc Soc Exptl Biol Med* 73 609 (1950)
- 41 McDermott W, Fry L C, Brobeck J R and Long C N H *Yale J Biol and Med* 23 52 (1950)
- 42 McDonald P R, Leopold I H, Vogel A W and Mulberger R D *Arch Ophthalmol* 49 400 (1953)
- 43 McEachern D *Bull N Acad Med* 27 3 (1951)
- 44 McGavack T T, Saccone A, Vogel M and Harris R *J Clin Endocrinol and Metabolism* 6 716 (1946)
- 45 McIntosh H W and Holmes C R *Lancet* 11 1061 (1951)
- 46 McLean C R, Hamilton J D and Fitzgerald J D I *Am J Pathol* 27 11 (1951)
- 47 McManus J F A, Cash J R, Carter J P, Alrich E M and Lehman E P *Federation Proc* 10 364 (1951)
- 48 McQuarrie I, Anderson J A and Ziegler M R *J Clin Endocrinol and Metabolism* 2 406 (1942)
- 49 McQuillan M T and Trikojus V M *Brit J Exptl Pathol* 27 247 (1946)
- 50 Meadows E C and Prudden J F *Surgery* 33 841 (1953)
- 51 Means J H *Lancet* 11 543 (1949)
- 52 Megel H and Gordon A S *Endocrinology* 48 591 (1951)
- 53 Meier R, Gysel H and Muller R *Schweiz Med Wochschr* 74 93 (1944)
- 54 Meier R, Schuler W and Desaulles P *Experientia* 6 469 (1950)
- 55 Meiklejohn K P *Vitamins and Hormones* 11 67 (1953)
- 56 Meites J *Proc Soc Exptl Biol Med* 78 692 (1951)
- 57 Meites J *Metabolism* 1 58 (1952)
- 58 Meites J *Proc Soc Exptl Biol Med* 81 307 (1952)
- 59 Meites J and Feng Y S L *Proc Soc Exptl Biol Med* 85 341 (1954)
- 60 Melampy R M, Cheng D W and Northrop L C *Proc Soc Exptl Biol Med* 76 24 (1951)
- 61 Melcher G W, Blunt J W and Ragan C *J Clin Invest* 31 649 (1952)
- 62 Menkin V *Am J Physiol* 166 518 (1951)
- 63 Merck and Co *The Cortisone Investigator* 1-10 Published by Merck and Co (1950-54)
- 64 Merritt H H *Yale J Biol and Med* 24 466 (1952)
- 65 Meyer R J *J Clin Endocrinol and Metabolism* 13 123 (1953)

- 766 Michael M Cummings M M and Bloom W I *Proc Soc Exptl Biol Med* 75 613 (1950)
- 767 Michael M Jr and Whorton C M *Proc Soc Exptl Biol Med* 76 754 (1951)
- 768 Michaelson I C *Arch Ophthalmol* 47 459 (1952)
- 769 Michaelson I C *Arch Ophthalmol* 48 144 (1952)
- 770 Migron C J *Proc Soc Exptl Biol Med* 80 571 (1952)
- 771 Migron C J and Cardner, I I *J Clin Endocrinol and Metabolism* 12 1513 (1952)
- 772 Migron C J Cardner I I Crigler J F Jr and Wilkins I *Endocrinology* 51 117 (1952)
- 773 Miescher K Ischer W H and Tschopp T *Nature* 142 435 (1938)
- 774 Miller L *Endocrinology* 54 431 (1954)
- 775 Mirand E A Reinhard M C and Coltz H I *Proc Soc Exptl Biol Med* 81 397 (1952)
- 776 Mirick G S *Bull Johns Hopkins Hosp* 88 332 (1951)
- 777 Mogabgab W J and Thomas I *J Lab Clin Med* 39 271 (1952)
- 778 Moll F C and Hawn C van Z *J Immunol* 70 431 (1953)
- 779 Molomut N Spain D M and Haber A *Proc Soc Exptl Biol Med* 73 416 (1950)
- 780 Money W I, Fager J Jaques A and Rawson R W *Endocrinology* 51 87 (1952)
- 781 Money W L Fager J and Rawson R W *Cancer Research* 12 206 (1952)
- 782 Money W I Kirschner I Kraitsz I Merrill P and Rawson R W *J Clin Endocrinol and Metabolism* 10 1982 (1950)
- 783 Money W L Kraitsz L Fager J Kirschner I and Rawson R W *Endocrinology* 48 682 (1951)
- 784 Monto R W Brennan M J Margulis R R and Smith R W *J Lab Clin Med* 36 1008 (1950)
- 785 Moon V H and Tershakovec G A *Proc Soc Exptl Biol Med* 79 63 (1952)
- 786 Moore B and Purinton C O *Am J Physiol* 5 182 (1901)
- 787 Moore C R *J Clin Endocrinol and Metabolism* 13 339 (1953)
- 788 Moore D F Lowenthal J Fuller M and Jaques L B *Am J Clin Pathol* 22 936 (1952)
- 789 More R H and Kobernick S D *Am J Pathol* 27 708 (1951)
- 790 Morel F *Compt rend soc biol* 146 202 (1952)
- 791 Morgan A F and Lewis E M *Federation Proc* 11 451 (1952)
- 792 Morgan A F and Summs H D *Science* 89 565 (1939)
- 793 Morgan J A *Surgery* 30 506 (1951)
- 794 Mosher H A *Arch Ophthalmol* 45 317 (1951)
- 795 Mote J R (ed) *Proc 1st Clin ACTH Conf Phila* (1950)
- 796 Mote J R (ed) *Proc 2nd Clin ACTH Conf Phila* (1951)
- 797 Muehrcke R C Lewis J L and Kark R M *Science* 115 377 (1952)
- 798 Muehrcke R C Staple T W and Kark R M *J Lab Clin Med* 40 169 (1952)
- 799 Mueller J F Weir D R and Heinle R W *Proc Soc Exptl Biol Med* 77 312 (1951)
- 800 Muhnos M G and Lomerantz L *J Nutrition* 19 493 (1940)
- 801 Muhnos M G and Lomerantz L *Am J Physiol* 132 368 (1941)

- 802 Muknos M C Spingarn C I and Etkin M I *Am J Physiol* 135 102 (1941)
- 803 Murphy J B and Sturm I *Science* 90 703 (1941)
- 804 Murphy J B and Sturm I *Cancer Research* 10 191 (1950)
- 805 Murray R and Brinkham S I *Proc Soc Exptl Biol Med* 78 70 (1951)
- 806 Muirhead C W Stebbins R B and Barton M N *Trans N Y Acad Sci* 9 791 (1947)
- 807 Nachl L M and Schneider J J *Endocrinology* 51 5 (1952)
- 808 Nagareda C S and Caunt R *Endocrinology* 48 560 (1951)
- 809 Nathanson I T and Kelley H M *New Engl J Med* 246 175-180 (1952)
- 810 Neal W B Jr Woodward L R Park A I Zubaran J M and Montalbetti J A *Arch Surg* 65 74 (1952)
- 811 Nelson A A and Woodward C *Arch Pathol* 48 357 (1949)
- 812 Nelson C T Fox C I and Freeman L B *Proc Soc Exptl Biol Med* 75 181 (1950)
- 813 Nelson C T Fox C I and Freeman L B *J Invest Dermatol* 18 113 (1952)
- 814 Nelson D H and Samuels I T *J Clin Endocrinol and Metabolism* 12 519 (1952)
- 815 Nelson D H Samuels I T Willardson D C and Tyler F H *J Clin Endocrinol and Metabolism* 11 1021 (1951)
- 816 Nelson I J *Proc Soc Exptl Biol Med* 77 589 (1951)
- 817 Newsom S W and Durrach M *Can J Physiol & Biochem* 32 37 (1954)
- 818 Nichols J and Davis C *Federation Proc* 12 103 (1953)
- 819 Nichols J and Gardner I I *J Lab Clin Med* 37 279 (1951)
- 820 Nichols J and Sheehan H I *Endocrinology* 51 362 (1952)
- 821 Nissim J A *J Endocrinol* 8 10 (1952)
- 822 Nissim J A *Endocrinology* 52 611 (1953)
- 823 Noble R I *The Hormones Vol 2 Academic Press New York 1950 p 65*
- 824 Noble R I *Can J Physiol & Biochem* 33 in press (1955)
- 825 Noble R I and Collip J B *Quart J Exptl Physiol* 31 700 (1942)
- 826 Noble R I and Crispin I J *Rec can Biol* 11 57 (1952)
- 827 Noble R I and Tolly C G *J Endocrinol* 5 203 (1944)
- 828 Norcross B M Lockie I M Constantine A C Talbot J H and Stein R H *Ann Internal Med* 36 751 (1952)
- 829 Novack A *Am J Physiol* 168 121 (1952)
- 830 Nowakowski H K and Puskel L *Acta Endocrin* 11 370 (1952)
- 831 O'Donnell W M and Fajans S S *Univ Mich Med Bull* 16 169 (1950)
- 832 O'Donnell W M Fajans S S and Weirbaum J C *Arch Internal Med* 68 28 (1951)
- 833 Oesterling M J and Long C N H *Science* 113 241 (1951)
- 834 Ojers G W Yasuna J M and Schrie H G *Am J Ophthalmol* 36 120 (1953)
- 835 Oleson J J *Arch Biochem* 29 413 (1950)
- 836 Olhagen B *Scand Med* 43 78 (1950)
- 837 O'Neal J W and Heinbecker I *Endocrinology* 53 60 (1953)
- 838 Oppenheim L and Bruger M *Circulation* 6 410 (1952)
- 839 Opsahl J C *Yale J Biol and Med* 22 115 (1949)
- 840 Ormsby H L Dempster C and van Rooven C F *Am J Ophthalmol* 34 1687 (1951)

- 841 Osnes M and Thorsen R K *Acta Med Scand* 145 44 (1953)
- 842 Overman J R and Hanan R *Proc Soc Exptl Biol Med* 82 477 (1953)
- 843 Pabst M I Sheppard R and Kuizenga M H *Endocrinology* 41 50 (1944)
- 844 Ladawer J and Gordon A S *Endocrinology* 51 57 (1952)
- 845 Page I W and Glendening M B *Proc Soc Exptl Biol Med* 82 466 (1953)
- 846 Page I H *Am J Physiol* 122 357 (1938)
- 847 Iage I H and Lewis I A *Am J Physiol* 164 61 (1951)
- 848 Panzenhagen H and Speirs R S *Blood* 8 536 (1953)
- 849 Purkes A S and Smith A U *J Physiol* 124 13 (1954)
- 850 Farmer I G Katonach F and August A A *Proc Soc Exptl Biol Med* 77 215 (1951)
- 851 Paschkis K I Ipstein D Cantarow A and Friedler G *J Clin Endocrinol and Metabolism* 12 939 (1952)
- 852 Paschkis K L Cantarow A Lberhard T and Boyle D *Proc Soc Exptl Biol Med* 73 116 (1950)
- 853 Paterson J C and Mitchell C A *Irish J Pathol* 52 260 (1951)
- 854 Payne R W *Endocrinology* 45 305 (1949)
- 855 Pearson O H and Eliel I I *J Am Med Assoc* 144 1349 (1950)
- 856 Pearson O H and Eliel L I *Recent Progr Hormone Research* 6 3,3 (1951)
- 857 Pearson O H Eliel I P and Hollander V I *J Clin Invest* 30 665 (1951)
- 858 Pearson O H Eliel I P and Talbot T R *Cancer Research* 10 235 (1950)
- 859 Pearson O H Mendelsohn M I and West C D *J Clin Endocrinol and Metabolism* 13 841 (1953)
- 860 Pearson O H Hollander V I West C D Whitmore W F and Randall H T *J Clin Invest* 31 653 (1952)
- 861 Peczenik O *Proc Roy Soc Edinburgh B* 62 50 (1944)
- 862 Pellegrino P C Morris G M and Trubowitz S *Proc Soc Exptl Biol Med* 74 330 (1950)
- 863 Perera G A *Proc Soc Exptl Biol Med* 76 583 (1951)
- 864 Perlman P I and Cassidy J W *Proc Soc Exptl Biol Med* 77 237 (1951)
- 865 Perlmutter M and Mufson M *J Clin Endocrinol and Metabolism* 11 77 (1951)
- 866 Perry W F *Endocrinology* 49 284 (1951)
- 867 Perry W F *Can J Med Sci* 30 36 (1952)
- 868 Perry W F Hawkins W W and Cumming G R *Am J Physiol* 172 259 (1953)
- 869 Persky L and Jacob S *Proc Soc Exptl Biol Med* 77 66 (1951)
- 870 Peschel E Black Schaffler B and Schlayer C *Endocrinology* 48 399 (1951)
- 871 Pick R and Stamler J *Federation Proc* 13 112 (1954)
- 872 Pihero S J Landau D and Gordon A S *Science* 112 559 (1950)
- 873 Pincus G *J Clin Endocrinol and Metabolism* 3 195 (1943)
- 874 Pincus G *Psychosomat Med* 12 275 (1950)
- 875 Pincus G and Elmadjian F *Ann Rev Physiol* 16 403 (1953)
- 876 Pincus G Romanoff L P and Carlo J *J Clin Endocrinol and Metabolism* 8 221 (1948)
- 877 Pincus J B Natelson S and Lugovoy J K *Proc Soc Exptl Biol Med* 78 24 (1951)
- 878 Pine, I Engel F L and Schwartz T B *J Electroenceph and Clin Neurophysiol* 3 301 (1951)

- 879 Pirani C L Stepto R C and Sutherland A *J Exptl Med* 93 217 (1951)
- 880 Plotz C M Howes F I Blunt J W Meyer K and Ragan C *Arch Dermatol and Syphilol* 61 919 (1950)
- 881 Plotz C M Howes F I Meyer K Blunt J W Lattes R and Ragan C *Am J Pathol* 26 709 (1950)
- 882 Polley H F and Mason H I *J Am Med Assoc* 143 1474 (1950)
- 883 Polley H F Ward I L Slocumb C H and Hench P S *J Lab Clin Med* 38 937 (1951)
- 884 Priestley J T Walters W Sprague R G and Salassa R M *Ann Surg* 134 464 (1951)
- 885 Prunty T T G Forsham P H and Thorn G W *Clin Sci* 7 109 (1948)
- 886 Quttnier H Wald N Sussmann I N and Antopol W *Blood* 6 513 (1951)
- 887 Rabinowitz M and Ratsimamanga A R *Compt rend soc biol* 144 1466 (1950)
- 888 Ragan C Howes F I Plotz C M Meyer K and Blunt J W *Proc Soc Exptl Biol Med* 72 718 (1949)
- 889 Rall F P *Endocrinology* 39 275 (1946)
- 890 Rall F P and Dumm M F *Endocrinology* 61 135 (1952)
- 891 Rall F P and Dumm M I *Vitamins and Hormones* 11 133 (1953)
- 892 Rall F I and Graef I *Endocrinology* 32 1 (1943)
- 893 Rall F P and Graef I *Endocrinology* 37 252 (1945)
- 894 Rambo J H T *Arch Otolaryngol* 55 554 (1950)
- 895 Random L and Michaux A *Compt rend* 183 105 (1926)
- 896 Ratsimamanga A R *Compt rend soc biol* 138 19 (1944)
- 897 Ratsimamanga A R and Giroud A *Compt rend soc biol* 138 19 (1944)
- 898 Rawlin A G *Ann Allergy* 10 410 (1952)
- 899 Real C H Vennig F H and Ripstein M P *J Clin Endocrinol and Metabolism* 10 815 (1950)
- 900 Rebuck J W and Mellinger R C *Ann N Y Acad Sci* 66 715 (1953)
- 901 Recant I Ott W H and Fichel E F *Proc Soc Exptl Biol Med* 75 264 (1950)
- 902 Recant L Hume D M Forsham P H and Thorn G W *J Clin Endocrinol and Metabolism* 10 187 (1950)
- 903 Reid E Smith R H and Young F G *Biochem J* 42 xiv (1948)
- 904 Reinhardt W O and Fitch C H *Science* 117 295 (1953)
- 905 Reinhardt W O Aron H and La C H *Proc Soc Exptl Biol Med* 67 19 (1944)
- 906 Reiss M *Intern Record of Med* 166 190 (1953)
- 907 Reiss M and Halberston J M *J Endocrinol* 6 569 (1950)
- 908 Rhoads C I *Cancer* 2 913 (1949)
- 909 Rice K K and Richter C P *Endocrinology* 33 106 (1943)
- 910 Rich A R Berthrong M and Bennett I L *Bull Johns Hopkins Hosp* 87 549 (1950)
- 911 Rich A R Cochran T H and McGoon D C *Bull Johns Hopkins Hosp* 83 101 (1951)
- 912 Rich A R Bennett I L Cochran T H Griffith P C and McGoon D C *Bull Johns Hopkins Hosp* 88 189 (1951)
- 913 Richards J B and Sayers G *Proc Soc Exptl Biol Med* 77 87 (1951)
- 914 Riedel B E and Rossiter R J *Can J Physiol & Biochem* 32 761 (1954)

- 915 Riedel B E Iogan J E Deluca H A and Ro siter R J *Endocrinology* 55 219 (1954)
- 916 Ritchie A C Shubik P Lane M and Ieroy I P *Cancer Research* 13 45 (1953)
- 917 Rivero Pontan J Pa chkis K I West J and Cantarow A *Endocrinology* 51 100 (1952)
- 918 Roberts E Karnofsky D A and Frankel S *Proc Soc Exptl Biol Med* 76 289 (1951)
- 919 Roberts I Ronzoni L and Frankel S *Cancer Research* 11 275 (1951)
- 920 Roberts, K I and Pitts R F *Endocrinology* 60 51 (1957)
- 921 Roberts K I and Pitts R F *Endocrinology* 52 324 (1953)
- 922 Roberts S *J Biol Chem* 200 77 (1952)
- 923 Roberts S and White A *Endocrinology* 48 741 (1951)
- 924 Robinson H J *Federation Proc* 10 332 (1951)
- 925 Robinson H J *In J M Yoffey, The Suprarenal Cortex* Academic Press New York 1953 p 105
- 926 Robinson H J and Smith A I *Ann N Y Acad Sci* 56 757 (1953)
- 927 Robinson H J Mason R C and Smith A I *Proc Soc Exptl Biol Med* 84 312 (1953)
- 928 Robinson S Kirkaid R C and Rhamy R K *J Appl Physiol* 2 399 (1950)
- 929 Robinson W D Wolfson W O and Duff I F *Ann Rheumatic Diseases* 11 313 (1952)
- 930 Robson H N and Duthie J J R *Brit Med J* 11, 971 (1950)
- 931 Robson H N and Duthie J J R *Brit Med J* 1 994 (1952)
- 932 Roche P Cummings M M and Hudgins P C *Am Rev Tuberc* 65 603 (1952)
- 933 Rockwell L M and Johnson P *J Invest Dermatol* 19 137 (1952)
- 934 Roemmelt J C Sartorius O W and Pitts R F *Am J Physiol* 159 124 (1949)
- 935 Rogoff J M Nixon F N and Stewart G N *Proc Soc Exptl Biol Med* 41 57 (1939)
- 936 Rogoff J M Quishnock J M Nixon N and Rosenberg A W *Proc Soc Exptl Biol Med* 73 163 (1950)
- 937 Ronzoni I Roberts E Frankel S and Ramasarma G B *Proc Soc Exptl Biol Med* 82 496 (1953)
- 938 Rose B *Recent Progr Hormone Research* 7 315 (1957)
- 939 Rosenberg E and Lewis R A *J Appl Physiol* 3 164 (1950)
- 940 Rosenberg F Cornfield J Bates R W and Anderson F *Endocrinology* 54 363 (1954)
- 941 Rosenbaum J D Papper S and Ashley M M *J Clin Invest* 31 637 (1952)
- 942 Rosenberg N Kunderman P J Vroman I and Moolten S F *Arch Surg* 63 1 (1951)
- 943 Rosenberg N Kunderman P J Vroman L and Moolten S F *Arch Surg* 66 593 (1953)
- 944 Rosenman R H Freed S C and Friedman M *Proc Soc Exptl Biol Med* 78 77 (1951)
- 945 Rosenthal R L Fless A D and Litwins J *Acta Haematol* 6 114 (1951)
- 946 Rosenthal R I Wald N Yager A and Litwins J *Proc Soc Exptl Biol Med* 75 740 (1950)

- 947 Rosvold H E Kaplan S J and Stevenson J A F *Proc Soc Exptl Biol Med* 80 60 (1953)
- 948 Rothman S and Dillaha C J *Arch Dermatol and Syphilol* 65 749 (1952)
- 949 Rowe C R Grimson K S Flowe B H Lyons C K Longino F H and Taylor H M *Surgery* 32 276 (1952)
- 950 Roy A C Kar A B and Sur R N *Acta Endocrinol* 14 254 (1953)
- 951 Ruwek H I Russek A S Doerner A A and Zohman B I *Arch Internal Med* 91 457 (1953)
- 952 Russell J A *Ann Rev Physiol* 13 327 (1951)
- 953 Russell J A and Wilhelm A L *J Biol Chem* 137 713 (1941)
- 954 Sala G Sereni F and Ballabio C B *Lancet* ii 1090 (1951)
- 955 Sala S R M *J Clin Endocrinol and Metabolism* 13 857 (1953)
- 956 Sames G I and Leatham J H *Proc Soc Exptl Biol Med* 78 231 (1951)
- 957 Saniberg A A Nelson D H Palmer J G Samuels L T and Tyler F H *J Clin Endocrinol and Metabolism* 13 629 (1953)
- 958 Sapirstein I A Brandt W I and Drury D R *Proc Soc Exptl Biol Med* 73 82 (1950)
- 959 Sartorius O W Calhoun D and Pitts R F *Endocrinology* 51 444 (1952)
- 960 Sayers G *Physiol Revs* 30 241 (1950)
- 961 Sayers G and Sayers M A *Ann N Y Acad Sci* 50 572 (1950)
- 962 Sayers G Burns T W Tyler F H Jager B V Schwartz T B Smith E L Samuels L T and Davenport H W *J Clin Endocrinol and Metabolism* 9 593 (1949)
- 963 Schaffenburg C Masson G M C and Corcoran A C *Proc Soc Exptl Biol Med* 74 358 (1950)
- 964 Scheinberg S R and Saltzstein H C *Arch Surg* 63 413 (1951)
- 965 Schenker V *Endocrinology* 53 345 (1953)
- 966 Scherr G H *Science* 116 685 (1952)
- 967 Schieve J F Scheinberg P and Wilson W P *J Clin Invest* 30 1527 (1951)
- 968 Schiffer F and Wertheimer L J *J Endocrinol* 6 147 (1947)
- 969 Schiller E *Brit J Ind Med* 10 1 (1953)
- 970 Schmidt L H and Squires W I *J Exptl Med* 94 501 (1951)
- 971 Schneebeli G L and Dougherty T F *Am J Physiol* 167 825 (1951)
- 972 Schneebeli G L Dougherty T F and Loewe S *Proc Soc Exptl Biol Med* 77 407 (1951)
- 973 Schoch E P and Glick D *J Invest Dermatol* 20 119 (1953)
- 974 Schrek R *Endocrinology* 45 317 (1949)
- 975 Schrek R *Proc Soc Exptl Biol Med* 76 557 (1951)
- 976 Schrek R *Proc Soc Exptl Biol Med* 77 709 (1951)
- 977 Schricker J A Fisher G R Tullner W W and Hertz R *Endocrinology* 51 394 (1952)
- 978 Schricker J A Hertz R and Tullner W W *Proc Soc Exptl Biol Med* 78 572 (1951)
- 979 Schuler W Desaulles P and Meier R *Experientia* 10 142 (1954)
- 980 Schulman I Lanman J T Laydal O E and Holt L E *Pediatrics* 8 34 (1951)
- 981 Schultz R B Winters R W and Krehl W A *Endocrinology* 51 336 (1952)
- 982 Schwartz T B and Engel F L *J Biol Chem* 180 1047 (1949)
- 983 Schwartz T B and Engel F I *Proc Soc Exptl Biol Med* 74 82 (1950)
- 984 Scott J L Jr and Engel F L *Endocrinology* 53 410 (1953)

- 985 Secker, J *J Physiol* 109 49 (1949)
- 986 Segaloff A and Horwitz B N *Science* 118 220 (1953)
- 987 Segaloff A and Many A S *Endocrinology* 49 390 (1951)
- 988 Seifter J Baeder D H and Begany A J *Proc Soc Exptl Biol Med* 72 277 (1949)
- 989 Seifter J, Worter P J and Fitch D R *Proc Soc Exptl Biol Med* 73 131 (1950)
- 990 Seifter J Ehrlich W E, Begany A J, and Warren G H *Proc Soc Exptl Biol Med* 75 337 (1950)
- 991 Seifter J Ehrlich W F Bieder D H Butt A J and Hau er, E A *Ann N Y Acad Sci* 66 693 (1953)
- 992 Sellers F A You R W Ridout J H and Best C H *Nature* 166, 514 (1950)
- 993 Selye H *J Immunol* 41 259 (1911)
- 994 Selye H *Endocrinology* 30 437 (1942)
- 995 Selye H *Brit Med J* 11 1129 (1949)
- 996 Selye H *Can Med Assoc J* 61 553 (1949)
- 997 Selye H *In Stress Acta Inc Montreal* 1950
- 998 Selye H *Can Med Assoc J* 64 489 (1951)
- 999 Selye H *Proc Soc Exptl Biol Med* 76 510 (1951)
- 1000 Selye H *Federation Proc* 11 144 (1952)
- 1001 Selye H *Proc Soc Exptl Biol Med* 82 328 (1953)
- 1002 Selye H *J Am Med Assoc* 162 1207 (1953)
- 1003 Selye H *Abstr 19th Intern Physiol Congr Montreal* p 750 (1953)
- 1004 Selye H *Recent Progr Hormone Research* 8 117 (1953)
- 1005 Selye H and Carey N *J Clin Endocrinol and Metabolism* 10 824 (1950)
- 1006 Selye H Hall C L and Rowley E M *Can Med Assoc J* 49 88 (1943)
- 1007 Selye H Hall O and Rowley F M *Lancet* 1 301 (1945)
- 1008 Seneeca H and Rockenbach J *Science* 116 14 (1952)
- 1009 Sensenbach W and Madison I *Am J Med* 13 652 (1952)
- 1010 Sensenbach W Madison I and Ochs L *J Clin Invest* 32 372 (1953)
- 1011 Shacter B Intenman C and Shumkin M B *J Natl Cancer Inst* 13 647 (1952)
- 1012 Shanaphy J F *Am J Obstet Gynecol* 64 1761 (1952)
- 1013 Shapiro R Taylor B and Taubenhaus M *Proc Soc Exptl Biol Med* 76 854 (1951)
- 1014 Sheehan H L Summers V K and Nichols J *Lancet* 1 312 (1953)
- 1015 Sheldon W H Cummings M M and Evans L D *Proc Soc Exptl Biol Med* 75 616 (1950)
- 1016 Sheldon W H Herman A and Evans I D *Am J Syphilis Gonorrhea Venereal Diseases* 36 77 (1952)
- 1017 Shepanek L A *Surg Gynecol Obstet* 96 200 (1953)
- 1018 Shick R M Baggenstoss A H and Polley H F *Proc Staff Meetings Mayo Clinic* 25 135 (1950)
- 1019 Shuman C R and Finestone A J *Proc Soc Exptl Biol Med* 73 248 (1950)
- 1020 Schwartzman G *Proc Soc Exptl Biol Med* 75 835 (1950)
- 1021 Schwartzman G *Am J Pathol* 27 714 (1951)
- 1022 Schwartzman G and Aronson S M *Ann N Y Acad Sci* 66 793 (1953)
- 1023 Schwartzman G and Fisher A *J Exptl Med* 95 347 (1952)
- 1024 Siebenmann R *Schweiz Z Pathol u Bakteriologie* 15 174 (1952)

- 1025 Silber R H and Porter C C *Endocrinology* 52 518 (1953)
- 1026 Simpson S A and Tait J F *Endocrinology* 50 150 (1952)
- 1026a Simpson S A and Tait J F *Recent Progr Hormone Research* 11 183 (1955)
- 1027 Simpson S A Tait J F and Bush I I *Lancet* ii 726 (1952)
- 1028 Simpson S A Tait J I Wettstein A Neher R von Fuw J and Reichstein T *Experientia* 9 333 (1953)
- 1029 Simpson S A Tait J F Wettstein A Neher R von Fuw J Schindler O and Reichstein T *Experientia* 10 132 (1954)
- 1030 Simek F M *Federation Proc* 10 217 (1951)
- 1031 Singer B and Venning E H *Endocrinology* 52 623 (1953)
- 1032 Sirek O A and Best C H *Proc Soc Exptl Biol Med* 80 594 (1952)
- 1033 Sissons H A and Hadfield C J *Brit J Surg* 39 172 (1951)
- 1034 Sjogren B *Acta Endocrinol* 10 40 (1952)
- 1035 Slessor A *J Clin Endocrinol and Metabolism* 11 700 (1951)
- 1036 Slocombe A G Hoagland H and Praglin J *Federation Proc* 2 149 (1950)
- 1037 Smith D J *Federation Proc* 10 749 (1951)
- 1038 Smith D I and d'Amour F F *J Pharmacol Exptl Therap* 106 429 (1952)
- 1039 Smith G H and Opsahl J C *Yale J Biol and Med* 23 361 (1951)
- 1040 Smith J M Murphy J S and Mirick G S *Proc Soc Exptl Biol Med* 78 50 (1951)
- 1041 Smith R H *Biochem J* 44 xlii (1949)
- 1042 Smith R W *Proc Soc Exptl Biol Med* 78 568 (1951)
- 1043 Smith R W and Steffenen E H *New Engl J Med* 245 972 1007 (1951)
- 1044 Smith R W Morgulis R R Brennan M J and Monto R W *Science* 112 795 (1950)
- 1045 Smith S G and Lasater T L *Proc Soc Exptl Biol Med* 74 427 (1950)
- 1046 Smith W W Smith F and Thompson E C *Proc Soc Exptl Biol Med* 73 529 (1950)
- 1047 Snelling C E Donohue W L Laski B and Jackson S H *Pediatrics* 8 22 (1951)
- 1048 Snyder J G and Wyman I C *Endocrinology* 49 205 (1951)
- 1049 Soffer L J Gabrielove J L and Dorrance W R *Proc Soc Exptl Biol Med* 76 763 (1951)
- 1050 Soffer L J Gabrielove J I and Jacobs M D *J Clin Invest* 28 1091 (1949)
- 1051 Soffer L J Gabrielove J I and Jailer J W *Proc Soc Exptl Biol Med* 71 117 (1949)
- 1052 Soffer L J Schwartzman G Schneiersson S S and Gabrielove J L *Science* 111 303 (1950)
- 1053 Sokoloff L Sharp J T and Kaufman L H *Am J Pathol* 27 706 (1951)
- 1054 Solomon D H and Shock N W *J Gerontol* 5 302 (1950)
- 1055 Splotorovsky M Gregory F J and Stoerk H C *Proc Soc Exptl Biol Med* 76 286 (1951)
- 1056 Somerville W *Brit Med J* ii 860 (1950)
- 1057 Somerville W Levine H D and Thorn G W *Medicine* 30 43 (1951)
- 1058 Sommers S C Wilson J C and Hartman F W *J Exptl Med* 93 505 (1951)
- 1059 Sourkes T L and Heneage P *Endocrinology* 49 601 (1951)
- 1060 Sourkes T L and Heneage P *Endocrinology* 50 43 (1952)
- 1061 Southam C M and Babcock V I *Proc Soc Exptl Biol Med* 78 105 (1951)

- 1065 Spain D C Fontant B J and Strauss M B *J Allergy* 23 212 (1952)
- 1066 Spain D M and Molomut N *Am Rev Tuberc* 62 337 (1950)
- 1067 Spain D M and Thallmer W *Proc Soc Exptl Biol Med* 76 320 (1951)
- 1068 Spain D M Molomut N and Haber A *Am J Pathol* 26 710 (1950)
- 1069 Spain D M Molomut N and Haber A *Science* 112 335 (1950)
- 1070 Spain D M Molomut N and Haber A *J Lab Clin Med* 39 383 (1952)
- 1071 Spait F H and Mednicoff I *Bull New Engl Med Center* 13 301 (1951)
- 1072 Sparrow E M *J Endocrinol* 9 101 (1953)
- 1073 Speirs R S *Science* 113 691 (1951)
- 1074 Speirs R S and Meyer R K *Endocrinology* 45 403 (1949)
- 1075 Speirs R S and Meyer R K *Endocrinology* 48 316 (1951)
- 1076 Spencer A G *Nature* 166 32 (1950)
- 1077 Spies T D and Stone R L *Lancet* ii 890 (1949)
- 1078 Spies T D Stone R I Garcia Lopez C Milanes F Lopez Toea R and Reboredo A *Lancet* ii 241 (1950)
- 1079 Sprague R C *Am J Med* 10 567 (1951)
- 1080 Sprague R G Mason H I and Power M H *Recent Progr Hormone Research* 6 315 (1951)
- 1081 Sprague R C Power M H and Mason H L *J Am Med Assoc* 144 1341 (1950)
- 1082 Sprague R G Power M H Mason H I and Cluxton H F *J Clin Invest* 28 812 (1949)
- 1083 Sprague R G Power M H Mason H I Albert A Mathieson D R Hench P S Kendall F C Blocumb C H and Polley H F *Arch Internal Med* 85 199 (1950)
- 1084 Sprechler M and Vesterdal J *Acta Endocrinol* 12 207 (1953)
- 1085 Stadie W C *Ann N Y Acad Sci* 64 641 (1951)
- 1086 Stadie W C *Physiol Revs* 34 52 (1954)
- 1087 Stadie W C and Haugaard N *J Biol Chem* 177 311 (1949)
- 1088 Stadie W C Haugaard N and Hills A G *J Biol Chem* 184 617 (1950)
- 1089 Stadie W C Haugaard N and Marsh J B *J Biol Chem* 198 780 (1952)
- 1090 Stadie W C Haugaard N and Vaughan M *J Biol Chem* 199 729 (1952)
- 1091 Stamler J and Pick R *Proc Soc Exptl Biol Med* 75 803 (1950)
- 1092 Stamler J Pick R and Katz L N *Circulation* 4 461 (1951)
- 1093 Stand R Anderson W and Allcroft W M *Biochem J* 28 640 (1934)
- 1094 Stebbins R B *Endocrinology* 49 25 (1951)
- 1095 Stebbins R B *Endocrinology* 49 671 (1951)
- 1096 Stien A S *Brit J Ophthalmol* 35 741 (1951)
- 1097 Steffann M and Rosenthal M C *Proc Soc Exptl Biol Med* 76 806 (1950)
- 1098 Steffensen L H Ivy H B and Nagle F O *Am J Ophthalmol* 35 933 (1952)
- 1099 Steffensen L H Olson J A Margulis R R and Smith R W *Am J Ophthalmol* 33 1033 (1950)
- 1100 Stein K F *Proc Soc Exptl Biol Med* 71 225 (1949)
- 1101 Stein K F and Martin C *Proc Soc Exptl Biol Med* 78 513 (1951)
- 1102 Stephens C A I Wallraff E B Borden A I Brodie L C Holbrook W P Hill D F Kent L J and Kemmerer A R *Proc Soc Exptl Biol Med* 74 275 (1950)
- 1103 Stepto R C Pirani C I Consolazio C F and Bell J H *Endocrinology* 49 755 (1951)

- 1101 Stepto R C Pirani C I Fisher J D and Sutherland K *Federation Proc* **11** 479 (1952)
- 1102 Stickney J M Heck F J and Watkins C H *Proc Staff Meetings Mayo Clinic* **25** 488 (1950)
- 1103 Stier R A Feinberg S M Malkiel S and Werle M D *J Allergy* **23** 390 (1957)
- 1104 Solomon D H and Humphreys S R *Blood* **6** 894 (1951)
- 1105 Stoerk H C *Proc Soc Exptl Biol Med* **62** 90 (1946)
- 1106 Stoerk H C *Federation Proc* **9** 345 (1950)
- 1107 Stoerk H C *Proc Soc Exptl Biol Med* **74** 798 (1950)
- 1108 Stoerk H C *Ann N Y Acad Sci* **56** 742 (1953)
- 1109 Stoerk H C and Porter C C *Proc Soc Exptl Biol Med* **74** 60 (1950)
- 1110 Stoerk H C and Solotorovsky M *Am J Pathol* **26** 708 (1950)
- 1111 Straube R L Patt H M Tyree E B and Smith D E *Proc Soc Exptl Biol Med* **71** 539 (1949)
- 1112 Sugura K *Cancer Research* **13** 431 (1953)
- 1113 Sugura K Stock C C Dobriner K and Rhoads C P *Cancer Research* **10** 244 (1950)
- 1114 Sulzberger M B and Witten V H *J Invest Dermatol* **19** 101 (1952)
- 1115 Sulzberger M B Witten V H and Smith C C *J Am Med Assoc* **151** 468 (1953)
- 1116 Sure B and Theis R M *Endocrinology* **24** 673 (1939)
- 1117 Sutherland E W *Ann N Y Acad Sci* **54** 693 (1951)
- 1118 Swann H G *Physiol Revs* **20** 493 (1940)
- 1119 Sweat M I Abbott W E Jefferies W McK and Bliss E L *Federation Proc* **12** 141 (1953)
- 1120 Swingle W W Collins F Barlow G and Fedor F J *Am J Physiol* **168** 118 (1957)
- 1121 Swingle W W Perlmutt J Seay P and Collins E *Am J Physiol* **169** 778 (1952)
- 1122 Swingle W W Barlow G Collins E Fedor E J Welch W J and Rampona J M *Endocrinology* **51** 353 (1952)
- 1123 Swingle W W Collins E Barlow G Fedor E J Ben M and Maxwell R *Proc Soc Exptl Biol Med* **81** 52 (1952)
- 1124 Swingle W W Fedor E J Ben M Maxwell R Baker C and Barlow G *Proc Soc Exptl Biol Med* **82** 511 (1953)
- 1125 Swingle W W Perlmutt J Collins E Seay P Fedor E and Barlow G *Proc Soc Exptl Biol Med* **78** 865 (1951)
- 1126 Swingle W W Barlow G Fedor E Ben M Maxwell R Collins E and Baker C *Am J Physiol* **173** 4 (1953)
- 1127 Syverton J T Werder A A Friedman J Roth F J Graham A B and Mira O J *Proc Soc Exptl Biol Med* **80** 173 (1952)
- 1128 Szego C M *Endocrinology* **50** 429 (1952)
- 1129 Szego C M and White A *Endocrinology* **44** 150 (1949)
- 1130 Szilagyi D E McGraw A B and Smyth N P D *Ann Surg* **136** 555 (1957)
- 1131 Tait J F Grundy H M and Simpson S A *Lancet* **1** 122 (1952)
- 1132 Talalay P Dobson M M Ebersole C M and Huggins C *Endocrinology* **50** 574 (1952)
- 1133 Taubenhaus M *Ann N Y Acad Sci* **56** 666 (1953)

- 1131 Taubenhaus M and Amromin G D *J Lab Clin Med* 36 7 (1950)
- 1132 Taubenhaus M Taylor B and Morton J B *Endocrinology* 51 183 (1952)
- 1133 Teicher R and Nelson C T *J Invest Dermatol* 19 205 (1952)
- 1134 Teilum G Imbäck H C and Simonson M *Acta Endocrinol* 5 181 (1950)
- 1135 Teng C Saxe I M Deane H W and Hastings A B *J Cellular Comp Physiol* 39 73 (1952)
- 1136 Tepperman J and Tepperman H M *Ann N Y Acad Sci* 54 707 (1951)
- 1137 Terry I I and London I *Proc Soc Exptl Biol Med* 73 251 (1950)
- 1138 Thiersch J B Conroy J, Stevens A R and Finch C A *J Lab Clin Med* 40 171 (1952)
- 1139 Thomas I *Ann N Y Acad Sci* 56 799 (1953)
- 1140 Thomas I and Good R A *Proc Soc Exptl Biol Med* 76 604 (1951)
- 1141 Thomas I and Mogabgab W J *Proc Soc Exptl Biol Med* 74 879 (1950)
- 1142 Thorn G W *J Clin Endocrinol and Metabolism* 13 614 (1953)
- 1143 Thorn C W Engel I I, and Lewis R A *Science* 94 348 (1941)
- 1144 Thorn C W Jenkins D and Iudlow J C *Recent Progr Hormone Research* 8, 171 (1953)
- 1145 Thorn G W Goetz I C Streeten D H I Dingman J F and Arons W I *J Clin Endocrinol and Metabolism* 13 604 (1953)
- 1146 Thorn C W Forsham I H Frawley T F Hill F R Roche M Staehelin D and Wilson D I *New Engl J Med* 242 783 824 865 (1950)
- 1147 Thorn C W Forsham P H Frawley T F Wilson D I Renold A F Fredrickson D S and Jenkins D *Am J Med* 10 595 (1951)
- 1148 Thorn G W Jenkins D Iudlow J C Goetz I C Dingman J F Arons W I Streeten H I and McCracken B H *New Engl J Med* 245 237 281 323 369 414 458 632 (1953)
- 1149 Thygeson P Geller H O and Schwartz A *Am J Ophthalmol* 34 885 (1951)
- 1150 Tipton S R *Am J Physiol* 127 710 (1939)
- 1151 Toolan H W *Proc Am Assoc Cancer Research* 1 56 (1953)
- 1152 Torda C and Wolff H C *Am J Physiol* 163 755 (1950)
- 1153 Treager H S Gabuzda C J Zamecheck N and Davidson C S *Proc Soc Exptl Biol Med* 75 517 (1950)
- 1154 Tuerkischer F and Wertheimer I *J Endocrinol* 4 143 (1945)
- 1155 Turkel J Zizine I Jeanjean A and Depraeter M *Presse med* 61 817 (1953)
- 1156 Turner I B and Crollman A *Am J Physiol* 167 462 (1951)
- 1157 Turner T B and Hollander D H *Bull Johns Hopkins Hosp* 87 505 (1950)
- 1158 Umbreit W W *Ann N Y Acad Sci* 54 561 (1951)
- 1159 Umbreit W W and Tonhazy N I *J Biol Chem* 191 249 (1951)
- 1160 Umbreit W W and Tonhazy N L *J Biol Chem* 191 257 (1951)
- 1161 Ungar G *Am J Physiol* 167 833 (1951)
- 1162 Ungar G *Lancet* II 742 (1952)
- 1163 Ungar G and Damgaard E *J Exptl Med* 93 89 (1951)
- 1164 Uotila U and Pekkarinen A *Acta Endocrinol* 6 23 (1951)
- 1165 Upton A C and Coon W W *Proc Soc Exptl Biol Med* 77 153 (1951)
- 1166 Vaillancourt de G Grokoest A W and Ragun C *Proc Soc Exptl Biol Med* 78 383 (1951)
- 1167 Van der Scheer J Bohnel F Clarke F H and Wyckoff R W G *J Immunol* 44 165 (1947)

- 1171 Van Ordstrand H S *Ann Internal Med* 35 1203 (1951)
- 1172 Van Putten I M Van Bekkum D W and Querido A *Acta Endocrinol* 12 159 (1953)
- 1173 Van Scott E J Rothman S and Greene C R *J Invest Dermatol* 20 111 (1953)
- 1174 Vaughan J H Bayles T B and Favour C B *Proc Soc Exptl Biol Med* 76 274 (1951)
- 1175 Venning E H Pattee C J McCall F and Browne J S L *J Clin Endocrinol and Metabolism* 12 1409 (1952)
- 1176 Verzar F and Ann N Y Acad Sci 54 116 (1951)
- 1177 Vilcek C A and Hastings A B *J Biol Chem* 179 673 (1949)
- 1178 Vogel A W and Leopold I H *Am J Ophthalmol* 36 690 (1953)
- 1179 Vogt M In J M Yoffey *The Suprarenal Cortex* Academic Press New York 1953 p 59
- 1180 Volk B W Lazarus S S and Tui C *Proc Soc Exptl Biol Med* 80 1 (1952)
- 1181 Wahlstrom R C and Johnson B C *Proc Soc Exptl Biol Med* 78 112 (1951)
- 1182 Walbach S B and Maddock C L *Arch Pathol* 53 54 (1952)
- 1183 Waldenstrom J *Acta Endocrinol* 5 235 (1950)
- 1184 Wall J W and Shure N *Arch Otolaryngol* 56 172 (1952)
- 1185 Wallach D P and Reineke E P *Endocrinology* 45 75 (1949)
- 1186 Ward L E Slocumb C H Polley H F Lowman L W and Hench P S *Proc Staff Meetings Mayo Clinic* 26 361 (1951)
- 1187 Watson D Bidmead D S and Hersley G D *Ann Rheumatic Diseases* 11 297 (1952)
- 1188 Wayne H L and Bayle J *J Clin Endocrinol and Metabolism* 13 1010 (1953)
- 1189 Weimer H W Boak R A Bogen E Drusch H E Miller J N Moshin J R and Carpenter C M *Am Rev Tuberc* 68 31 (1953)
- 1190 Weir D R and Heinle R W *Proc Soc Exptl Biol Med* 75 655 (1950)
- 1191 Weir D R Heinle R W and Welch A D *Proc Soc Exptl Biol Med* 72 457 (1949)
- 1192 Weisberger A S and Levine B *J Clin Invest* 31 671 (1952)
- 1193 Weisman P A Wight A Quinby W C and Cannon B *Plastic and Reconstruct Surg* 8 417 (1951)
- 1194 Welbourn R B and Code C F *Gastroenterology* 23 356 (1953)
- 1195 Wells B B and Kendall E C *Proc Staff Meetings Mayo Clinic* 15 524 (1940)
- 1196 Welt I and Wilhelm A E *Yale J Biol and Med* 23 99 (1950)
- 1197 Wentworth J H and Billows J A *Radiology* 59 559 (1952)
- 1198 Werder A A Friedman J MacDowell E C and Syvertson J T *Cancer Research* 13 158 (1953)
- 1199 Werner S C *J Clin Endocrinol and Metabolism* 13 1332 (1953)
- 1200 West C D Pearson O H and Kappas A *J Clin Endocrinol and Metabolism* 13 841 (1953)
- 1201 West C D Hollander V P Whitmore W F Randall H T and Pearson O H *Cancer* 5 1009 (1952)
- 1202 Wheeler C E Cawley E P and Curtis A C *J Invest Dermatol* 20 385 (1953)

- 1203 White A Hoberman H D and Szego C M *J Biol Chem* 174 1049 (1949)
- 1204 White M R Linker A J and Schubert J *Proc Soc Exptl Biol Med* 80 603 (1952)
- 1205 Whitney J L and Bennett I L *Endocrinology* 60 657 (1957)
- 1206 Wick A N, Drury D R and Mackay E M *Ann N Y Acad Sci* 64 684 (1951)
- 1207 Weiland P Heer J Schmidlin J and Miescher K *Helv Chim Acta* 34 351 (1951)
- 1208 Wilhelmy C M Walderman E B and McGuire T F *Proc Soc Exptl Biol Med* 77 379 (1951)
- 1209 Wilkins I Lewis R A Klein R and Rosenberg L *Helv Paediat Acta* 6 418 (1950)
- 1210 Wilkins I Lewis R A Klein R, and Rosenberg E *Bull Johns Hopkins Hosp* 86 249 (1950)
- 1211 Wilkins L Lewis R A, Klein, R and Rosenberg I *Am J Diseases Children* 80 883 (1950)
- 1212 Wilkins L Crigler J F Silverman S H Gardner L I and Migeon C J *J Clin Endocrinol and Metabolism* 12 1015 (1952)
- 1213 Wilkins L Lewis R A Klein R Gardner L I Crigler J F Jr Rosenberg E, and Migeon C J *J Clin Endocrinol and Metabolism* 11 1 (1951)
- 1214 Williams W Whismant C and Litts W T *Am J Physiol* 170 57 (1957)
- 1215 Willis, G C *Can Med Assoc J* 69, 17 (1953)
- 1216 Winter C A and Flataker I *Federation Proc* 9 137 (1950)
- 1217 Winter C A and Flataker I *J Pharmacol Exptl Therap* 105 358 (1957)
- 1218 Winter C A Hollings H L and Stebbins R B *Endocrinology* 52 123 (1953)
- 1219 Winter C A Silber R H and Stoerk H C *Endocrinology* 47 60 (1950)
- 1220 Winternitz W W and Long C V H *Proc Soc Exptl Biol Med* 81 683 (1952)
- 1221 Winters R W Schultz R B and Krehl W A *Endocrinology* 50 317 (1957)
- 1222 Winters R W Schultz R B and Krehl W A *Endocrinology* 50 388 (1957)
- 1223 Winters R W Schultz R B and Krehl W A *Proc Soc Exptl Biol Med* 79 693 (1952)
- 1224 Wolfson W Q *J Clin Endocrinol and Metabolism* 13 175 (1952)
- 1225 Wolfson W Q Beierwaltes W H Robinson W D Duff I F Jones J R Knorpp C T and Lya M *J Lab Clin Med* 36 1005 (1950)
- 1226 Wood J W *Federation Proc* 12 159 (1953)
- 1227 Woodbury D M *J Pharmacol Exptl Therap* 105 46 (1952)
- 1228 Woodbury D M and Davenport V D *Am J Physiol* 157 234 (1949)
- 1229 Woodbury D M and Sayers G *Proc Soc Exptl Biol Med* 75 398 (1950)
- 1230 Woodbury D M Ghosh B N and Sayers G *J Clin Endocrinol and Metabolism* 11 761 (1951)
- 1231 Woodbury D M Cheng C P Sayers G and Goodman L S *Am J Physiol* 160 217 (1950)
- 1232 Woodbury D M Emmett J W Hinckley G V Jackson N R Newton J D Bateman J H Goodman L S and Sayers G *Proc Soc Exptl Biol Med* 76 65 (1951)
- 1233 Woodruff M F A and Boswell T *J Endocrinol* 10 86 (1953)
- 1234 Woods A C *Am J Ophthalmol* 33 1325 (1950)
- 1235 Woods A C *Am J Ophthalmol* 34 913 (1951)
- 1236 Woods A C and Wood R M *Arch Ophthalmol* 47, 477 (1952)

- 1237 Woolley G W *Ann N Y Acad Sci* 50 616 (1949)
- 1238 Woolley G W *Proc Soc Exptl Biol Med* 74 286 (1950)
- 1239 Woolley G W *Cancer Research* 11 291 (1951)
- 1240 Woolley G W and Little C C *Cancer Research* 5 193 (1945)
- 1240a Woolley G W and Peters B A *Proc Soc Exptl Biol Med* 82 286 (1953)
- 1241 Wragg I E and Spears R S *Proc Soc Exptl Biol Med* 80 680 (1952)
- 1242 Wrong N M and Smith R C *Can Med Assoc J* 68 50 (1952)
- 1243 Wyman G J *Am J Ophthalmol* 35 1706 (1952)
- 1244 Wyman L C Fulton G P and Shulman M H *Ann N Y Acad Sci* 56 643 (1953)
- 1245 Wyman L C Fulton G P Sudak F N and Patterson G N *Proc Soc Exptl Biol Med* 84 280 (1953)
- 1246 Yoffey J M (ed) *In The Suprarenal Cortex* Academic Press New York 1953 p 31
- 1247 Yoffey J M Metcalf W K Herdan G and Nairn V *Brit Med J* 1 660 (1951)
- 1248 Zarrow M N and Baldini J T *Endocrinology* 50 555 (1952)
- 1249 Zarrow M N and Money W I *Endocrinology* 44 345 (1949)
- 1250 Zarrow M N and Zarrow I G *Anat Record* 108 112 (1950)
- 1251 Zarrow M N and Zarrow I G *Proc Soc Exptl Biol Med* 76 620 (1951)
- 1252 Zeckwer I T *Endocrinology* 52 39 (1953)
- 1253 Zierler K I and Ishenthal J L Jr *Am J Med* 4 186 (1948)
- 1254 Zilversmit D B Stern T N and Overman R R *Am J Physiol* 164 31 (1951)
- 1255 Zimmerman H J Parrish A E and Alpert L K *Proc Soc Exptl Biol Med* 73 81 (1950)
- 1256 Zimmerman H J Parrish A E and Alpert L K *Am J Med* 10 235 (1951)
- 1257 Zingg W and Perry W F *J Clin Endocrinol and Metabolism* 13 712 (1953)
- 1258 Zintel H A Wolferth C C Jeffers W A Hafkenschiel J H and Lukens F D W *Ann Surg* 134 351 (1951)
- 1259 Zondek B and Burstein S *Endocrinology* 50 419 (1952)
- 1260 Zubiran J M Kark A E Montalbetti J A Morel C J L and Dragstedt I R *Arch Surg* 65 809 (1952)
- 1261 Zubirán S and Gómez Mont F *Vitamins and Hormones* 11 97 (1953)
- 1262 Zuckerman S *In J M Yoffey The Suprarenal Cortex* Academic Press New York 1953 p 69
- 1263 Zweifach B W Shorr E and Block M M *Ann N Y Acad Sci* 56 676 (1953)

CHAPTER XV

Clinical Endocrinology

BY R. E. PASCHKIS AND A. E. RAHOFF

CONTENTS

	<i>P a g e</i>
I Introduction	821
II Anterior Pituitary Gland	822
A General Remarks	822
B Acromegaly and Gigantism	823
C Hypopituitarism	825
III Thyroid Gland	827
A Diagnostic Procedures	827
B Thyrotoxicosis	831
C Cretinism	836
D Goiter	838
E Thyroiditis	839
IV The Adrenal Gland	840
A Adrenocortical Hypofunction	840
B Addison's Disease	840
C Cushing's Syndrome	843
D The Adrenogenital Syndrome	848
V Ovary	852
A Genetic Abnormalities	853
B Infantile Ovaries with Eunuchoidism	854
C Pituitary Hypogonadism	855
D Nonneoplastic Cysts of the Ovary	855
E Ovarian Deficiencies of Psychogenic Origin	857
F Metabolic Disturbances Affecting Ovarian Function	859
G Effect of Adrenal Hyperfunction on the Ovaries	859
H Ovarian Hyperfunction	860
I Corpus Luteum Cysts	860
J Pseudocyesis	860
VI Placenta	861
A Pregnancy Complications	862
VII Testes	864
A Hypogonadism	865
B Testosterone as a Gonadotropin	867
C Testicular Tumors	867
VIII Puberty	868
References	870

I Introduction

The task of reviewing, within the framework of this book, recent progress in clinical endocrinology is not easy. In view of the extensive literature we had to impose upon ourselves certain limitations. It would hardly appear likely that the student or the practicing physician would turn for detailed practical guidance, to a book dealing quite predominantly with the chemistry and physiology of the hormones. Rather the biochemist and physiologist would seek here information on the clinical applications of the scientific achievements in his field. In fact, the line between physiological and clinical investigations is being reduced almost to the vanishing point. We are therefore presenting advances in principles and approaches rather than details of procedures or of therapeutic results.

II Anterior Pituitary Gland

A GENERAL REMARKS

In the not so distant past the anterior pituitary was supposed to secrete some twenty different hormones. For every and any effect of crude pituitary extracts a separate hormone entity was postulated and a name coined. Chemical isolation procedures have yielded six hormones and a better knowledge of metabolic interrelations has led to an understanding of multiple effects of individual single hormones.

The anterior pituitary then produces six hormones: growth hormone (also called somatotrophic hormone), thyrotrophic hormone (TSH), adrenocorticotrophic hormone (ACTH) and the three gonadotrophic hormones: follicle stimulating hormone (FSH), luteinizing or interstitial cell stimulating hormone (LH-ICSH) and luteotrophic hormone (LTH), identical with lactogenic hormone.

At present the trend appears to reverse itself. Observations have been recorded suggesting the existence of two adrenocorticotrophic hormones (one responsible for adrenocortical growth, another for secretory function) (8, 125) as well as of two thyrotrophic hormones (73). Regarding growth hormone, a separation of the diabetogenic factor from the protein anabolic growth hormone has been reported (169). These problems are under current investigation and cannot as yet be evaluated from the standpoint of the physiologist or clinician. Even if the fractionations and chemical separations mentioned above should be borne out by further work, the question will have to be answered whether the pituitary gland at any time secretes these factors separately or whether it secretes a complex hormone with different actions residing in different parts of the molecule, parts which could be separated *in vitro*.

Regarding ACTH, various preparations have been made and have been

come available for clinical use. The action of these preparations is qualitatively identical. A highly purified preparation designated ACTA (Corticotropin A) is much less subject to destruction than is ACTH (the crude preparation) when administered intramuscularly. Actide corticotropin (Corticotropin B) is a preparation obtained by peptic hydrolysis of ACTH. It evidently represents the active peptides of the larger ACTH molecule. It is readily inactivated or destroyed in the extravascular tissues and at present appears to offer little advantage for clinical use (239).

As far as the six hormone entities are concerned it appears that their secretion is to a large extent independent and that the gland does not function as a unit. There is no panhyperpituitarism, hyperfunction of the entire anterior pituitary, nor is there a concerted hypofunction, panhypopituitarism, except that due to destruction of the glandular tissue. As a matter of fact, interrelations of secretion of individual pituitary hormones appear to be the result of retrograde action upon the pituitary from secretions of the secondary glands. Examples of this type of interrelation may be the dependence of growth hormone secretion upon thyroid hormone secretion and the release of LH through changes in estrogen levels.

The independence of secretion of individual pituitary hormones has become more intelligible through the discovery of individual cell types responsible for elaboration of the individual hormones. Traditionally the hormone secreting cells of the anterior pituitary are classified on the basis of the staining properties of their granules and of the morphologic characteristics of the Golgi apparatus as acidophiles (eosinophiles) and basophiles. The elaboration of FSH, TSH, and ACTH is tentatively ascribed to the basophiles and that of the other three hormones to the eosinophiles. It has, however, become evident that basophilic staining is the common property of specialized cells, differentiated by histochemical reactions. A basophilic FSH producing cell is different from the basophilic TSH producing cell (78, 79, 165, 166). No similar recent differentiations have been reported within the eosinophilic cells, but an older observation of carminophilic cells in the cat pituitary, believed responsible for LH production, points in the same direction (60).

B. ACROMEGALY AND GIGANTISM

The clinical picture of gigantism and acromegaly was outlined in *The Hormones*, Volume II. More information has accumulated regarding the natural history of acromegaly. The excessive secretion of growth hormone by the eosinophilic cells occurs periodically. Periods of excessive secretion are followed by periods of relative quiescence. Such cycles may be repeated several times in the course of years. It is of more than academic interest to recognize the active phase of the disease earlier than in retro

spect by increased growth phenomena such as enlargement of feet or separation of teeth. Bio assay of growth hormone in the blood has been reported in one case (115) but assay methods which would lend them selves readily to clinical work have yet to be developed. Increased levels of serum inorganic phosphate have been found in the active phase of acromegaly, the levels return to normal under treatment with sex steroids as well as with radiation therapy (115).

Diabetes has long been known to be a frequent complication of acromegaly. In many respects it appears to be the human equivalent to the growth hormone-induced diabetes of dogs. Even the reversibility of early stages of growth hormone-induced diabetes in dogs has been paralleled in diabetes of acromegaly. In an acromegalic, observed and studied repeatedly for several years, a diabetic glucose tolerance made its appearance and reverted to normal following intensive roentgen irradiation of the pituitary (152). It has been pointed out that diabetes in acromegaly is more frequent in acromegalics in whose families non acromegalic members have diabetes. This would tentatively suggest that an inherited 'vulnerability' of the islets (doglike islets) is a prerequisite for the growth hormone to exert its diabetogenic effect. The bearer of 'nonvulnerable' islets would respond much like those species (*e.g.*, the rat) in whom diabetes cannot readily be induced by growth hormone.

It has long been known that goiter is a frequent occurrence in acromegalics. Also in many cases a high BMR is recorded (39). In view of the well known stimulation of the thyroid by anterior pituitary TSH the obvious assumption was that in this type of hyperpituitarism an increased secretion of TSH was associated with the increased secretion of growth hormone and that the former caused goiter and increased function of the thyroid. The situation is undoubtedly more complicated and the whole story has yet to be told. Cushing and his collaborators pointed out early that the histologic appearance of goiters of acromegalics differed markedly from that in thyrotoxicosis; also thyroidectomy caused regression of the elevated BMR and of other manifestations only in exceptional cases of acromegaly (39). More recently it has been found that the I 131 uptake of the thyroid gland of acromegalics was normal regardless as to whether there was a goiter or whether the BMR was elevated or normal (136). These observations have suggested, as did the older clinical observations of Cushing and associates that the high BMR of many acromegalics may be due to factors other than increased thyroid function (136). TSH levels in the serum of acromegalics have been found elevated regardless of the presence or absence of goiter and of the levels of the BMR (42). Neither the significance of this finding in the pathophysiology of acromegaly nor

the cause of excessive secretion of TSH (believed to be secreted by the basophilic anterior pituitary cells) is at present understood

Treatment In the opinion of most clinicians indication for surgical treatment is not the demonstrable evidence for presence of an adenoma but rather a rapidly progressive encroachment of the optic nerves. Other cases should be subjected to roentgen ray treatment, its effectiveness has been greatly improved with the use of higher doses of radiation than were previously employed (112). If there is no evidence of increased hormonal activity nor evidence of expansive growth of an adenoma treatment should be withheld. Treatment with androgens or with androgens and estrogens has been tried in an attempt to suppress growth hormone secretion through the suppressive action of the steroid hormones upon the anterior pituitary. With this form of therapy return of the elevated serum inorganic phosphorus levels to normal as well as improvement of the diminished carbohydrate tolerance has been observed but it is as yet too early to judge the practical usefulness of steroid hormone therapy in acromegaly (115).

C. HYPOPITUITARISM

Destruction of the anterior pituitary resulting from tumor (chromophobe adenoma, parasellar tumors *e.g.* craniopharyngioma) or post partum necrosis causes panhypopituitarism. Depending on the amount of pituitary tissue destroyed the severity of the disorder may vary from complete loss of pituitary function to milder forms. It is interesting that in the incomplete forms not all pituitary functions are diminished or abolished in the same proportions (151). It is our impression that the gonadotropin secretion is most sensitive and is usually lost first. The various combinations of functional deficiencies seen in tumor cases in which the nontumorous part of the gland is exposed to compression from the tumor cannot be explained readily by topographic arrangement of the cells responsible for the respective hormone secretion but perhaps rather by differences of vulnerability of these cells to the pressure exerted on all.

The essential findings are described in Volume II. It should be added that sodium loss evidently due to adrenocortical salt hormone deficiency is found in severe cases (151) this is of interest because it has been stated that in rodents salt hormone function of adrenal cortex is not governed by the anterior pituitary. The lack of response of the adrenal cortex to ACTH mentioned in Volume II pertains only to single intramuscular injections. With repeated injections and especially with intravenous drip infusion the atrophic adrenal can be stimulated and evidence of function be obtained (rise of 17 ketosteroid and 11 oxysteroid excretion eosinopenia). We have been able to stimulate the adrenal cortex 18 years after

post partum necrosis (152). The therapeutic regime as outlined in Volume II now includes cortisone and hydrocortisone.

1 *Single Hormone Hypopituitarism*

In contradistinction to panhypopituitarism due to destruction of the anterior pituitary the secretion of individual single hormones may be deficient, in the presence of normal secretion of the others.

Hypogonadotropism occurs both in males and females and causes clinical pictures indistinguishable from that due to primary gonadal failure. The distinction is made chiefly by assay of urinary levels of gonadotropin excretion which is high in primary gonadal failure and absent in hypogonadotropism. This will be further discussed in the chapters on Testes (VII) and Ovaries (V).

Hypothyrotropism is indistinguishable clinically from hypothyroidism due to primary thyroid failure. A distinction based on thyrotropin assay, analogous to the gonadotropin studies mentioned above, can be attempted only in laboratories equipped for TSH assays. In view of the difficulties of this assay procedure the two types of hypothyroidism are usually differentiated by the response to injections of TSH (102, 159). An increase of I 131 uptake, protein bound iodine (PBI), or BMR can be expected only if the thyroid gland is intact, and can therefore be accepted as evidence of hypothyrotropism.

There is at present no information as to whether these single hormone deficiencies of the anterior pituitary are the result of an anatomical derangement or whether the lack of secretion of the respective hormones is due only to a functional disturbance.

The growth hormone deficiency causing pituitary dwarfism of the Lorain-Levy type does not belong strictly in the category of single hormone deficiencies because in this disorder growth hormone deficiency is usually associated with gonadotropin deficiency and consequent hypogonadism. In exceptional cases however gonadal structure and function are normal (88, 180) and these cases legitimately can be considered under the heading of single hormone deficiencies. As is discussed below under the heading of pituitary dwarfism there is evidence for morphological changes in the eosinophilic cells of the anterior pituitary.

2 *Pituitary Dwarfism*

Dwarfism due to lack of growth hormone secretion is seen in children in whom the anterior pituitary is destroyed or encroached upon by tumors. In most instances craniopharyngiomas. Almost always the growth deficiency in these cases is associated with deficiencies of the tropic hormones, in other words the growth deficiency is part of panhypopituitarism. In

cases of pituitary dwarfism without clinically or radiologically demonstrable pituitary lesion (Lorain Levy type) the diagnosis is difficult. In most cases hypogonadism is present (but of course recognizable only after the age of puberty) but normal sexual development and even fertility have been recorded in exceptional cases (88-180). Delayed closure of the epiphyses appears to be typical. Increased insulin sensitivity might be expected in view of the anti-insulin action of growth hormone; however few observations are available and it would appear far from certain whether a normal insulin sensitivity would rule out growth hormone deficiency in the presence of normal adrenocortical function. The pituitary dwarfism of the Lorain Levy type may be compared with the dwarfism of the dwarf Silver mouse. In the latter dwarfism and hypogonadism occur as a recessive hereditary trait (188-189). The pituitary shows absence of eosinophilic cells. A marked diminution of the number of eosinophilic cells and degenerative changes in the cytoplasm have been reported in a case of pituitary dwarfism in a woman (88).

Whereas most growth hormone preparations have been found to be entirely ineffective in man inducing neither N retention nor growth in length (in spite of good potency in the rat (181)) positive results have been obtained recently by several investigators (114-184). Therapeutic growth hormone preparations are not yet commercially available.

II Thyroid Gland

A DIAGNOSTIC PROCEDURES

A considerable amount of work during the last several years has been concerned with diagnostic methods based on the role of the thyroid gland in iodine metabolism. The interest in these aspects has overshadowed that in the older procedures especially in the determination of the BMR. Conditions other than those of abnormal thyroid function are known to effect the oxygen consumption. In clinical diagnostic use one of the most common and most disturbing factors is that of emotional unrest causing high oxygen consumption probably by release of epinephrine. Techniques have been introduced for performing the test in barbiturate induced sleep or anesthesia under such conditions the high oxygen consumption resulting from nervous factors is reduced to normal values whereas the high BMR due to hyperthyroidism is not appreciably affected (13-172).

The use of $\text{I } 131$ tracer studies and PBI determinations has added materially to the knowledge of the pathophysiology of thyroid disease and these procedures have proved invaluable as diagnostic tools. Obviously a study of these parameters of thyroid function gives no information regarding the target response of the body tissues. This is highlighted by the unex-

plained syndrome of "hypometabolism" in which the basal oxygen consumption may be as low as in severe hypothyroidism, but in which the manufacture and release of thyroid hormone is entirely normal, as are all other body functions dependent on thyroid hormone.

1. Tracer Studies

The physiological basis for the diagnostic use of I 131 tracers has been discussed in Volume II. In many clinics the uptake is determined 24 hours after an oral tracer dose by measuring the radiation over the thyroid region. It has generally been confirmed that the uptake is high (more than 40%) in hyperthyroidism and low (less than 10%) in hypothyroidism. The tracer dose employed depends largely upon the sensitivity of the measuring instruments and of course on the particular parameters to be studied. With the use of scintillation counters the dose can be as low as 1 to 10 microcuries (3). Urinary excretion studies can supplement the measurements over the thyroid being merely related to the latter. Most of the iodine not taken up by the gland is rapidly excreted in the urine. It should be remembered that conditions other than thyrotoxicosis cause increased uptake of I 131: the high uptake of iodine deficiency goiter long postulated on the basis of animal experiments, has now been demonstrated in an endemic goiter population not exposed to iodine prophylaxis such as the use of iodized table salt (195). The high uptake in some goitrous cretins is discussed on page 837. Drug induced situations of increased iodine starvation also may cause a temporary rise of uptake considerably above normal. This is the case in individuals in whom the iodine trapping mechanism was blocked by thiocyanate with development of goiter. Following withdrawal of thiocyanate the uptake by the iodine starved goitrous gland is very high. A similar situation occurs after withdrawal of thiouracils in experimental animals but has not been definitely observed in man (41).

Conversely, low uptakes occur in conditions other than hypothyroidism. Antithyroid drugs (thiourea derivatives, imidazoles) as well as thiocyanate block the uptake, though in different ways. Thiocyanate truly blocks iodide uptake under ordinary conditions; only the presence of very large amounts of iodide would force the latter into the gland through the block. The antithyroid drugs do not materially interfere with the iodide concentrating mechanism but inhibit incorporation of molecular iodine into the amino acid. Since 24 hours after the tracer dose mainly the protein bound moiety remains, the result at this time interval is one of decreased uptake. Low uptake values are also obtained during and for several weeks after medication with thyroid hormone evidently through suppression of TSH secretion from the anterior pituitary. Administration of large amounts of iodine (either as iodide or as organically bound compound

(as used in roentgenographic studies of cholecystograms uterosalpingograms urograms bronchograms encephalograms) also causes low tracer uptakes. Two factors may be involved. The iodide may cause a real block perhaps by inactivating TSH. Furthermore if the iodide pool (circulating and extracellular) is considerably increased the tracer is diluted much more than in an average pool the uptake of the tracer may thereby be smaller owing to dilution. In renal disease the urinary excretion of iodide is diminished both because of impaired glomerular filtration and increased tubular reabsorption. Tracer studies based on urinary excretion of I 131 would give erroneous results. The increased I 127 pool also causes a greater dilution of the administered tracer and the accumulation of I 131 in the thyroid is delayed. However, values obtained by direct measurement over the thyroid at 26 hours are within normal range (157).

The physiological and pharmacological factors influencing I 131 uptake are of interest not only because of the light they throw on the pathophysiology of thyroid disease but also because they are pitfalls in diagnostic interpretation of the results of such studies. Another interesting finding is the almost complete obliteration of I 131 uptake in acute thyroiditis without other evidence of hypothyroidism.

Several procedures, using I 131 tracers have been introduced which add materially to the information obtained by determining the uptake after 24 hours. By repeated measuring of activity over the neck following a tracer dose the concentration of I 131 in the thyroid is found to be a linear function of the square root of time. The slope of this curve is called the accumulation gradient (9-74).

The mathematical relationship is simple but rather obscure in terms of biological significance. The linear relationship just mentioned is found in euthyroid and hypothyroid individuals but not in hyperthyroidism. The method is valuable inasmuch as it establishes the accumulation gradient within 1 to 2 hours after administration of the tracer and permits study of factors influencing the iodine accumulation of the thyroid (168). For example the activity of antithyroid drugs has been screened by this method (200).

Determination of thyroidal and renal plasma I 131 clearance rates has been recommended as a routine diagnostic test. Whereas determinations of the 24 hour uptake fail to correlate with the clinical condition in 7% to 10% of cases the claim is made that the clearance rates show no overlapping of values in normal and hyperthyroid individuals (17). It will be interesting to see whether such a sharp dividing line between euthyroid and hyperthyroid individuals will be confirmed on a larger material which will include mild cases of hyperthyroidism. It may appear questionable to some whether there is biologically a sharp line of demarcation between a thyroid

gland with 'top normal' function and that functioning slightly in excess of normal

Another parameter of thyroid function which can be studied by the use of I 131 tracers is that of release of thyroid hormone from the gland. By direct determination of the radiation over the thyroid for several days a measure for the secretion of thyroid hormone can be obtained (9).

This is also referred to as 'physiological decay,' and the time during which the gland loses 50% of its activity (corrected for physical decay) is called the biological half life. For more precise determinations it is necessary to administer antithyroid drugs (thiouroid derivatives or imidazole derivatives) after the tracer dose in order to block uptake of that iodide moiety originating from the secreted thyroid hormone (9). Because of the necessary length of observation period this method is not generally used in clinical practice. Measurements of protein bound iodine I 131 following a tracer dose reflect the secretory activity of the thyroid (186). Under certain circumstances the values of PBI₁₃₁ do not correlate with the clinical physiological status of the individual. In some cases of postoperative myxedema the PBI₁₃₁ values have been found high in the presence not only of clinical manifestations of hypothyroidism but also of low PBI₁₂₇ values. This has been thought to indicate a very small but very active thyroid remnant with rapid turnover rate but small total output (19). By determining total plasma I 131 and protein bound plasma I 131 a conversion ratio has been calculated by the formula conversion ratio =

$$\frac{\text{PBI}_{131}}{\text{total plasma I 131}} \times 100$$

This ratio is believed to be related to the rate at which the gland utilizes iodine for hormone synthesis and release. There is some doubt as to the validity of such a simple formula on physiological grounds; correlation with the clinical condition has been found only in hyperthyroidism whereas the conversion ratio of hypothyroid and euthyroid individuals did not differ materially (183).

A further application of the tracer technique made possible through the introduction of the highly sensitive scintillation counter is the scanning procedure (68, 69). One or several scintillation counters are moved by a motor driven device across the thyroid region; the impulses are transmitted to a stylus which records them on graph paper. Such 'scintigrams' record the outlines of the thyroid gland and have been used for determining thyroid weight as a basis for calculating therapeutic doses (69). Inactive and hyperactive nodules can be demonstrated inasmuch as the scintigram will show regions of diminished or absent density or regions of increased density (68). The procedure has furthermore been used as an aid in differential diagnosis of masses of the neck and by scanning body regions other than the neck for detection of thyroid cancer metastases (10, 195).

2 Protein Bound Iodine (PBI) (207)

Methods for determination of PBI have been improved a discussion of these methods is beyond the scope of this review. In interpreting PBI values as indicative of abnormal thyroid function, sources of error other than those of a technical nature must be kept in mind. The most important error results from the presence in the body fluids of large amounts of nonhormonal iodine. This may be the result of ingestion of large amounts of iodide organic iodine compounds used as contrast media in various diagnostic procedures (cholecystography, pyelography, bronchography, encephalomyelography, uterine alpingography) cause even greater and more prolonged spuriously high PBI values. It appears that in the presence of large amounts of iodide in the serum some iodine will come down with the protein precipitate causing high yields in this fraction. High PBI levels are regularly found in pregnancy (129). This finding poses an intriguing question. Evidently the normal pregnant woman exhibits no signs of hyperthyroidism. Is this because the body needs more thyroid hormone in pregnancy? Spuriously low PBI levels have been obtained during the use of mercurial diuretics, during the technical procedures of digestion insoluble mercuric iodates or iodides are formed (139).

B THYROTOXICOSIS

No progress has been made regarding the etiology of thyrotoxicosis.

Advances in diagnostic methods are discussed on page 827. Diagnosis is not difficult in fairly typical cases certain types of patients pose a more difficult diagnostic problem in which the use of the newer procedures outlined above are of particular value. One is the thyrocardiac individual presenting cardiac failure frequently without a history typical of thyrotoxicosis nor are the clinical manifestations unequivocal. Frequently a failure to respond to digitalis first arouses suspicion. Signs and symptoms such as auricular fibrillation, tachycardia and nervousness can be the result of the cardiac condition and therefore help little in establishing the diagnosis of thyrotoxicosis. In the presence of cardiac failure and passive congestion a true basal metabolic rate cannot be determined. PBI and tracer studies have been of great diagnostic value in these cases. Control of the thyrotoxicosis will improve the cardiac condition significantly. Whether thyrotoxicosis can cause failure and consequent passive congestion of an otherwise entirely normal heart is debatable. The fact that the majority of thyrocardiac patients are in the older age group would lend support to the assumption that thyrotoxicosis causes failure only of a damaged heart. The argument loses practical significance if one keeps in mind that the pre-existing cardiac damage (most frequently arteriosclerotic in origin) may be subclinical and may revert to a symptomless stage after the thyrotoxic component is controlled.

Another group of thyrotoxic patients in whom the diagnosis can be established only with the aid of the tests described is that of certain severely agitated psychoneurotics. It is generally believed that psychic factors can precipitate or aggravate thyrotoxicosis, equally thyrotoxicosis is apt to aggravate or bring into the open psychoneurotic tendencies. Regardless of what is cause or effect in an individual case control of thyrotoxicosis if present is of greatest importance. Sleeping BMR, PBI, and tracer studies will help to establish the presence or absence of thyrotoxicosis.

Therapy. No basically new approach in the treatment of thyrotoxicosis has evolved. However a great amount of experience has accumulated with the use of the methods outlined in Volume II.

Numerous antithyroid drugs have been tested. Those which have proved most useful and are generally used, are two thouracil derivatives propylthiouracil and methylthiouracil and an imidazole derivative 1-methyl-2-mercaptoimidazole (trade name Tapazole). They can be used in preparation for surgery or for definitive medical treatment. If given preoperatively the increased vascularity and friability of the hyperplastic gland may cause technical difficulties at operation. Iodide (e.g. in the form of liquor iodi comp. Lugol's solution) causes involution of the hyperplastic gland with decreased vascularity and friability iodide is therefore administered together with the antithyroid drug for a few days prior to operation. The mechanism of action of iodide under the conditions is but poorly understood. It may enter the gland but is not oxidized and rapidly leaves the gland again. It may possibly inactivate endogenous TSH. If the antithyroid drugs are used for definitive medical treatment they should be administered for long periods of time. It is advisable to continue maintenance doses for 8 to 12 months after a euthyroid condition has been attained. Then treatment is discontinued. Fifty to sixty per cent of the patients remain in remission the others relapse. It has been possible in some instances to keep up maintenance treatment for several years if for any reason the other forms of therapy surgery or radiation were inadvisable.

Toxic reactions are observed in a small number of cases both with propyl and methyl thiouracil and with Tapazole. The total incidence is less than 2% in a few instances agranulocytosis has occurred.

Experience with radioactive iodine has also been greatly extended. There still is considerable difficulty in determining the dose which will reduce thyroid activity to normal but not cause hypothyroidism (see also p. 830). In many cases there occurs an exacerbation of the thyrotoxicosis during the first two to three weeks following the administration of I 131 owing to the liberation of stored hormone incident to the destruction of thyroid tissue. This exacerbation is rarely serious but thyroid crisis has

been observed. We have seen the most gratifying results of I 131 therapy in the thyrocardiac patient but the temporary exacerbation of the thyrotoxicosis may cause the patient to go through a period of more severe cardiac failure.

The possibility, that the radiation from therapeutic doses of I 131 may be carcinogenic is considered remote by most investigators. However thyroid cancer has been observed in rats treated with I 131 (6). Furthermore the possibility of late development of radiation induced fibrosis has to be considered. Many clinics therefore prefer not to treat thyrotoxicosis with I 131 in individuals younger than 35 to 40 years.

1 *Ophthalmopathy*

Interest in the ophthalmopathic facet of Graves disease continues. There appears to be no question that the severe ophthalmopathy is caused by a pituitary factor as outlined in Volume II. In the older experiments in which exophthalmos had been induced in experimental animals with pituitary extracts this action had been ascribed to TSH or to a pituitary factor closely allied to TSH. Dobyns and Steelman have succeeded in separating the exophthalmos producing principle from the thyroid stimulating hormone (49). This still leaves open the question whether the pituitary actually secretes the two factors separately, or whether the separation is a chemical one only. The most severe forms of ophthalmopathy are usually observed following thyroidectomy for thyrotoxicosis; occasionally they occur during treatment with antithyroid drugs. As long as the thyrotoxicosis is uncontrolled the ophthalmic picture is usually less severe; in exceptional cases the dissociation of thyrotoxicosis and ophthalmopathy is complete, the latter occurring in euthyroid individuals or in spontaneous myxedema, in either case without any indication of a preceding hyperthyroid episode.

The early changes in the retrobulbar tissue both in exophthalmos produced experimentally by injections of TSH and in human cases are those of edema; the edema fluid is rich in mucopolysaccharides (127). The muscle changes are swelling, edema and lymphocytic infiltration (pseudo hypertrophy). Whereas these muscular changes occur quite predominantly in the extrinsic muscles of the eye they are demonstrable to a minor extent also in skeletal muscles.

Occasionally severe ophthalmopathy occurs simultaneously with pretibial localized myxedema (38, 224). The cutaneous localized infiltration shows the same chemical characteristics as the retrobulbar tissue (227). Injection of hyaluronidase into such lesions has been reported to cause resolution, evidently owing to the specific depolymerizing action of this enzyme on the mucopolysaccharides (77, 142) in one case of our own

observation we were not able to show any changes following local hyaluronidase injection (132). Similarly, hyaluronidase has been injected into the retrobulbar tissue in cases of severe ophthalmopathy, but results have been unsatisfactory (77-107). Because TSH was believed to be the active factor from the anterior pituitary, several investigators have studied the TSH levels in blood or the urinary excretion of TSH in ophthalmopathic cases (5-42, 161). In most instances, high blood levels or high urinary excretion was found (1), but this appears not to be invariably the case (42, 161). If the *in vitro* separation of an exophthalmos inducing factor from the thyroid stimulating moiety is confirmed, the TSH studies just described would suggest that TSH secretion usually, but not invariably, parallels that of exophthalmos inducing factor.

The question whether the ophthalmopathy associated with Graves' disease is always and in all cases due to a pituitary factor (so called thyrotropic ophthalmopathy) can as yet not be definitely answered. Most investigators favor a unitary etiology for both the severe ophthalmopathy and the ocular manifestations of classical Graves' disease. However, the occurrence of two etiologically different types of ophthalmopathy has been suggested, thyrotropic and thyrotoxic in nature (143). In the latter the ocular changes are supposed to be the direct effect of excessive thyroid hormone excretion. If this type actually exists, the eye manifestation would be expected to subside after the thyrotoxicosis is brought under control. In many instances, improvement of the eyes following control of the thyrotoxicosis is more apparent than real; the lid retraction (a definitely thyrotoxic manifestation) subsides and with weight gain the cheeks fill out, giving the eyes a more normal appearance, whereas the exophthalmos may not recede or actually increase even if slightly. However, occasional observations do suggest a real regression of exophthalmos and of paresis of extrinsic ocular muscles.

It is most important to prevent the development of the most severe or malignant forms. Since they occur mostly postoperatively, thyroidectomy should be advised against in thyrotoxic patients in whom severe ophthalmopathy, including swelling of the lids, pain, extensive muscle paresis and edema of the conjunctiva is present. In rare instances, severe deterioration of the ophthalmopathy has been observed during thiouracil therapy. In such cases the thyrotoxicosis should be brought under control very gradually by relatively small doses of antithyroid drugs (propyl thiouracil, Tapazole) or by small repeated doses of I 131 or by external roentgen radiation.

The treatment of the ophthalmopathy is directed to the anterior pituitary. If at the time of treatment the patient is euthyroid or hypothyroid (regardless of whether this was preceded by thyrotoxicosis or not), desic-

cated thyroid given to tolerance is frequently beneficial. Results appear better, the earlier treatment is started. Thyroid hormone probably acts by suppressing secretion of TSH (or an exophthalmos factor) from the anterior pituitary. Estrogens in large doses have been used successfully in severe cases; they also probably act by suppressing the anterior pituitary. Roentgen ray irradiation to the pituitary was up to recently, used mostly as an adjunct treatment of questionable value combined with either thyroid or estrogen medication; recently, however, cases have been reported in which roentgen radiation to the pituitary alone yielded good results (15).

2 *Thyrotoxic Myopathy* (141, 167-240)

It has long been known that some muscle wasting is a common occurrence in thyrotoxicosis. The increased (endogenous) creatinuria as well as the increased creatinuria following an exogenous creatine load (decreased creatine tolerance) is probably the result of the muscle dysfunction induced by the excess circulating thyroid hormone. In rare instances of thyrotoxicosis muscle wasting is very much more pronounced and may be the presenting feature of the disease; these cases are referred to as thyrotoxic myopathy. Doubt has been voiced as to whether this syndrome represents only a quantitative exaggeration of the more moderate muscle involvement commonly seen in many severe thyrotoxic patients; it has been suggested that a qualitatively different but as yet unknown process is involved. Good recovery is observed when the thyrotoxicosis is brought under control. Diagnosis may be difficult inasmuch as the myopathy can dominate the picture to the point of masking the thyrotoxicosis; careful history and examination and appropriate tests will aid in establishing the diagnosis. When the presence of thyrotoxicosis and of myopathy has been established, the cases have to be differentiated from myasthenia gravis. The latter disease is occasionally associated with thyrotoxicosis. Whether there is any causal relationship is as yet unknown. Commonly the response to prostigmine is used to differentiate between myasthenia gravis as associated with thyrotoxicosis and thyrotoxic myopathy. The former does and the latter does not respond to the drug. However the specificity of the response to prostigmine has been questioned and cases responding to a test dose of prostigmine have been reported as thyrotoxic myopathies. This must remain a debate in terminology until other tests characteristic for myasthenia gravis are also employed in the study of these cases (response to curare, electromyogram, etc.). The review of clinical problems in neuromuscular physiology by Denny Brown is helpful in understanding these complex problems (46). In patients simultaneously afflicted with myasthenia gravis and with thyrotoxicosis the myopathy has been reported to show improvement, deterioration, or no change after

control of the thyrotoxicosis. It is trite to state that more studies of this problem are needed.

Cases are on record of simultaneous occurrence of thyrotoxicosis and periodic familial paralysis (141, 228). As far as can be judged from the reported cases the periodic paralysis appears to be improved but not abolished following control of the thyrotoxicosis.

3. *Thyroid Crisis*

The exact nature of thyroid crisis or thyroid storm has yet to be completely unraveled. The factors most commonly precipitating crisis are (1) any operations performed upon untreated or incompletely controlled thyrotoxicosis cases (surgical storm) and (2) severe infections as well as major upsets (medical storm). Whether these factors act as 'stressors' through the adrenal cortex (see below) is at present a matter for speculation. There seems, however, little doubt that thyroid crisis represents thyrotoxicosis at peak severity. Why, in this condition, the heat regulating processes of the body break down, with the skin becoming hot and dry rather than dripping with perspiration, is not understood. It has long been suggested that an acute adrenocortical insufficiency occurs during thyroid storm and may in part be responsible for the fatal outcome. Treatment with cortisone has been successful and in several cases the development of post-operative 'surgical' storm was prevented by early treatment with ACTH or with cortisone, when an emergency operation became necessary in uncontrolled thyrotoxicosis (209).

C. CRETINISM

Cretinism is clinically subdivided into endemic and sporadic cretinism. *Prima facie* this is a geographical classification of cretinism occurring in goiter areas (iodine deficiency areas) or in nongoitrous regions. The endemic cretin is identified with the bearer of a degenerated thyroid, usually but not always goitrous, incapable of manufacturing thyroid hormone. The sporadic cretin on the other hand is identified with a congenital defect—a thyroid aplasia.

Tracer studies with I 131 on cretins believed to have congenital thyroid aplasia have in some instances yielded some small uptake in the region of the thyroid indicating the presence of some thyroid tissue. This confirms old histological reports in such cases in which small amounts of thyroid tissue rather than complete absence was reported. It will be interesting to see whether studies including the response to ISH of a larger material will detect early cases of TSH hypopituitarism in sporadic cretins.

The occurrence of goitrous cretins outside areas of goiter endemics has been described. Studies reported in some of these cases raise intriguing

problems regarding the pathophysiology. The uptake of I 131 is rapid and of a high order (81 122 137 199) such as is usually seen in the thyrotoxic gland and has been found in endemic iodine deficiency goiter (198). The utilization of the iodine by the thyroid gland appears to differ in different cases. In Stanbury's cases the rapid purging of the gland by thiocyanate indicates that the iodide was not oxidized, and therefore not incorporated into organic compounds (199). The deficiency in these cases was similar to that induced by thiourea and imidazole compounds. In contradistinction in McGirr's cases thiocyanate failed to purge the gland suggesting oxidation of iodide with subsequent organic binding. This was further supported by the presence of protein precipitable I 131 in the blood. It was assumed that this was an abnormal compound which was not identified (137). In Hamilton's cases analysis of thyroid tissue following administration of I 131 yielded radioactive fractions corresponding to thyroxine and diiodotyrosine. These studies were done before chromatographic analysis of these compounds had been introduced. In these cases it was assumed that the thyroid gland was capable of synthesizing thyroid hormone but not of releasing it (81).

Hurvital has claimed that some goitrous cretins subsequently and without treatment become euthyroid or even hyperthyroid (94 96). These cases have not been reported in great detail and from the data recorded the diagnosis of cretinism may be questioned.

It is now generally recognized that treatment of cretins even when started early and continued uninterruptedly frequently fails to bring about normal mental development. Mental development improves during therapy, but in many instances does not reach normality. The nature of the changes underlying the lack of mental development is not known. Recent observations indicate the presence of electroencephalographic abnormalities in untreated cretins (43 217). It has been suggested that the EEG pattern becomes normal under thyroid medication only in those cases in whom mental development is fully restored and not in those who remain mentally deficient in spite of treatment (217). This has not been borne out regularly in our as yet limited experience (471).

On the basis of the concept of a reciprocal relationship of the pituitary tropic hormones with those of the secondary glands one would expect a high TSH secretion from the anterior pituitary in severe hypothyroidism including congenital thyroaplasia (an analogy could be drawn between the latter and ovarian agenesis associated with high FSH excretion). However this is not always the case. In a study of cretins Paschalis and D'Angelo found absence of TSH in the blood of an untreated cretin 11 years of age. TSH becoming detectable only after thyroid medication. This suggests that the anterior pituitary which has not been exposed to

thyroid hormone may be incapable of manufacturing and/or secreting TSH. In untreated cretinous babies the expected high TSH levels have been found. Studies regarding secretion of hormones other than TSH by such pituitaries will be of interest.

Familial incidence of sporadic cretinism has been reported (22) (familial incidence is of course common in endemic cretinism). On the other hand, the occurrence of cretinism in one of two identical twins has been reported unfortunately very briefly and without detailed data (175).

D. COITER

Iodine deficiency is firmly established as the most common cause of goiter (see Chapter V by Salter, in Volume II of *The Hormones*). Interesting historical aspects are touched upon by Salter (177). Certain aspects of the pathophysiology of human iodine-deficiency goiter had not been studied with modern methods because iodine prophylaxis in the form of iodized table salt has been in general use for several decades in the goiter regions and goiter belts of Europe and North America. Recently such studies were carried out in an endemic goiter area in the Argentine Andes where the natural conditions of the iodine-deficiency goiters had heretofore never been interfered with by administration of extra iodide. The great avidity of these goiters for iodine was demonstrated by the very high uptake of tracer doses of I 131 (198). Further results of this study will be awaited with interest.

Colloid goiters clinically and pathologically indistinguishable from those occurring in iodine deficiency regions are observed sporadically in regions such as the Atlantic seaboard where iodine supply in water and food is adequate. Endemic occurrence of goiter in locations not deficient in iodine has long been recorded (*e.g.* Derbyshire goiter in England) (138). Following the observations of goitrogenic action of cabbage brussels seeds and other vegetable matter in rabbits and rat, Astwood isolated an active goitrogen 1:5:2 vinyl thiocyanazolidone from rutabaga and studied the goitrogenic action of a great number of vegetables (6, 7). The chemical composition as well as the goitrogenic action is similar to that of thiourea and its derivatives. The subject has been reviewed by Greer (75) and by Fertman *et al.* (54). Only very few observations in human beings are recorded in whom the presence of a goiter could with great probability be traced to the ingestion of large amounts of goitrogenic vegetables (55, 138). Means has coined the term *struma cibaria* for this type of goiter (138). Careful nutritional histories should be obtained in all goiter cases especially outside iodine deficiency areas. However it appears very doubtful to us that the majority of colloid goiters occurring in regions with adequate iodine supply can be explained on this basis.

The etiology of most cases is still obscure; an endogenous deficiency in iodine utilization may be thought of hypothetically (76).

In the medical treatment of colloid goiter medication with desiccated thyroid has been revived (76). Thyroid medication evidently acts by suppressing endogenous TSH secretion. The latter was previously increased causing enlargement of the thyroid because the normal size gland has for some reason been unable to produce adequate amounts of thyroid hormone.

F. THYROIDITIS

Whereas the etiology of acute nonsuppurative thyroiditis remains obscure in most instances, an interesting physiopathologic aspect has been detected. The uptake of tracer doses of I 131 is greatly diminished or absent. No other manifestations of hypothyroidism are present. PBI values are normal or even slightly elevated. The reason for the depressed I 131 uptake is not evident; administration of TSH will increase the uptake though to a lesser degree than in euthyroid individuals in thyrotoxicosis factitia or in hypopituitarism. Curiously, administration of TSH brings about an improvement, at least a temporary one, of the condition. Various hypotheses have been offered in explanation of the low I 131 uptake and the response to TSH, but none is entirely satisfactory (176).

Milder cases last only a few days, but in more severe cases fever, pain, and swelling may continue for several weeks. In the absence of any knowledge regarding the etiology (bacteriological cultures have been almost uniformly sterile), treatment is symptomatic. Antibiotics have been without effect in almost all instances. There have been reports of beneficial action of external radiation and of thiouracil therapy; more recently, cortisone or ACTH has been used successfully; their beneficial action is probably due to the anti-inflammatory action of the adrenal steroids (230).

1. Subacute Thyroiditis (36a)

Attention has been drawn to this form of thyroiditis, which is not infrequently confused with Riedel's struma. A history of an acute phase is frequently obtained, but sometimes the onset is insidious and fever and severe general constitutional manifestations are absent. The gland is enlarged and very hard; it may or may not be tender. Even in the absence of fever, the sedimentation rate is elevated, and the I 131 uptake quite as low as in acute thyroiditis. Extensive fibrosis of the surrounding structures characteristic of Riedel's struma is absent. The histological picture (surgical or needle biopsy) is characterized by pseudotubercles and

giant cells. In contradistinction to Riedel's struma, subacute thyroiditis is self limited and responds rather well to roentgen ray therapy, operations are unnecessary. The etiology is as obscure as that of other forms of thyroiditis.

IV The Adrenal Gland

A ADRENOCORTICAL HYPOFUNCTION

Chronic hypoadrenocorticalism in human beings is due to (1) Destruction of the adrenal cortex by chronic disease (Addison's disease), (2) adrenalectomy performed for hypertension or advanced cancer of prostate or breast, (3) hypopituitarism resulting in lack of stimulation of the adrenal cortex by ACTH, (4) salt losing adrenogenital syndrome' (see p 831). Acute adrenocortical insufficiency probably plays a role in Waterhouse-Friderichsen syndrome (acute fulminant bacteremia, mostly meningococemia, with extensive hemorrhage into the adrenal cortices), and perhaps as a phenomenon of adrenocortical exhaustion in severe infection without hemorrhage into the adrenals.

B ADDISON'S DISEASE

'Cortical atrophy' so called and tuberculosis are the causes of Addison's disease in the vast majority of cases. The term atrophy is a misnomer unfortunately ingrained in the clinical literature. The anatomical changes in the adrenal cortex in these cases are the result of cytotoxic damage, necrosis and fibrosis. In a small number of cases histoplasmosis involving the adrenal cortex causes Addison's disease, cases of amyloidosis (84-225), leucemic infiltration and extensive tumor metastases are rare occurrences (29). It is believed that severe clinical manifestations occur only after 90% of the adrenal cortex is destroyed.

1 Diagnostic Procedures

The absence of response to ACTH has been widely employed, with ACTH given either in a single intramuscular injection or by intravenous drip infusion. The end points studied are the decrease in circulating eosinophils and the rise in urinary 17 ketosteroids and 11 oysteroids after the more intensive stimulation by continuous infusion. The eosinophil response to epinephrine however cannot be used as a test for adrenal responsiveness since it is very doubtful whether in man epinephrine in doses employed actually fires the pituitary-adrenal system (213). The most significant tests are those based on the disturbance of electrolyte metabolism. The response to NaCl deprivation combined with potassium administration (Cutler-Power-Wilder) still appears to be the most reliable and simple procedure if it is carried out with due safeguards for the possi-

ble precipitation of a crisis (40). The deranged Na and K metabolism can be demonstrated by testing the composition of sweat (35-126) or saliva (57-226). These studies have yielded interesting results regarding hormonal control of electrolyte balance but they would at present appear less suitable than urine studies for diagnostic purposes. Tests designed to demonstrate the abnormal electrolyte metabolism are the most significant not only because most of the clinical manifestations of Addison's disease are due to the abnormal electrolyte metabolism but also because only the "salt hormone deficiency" is a constant finding in all cases of Addison's disease. Disturbances of carbohydrate metabolism are present in the majority of cases in varying severity. Severe hypoglycemia after prolonged fast and severe hypoglycemia following the hyperglycemia induced by intravenous glucose administration are indicative of the sugar hormone deficiency. In many instances this deficiency is not severe enough to be demonstrable by these procedures. Increased sensitivity to insulin (hypoglycemia irresponsiveness) is present in the majority of cases. Because most Addison patients lapse into hypoglycemic coma when tested with intravenous administration of insulin by the standard procedures modifications of the test have been devised. Thorn suggests giving insulin in increasing doses with the meals (214). Engel has devised an insulin glucose test in which the glucose administration prevents severe hypoglycemic manifestations (52). Cases of dissociated Addison's disease are of considerable interest in the presence of the classical picture with pigmentation, hypotension, weakness and demonstrable defect in Na and K metabolism the response to insulin is normal and low normal urinary excretion of glucocorticoids is present (152). Similarly the very low or absent 17 ketosteroid excretion is not always found in some instances values in the lower range of normal excretion have been reported (131). These dissociated cases cannot be adequately interpreted at this time. Even if in man the different hormones are secreted by different zones of the cortex as has been suggested in the rat and mouse it is difficult to visualize a zonal destruction in the cortex. No autopsy reports have come to our attention which would throw light on this problem if indeed the pathologist will hold the answer. Also follow up studies of such patients will be desirable in order to see whether eventually the dissociated disturbance will in later stages of the disease embrace all hormones resulting in what may be called panhypocorticalism.

2 Addison's Disease and Pregnancy

Before adrenocortical hormones became available for treatment of Addison's disease the occurrence of pregnancy in Addison patients was extremely rare and almost invariably fatal (25). With present therapy

several women have been carried through pregnancy successfully and have given birth to viable children. The most critical periods are the first trimester, during which profuse vomiting may precipitate crisis and the delivery with the incident stress and blood loss. Adrenalectomized pregnant rats have been known not only to tolerate pregnancy but to be maintained easier and better. This was ascribed to the deoxycorticosterone like action of progesterone. However in human beings there is evidence that the placenta produces corticoids (and probably ACTH). The *c* compounds are not produced in amounts sufficient for complete maintenance but may contribute to make maintenance after the third month easier (2, 99, 150).

3. Treatment of Addison's Disease

The use of deoxycorticosterone acetate and salt was discussed in Volume II of *The Hormones*. The compound is now available in 'buccal tablets' resorption is through the mucosa of the oral cavity into the major circulation. Dosage is of necessity less accurate than with administration by injection or pellet implantation; the degree of accuracy or inaccuracy depends entirely upon the intelligent and meticulous co-operation of the patient. Recently deoxycorticosterone trimethylacetate was introduced. This is a 'long acting' preparation permitting of maintenance by one intramuscular injection about every four weeks (215). The required dose of deoxycorticosterone and of salt must be determined empirically for every case; overdosage must be avoided because it causes hypertension, edema, and cardiac failure. Standardization is carried out with the use of body weight, blood pressure, hematocrit and serum electrolyte levels as criteria. One of the most valuable and sensitive effects is that on heart size as determined by roentgenograms. Cortisone is used as an adjunct to the treatment with deoxycorticosterone. Even when treatment with the latter succeeds in normalizing the patient by all objective criteria additional therapy with small doses of cortisone (12.5 to 20 mg. per day) frequently adds to the sense of well being. Whether this is due to the action on carbohydrate metabolism or rather to the cerebral euphoria, the action of cortisone is not immediately apparent. Perhaps the fact may be significant in this respect that the abnormal electroencephalographic pattern of Addison's disease is corrected by cortisone therapy but not by the use of deoxycorticosterone. Perhaps the most important place of cortisone is in prevention of crisis. Patients can be instructed to take extra cortisone if they have a sore throat or any other febrile episode even before they report to their physician. Corticosterone (Compound B of Kendall) has been investigated as a therapeutic agent in Addison's disease but this steroid has not been available for large scale clinical testing (34).

The fact that glycyrrhizic acid, the active ingredient of Liquorice is capable of correcting the electrolyte disturbance of Addison's disease is at this time at least less of practical than of theoretical interest (24-71)

Addisonian crisis is treated by intravenous infusion of glucose saline plasma antibiotics and aqueous adrenocortical extract and by intramuscular injections of cortisone and deoxycorticosterone acetate

C. CUSHING'S SYNDROME

1 Pathogenesis

Cushing's syndrome has been recognized as a form of hyperadrenocorticism as was described in Volume II page 692. The adrenals are bilaterally diffusely hyperplastic in some cases whereas in others an adrenocortical cancer is present. In a small number of cases no morphological changes are found in the adrenals if no ovarian tumor is present an excessive function of adrenal of normal size and structure may be assumed. This however is an assumption since it is conceivable that normal amounts of hormone might be secreted by the glands and the picture of hyperfunction be simulated by abnormal metabolism of the hormone. The tumors of the ovary mentioned above are probably adrenal rest tumors such tumors are rare and most of those reported have caused adrenogenital syndrome rather than Cushing's the occurrence of the latter however has been established.

Cases of hyperplasia may be the link between the newer concept of Cushing's syndrome as hyperadrenocorticism and Cushing's original thesis that these cases were due to basophilic hyperpituitarism. Whether or not a basophilic adenoma is present excessive secretion of ACTH by the pituitary basophils would explain hyperplasia and hyperfunction of the adrenal cortex. Older reports of increased ACTH levels in the blood of such patients are inconclusive because of the inadequacy of the methods employed. With the use of Sayer's method increased levels of ACTH have now been demonstrated in some instances (23-70) but not in others (208a, 209a). Correlation with anatomical findings in adrenals and pituitary will be desirable. If the findings of increased ACTH levels in the blood of Cushing cases with adrenocortical hyperplasia are confirmed in a larger number of cases a further problem arises. We have reason to assume that hypersecretion of cortical hormones will suppress ACTH secretion from the anterior pituitary making endogenously caused adrenocortical hypersecretion self limited. This would of course not apply to adrenocortical carcinomas which may be autonomous and continue to function at high levels in spite of the retrograde suppression of ACTH secretion the contralateral adrenal in such cases is however atrophic because of this mechanism. In cases of hyperplasia the question therefore

arises why the excessive ACTH should persist, and not be suppressed by the liberated adrenal steroids. There is at present no answer, one could suggest the hypothesis that the control mechanism in such cases is 'set' at a higher level, either in the pituitary or in the hypothalamus.

The latter has been implicated in the pathogenesis of Cushing's syndrome (63), and whereas the evidence is far from conclusive the experimental data on hypothalamic control of ACTH secretion (26, 56, 93, 134) should stimulate further study of clinical material.

In a discussion of the pathogenesis mention should be made of the Cushing like pictures which have been observed (1) after therapeutic administration of large doses of cortisone (197) and (2) in certain forms of hepatic cirrhosis (21). From a purely therapeutic angle, the occurrence of Cushing like features in patients treated with cortisone is an undesirable 'side effect'. However these cases are of considerable interest inasmuch as they would seem at first sight to nullify the concepts of the pathophysiology of Cushing's syndrome as first proposed by Albright (1) and Kepler (110). According to this concept Cushing's syndrome would represent 'panhypercorticalism' some of the manifestations would be the result of excessive secretion of deoxycorticosterone like salt hormone (hypertension cardiac enlargement, edema) with others to be ascribed to the excess sugar hormones' (diabetes obesity deossification purple striae) and yet others (acne hirsutism, plethora) to increased androgen secretion. After all these manifestations were induced by excess cortisone administration it appeared doubtful to some investigators whether the concept of Cushing's syndrome as 'multihormone excess or panhypercorticalism' could be upheld. In view of the fact that "single hormone" hypersecretion as well as hyposecretion appears to occur and in view of the fact that the adrenal gland can perform remarkable interchanges in the steroid structure the induction of Cushing's syndrome by cortisone administration does not appear conclusively to invalidate the previous concepts.

The observation of Cushing like pictures in certain forms of hepatic cirrhosis is of interest. This has been explained by assuming that the diseased liver is deficient in inactivating or destroying endogenous cortical hormones. The urinary excretion of formaldehydogenic steroids was found increased in these cases—a finding compatible with the explanation (21). Hepatic inactivation of cortisone has been demonstrated *in vitro* and *in vivo*.

To return to the pituitary basophils the only constant finding in all cases of Cushing's syndrome has been that of hyalinization of many basophils these changes are usually referred to as 'Crooke's changes'. The physiological significance of this finding was not apparent the hyaliniza-

tion was interpreted as possibly indicative of excessive ACTH secretion, or as the late Dr Kepler suggested as an effect of the excessive adrenocortical secretion on the pituitary (109). The latter explanation may now be accepted in view of the fact that identical changes have been found in the pituitaries of subjects treated with ACTH or with cortisone for various conditions (66, 121).

2 *Hormone Secretion*

Following the development of methods for determining adrenocortical 11-17 hydroxy compounds in blood, high levels were found in cases of Cushing's syndrome (156). Urinary excretion studies have been carried out with various methods. 17-ketosteroids have been found to be within normal range, or moderately elevated (131). This would correlate well with the moderate degree of virilizing manifestations seen in Cushing's syndrome in contradistinction to the marked degree observed in the various forms of adrenogenital syndrome. Corticoid excretion determined by bioassay, is elevated and this finding correlates well with isolation studies which have yielded large amounts of 17-hydroxycorticosterone (hydrocortisone). Reports of studies using either reducing methods or formaldehyde generation are less consistent: excretion levels have been found high in some and within normal range in other cases; most of these studies were carried out without β -glucuronidase hydrolysis. On the other hand, the enzyme appears to liberate substances not related to corticosteroids which react both in the reducing and formaldehydogenic procedures. Even with technical refinements of the formaldehydogenic method (90, 237) there are still considerable sources of error in this procedure. With the use of methods based on the Porter-Silber reaction (30, 64, 173, 179, 223) more consistent results are obtained, even though (or perhaps because) this reaction determines a more limited range of compounds (17-hydroxycorticosteroids). Determination of the 17-hydroxycorticosteroids and especially of the enzyme-liberated Porter-Silber positive compounds (17-hydroxycorticosteroids) has yielded high values in cases of Cushing's syndrome (173, 178); additional information can be obtained by separate determination of the free and the glucuronidase-liberated 17-hydroxysteroids (33).

3 *Metabolic Changes*

The retention of sodium and loss of potassium in Cushing's syndrome is a well established fact (see Volume II). Eventually this may lead to a severe hypokalemic hypochloremic alkalosis. To what extent the electrolyte changes are due to the action of C_{11} -oxygenated compounds alone or to a concomitant increased secretion of salt hormone is as yet unsettled.

Studies for the presence of aldosterone¹ will shed light on this problem. An interesting abnormality of the electrolyte metabolism pertains to the response to deoxycorticosterone acetate. Whereas in the normal individual as well as in the Addisonian injection of this compound together with intravenous administration of sodium chloride is followed by sodium and chloride retention, sodium and chloride loss occurs under the same conditions in cases of Cushing's syndrome (191, 192). There is at present no adequate explanation for the paradoxical action.

The disturbance of carbohydrate metabolism is clearly the result of excess secretion of C_{21} oxygenated steroids ('sugar hormones'). As would be expected the resulting diabetes is relatively insulin resistant. Differences of severity of impairment of carbohydrate metabolism ranging from severe diabetes to 'latent' diabetes (impaired glucose tolerance), may in part at least be due to differences of β cell 'reserve capacity'. The diabetogenic action of the C_{21} oxygenated corticosteroids is due to increased gluconeogenesis and to decreased glucose utilization. An interesting aspect of disturbed carbohydrate metabolism in Cushing's syndrome has received some attention recently. It has been shown previously that in normal individuals the fasting blood values for glucose, lactate and pyruvate were in the ratio of 100:10:1; this ratio is maintained following ingestion of glucose. In diabetics the rise of blood sugar during a glucose tolerance test is not associated with a proportional rise of lactate and pyruvate; these metabolites either fail to rise or this rise lags appreciably. In contradistinction in some cases of Cushing's syndrome the fasting levels of lactate and pyruvate were disproportionately high, and following glucose load the diabetic glucose curve was associated with high levels of lactate and pyruvate (89-111). This has been interpreted to indicate that in this form of 'steroid diabetes' the islets respond to the glucose load with good insulin secretion in spite of the diabetic shape of the glucose tolerance curve. High blood pyruvate levels following glucose load have also been reported after administration of cortisone and ACTH (111). If the high blood levels indicate increased formation of pyruvate this could conceivably also have some bearing on the fat formation via acetate and help to explain the obesity observed both in Cushing's syndrome and following cortisone and ACTH therapy.

¹ A number of important papers regarding the physiologic and therapeutic action of aldosterone have appeared after this manuscript was submitted. (R. Gaunt *et al.* *Endocrinol.* 55: 236 (1954); F. T. G. Prunty *et al.* *Lancet* III: 620 (1954); G. L. Farrell *et al.* *Proc. Soc. Exptl. Biol. Med.* 87: 141 (1954); J. A. Luetscher Jr. *et al.* *J. Clin. Invest.* 33: 276 (1954); J. A. Luetscher Jr. *et al.* *J. Clin. Endocrinol. and Metabolism* 14: 1086 (1954).)

4 Diagnosis

There has been no significant new contribution with regard to diagnosis. The frequency of occurrence of signs and symptoms as well as of abnormal laboratory data has been compiled by Plotz *et al* (163). Determination of urinary 17 ketosteroids is of limited diagnostic value because values are within normal range in many cases. Determination of 17 hydroxysteroids promises to be of great significance.

After the diagnosis of Cushing's syndrome is made it is obviously of greatest importance to determine presence or absence of a cortical tumor. An interesting biological test is based upon the response to exogenous cortisone. In the normal individual as well as in cases of adrenocortical hyperplasia administration of cortisone suppresses 17 ketosteroid excretion probably by suppressing ACTH secretion from the anterior pituitary with consequent suppression of secretion of the adrenal cortex. In cases of adrenocortical cancer cortisone administration is not followed by a decrease of 17 ketosteroid excretion evidently because the adrenocortical cancer is autonomous and continues to function in the absence of ACTH stimuli (97). In a case of Cushing's syndrome which one of us (K. E. P.) discussed in a CPC conference injection of a test dose of insulin was followed by profound and prolonged hypoglycemia usually seen in Addison's disease and in panhypopituitarism. This was interpreted as indicating the presence of an autonomous adrenocortical cancer associated with the typical atrophy of the other adrenal.

The biological tests (response to cortisone response to insulin) will have to be confirmed in a larger number of cases before the reliability and practical usefulness can be evaluated.

At present direct evidence for the presence of a tumor must still be sought in every case. To the older methods of roentgenography following urography and periadrenal air insufflation roentgenography following introduction of air into the retrorectal space (retroperitoneal pneumography) has been added (203). This procedure appears to be safe and has yielded good pictures in several cases. As before exploratory operation has to be resorted to in doubtful cases.

5 Course of Disease

The natural history of Cushing's syndrome has been critically reevaluated in a study by Plotz *et al* based on 33 observed cases and analysis of 189 cases reported in the literature (163). The rare occurrence of spontaneous remissions has been reemphasized by Knowlton (117) this may be important in trying to evaluate results of therapy.

6 Treatment

The treatment of choice in cases of tumor is clearly surgical removal of the tumor.

As mentioned in Volume II treatment in cases due to adrenocortical hyperplasia had not been very satisfactory. Roentgen irradiation of the pituitary is now being carried out with much larger doses (3000 to 5000 tissue r) than were employed previously. Good results with remission of long duration have been obtained in two series of cases (51, 103). However even these large radiation doses fail to bring about any improvement in some cases (33, 103). Partial hypophysectomy has been carried out in a few cases (128). B. Arner *et al.* report a good result following exposure and electrocoagulation of the pituitary (4). Beneficial results of subtotal and of total adrenalectomy have been reviewed (196). 2,2-bis (para-chlorophenyl) 1,1-dichloroethane (D D D) has been found to cause atrophy of the zona fasciculata of the adrenals of dogs rendering them insulin sensitive (146). It is also active in rats increasing their insulin sensitivity and alleviating alloxan diabetes but without causing adrenocortical atrophy (27, 28). Similar observations have been made in rabbits (161). However, in two cases of Cushing's syndrome this drug failed to exert any effect on corticoid excretion and on carbohydrate metabolism (33, 182).

D THE ADRENOGENITAL SYNDROME

As indicated in Volume II the clinical manifestations are due to increased secretion of androgenic 17 ketosteroids from the adrenal cortex. The anatomical change in the latter is either diffuse bilateral hyperplasia or tumors of one cortex. The picture will differ depending on the age and sex of the individual as summarized in Table I.

Excretion of 17 ketosteroids is increased above the levels normal for the respective age group. Higher values are found in tumor cases but there is some overlap. However excretion of more than 150 mg/24 hours indicates the presence of adrenocortical cancer. Tumor cases excrete proportionately more β 17 ketosteroids (mainly dehydroepiandrosterone) than do hyperplasia cases. In the latter α and β 17 ketosteroids are increased in approximately normal ratio the β fraction comprising up to 25% of the total. Separation into α and β fraction can be carried out by Digitonin precipitation or with the use of color reaction based on the Pettenkofen reaction. However cases of adrenocortical cancer with low β fractions have been recorded (62, 132) and in our experience also the differentiation of tumor and hyperplasia by determination of the α - β ratio is not reliable. Increased 17 ketosteroid values have recently also been found in blood in adrenogenital syndrome (61). Recently high urinary excretion values of pregnanetriol in adrenogenital syndrome have been

TABLE I
SYMPTOMS AND GENITAL SYNDROME

Age of onset	Anatomy of adrenal	Clinical manifestations		Remarks
		Female	Male	
1 Fetal	Diffuse hyperplasia	Female pseudohermaphrodite Persistent urogenital sinus Large clitoris Vulva in addition features like #2	Macrogenitosoma precocious Manifestations not conspicuous at birth	Final incidence in some instances Type 1 Effects of excess androgens only Type 2 Salt losing Effects of excess androgens plus salt hormone deficiency Type 3 Effects of excess androgens plus sugar hormone deficiency Tumor more frequent than in hyperplasia Similar syndrome in females from virilizing ovarian tumors in males from Leydig cell tumors and teratomas
2 Childhood	Tumor or diffuse hyperplasia	Heterosexual precocious puberty Large clitoris Body and facial hair Deep voice Advanced bone age Rapid growth in length Amenorrhea Absent mammary tissue	Isosexual precocious puberty Large penis Small testes Body and facial hair Deep voice Advanced bone age Rapid growth in length Azoospermia	Same as #2
3 Adult	Tumor or diffuse hyperplasia	Large clitoris Deep voice Body and facial hair Amenorrhea Atrophy of breasts		

found (20, 132-140), perhaps the high pregnanediol excretion reported in earlier studies (63, 221) actually was due to the presence of the triol in the precipitate. The possibility that the pregnanetriol is a metabolite of 17 hydroxyprogesterone, probably a precursor of hydrocortisone, has been discussed (20).

The pathogenesis of the diffuse hyperplasia poses a challenging problem. There is no evidence to suggest a primary hypothalamic or pituitary disorder causing excessive secretion of ACTH. However, high blood levels of ACTH have been reported (208). It is not immediately apparent how continuous stimulation by excessive ACTH causes excess androgen secretion only, and not panhypercorticism as seen in Cushing's syndrome. A very interesting hypothesis has been proposed by F. Bartter *et al.* (14). According to this schema the primary abnormality of the adrenal cortex in these cases is the decreased ability to produce C_{11} oxygenated steroids ("sugar hormones") or, with other words, a low target responsiveness of 11 oxysteroid production to ACTH. It has been subsequently suggested that the 'block' might be at the conversion of 17 hydroxyprogesterone to hydrocortisone and that perhaps the high excretion of pregnanetriol (probably a metabolite of 17 hydroxyprogesterone) is due to this block (20). The low levels of circulating 11 oxysteroids take the brakes off the anterior pituitary's ACTH secretion, the larger amounts of ACTH now in circulation are adequate to maintain a fair 11 oxysteroid production but the 17 ketosteroid production now becomes excessive because the sensitivity of the processes responsible for their production is normal; this hypothesis assumes that the adrenal androgens are much weaker inhibitors of ACTH secretion than the C_{11} oxysteroids. A number of observations in patients with adrenogenital syndrome would be explicable on the basis of Bartter's schema: ACTH administration fails to induce eosinopenia, nitrogen loss and increased urinary levels of reducing steroids in cases of adrenogenital syndrome. Also ACTH administration was not followed by Na retention or K loss (14-124). Low blood levels of C_{11} -17 oxygenated steroids have been reported in several cases (105); this is in apparent discrepancy with the normal urinary values of reducing or formaldehydogenic steroids found in other cases (14). The discrepancy may be more apparent than real since methods determining reducing and formaldehydogenic steroids are not entirely specific. On the other hand it is curious and not immediately understandable that individuals with low secretion of C_{11} - C_{17} oxysteroids should have no manifestations referable to this deficiency. Furthermore studies indicate that these adrenals are incapable of increasing their 11 oxysteroid production under additional ACTH stimulation (14-124) yet the individuals appear to tolerate infections and other stresses quite well. It would appear that

the steroid deficiency differs in degree in different cases being relatively mild and subclinical in most instances in others severe deficiency with hypoglycemia has been observed (231-235). Hypertension has been recorded in a number of cases (236).

In some cases of adrenogenital syndrome a disturbance of electrolyte metabolism similar to that present in Addison's disease is found (12-36). In this type designated as the salt losing type of adrenogenital syndrome inability to retain sodium lead to Addisonian crisis. From a clinical standpoint the most difficult cases are those occurring in very young infants in boys there may be little in the appearance of the patient to suggest virilizing adrenocortical hyperplasia because manifestations may not appear before the end of the first year of life. An assay of urinary 17 keto-steroid excretion will permit of early diagnosis.

In some cases an aplasia of the zona glomerulosa in addition to the hyperplasia of the inner zone of the cortex was found (18). In view of the fact that at least in rodents the zona glomerulosa is believed to elaborate salt hormone it was suggested that in these cases salt loss was due to a deficiency of salt hormones (234). Another explanation was suggested based on the observation that these patients responded to ACTH injection with sodium loss instead of with the sodium retention observed in normal individual. It is possible that these adrenals elaborate an abnormal hormone which causes Na loss. As yet such a compound has not been demonstrated. The effect of this hypothetical factor has to be overcome with large doses of deoxycorticosterone acetate and salt until cortisone successfully suppresses its secretion (101-124).

The familial incidence of congenital adrenocortical hyperplasia causing pseudohermaphroditism in girls and macrogenitosomia precox in boys has been known for some time. Bentinck and co-workers have analyzed all reported instances they point out that in no instance has the abnormality been found in more than one generation. Generally they believe that the evidence available militates against the assumption of a genetic factor (16).

The differentiation of cases due to adrenocortical tumor from those resulting from diffuse hyperplasia is of course of greatest practical importance. As mentioned above neither the urinary excretion level of total 17 keto-steroids nor the ratio of urinary alpha and beta 17 keto-steroid give reliable information. Whether pregnanetriol excretion in hyperplasia will prove a reliable differentiating feature from the tumor cases will have to await study of a larger material. Except in the congenital form which is invariably due to hyperplasia a tumor must always be suspected. The visualization of the adrenal glands on roentgenogram following air injections has been greatly improved by the use of retro-

found (20, 132, 140), perhaps the high pregnanediol excretion reported in earlier studies (63, 221) actually was due to the presence of the triol in the precipitate. The possibility that the pregnanetriol is a metabolite of 17 hydroxyprogesterone probably a precursor of hydrocortisone has been discussed (20).

The pathogenesis of the diffuse hyperplasia poses a challenging problem. There is no evidence to suggest a primary hypothalamic or pituitary disorder causing excessive secretion of ACTH. However, high blood levels of ACTH have been reported (208). It is not immediately apparent how continuous stimulation by excessive ACTH causes excess androgen secretion only, and not panhypercorticalism as seen in Cushing's syndrome. A very interesting hypothesis has been proposed by F. Bartter *et al.* (14). According to this schema the primary abnormality of the adrenal cortex in these cases is the decreased ability to produce C_{11} oxygenated steroids (sugar hormones) or with other words a low target responsiveness of 11α oxysteroid production to ACTH. It has been subsequently suggested that the 'block' might be at the conversion of 17 hydroxyprogesterone to hydrocortisone and that perhaps the high excretion of pregnanetriol (probably a metabolite of 17 hydroxyprogesterone) is due to this block (20). The low levels of circulating 11α oxysteroids take the brakes off the anterior pituitary's ACTH secretion: the larger amounts of ACTH now in circulation are adequate to maintain a fair 11α oxysteroid production but the 17 ketosteroid production now becomes excessive because the sensitivity of the processes responsible for their production is 'normal', this hypothesis assumes that the adrenal androgens are much weaker inhibitors of ACTH secretion than the C_{11} oxysteroids. A number of observations in patients with adrenogenital syndrome would be explicable on the basis of Bartter's schema: ACTH administration fails to induce eosinopenia, nitrogen loss, and increased urinary levels of reducing steroids in cases of adrenogenital syndrome. Also ACTH administration was not followed by Na retention or K loss (14, 124). Low blood levels of C_{11} 17 oxygenated steroids have been reported in several cases (108); this is in apparent discrepancy with the normal urinary values of reducing or formaldehydogenic steroids found in other cases (14). The discrepancy may be more apparent than real since methods determining reducing and formaldehydogenic steroids are not entirely specific. On the other hand it is curious and not immediately understandable that individuals with low secretion of C_{11} C_{17} oxysteroids should have no manifestations referable to this deficiency. Furthermore studies indicate that these adrenals are incapable of increasing their 11α oxysteroid production under additional ACTH stimulation (14, 124) yet the individuals appear to tolerate infections and other stresses quite well. It would appear that

stance adrenergic in nature which passes along the portal vessels from the hypothalamus to the adenohypophysis (130)

Not only organic lesions but also various stress reactions both physical and emotional may influence this mechanism markedly and in at least several different ways to produce several syndromes of pituitary ovarian dysfunction

A GENETIC ABNORMALITIES

1 Ovarian Agenesis (*Turner Syndrome (218) Turner Varney Syndrome*)

In this condition there is a congenital absence of the germ cells and failure of the follicular epithelium to develop, so that the ovary consists of a thin band of fibrous tissue lined by germinal epithelium lying in the mesosalpinx parallel to the Fallopian tube. It is not known whether this condition is due to failure of production of the germinal cells or to failure of the germinal cells to migrate from their site of origin in the hind gut into the gonad.

The question has been raised (232) as to whether all of these patients are in reality females as their external genital development denotes. Since in the male the Mullerian system appears to regress under the influence of androgen from the developing testes is it not possible that many of the essentially gonadal individuals are genetic males. On the other hand if this were so some other explanation would be required for the occasional male who presents a similar syndrome (p 865). Experimental observations made in young rabbit embryos by Jost (106) and in mice by Wells (229) support this hypothesis.

Many other genetic and endocrine features in this syndrome require clarification. The shortness of stature is hardly to be expected in view of the congenital hypogonadism. There is a remarkable tendency for most of these people to range in height from 54 to 56 inches. The skeletal proportions are usually normal rather than eunuchoid. The bone age is generally within the normal range occasionally being moderately delayed. Pubic hair is present in some but absent in other cases. Webbing of the neck is often noted. Among the other congenital defects sometimes seen are a receding chin, hypertension, coarctation of the aorta, osteoporosis with resulting orthopedic abnormalities and polydactylism. Some patients may show no congenital defects except for the hypogonadism and shortness of stature.

Gonadotropin (FSH) excretion is uniformly elevated and indeed serves as an important point in differential diagnosis from pituitary infantilism.

* A recent report by Wilkins *et al.* *J Clin Endocrinol and Metabolism* 14: 170 (1954) indicates that the chromosomal sex is male in most cases.

peritoneal pneumography (203) Air is injected into the retrorectal space and spreads upward retroperitoneally outlining all retroperitoneal structures. This procedure appears to be technically simpler, and above all safer than the perirenal or insufflation. If for some reason adequate visualization is not obtained, surgical exploration is necessary.

Treatment in tumor cases consists of course in surgical removal of the tumor. Treatment of hyperplasia cases was very unsatisfactory up to the introduction of cortisone therapy (232, 233, 236). Treatment with cortisone suppresses the function of the adrenal cortex, evidently by suppressing ACTH secretion from the anterior pituitary. The inhibition of androgen production causes arrest of the virilization in females as well as of the rapid growth and bone maturation with premature closure of epiphysis. Development is normalized and a normal female pubertal development in girls with menarche and breast development takes place. Evidently exogenously administered cortisone adequately replaces the suppressed secretion of the other cortical hormones.

1. *Treatment of the Salt Losing Type (36, 232)*

Treatment with cortisone alone even with administration of extra salt, is usually inadequate for maintaining electrolyte metabolism. Deoxycorticosterone acetate (DCA) is used in combination with cortisone the latter chiefly for suppressing androgenic 17 ketosteroid excretion the former for correction of the inability to retain sodium. It has been found that large doses of deoxycorticosterone and NaCl are necessary. This has been tentatively considered supportive evidence for the theory that the salt loss is due not to a deficiency of salt hormone but to the production of a Na losing adrenal compound the action of which has to be overcome by DCA. The requirement for the latter is appreciably reduced after cortisone therapy has brought about suppression of adrenocortical activity.

V. Ovary

Normal cyclic function of the ovaries with regard to both ovulation and hormone secretion involves not only a complex but also a very labile mechanism involving particularly (1) neurohumoral factors regulating the cyclic release of three gonadotropic substances (FSH, LH and LTH) (2) a precise sequential response of the germ cells to gonadotropic stimulation and (3) the cyclic secretion of steroid hormones by the granulosa and theca cells in proper amounts and at the proper time to favor a self continuing push pull relationship between the gonad and the hypophysis. The hypothalamus is undoubtedly an important relay station in the neurohumoral control of the pituitary ovarian axis and present evidence suggests that gonadotropin release is mediated by a neurohumoral sub

closure of the epiphyses will be hastened the secondary sex characters will develop as will the vagina and uterus and the vasomotor symptoms will be controlled but spontaneous ovarian function rarely follows the withdrawal of substitutional therapy

C PITUITARY HYPOGONADISM

Pituitary hypogonadism in women may be associated with pituitary dwarfism (p 826) or may be a single gonadotropic deficiency The skeletal development may therefore range from marked dwarfism to eunuchoidism depending apparently upon the degree of growth hormone deficiency In some instances with moderate retardation of growth the clinical picture may resemble that of ovarian agenesis but can be distinguished from the latter by absence of FSH in the urine Other congenital defects are usually absent the bone age is more markedly retarded and the urinary 17 ketosteroids are lower (often less than 10 mg per 24 hours) (232) As indicated above the differential diagnosis from a primary ovarian deficiency is of importance because of the possibility of a response to gonadotropic therapy

D NONNEOPLASTIC CYSTS OF THE OVARY

1 *Follicle Cysts of the Ovary*

The etiologic factors in follicular cystic disease of the ovary appear to be many and varied including genetic abnormalities inflammatory lesions of the ovary degenerative processes and endocrine factors Usually the physiologic disturbance results in a menstrual dysfunction and impairment of ovulation Estrogen secretion varies the follicular fluid in some of the large follicle cysts of the ovary contains small to moderate amounts of estrogenic activity whereas others are hormonally inactive Urinary excretion of estrogens also vary from diminished titers in most cases to excessive levels in occasional instances The endometrial pattern is likewise diverse ranging from an early proliferative type to one of cystic hyperplasia A secretory pattern with apparently normal ovulation may occur in the patient with one or a few follicle cysts of the hormonally inactive type in which the ovaries are otherwise normal

Little is known of the endocrine factors which may cause large follicle cysts of the ovary Occasionally these may be induced by too vigorous gonadotropic therapy (44) suggesting that a disturbance of the pituitary ovarian endocrine mechanism may be the etiologic factor in some cases In this respect it is of interest that the use of estrogens or androgens given in full dosage for several weeks may cause rapid regression of simple follicle cysts

in questionable case. Estrogens in the urine are very low. Despite the pattern of high gonadotropins and low estrogens vasomotor symptoms do not occur in these patients as they so often do in acquired types of prepubertal hypogonadism. The 17 ketosteroid excretion varies considerably; in some cases it is normal and in others it is quite low.

As a rule, the diagnosis of this condition is not established with certainty until the postpubertal years when the hypogonadism becomes evident. However, in instances in which it is suspected on the basis of dwarfism together with other congenital stigmata, the diagnosis can be made with certainty if there is a high urinary gonadotropic excretion, a feature distinguishing this condition from pituitary and other types of dwarfism. Wilkins (232) has noted that urinary FSH was absent or low during early childhood but rose as puberty was approached. The administration of estrogens produces a gratifying development of the breasts, secondary sex characters and growth of the vagina suitable for sexual purposes. This should be continued in cyclic fashion to maintain development but in moderate dosage since otherwise marked endometrial hyperplasia with troublesome bleeding can occur. We have not noted a stimulation of skeletal growth with estrogens even when treatment has been started in late childhood.

B. INFANTILE OVARIES WITH EUNUCHOIDISM

In this syndrome, the ovaries are of the elongated ovoid (infantile) type containing apparently normal numbers of primordial follicles. The follicles fail to grow and mature despite adequate FSH secretion as evidenced by high urinary titers. The irresponsiveness of the ovary to gonadotropic stimulation is believed to be due to a genetic defect although the possibility of a severe childhood infection affecting the ovaries must also be considered. These patients show the characteristic stigmata of prepubertal hypogonadism: they are relatively tall with eunuchoid proportions, the secondary sex characteristics and reproductive organs are immature, they have primary amenorrhea and hot flashes. The latter symptoms, as well as the high urinary FSH titers, distinguish them from the prepubertal eunuchoid hypogonad of pituitary origin. Furthermore, these patients rarely improve spontaneously, whereas the pituitary hypogonad may sometimes go into a delayed puberty in her late teens or early twenties and may eventually attain normal reproductive function. The latter may be encouraged or hastened by the cyclic administration of gonadotropic hormones with follicle stimulating activity, whereas the primary ovarian eunuchoid can be treated only symptomatically with estrogens or with estrogens and progesterone in cyclic fashion. The

closure of the epiphyses will be hastened the secondary sex characters will develop as will the vagina and uterus and the vasomotor symptoms will be controlled but spontaneous ovarian function rarely follows the withdrawal of substitutional therapy

C PITUITARY HYPOGONADISM

Pituitary hypogonadism in women may be associated with pituitary dwarfism (p 826) or may be a single gonadotropic deficiency The skeletal development may therefore range from marked dwarfism to eunuchoidism depending apparently upon the degree of growth hormone deficiency In some instances with moderate retardation of growth the clinical picture may resemble that of ovarian agenesis but can be distinguished from the latter by absence of FSH in the urine Other congenital defects are usually absent the bone age is more markedly retarded and the urinary 17 ketosteroids are lower (often less than 10 mg per 24 hours) (232) As indicated above the differential diagnosis from a primary ovarian deficiency is of importance because of the possibility of a response to gonadotropic therapy

D NONNEOPLASTIC CYSTS OF THE OVARY

1 *Follicle Cysts of the Ovary*

The etiologic factors in follicular cystic disease of the ovary appear to be many and varied including genetic abnormalities inflammatory lesions of the ovary degenerative processes and endocrine factors Usually the physiologic disturbance results in a menstrual dysfunction and impairment of ovulation Estrogen secretion varies the follicular fluid in some of the large follicle cysts of the ovary contains small to moderate amounts of estrogenic activity, whereas others are hormonally inactive Urinary excretion of estrogens also vary from diminished titers in most cases to excessive levels in occasional instances The endometrial pattern is likewise diverse ranging from an early proliferative type to one of cystic hyperplasia A secretory pattern with apparently normal ovulation may occur in the patient with one or a few follicle cysts of the hormonally inactive type in which the ovaries are otherwise normal

Little is known of the endocrine factors which may cause large follicle cysts of the ovary Occasionally these may be induced by too vigorous gonadotropic therapy (44) suggesting that a disturbance of the pituitary ovarian endocrine mechanism may be the etiologic factor in some cases In this respect it is of interest that the use of estrogens or androgens given in full dosage for several weeks may cause rapid regression of simple follicle cysts

2 *The Stein Leventhal Syndrome*

A particular type of microfollicular cystosis of the ovary which is of interest because of its endocrine stigmata is that seen in the Stein Leventhal syndrome (201, 202, 123). This condition is characterized by a menstrual dysfunction (usually amenorrhea but sometimes irregular or excessive bleeding), infertility, and hirsutism of varying degree, together with bilateral cystic enlargement of the ovaries. The condition generally begins in the late teens or early twenties, usually after prior normal menstrual function. Although the hirsutism may become quite marked in some cases, the clitoris is rarely appreciably enlarged or the voice deepened. The 17 ketosteroid excretion is generally within the normal range, occasionally being slightly elevated (up to 20 mg per 24 hours). A moderately high β fraction (20% to 40%) is often noted. Estrogens are present but usually in moderately low titers, and the vaginal smear shows a persistent early follicular reaction. FSH is often not demonstrable in the urine but high titers occur sporadically, especially early in the condition.

The appearance of the ovaries on exploration is quite characteristic. They are enlarged, flattened, and pale and have a thickened gray capsule; they resemble oysters. The cut surface reveals many small cystic areas of varying size. On histologic examination the significant abnormalities are the marked fibrous thickening of the ovarian capsule, the presence of numerous follicle cysts, with a paucity of normal growing follicles, and hyperplasia of the theca surrounding the cystic follicles. The latter finding is not consistently present.

A small type of Stein Leventhal ovaries is sometimes noted, in which the ovaries are only slightly enlarged and ovoid but have a markedly thickened capsule, microfollicular cysts, and theca hyperplasia. These patients often have excessive and irregular bleeding rather than amenorrhea.

The etiology of this condition is not known. There is little to support an inflammatory basis for the excessive fibrosis of the capsule which could secondarily produce follicular cystosis. A genetic dysfunction likewise seems improbable in view of the previous apparently normal menstrual history and absence of other stigmata in most cases. It seems more likely that the condition is due to an endocrine dysfunction. One wonders whether the hyperplastic theca may not be secreting increased amounts of an unusual steroid, perhaps not excreted as a 17 ketosteroid, which is responsible for the mild masculinization and inhibition of FSH. In this regard, it is to be noted that hirsutism has been observed in association with ovarian hyperthecosis (162, 167, 204, 211).

This condition may be differentiated (72) from an adrenogenital syndrome on the basis of lack of true masculinization and normal 17 ketoster

oids and from a Cushing's syndrome by lack of the typical habitus, purple striae and metabolic disturbances. A masculinizing tumor of the ovary may also be differentiated by lack of true masculinization and the fact that the ovarian enlargement is bilateral and 'cystic'.

Generous wedge resection of the ovaries frequently results in prompt and striking improvement of this syndrome, with return to normal ovulatory cycles in a high percentage of cases as indicated by basal temperature charts, endometrial biopsies and improved fertility (202). This improvement in endocrine function following removal of ovarian tissue is difficult to explain.

Hormonal therapy is not often of help in this condition. In its early stages occasional cases benefit from cyclic progesterone therapy or treatment with cortisone (p. 860).

E. OVARIAN DEFICIENCIES OF PSYCHOGENIC ORIGIN

It is well known that nervous or emotional disturbances may be followed by impairment of ovarian function resulting in anovulatory infertility, secondary amenorrhea, dysfunctional uterine bleeding, cystic mastitis or other manifestations of disturbed ovarian function. The clinical findings together with hormonal and other endocrine studies of such patients indicate that there are several patterns or syndromes of ovarian hypofunction which may occur on a psychogenic basis (153, 171, 206).

1 Hypogonadotropism. Inhibition of pituitary gonadotropic secretion with secondary ovarian deficiency appears to be most common in such disturbances. In its mildest form the menses may occur at fairly regular intervals but there is interference with normal ovulation causing relative infertility. This may progress to dysfunctional bleeding or to secondary amenorrhea depending upon the degree of inhibition of the pituitary-ovarian mechanism.

2 Inhibition of gonadotropins plus other pituitary hormones. These patients generally have secondary amenorrhea with absence of gonadotropin excretion in the urine plus clinical and metabolic evidences of mild hypothyroidism. In severe cases the clinical picture is that of anorexia nervosa with loss of weight, tiredness, hypotension and moderately decreased 17 ketosteroids. In occasional instances the clinical manifestations may be so marked as to require differentiation from Simmonds disease (57).

3 Normal FSH with ovarian deficiency. This is the pattern seen in patients with so-called hypothalamic amenorrhea as described by Klinefelter, Albright and Griswold (116). It has been presumed by these authors that the ovarian hypofunction results from lack of LH secretion to synergize with FSH to stimulate the ovary. The clinical

manifestations are those of secondary amenorrhea with anovulatory infertility. In our experience, this group is not as commonly seen as those described under the first classification although the clinical picture is often the same.

4 Primary ovarian deficiency with high FSH. This syndrome is characterized clinically by secondary amenorrhea and infertility and usually accompanied by hot flashes and other menopausal symptoms. It generally occurs after an acute psychic trauma and is often of long duration and highly resistant to treatment.

From the endocrine patterns which have been described above, it would appear that psychogenic factors may influence ovarian function by way of the pituitary, the hypothalamus, or the ovary directly. Unfortunately, very little is known of the intermediary mechanisms by which psychogenic factors influence these structures.

The therapeutic approach to psychogenic ovarian hypofunction is often difficult. In milder cases readjustment of environmental factors and superficial psychotherapy may be all that is required. For this reason, many such cases eventually undergo spontaneous correction. On the other hand, in many other instances the syndrome may be prolonged and may fail to respond even to intensive psychotherapy. Such patients are also known to be very resistant to endocrine therapy of various types. The use of gonadotropic hormones in patients with low FSH only rarely results in stimulation of ovarian function. Preparations containing follicle stimulating activity should be tried in cyclic fashion in the milder cases. In the more severe cases substitutional therapy with estrogen followed by estrogen and progesterone may bring on cyclic bleeding. These patients seem to require much higher doses of estrogen and progesterone to obtain an endometrial response and bleeding than patients with other types of amenorrhea (67, 171). The repeated induction of cyclic bleeding is occasionally followed by resumption of spontaneous ovarian function. This approach with substitutional therapy is of course the only plan of endocrine therapy indicated in patients with a primary ovarian deficiency.

Low dosage irradiation to the pituitary or ovaries is not usually helpful in these cases of psychogenic ovarian hypofunction (170) and is reserved for cases of infertility in which all other methods of treatment have failed. Better results are obtained in patients with a gonadotropic deficiency than in those with primary ovarian failure. The effects of the irradiation therapy appear to be mediated by way of the ovary rather than the pituitary (170).

F METABOLIC DISTURBANCES AFFECTING OVARIAN FUNCTION

In some patients excessive gain of weight may result in ovarian hypofunction with secondary amenorrhea or oligomenorrhea. This is usually due to inhibition of pituitary FSH secretion although the mechanism by which this occurs is not known. It is remarkable to observe in some patients the manner in which menstrual function may return and then disappear again depending upon their gain or loss of weight by dieting.

In this respect it is also to be noted that some young women may develop a very severe or resistant type of secondary amenorrhea following rapid loss of weight below their usual normal by severe dietary restriction (96). In these patients there is also an FSH deficiency. Unfortunately in some cases ovarian function does not resume spontaneously following a return to normal diet and normal weight. It is believed that diets deficient in protein and certain members of the B complex are particularly likely to cause ovarian dysfunction.

Disturbances in liver function may indirectly affect the function of the ovaries because of failure of normal inactivation of the ovarian hormones. Thus in hepatitis and in cirrhosis of the liver disturbances of ovarian function causing menorrhagia, polymenorrhea or cystic mastitis are sometimes encountered (98).

G EFFECT OF ADRENAL HYPOFUNCTION ON THE OVARIES

It is of course well known that marked degrees of adrenal cortical hyperfunction will inhibit ovarian function by depressing the gonadotropins. In the adrenogenital syndrome this may be attributed primarily to the androgens which are produced in excess. In some cases there is also increased estrogen secretion by the adrenal (99) the effects of which are masked by the greater increase in the androgens. However in some cases of Cushing's syndrome in which the 17 ketosteroids are not elevated and the total estrogen excretion is diminished the inhibition of gonadotropic function may be attributed to other steroids of the adrenal cortex.

The histologic picture of the ovaries in patients with an adrenogenital syndrome due to an adrenal cortical tumor has been of particular interest to us. In two such cases in which ovarian biopsies were obtained there was a striking resemblance of the histologic findings in the ovary to those seen in the Stein-Leventhal syndrome. Although the ovaries were not enlarged there was marked thickening of the capsule, microfollicular cystosis and hyperplasia of the theca cells.

Another type of ovarian dysfunction which appears to be related to a mild degree of adrenal cortical hyperfunction or dysfunction is seen in patients with mild hirsutism, anovulatory infertility, oligomenorrhea or

secondary amenorrhea and 17 ketosteroid excretion which is only slightly elevated above the normal, ranging from 15 to 22 mg per 24 hours. Such patients will sometimes respond favorably to the administration of cortisone in a dosage of 20 to 30 mg daily by mouth. Jones *et al* (105) have drawn attention to the use of cortisone in stimulating greater follicular activity in patients of this type. The mechanism by which cortisone produces this favorable action would appear to be similar to that described by Wilkins in patients with congenital adrenal hyperplasia, namely the depression of adrenal androgens by cortisone with subsequent release of FSH.

II OVARIAN HYPERFUNCTION

The most common and the most perfect example of ovarian hyperfunction is that which is seen in early pregnancy, during which the corpus luteum is stimulated to produce maximal amounts of estrogen and progesterone by the rising titer of chorionic gonadotropin secreted by the cytotrophoblast. Even in states of anterior pituitary hyperfunction the hypophysis is not capable of secreting sufficient luteinizing and luteotropic hormone to maintain and stimulate the corpus luteum to a degree comparable to that seen in early gestation. There are however several syndromes in which this may occur to a lesser extent.

I CORPUS LUTEUM CYST

In some instances in which there has been excessive hemorrhage into a corpus luteum this structure may become cystic and persist with functional activity beyond the usual two week period so regularly noted in the human cycle. Menstruation may be delayed because of a persistently high estrogen and progesterone level. There may be signs and symptoms of early pregnancy or the clinical picture may suggest an ectopic pregnancy. Also, in some cases, the pregnancy test may be weakly positive, suggesting increased secretion of LH and LTH.

Rare patients may often or regularly have a slightly prolonged corpus luteum phase, lasting for 16 to 18 days as indicated by basal temperature charts. This occurs occasionally in patients who are particularly anxious for pregnancy but who do not develop a true pseudocyesis syndrome.

I PSEUDOCYESIS

The pseudocyesis syndrome is a striking example of the influence of psychogenic factors upon the pituitary-ovarian mechanism to produce a state of ovarian hyperfunction. This condition occurs primarily in women who have a great desire to become pregnant but who have been unable to do so or sometimes in women who have a great fear that they have become pregnant illegitimately or against their wishes. Endocrine studies (38-59)

indicate that they have persistent corpus luteum function characterized by hyperhormonal amenorrhea associated with a secretory endometrium, persistently good levels of estrogen and pregnanediol plus such other evidences of ovarian hyperfunction as increased glandular tissue of the breasts pigmentation of the nipples softening of the cervix and slight enlargement of the uterus. FSH in the urine is generally absent but assays for prolactin have been positive. These findings explain the milky secretion which may often be expressed from the nipples of these patients. It would seem that psychogenic factors have in some way interfered with the pituitary hypothalamic mechanism so as to favor continued secretion of LH and LTH and persistence of the corpus luteum. The endocrine disturbance together with the other psychogenic manifestations accounts for the full blown syndrome of pseudocyesis. Psychiatric studies on these patients indicate that the psychosis is a type of hysteria. Psychotherapy is only of limited value in the treatment of this condition which also has a great tendency to be recurrent. Testosterone is sometimes helpful apparently by inducing regression of the corpus luteum. This is generally followed by a menstrual flow suggesting to the patient that she is no longer pregnant.

VI Placenta

The human placenta is a versatile and prodigious producer of both protein and steroid hormones. There is well-documented evidence that the placenta secretes large amounts of chorionic gonadotropin estrogens and progestogens. There is also good evidence suggesting that the placenta secretes glucocorticoids (220-100-150). Studies of the ketosteroids in the urine of pregnant women (48) show both qualitative and quantitative differences from those of nonpregnant women suggesting that the placenta produces other steroid precursors or that pregnancy induces alterations in steroid metabolism or both. Evidence for the secretion of a protein hormone containing ACTH activity has also been presented (101).

There is increasing evidence (205-238) that the cytotrophoblast (Langhans layer) is the site of origin of chorionic gonadotropin and possibly of other protein hormones whereas histochemical studies suggest that the syncytial cells are the site of formation of the steroid hormones.

Comparatively little is known concerning the functions of the placenta in gestation. Chorionic gonadotropin appears in the blood and urine in rapidly rising titers in the early days of pregnancy and apparently maintains the corpus luteum of pregnancy. During the last half of pregnancy the chorionic gonadotropin levels are much lower. The function of this hormone during the placental phase is not known; it has been suggested that it stimulates the production of estrogens and progesterone by the syncytial cells (190). The early appearance of chorionic gonadotropin in

blood and urine and its high concentration make it possible to use its assay as the basis for the more reliable pregnancy tests including the Aschheim Zondek, Friedman, South African frog, American male frog, and rat-ovary hyperemia tests.

Estrogens and progesterone are secreted in increasing amounts as pregnancy progresses. It has been generally believed that during the first trimester the bulk of these steroids are of corpus luteum origin and that the corpus luteum is essential for the maintenance of pregnancy during this period. More recent studies (118) would indicate that in the human being the corpus luteum is essential for a much briefer period of time and that removal of the corpus luteum during the first weeks of pregnancy does not necessarily result in termination of the pregnancy or diminution in pregnanediol excretion, thus suggesting that the trophoblast may adequately take over this function very early.

Estrogen and progesterone appear to be essential for growth and quiescence of the uterine muscle during gestation (174). They also induce many other changes favorable for mother and fetus such as the further growth and development of the breasts, softening of the cervix, thickening of the vaginal epithelium, and increased vaginal acidity. With regard to some of these changes the action of the two hormones appears to be synergistic, whereas in others they are antagonistic.

The intermediary metabolism of estrogen and progesterone during pregnancy is discussed in Volume I of *The Hormones*.

A. PREGNANCY COMPLICATIONS

1. Abortion

Several patterns of hormonal imbalance have been observed (104) in patients with sporadic or habitual abortion. The latter condition has been particularly useful for such studies, since the hormonal excretion patterns have been studied from early gestation to the time of abortion. In one type the titer of chorionic gonadotropin is low from the very early days of pregnancy and progressively falls. Estrogen and pregnanediol excretion is also diminished. In these pregnancies the fetus is often resorbed and there is reason to suspect a genetic defect. In another type pregnanediol and estrogen levels are diminished in spite of normal titers of chorionic gonadotropin, suggesting poor response by the corpus luteum or syncytiotrophoblast. In this group substitutional therapy with large doses of progesterone and estrogens appears to be particularly indicated. In a third type abortion may occur repeatedly in spite of normal urinary levels of all three hormones, suggesting a uterine factor or possibly a disturbance of other unknown hormonal factors. Diminished thyroid function must be

considered it has been suggested (158) that failure of the serum protein bound iodine to rise during pregnancy may be of significance in this regard

2 *Hydatidiform Mole and Chorionepithelioma*

In many instances these conditions are associated with unusually high titers of chorionic gonadotropin which may be helpful in their diagnosis. Since hydatidiform mole is most often suspected during early pregnancy when the titer is normally quite high a high serum chorionic gonadotropic titer will be indicative of hydatidiform mole only if it is several times higher than the peak values for normal pregnancy or if a moderately high level is sustained significantly beyond the period when the titer normally drops (12th to 16th weeks). A rising titer of chorionic gonadotropin following expulsion of a mole or after an abortion or normal delivery is presumptive evidence of chorionepithelioma provided the possibility of another pregnancy has been ruled out. Patients who have expelled a mole or have had surgery for a chorionepithelioma should have repeated follow up assays over a long period of time in order to detect recurrence or metastases. It should however, be emphasized that occasionally chorionepitheliomas produce little or no chorionic gonadotropin particularly in their early stages (212). The secretion of estrogens and progesterone by moles and chorion epitheliomas varies much more than that of chorionic gonadotropin the assay of the former is therefore less useful from a diagnostic standpoint. Some of these tumors may secrete very large amounts of chorionic gonadotropic hormone with but little if any estrogens or progesterone whereas others may secrete considerable amounts of all three hormones (148).

3 *Toxemias of Pregnancy*

There are many reasons to suspect a placental factor possibly of hormonal nature of etiologic significance in the toxemias of pregnancy but satisfactory evidence for such an endocrine factor has not yet been obtained. Results of hormonal excretion studies in toxemias of pregnancy are quite controversial. The pattern reported by the Smiths (190) in many instances namely increased urinary chorionic gonadotropin and diminished urinary pregnanediol and estrogen as well as changes in the partition of the urinary estrogens has in part been confirmed by some workers. On the other hand these alterations are often absent in patients with severe pregnancy toxemia (155, 160, 193).

An increase in the urinary excretion of corticoids in toxemia of pregnancy has been reported (47, 148, 216) this finding has been tentatively correlated with the sodium and water retention observed in toxemia. The increase above that occurring in normal pregnancy is probably of questionable significance especially in view of the methods employed in these studies.

Even if such an increase will be confirmed with the use of more specific methods such as those based on the Porter-Silber reaction, the question will still remain what is cause, and what effect? The occurrence of a severe toxemia could well cause a non specific rise of adrenocortical secretion. It has been suggested that increased production of a mineral corticoid might play a role in this condition (147). *This entire problem appears to be quite unsettled and further advances will depend in part on better assay methods for these steroids, particularly for the mineralocorticoids.* Similarly the question of the use of cortisone or ACTH in the treatment of late pregnancy toxemias is controversial (147, 211).

VII Testes

The hypophyseal testicular interrelationships have been discussed in Volume II. Several problems in this sphere remain unsolved. Although FSH is believed to be primarily responsible for stimulation of spermatogenesis, it is uncertain whether this occurs without the synergism of the other gonadotropins or of androgens. It is equally uncertain whether luteotropin (LTH prolactin) plays any role in the male, perhaps as a synergist of LH (ICSH) in stimulating Leydig cell function. The question of a 'second' testicular hormone awaits final answer. The existence of such a hormone has been postulated on the basis of analogy with the female Interstitial tissue (and corpus luteum) in the female and interstitial tissue (Leydig cells) in the male producing progesterone and testosterone respectively is regulated by LH (ICSH). Ovarian follicles and estrogen secretion in the female and spermatogenic tubules in the male are governed by FSH. Is there a tubular hormone in the male corresponding to the follicular estrogen?

In addition to such general considerations certain clinical observations suggest the existence of a tubular hormone. In Klinefelter's syndrome (Volume II p. 679) hyalinization of the seminiferous tubes and intact Leydig cell function are associated with excessive urinary FSH excretion. This suggests that under normal circumstances the tubules secrete a hormone the absence of which releases pituitary FSH secretion. It has been proposed that this hormone might be an estrogen. Estrogens have been extracted from the testis of human beings as well as certain animal species and are believed by some to be secreted by the Sertoli cells. This assumption is based chiefly on the findings in Sertoli cell tumors of the testis which induce feminization in dogs and in man. From Sertoli cell tumors of dogs estrogens have been extracted. However Nelson has questioned the origin of these testicular tumors from Sertoli cells and believes them to be Leydig cell tumors (144). If androgens and estrogens are according to

Nel on produced in the Leydig cell the postulated tubular hormone still goes begging

A HYPAGONADISM (91)

Hypogonadism is discussed in Volume II of *The Hormones*

In view of the fact that in many instances nothing is known regarding the etiology of the various syndromes hypogonadism may be classified, on a symptomatological basis as follows Failure (1) of both tubular (spermatogenic) and Leydig cell function (2) of tubular function with intact Leydig cell function and (3) of Leydig cell function with intact tubular function All three types can at least theoretically be caused (a) by a primary testicular deficiency and (b) by a testicular deficiency secondary to pituitary gonadotropic failure

1 Failure of Tubular and Leydig Cell Function

This type is represented by the surgical castrate (eunuch) and by the eunuchoid in whom testicular failure is due to an anlage defect (functional castrate 86), or due to impaired blood supply incident to operations for cryptorchidism or hernia or due to bilateral orchitis

These cases may be indistinguishable in their physical appearance and in their functional deficiencies from cases of pituitary hypogonadotropism Differentiation of primary testicular eunuchoidism and eunuchoidism secondary to hypogonadotropism is readily accomplished by assay of urinary gonadotropins (FSH assay) which yields very high values in the former whereas gonadotropin excretion is absent in the latter

The castrates and the primary testicular eunuchoid are treated with testosterone In hypogonadotropic eunuchoids treatment with gonadotropic hormones (chorionic gonadotropin and FSH containing pituitary preparations) has been successful

Whether an anlage defect of the testis ever is associated with a syndrome analogous to that of ovarian agenesis (Turner's syndrome) is as yet uncertain As is discussed elsewhere (p 83) the absence of estrogen effects and of germ cells and the presence of high urinary gonadotropin (FSH) levels in ovarian agenesis are associated with features which are different from those of the prepubertal castrate shortness of stature frequent occurrence of other congenital malformations including webbing of the neck usually no delay of closure of the epiphyses None of the few cases reported as Turner's syndrome in the male (194) appear to us convincingly analogous to the female syndrome The suggestion has been made (p 83) that some patients with ovarian agenesis may in fact be genetic males and thus represent instances of testicular aplasia

Even if such an increase will be confirmed with the use of more specific methods, such as those based on the Porter-Silber reaction, the question will still remain what is cause, and what effect? The occurrence of a severe toxemia could well cause a non specific rise of adrenocortical secretion. It has been suggested that increased production of a mineral corticoid might play a role in this condition (147). This entire problem appears to be quite unsettled and further advances will depend in part on better assay methods for these steroids particularly for the 'mineralocorticoids'. Similarly the question of the use of cortisone or ACTH in the treatment of late pregnancy toxemias is controversial (147, 211).

VII Testes

The hypophyseal testicular interrelationships have been discussed in Volume II. Several problems in this sphere remain unsolved. Although FSH is believed to be primarily responsible for stimulation of spermatogenesis it is uncertain whether this occurs without the synergism of the other gonadotropins or of androgens. It is equally uncertain whether luteotropin (LTH prolactin) plays any role in the male perhaps as a synergist of LH (ICSH) in stimulating Leydig cell function. The question of a 'second' testicular hormone awaits final answer. The existence of such a hormone has been postulated on the basis of analogy with the female Interstitial tissue (and corpus luteum) in the female and interstitial tissue (Leydig cells) in the male producing progesterone and testosterone respectively, is regulated by LH (ICSH). Ovarian follicles and estrogen secretion in the female and spermatogenic tubules in the male are governed by FSH. Is there a tubular hormone in the male corresponding to the follicular estrogen?

In addition to such general considerations certain clinical observations suggest the existence of a tubular hormone. In Klinefelter's syndrome (Volume II p. 679) hyalinization of the seminiferous tubes and intact Leydig cell function are associated with excessive urinary FSH excretion. This suggests that under normal circumstances the tubules secrete a hormone the absence of which releases pituitary FSH secretion. It has been proposed that this hormone might be an estrogen. Estrogens have been extracted from the testis of human beings as well as certain animal species and are believed by some to be secreted by the Sertoli cells. This assumption is based chiefly on the findings in Sertoli cell tumors of the testis which induce feminization in dogs and in man. From Sertoli cell tumors of dogs estrogens have been extracted. However Nelson has questioned the origin of these testicular tumors from Sertoli cells and believes them to be Leydig cell tumors (144). If androgens and estrogens are according to

(119-133-151) The evidence for the latter varied somewhat in individual cases: eunuchoid body proportions, sparseness or absence of pubic and axillary hair, poor beard growth, high pitched voice, undersized penis as well as diminished urinary 17 ketosteroid values have been reported. Testicular biopsy specimens showed diminished number and pycnotic appearance of Leydig cells and production of mature sperm in the tubules. In some instances an ejaculate could not be obtained, in one case the fructose content of the ejaculate was very low (119), the fructose content of the semen is considered to be a measure of androgen secretion (120). In other cases the ejaculate was normal and fertility was claimed for one of these cases. McCullagh has applied in his cases an assay method believed to show LH (ICSH) in urinary extract. In three of his five cases LH was not demonstrable, suggesting a pituitary deficiency limited to this one hormone. Some response was obtained by treatment with chorionic gonadotropin; this would add further support to the assumption of a pituitary deficiency (133).

B TESTOSTERONE AS A GONADOTROPIN

Several older studies are on record indicating that under narrowly circumscribed experimental conditions testosterone can exert a gonadotropic function. In the male hypophysectomized rat treatment with testosterone initiated immediately after hypophysectomy will maintain spermatogenesis and prevent testicular atrophy, but treatment with testosterone will not repair the previously atrophied testicle nor restore spermatogenesis once the posthypophysectomy changes have taken place (145-187).

There are for obvious reasons no observations in man which would duplicate the experimental conditions in the rat. However in several cases of hypopituitarism sperm production occurred under treatment with testosterone (93-113). In contradistinction to the hypophysectomized animal the possibility cannot be ruled out in such a case that testosterone therapy induced the pituitary to secrete gonadotropins. Kinsell rejects this possibility because no rise of urinary gonadotropin titer was observed during androgen therapy (113).

C TESTICULAR TUMORS

Tumors of the testis will be discussed briefly only with regard to endocrine aspects. Leydig cell tumors secrete large amounts of androgens and 17 ketosteroids and androsterone sulfate has been isolated from the urine. In prepubertal boys these tumors cause precocious pseudopuberty (p. 869). It has been pointed out that probably because of the protein anabolic action of the androgen secreted by the tumor tumor cachexia may be absent even in advanced cases with widespread metastasis (133). It is

2 *Impairment of Tubular Function with Intact Leydig Cell Function*

Typically the individuals are normally developed males, with normal secondary sex characteristics, normal libido and potency, and normal 17 ketosteroid excretion. The size of the testes varies from normal to markedly diminished. The sperm content of the ejaculate varies from oligospermia to azospermia. Testicular biopsy specimens show normal Leydig cells; the germinal epithelium may be well developed, with spermatogenesis progressing to the spermatide stage ('spermatogenesis'), but with absent sperm maturation ('spermiogenesis'). In other cases the tubules are completely denuded with only Sertoli cells left. Pericanalicular fibrosis is present in varying degrees in most but not in all cases of the latter type. Urinary gonadotropin excretion is normal in some and very high in other cases, but a definite correlation of the FSH excretion pattern with the histological picture of the testis is as yet not possible. There is no evidence that any of these cases is due to a pituitary gonadotropic deficiency. The etiology is unknown.

It has been claimed that seminal failure can be recognized already in the histologic picture of the pubertal testis by marked retardation of tubular development (31).

The Klinefelter syndrome discussed in Volume II, characterized by hyalinization of the tubules, normal Leydig cells, gynecomastia, and high FSH excretion, would appear to be a special entity within the group of seminal failure without Leydig cell failure (45-116). Cases have been reported which show features of Klinefelter syndrome associated with varying degrees of Leydig cell failure (85-87).

Treatment. Cases with azospermia and extensive fibrosis and/or hyalinization of the tubules as seen in the biopsy specimen are obviously beyond any possibility of repair. However, results are also unsatisfactory in most instances of milder forms with oligospermia and maturation arrest. Various gonadotropic preparations have been used on a purely empirical basis. One would expect results if any from preparations containing adequate amounts of FSH such as pituitary gonadotropins or Synapoidin. Chorionic gonadotropin has also been used and occasional good results have been claimed with any of the preparations. Treatment with large doses of testosterone has also been suggested; complete suppression of sperm production is reported to be followed by a rise of viable sperm cells in the ejaculate to values permitting fecundation ('rebound effect') (82). Confirmation has to be awaited.

3 *Leydig Cell Deficiency with Normal Tubular Function*

Recently three reports have appeared concerning patients who had spermatogenesis in the presence of evidence for impaired Leydig cell function

of the sex steroids. These amounts are too small to induce somatic changes as evidenced by the absence of secondary sex characteristics in normal children and by the absence of detectable somatic changes in prepubertal castrates. However the fact that the prepubertal castrate (80) as well as the child with ovarian agenesis (18a) may secrete appreciable amounts of FSH indicates that the child's gonad is not wholly inactive. The pubertal spurt of sex steroid secretion from the gonad is caused by a spurt of gonadotropin secretion from the anterior pituitary. This can be shown by urinary gonadotropin assay and is evident from the fact that administration of exogenous gonadotropin causes sex hormone secretion and development of secondary sex characteristics years before the physiological onset of puberty.

The anterior pituitary probably is triggered into gonadotropin release at the time of puberty by an hypothalamic mechanism presumably by the secretion of a neurohumeral agent which reaches the anterior pituitary by way of the portal venous system of the pituitary stalk. Just as the gonad is capable of responding to gonadotropic stimulation long before the physiological onset of puberty, the anterior pituitary of animals (181) and human beings (11) contains appreciable amounts of gonadotropic hormones without releasing them. The release of gonadotropic hormones from the anterior pituitary through the hypothalamic mechanism has been demonstrated for the LH release in the rabbit (130); the fact that certain organic lesions of the hypothalamus in children cause precocious puberty suggests that a similar mechanism operates in the initiation of the pubertal chain of events in the human being.

Precocious puberty is found in cases of hypothalamic lesions (tumors, encephalitis, hydrocephalus); perhaps the precocious puberty associated with polyostotic fibrous dysplasia (Albright's disease) is due to cerebral changes resulting from deformities of the base of the skull. In the majority of cases of precocious puberty no organic lesion is present; these cases are referred to as constitutional precocious puberty. It may be assumed that for some reason the intrinsic triggering mechanism in the hypothalamus is set off abnormally early in life. Because in precocious puberty both in that due to organic lesions and in the constitutional type the entire physiological pubertal mechanism is duplicated, the precocious development includes that of the gonad involving both its endocrine secretions and maturation of ova and sperm. Pregnancy has been reported in a girl $5\frac{1}{2}$ years old.

Tumors or hyperplasia of the adrenal cortex, tumors of the testis or ovary, and teratomas containing testicular tissue cause precocious development of secondary sex characteristics, either isosexual or heterosexual, but not maturation of the gonad. This type of precocity is called precocious pseudopuberty.

interesting that the estrogen excretion may be increased simultaneously in such cases (222). This finding would add weight to the theory that the Leydig cell is the source of both androgens and estrogens (p. 861). However, too few cases have been adequately and completely studied to permit any generalizations or final conclusions, unfortunately reports on this rare tumor continue to appear in the literature with little information other than a histologic study of the tumor.

Tumors of the testis believed to originate from Sertoli cells, have been found in man (210) and more commonly in the dog (92). These tumors secrete estrogens and cause gynecomastia. Nelson questions their origin from Sertoli cells and believes them to be a variety of the Leydig cell tumor (144) (see also p. 861).

Certain malignant tumors of the testis contain chorion like tissue (choriocarcinoma) they are characterized by excretion of large amounts of gonadotropins which are biologically identical with chorionic gonadotropins of pregnancy. Pregnancy tests' (Asheim Zondek Friedman, rat ovary hyperemia test) carried out with urine of such cases are positive. Increased urinary estrogen levels and the occurrence of gynecomastia in some cases suggest that the choriocarcinoma duplicates not only the gonadotropin production but also the estrogen production of normal chorionic tissue. There is suggestive evidence that in some cases of choriocarcinoma of the testis there is increased urinary excretion not only of chorionic gonadotropin but also of FSH. Because of the technical difficulties of assaying the two gonadotropins separately in a mixture of both the data should at present not be considered final. The presence of large amounts of urinary FSH in such cases has been tentatively explained as the result of destruction of testicular tissue (castration by tumor) but in view of the fact that the same hormone pattern occurs in unilateral tumors with intact second testis this explanation is not wholly satisfactory. Similar observations of increased urinary FSH levels have been reported in other testicular tumors. Regarding the significance of this finding the same uncertainty has to be recorded as that mentioned above in the case of choriocarcinomas. No essentially new facts have been reported recently a good review of the subject appeared in 1946 (219).

VIII Puberty (149)

In our population the onset of puberty normally takes place at the age of about 11 to 12 years in girls and 12 to 13 years in boys. From this mean age of onset there occurs considerable deviation with onset delayed up to 18 to 20 years or advanced to as early as 1 to 2 years of age (precocious puberty). The onset of pubertal changes is brought about by the beginning of the secretions of estrogens and androgens respectively. The prepubertal gonad appears to be not completely inactive but to secrete small amounts

- 35 Conn J W and Louis I H *J Clin Endocrinol* 10 1 (1950)
- 36 Engler J F Jr Silverman S H and Wilkins I *Pediatrics* 10 39 (1952)
- 36a Crile C and Hazard J B *J Clin Endocrinol* 11 1123 (1951)
- 37 Culiner A and Shappel S *J Obstet and Gynaecol Brit Empire* 56 439 (1949)
- 38 Curtis A C Cawley L I and Johnwick E B *Arch Dermatol and Syphilol* 60 318 (1949)
- 39 Cushing H and Davidoff I M *Arch Internal Med* 39 673 (1927)
- 40 Cutler H H Power M H and Wilder R M *J Am Med Assoc* 111 117 (1938)
- 41 D Angelo S A Paschke K E Cantarow A Siegel A N and Rivero Fontan J L *Endocrinology* 49 621 (1951)
- 42 D Angelo S A Paschke K E Cordon A S and Cantarow A *J Clin Endocrinol* 11 1237 (1951)
- 43 D Avignon M and Melin K A *Acta Paediat* 40 368 (1951)
- 44 Davis M E *In J V Meigs and S H Sturge Progress in Gynecology* Grune & Stratton New York 1946 Vol I
- 45 de la Balze F A Arrillaga F C Irazu J and Mancini R E *J Clin Endocrinol and Metabolism* 12 1426 (1952)
- 46 Denny Brown D *Am J Med* 15 368 (1953)
- 47 Devis R and Devis Vanden F *J Clin Endocrinol* 9 1436 (1950)
- 47a DiGeorge A and Paschke K L Unpublished observations
- 48 Dobriner K Lieberman S Rhoads C P and Taylor H C *In The Normal and Pathologic Physiology of Pregnancy* Williams & Wilkins Baltimore 1948
- 49 Doherty B M and Steelman S I *Endocrinology* 52 60 (1953)
- 50 Dohan F C Jose E Eiman J W Richardson F M and Zintel H A *J Clin Endocrinol and Metabolism* 13 415 (1953)
- 51 Dohan C F Rose L Boucot N and Chamberlain R *Trans & Studies College Physicians Phila* 20 150 (1953)
- 52 Engl F I and Scott J I Jr *J Clin Invest* 29 151 (1950)
- 53 Farquharson R F *Simmonds Disease* Charles C Thomas Springfield Ill 1950
- 54 Fertman M B and Curtis G M *J Clin Endocrinol* 11 1361 (1951)
- 55 Fisher G Epstein D and Paschke K E *J Clin Endocrinol and Metabolism* 12 1100 (1952)
- 56 Fortier C *Ciba Colloq Endocrinol* 4 148 (1952)
- 57 Frawley T F and Thorn G W *Proc 2nd Clin ICTH Conf* 1 115 (1951)
- 58 Fried P H Rakoff A J Schopback R R and Kaplan A *J Am Med Assoc* 145 1329 (1951)
- 59 Fried P H and Rakoff A E *J Clin Endocrinol* 12 371 (1952)
- 60 Friedgood H B *In The Chemistry and Physiology of Hormones* Am Assoc Advancement of Science Washington D C 1944
- 61 Gardner I I *Proc Soc Exptl Biol Med* 83 201 (1953)
- 62 Genell S *Acta Endocrinol* 7 133 (1951)
- 63 Genitts A E and Bornstein I P *J Am Med Assoc* 119 704 (1942)
- 64 Glenn E M and Nelson D H *J Clin Endocrinol and Metabolism* 13 911 (1953)
- 65 Goldberg R C and Chaikoff I I *Arch Pathol* 53 27 (1952)
- 66 Golden A Bonly P K and Sheldon W H *Proc Soc Exptl Biol Med* 74 450 (1950)
- 67 Goldzieher M A and Goldzieher J W *J Clin Endocrinol* 12 42 (1952)

REFERENCES

- 1 Albright F *Harvey Lectures* (1942-43) p 123
- 2 Allen H C Jr, Kelly F J and Greene J A *J Clin Endocrinol* **12** 1356 (1952)
- 3 Allen H C Jr, Libby R I and Cassen B *J Clin Endocrinol* **11** 492 (1951)
- 4 Arner B, Luft R, Olsson H and Sjogren B *J Clin Endocrinol and Metabolism* **13** 1101 (1953)
- 5 Ashoe Hansen G, Iversen K and Wichman R *Acta Endocrinol* **11** 3,6 (1952)
- 6 Astwood E B *Ann Internal Med* **30** 1087 (1949)
- 7 Astwood E B, Greer M A and Ettlinger M G *J Biol Chem* **181** 121 (1949)
- 8 Astwood E B, Raben M S and Payne R W *Recent Progr Hormone Research* **7** 1 (1952)
- 9 Astwood E B and Stanley M M *West J Surg* **55** 625 (1947)
- 10 Auger, H O *Am J Roentgenol Radium Therapy Nuclear Med* **70** 605 (1953)
- 11 Bahn R C, Lorenz H, Bennett W A and Albert A *Endocrinology* **52** 605 (1953)
- 12 Barnett H L and McNamara H *J Clin Invest* **28** 1498 (1949)
- 13 Bartels E C *J Clin Endocrinol* **9** 1190 (1949)
- 14 Bartter F C, Albright F, Forbes, A P, Leaf A, Dempsey F, and Carroll E *J Clin Invest* **30** 237 (1951)
- 15 Beierwualtes W H *J Clin Endocrinol and Metabolism* **13** 1090 (1953)
- 16 Bentinck R C, Hinman F, Lissner H and Traut H F *Postgraduate Med* **11** 301 (1952)
- 17 Berson S A, Yalow R S, Sorrentino J and Roswitt B *J Clin Invest* **31** 141 (1952)
- 18 Blackman S S Jr *Bull Johns Hopkins Hosp* **78** 180 (1946)
- 19 Blom I S and Terpstra J *J Clin Endocrinol* **13** 949 (1953)
- 20 Bongiovanni A M *Bull Johns Hopkins Hosp* **92** 244 (1953)
- 21 Bongiovanni A M and Eisenmenger W J *J Clin Endocrinol* **11** 152 (1951)
- 22 Bornstein I P, Bower I E and Murphy J *Am J Med Sci* **205** 114 (1943)
- 23 Bornstein J and Treshella P *Lancet* **ii** 678 (1950)
- 24 Borst J G G, ten Holt S I, de Vries I A and Molhuysen J A *Lancet* **i** 657 (1953)
- 25 Brant F *Am J Surg* **79** 645 (1950)
- 26 Brobeck, J R *Ciba Colloquia Endocrinol* **4** 124 (1952)
- 27 Brown J H U *Proc Soc Exptl Biol Med* **83** 59 (1953)
- 28 Brown J H U *Endocrinology* **53** 116 (1953)
- 29 Butterly J M, Fishman L, Secklen J and Steinberg H *Ann Internal Med* **37** 930 (1952)
- 30 Carroll K K, McAlpine H T and Noble R I *Can Med Assoc J* **65** 363 (1951)
- 31 Charny C W, Conston A S and Meranze D S *Fertility and Sterility* **4** 518 (1953)
- 32 Chart J J, Shipley E G and Gordon E S *Proc Soc Exptl Biol Med* **78** 224 (1951)
- 33 Cloud L P, Schneider J J and Paschkis K E Unpublished observations
- 34 Conn J W, Fajans S S, Louis L H and Johnson B *Proc 2nd Clin ICTH Conf* **1** 221 (1951)

- 109 Kepler E J *J Clin Endocrinol* 5 70 (1945)
- 110 Kepler E J Sprague R G Mason H I and Lower M H *Recent Progr Hormone Research* 2 315 (1948)
- 111 Kerppola W *Acta Med Scand* 145 357 (1953)
- 112 Kerr H D *Am J Roentgenol and Radium Therapy* 60 348 (1948)
- 113 Kinsell I W *J Clin Endocrinol* 7 781 (1947)
- 114 Kinsell I W Margen S Partridge J W Michaels G D Balch H F and John J P *J Clin Endocrinol and Metabolism* 14 110 (1954)
- 115 Kinsell I W Michaels C D Jr C H and Larsen W F *J Clin Endocrinol* 10 1013 (1948)
- 116 Klinefelter H F Jr Albright F and Griswold G C *J Clin Endocrinol* 3 579 (1943)
- 117 Knowlton A I *Brit N Y Acad Med* 29 441 (1953)
- 118 Koff A K and Tulsky A S *Surg Clin N Amer* 33 3 (1953)
- 119 Landau R I *J Clin Endocrinol and Metabolism* 13 510 (1953)
- 120 Landau R L and Loughhead R *J Clin Endocrinol* 11 1411 (1951)
- 121 Laqueur G I *Science* 112 479 (1950)
- 122 Lerman J Jones H W and Calkins F *Ann Internal Med* 25 677 (1946)
- 123 Leventhal M I and Cohen M R *Am J Obstet Gynecol* 61 1034 (1951)
- 124 Lewis R A and Wilkins I *J Clin Invest* 28 394 (1949)
- 125 Liddle G W Rinfret A P Richard J and Forsham P H *J Clin Endocrinol and Metabolism* 13 842 (1953)
- 126 Locke W Talbot N B Jones H S and Worcester J *J Clin Invest* 30 325 (1951)
- 127 Ludwig A W Boas N F and Soffer I J *Proc Soc Exptl Biol Med* 73 137 (1950)
- 128 Luft R and Olivecrona H *J Neurosurg* 10 301 (1953)
- 129 Man F B Heineman M Johnson C L Leary D C and Peters J P *J Clin Invest* 30 137 (1951)
- 130 Markee J F Everett J W and Sawyer C H *Recent Progr Hormone Research* 7 139 (1952)
- 131 Mason H L and Engstrom W W *Physiol Revs* 30 321 (1950)
- 132 Mason H I and Kepler L J *J Biol Chem* 161 235 (1945)
- 133 Masson P *Rev can biol* 1 570 (1942)
- 134 McCann S M *Am J Physiol* 175 13 (1953)
- 135 McCullagh E P Beck J C and Schaffenburg C A *J Clin Endocrinol and Metabolism* 13 489 (1953)
- 136 McCullagh E P Gold A and McKendry J B R *J Clin Endocrinol* 10 687 (1950)
- 137 McGirr E M and Hutchinson J H *Lancet* II 1117 (1953)
- 138 Means J H *The Thyroid and Its Diseases* 2nd ed Lippincott Philadelphia 1948
- 139 Meyers T H and Man E B *J Lab Clin Med* 57 867 (1951)
- 140 Miller A M and Dorfman R I *Endocrinology* 46 514 (1950)
- 141 Millikan C H and Haines S F *Arch Internal Med* 92 5 (1950)
- 142 Mills E S and Forsey R R *Trans Assoc Am Physicians* 62 70 (1949)
- 143 Mulvancy T H *Am J Ophthalmol* 27 589 (1944)
- 144 Nelson W V and Heller G G *Recent Progr Hormone Research* 3 197 (1948)
- 145 Nelson W O *Inst Record Suppl* 67 36 (1937)
- 146 Nichols J and Gardner I I *J Lab Clin Med* 37 279 (1951)

- 68 Goodwin W E, Bauer F K, Barrett T F and Cassen B *Am J Roentgenol Radium Therapy Nuclear Med* **68** 963 (1952)
- 69 Goodwin W I, Cassen B and Bauer F K *Radiology* **61** 88 (1953)
- 70 Gray C H and Larrott D M *Ciba Colloq Endocrinol* **5** 153 (1953)
- 71 Green J, Isler H, Frenkel M, Rammunga I L and Willebrands A F *J Clin Invest* **31** 57 (1952)
- 72 Greenblatt R B *Postgraduate Med* **9** 492 (1951)
- 73 Greer M A *J Clin Endocrinol* **12** 1259 (1952)
- 74 Greer M A *J Clin Invest* **30** 301 (1951)
- 75 Greer M A *Physiol Revs* **30** 513 (1950)
- 76 Greer M A and Astwood E B *J Clin Endocrinol* **13** 1312 (1953)
- 77 Grynkewich S L, Laughlin R M, Herold F T and Carmel W J *Am J Med Sci* **222** 142 (1951)
- 78 Halmi N S *Endocrinology* **47** 289 (1950)
- 79 Halmi N S *Endocrinology* **50** 140 (1952)
- 80 Hamilton J *Recent Progr Hormone Research* **3** 250 (1948)
- 81 Hamilton J, Soley M H, Reilly W A and Lichorn K B *Am J Diseases Children* **66** 495 (1943)
- 82 Heckel N J *J Clin Endocrinol* **11** 235 (1951)
- 83 Heinbecker P and Pfeifferberger M Jr *Am J Med* **9** 3 (1950)
- 84 Heller E L and Camarata S J *Arch Pathol* **49** 601 (1950)
- 85 Heller C C and Nelson W O *Recent Progr Hormone Research* **3** 229 (1948)
- 86 Heller C G, Nelson W O and Roth A A *J Clin Endocrinol* **3** 573 (1943)
- 87 Heller C G and Nelson W O *J Clin Endocrinol* **8** 345 (1948)
- 88 Hower T F *J Endocrinol* **3** 387 (1942-1944)
- 89 Hills O W, Power M H and Wilder R M *Diabetes* **1** 351 (1952)
- 90 Hollander V P, Di Mauro S and Pearson O H *Endocrinology* **49** 617 (1951)
- 91 Howard R P, Sniffen R C, Simmons F A and Albright F *J Clin Endocrinol* **10** 121 (1950)
- 92 Huggins C and Moulder I V *Cancer Research* **5** 510 (1945)
- 93 Hume D M *Ciba Colloq Endocrinol* **4** 87 (1952)
- 94 Hurvthal L M *Med Clin N Amer* (January 1947)
- 95 Hurvthal L M, Bruns H J and Musulin N *J Clin Endocrinol* **9** 1245 (1949)
- 96 Hurvthal L M and Musulin N *Am J Med* **1** 53 (1946)
- 97 Jailer J W Quoted by Knowlton in reference 117
- 98 Jailer J W *J Clin Endocrinol* **9** 557 (1949)
- 99 Jailer J W and Knowlton H I *J Clin Invest* **29** 1430 (1950)
- 100 Jailer J W *Proc 2nd Clin ACTH Conf* **1** 19 (1951)
- 101 Jailer J W *Trans N Y Acad Sci* [2] **13** 262 (1951)
- 102 Jefferies W McK, Levy R P, Palmer W G, Storaasli J P and Kelly L W *New Engl J Med* **249** 876 (1953)
- 103 Johnson S C *Acta Med Scand* **144** 165 (1952)
- 104 Jones G E S and Delf E *J Am Med Assoc* **146** 1212 (1951)
- 105 Jones G E S, Howard J E and Langford H *Fertility and Sterility* **4** 49 (1953)
- 106 Jost A *Arch anat microscop et morphol expil* **36** 271 (1947)
- 107 Kaftin M *Am J Ophthalmol* **33** 962 (1950)
- 108 Kelley A C, Ely R S and Raile R B *Pediatrics* **12** 541 (1953)

- 183 Shelton G I Moore M C Kapps A and Clark D I *J Clin Endocrinol* **11** 91 (1951)
- 184 Shorr T Carter A C Kennedy B J and Smith R W Jr *Trans Assoc Am Physicians* **66** 114 (1953)
- 185 Silver H K *Pediatrics* **8** 368 (1951)
- 186 Silver S Fieber M H and Yohalin S B *Am J Med* **13** 125 (1952)
- 187 Simpson M I and Evans H M *Endocrinology* **39** 281 (1946)
- 188 Smith P E, and MacDowell I C *Anat Record* **46** 249 (1930)
- 189 Smith P E and MacDowell I C *Anat Record* **50** 85 (1931)
- 190 Smith G V and Smith O W *Physiol Revs* **28** 1 (1948)
- 191 Soffer I J Diseases of the Endocrine Glands Philadelphia 1951 Lea & Febiger Philadelphia
- 192 Soffer I J Lesnick C Sorkin S / Sobotka H H and Jacobs M J *Clin Invest* **23** 51 (1944)
- 193 Sommerville I T In John Hammond Toxemias of Pregnancy Human and Veterinary Blackiston Philadelphia 1950
- 194 Sougin Mibashan R and Jackson W P U *Brit Med J* **2** 311 (1953)
- 195 Specht N W Bauer F K and Adams R M *Am J Med* **14** 766 (1953)
- 196 Sprague R G *Iroc Poy Soc Med* **46** 1010 (1953)
- 197 Sprague R G Mason H I and Power M H *Recent Progr Hormone Research* **6** 315 (1951)
- 198 Stanbury J B Brownell G Riggs D S Perinetti H del Castillo E and Itolz J *J Clin Endocrinol* **12** 191 (1952)
- 199 Stanbury J B and Hedge A N *J Clin Endocrinol* **10** 1471 (1950)
- 200 Stanley M M and Astwood E B *Endocrinology* **41** 66 (1947)
- 201 Stein I F Cohen M R and Elson R *Am J Obstet Gynecol* **58** 267 (1949)
- 202 Stein I F and Ieventhal M I *Am J Obstet Gynecol* **29** 181 (1935)
- 203 Steinback H I Lyon R P Smith D R and Miller F R *Radiology* **69** 167 (1952)
- 204 Sternberg W H Segaloff A and Gaskill C J *J Clin Endocrinol and Metabolism* **13** 139 (1953)
- 205 Stewart I H *Am J Obstet Gynecol* **61** 990 (1951)
- 206 Sturgis S H In Progress in Gynecology Grune & Stratton New York 1950 Vol 2
- 207 Sunderman F W and Sunderman F W Jr *Am J Clin Pathol* **24** 885 1954
- 208 Sydnor K L Kelley V C Paile R B Eh R S and Savers G *Proc Soc Exptl Biol Med* **82** 695 (1953)
- 208a Sydnor K I Savers G Brown H and Tyler T H *J Clin Endocrinol and Metabolism* **13** 891 (1953)
- 209 Szilagyi D F McGraw A B and Smyth N P D *Ann Surg* **136** 555 (1952)
- 209a Taylor A B Albert A and Sprague R G *Endocrinology* **45** 335 (1949)
- 210 Teilum G *J Clin Endocrinol* **9** 301 (1943)
- 211 Tew M B and McAlpins H T *Can Med Assoc J* **63** 287 (1950)
- 212 Thompson R Gross S and Strauss R *Am J Obstet Gynecol* **61** 80 (1951)
- 213 Thorn C W *J Clin Endocrinol and Metabolism* **13** 614 (1953)
- 214 Thorn G W Forshum P H and Emerson K The Diagnosis and Treatment of Adrenal Insufficiency 2nd ed Charles C Thomas Springfield Ill 1951
- 215 Thorn G W Jenkins D Arons W L and Frawley T F *J Clin Endocrinol and Metabolism* **13** 957 (1953)

- 147 Page I W The Hypertensive Disorders of Pregnancy Charles C Thomas
Springfield Ill 1953
- 148 Parvonen S Soiva K and Vartiainen S *Acta Obstet Gynecol Scand* 29 5
(1950)
- 149 Paschkis K I *Med Clin Amer* 36 1711 (1952)
- 150 Paschkis K I Rupp J and Schneider J J *Trans Studies College Physi-
cians Phila* 19 149 (1952)
- 151 Paschkis K I, and Cantarow A *Ann Internal Med* 34 669 (1951)
- 152 Paschkis, K I and Rupp J Unpublished observation
- 153 Paschkis K I Rakoff A E and Cantarow A *Clinical Endocrinology*
Hoebner New York 1953
- 154 Pasquallini R Q *J Clin Endocrinol and Metabolism* 13 178 (1953)
- 155 Pearlman W H Pearlman M R J and Rakoff A I *Am J Obstet Gynecol*
66 370 (1953)
- 156 Perkoff C T Sandberg A A Nelson D H and Tyler F H *Arch Internat
Med* 93 1 (1954)
- 157 Perry W F and Hughes J F *J Clin Invest* 31 457 (1952)
- 158 Peters J P Mann L P and Heineman M *In The Normal and Pathological
Physiology of Pregnancy* Williams & Wilkins Baltimore 1948
- 159 Pickering D F and Miller F R *Am J Diseases Children* 85 135 (1953)
- 160 Pigeaud H and Barthiault R *Gynecol et obstet* 50 341 (1951)
- 161 Pincus I J Personal communication
- 162 Plate W P *Acta Endocrinol* 8 17 (1951)
- 163 Plotz C M Knowlton A I and Ragin C *Am J Med* 13 597 (1952)
- 164 Purves H D and Griesbach W I *Brit J Exptl Pathol* 30 23 (1949)
- 165 Purves H D and Griesbach W I *Endocrinology* 49 244 (1951)
- 166 Purves H D and Griesbach W I *Endocrinology* 49 427 (1951)
- 167 Quinn I J and Worcester R I *J Clin Endocrinol* 11 1561 (1951)
- 168 Raben M S and Astwood J B *J Clin Invest* 28 1317 (1949)
- 169 Raben M S and Westermeyer V W *Proc Soc Exptl Biol Med* 80 83
(1952)
- 170 Rakoff A E *Fertility and Sterility* 4 263 (1953)
- 171 Rakoff A E *Med Clin Amer* 32 1509 (1948)
- 172 Rapport R I Curtis C M and Simcox S J *J Clin Endocrinol* 11 1549
(1951)
- 173 Reddy Wm J Jenkins D and Thorn S W *Metabolism* 1 511 (1952)
- 174 Reynold S R M *Physiology of the Uterus* 2nd ed Hoeber New York
1949
- 175 Reynolds I Corrigan K F and Hayden H S *Harper Hosp Bull* 8 4
(1950)
- 176 Robbins J Rall J L Trunell J B and Rawson R W *J Clin Endocrinol*
11 1116 (1951)
- 177 Salter W T *Science* 109 453 (1949)
- 178 Sandberg A A Nelson D H Glenn F M Tyler F H and Simuels I T
J Clin Endocrinol and Metabolism 13 1445 (1953)
- 179 Schneider J J Unpublished
- 180 Schwarzer K *Ergebn innere Med u Kinderheilk* 58 55 (1949)
- 181 Severinghaus A F *Research Publ Assoc Research Nervous Mental Dis*
17 69 (1938)
- 182 Sheehan H I Summers V K and Nichols J *Lancet* 266 312 (1953)

Author Index

Numbers in parentheses are reference numbers and are included to assist in locating references when the authors' names are not mentioned in the text. Numbers in italics indicate the page on which the reference is listed at the end of the chapter.

A

- Abbott W E 694 (1119) 696 (1119) 815
 Abderhalden, L 475 (1) 499
 Abdon N O 129 (1) 144
 Abel J J 190 185 425
 Abelin I 436 (9) 445 (2) 474 (6) 476 (3 4 7 8) 453 (5) 499 500
 Abelman M A 700 (218 219) 790
 Abercrombie W F 459 (10) 500
 Åberg B 11 (2 3 5) 12 (3) 13 (4) 14 (2 4) 15 (3 4) 16 25 (5) 48
 Abernathy R 779 (1 2) 784
 Aboud I G 377 383
 Abahams V C 425
 Abramowitz A A 100 (2) 144
 Abrams G D 258 (1) 259 (1) 296
 Abus M E 271 379
 Aburas W B 713 (5) 785
 Acevedo D 172 (1) 677
 Acher P 475 (75) 404 414 (12) 425 427
 Ackermann I G 727 (653) 803
 Adams A 231 230
 Adams C H 696 (350) 766 (350) 794
 Adams E 287 (2) 293 (2) 296 743 (4) 785
 Adama J E 718 (78) 787
 Adams R M 80 (195) 875
 Additt F T 23 (264) 54
 Adezati L 672 (51) 679
 Adler M 74 (25) 785
 Adlersberg D 721 (103) 725 (103) 732 (5-9) 785 787
 Adolph E F 425
 Adrouny G A 25 (3) 296
 Aebi H 477 (11) 500
 Agnello E J 556 (201) 584
 Agnoletto A 482 (12) 500
 Agnoletto C 482 (12) 500
 Agolini C 672 (2) 674 (2) 677
 Agosin M 751 (10) 785
 Agrell I 75 (1-3) 85
 Ahlmark A 670 (3) 677
 Ahlquist R I 154 144
 Aikawa J K 706 (11) 785
 Ainslie J D 779 (12) 785
 Aird P B 713 (13) 785
 Albaum H G 724 (14) 785
 Albert A 331 (9-12) 342 (243) 347 379 388 439 (541) 440 (14) 445 (18) 449 (21 541) 450 (19) 451 (15 16 17 378) 452 (16 17 61 325) 453 (61) 454 (15 378 338) 455 (18 543) 490 (446) 500 501 507 510 512 572 (226) 574 (226) 584 705 (1080) 707 (737 1080) 708 (1080) 709 (1080) 721 (1080) 724 (1080) 732 (1080) 753 (1080) 765 (1080) 782 (15) 784 (371) 785 795 805 814 843 (209a) 869 (11) 870 875
 Albert S 556 800
 Albright E C 450 (144 381) 484 (380) 503 508
 Albright F 154 155 160 168 170 718 (59) 783 (59) 786 844 850 (14) 857 862 (91) 865 (116) 870 872 873
 Albright F J 703 (339) 794
 Aldrich T B 401 (116) 403 (116) 428
 Alexander C R 779 (708) 804
 Alexander J A 707 (345) 721 (343) 753 (343) 794
 Alexandrowicz J S 61 (4-7) 67 (7) 83 109 110 139 (b) 144
 Alibrandi A 488 (22) 500
 Alimnosa L M 552 (215) 554 (47) 563 (47) 579 586

- 216 Tobian I, Jr *J Clin Endocrinol* 9 319 (1949)
- 217 Topper A *Am J Diseases Children* 81 233 (1951)
- 218 Turner, H H *In* S Soeldin *Progress in Clinical Endocrinology* Crane & Stratton New York 1950
- 219 Twombly G A *Surgery* 16 181 (1944)
- 220 Venning E H *In* *The Normal and Pathological Physiology of Pregnancy* Williams & Wilkins Baltimore 1948
- 221 Venning E H Weil P C and Browne J S L *J Biol Chem* 128 cvii (1939)
- 222 Venning I H *Rev can biol* 1 571 (1942)
- 223 Vestergaard I *Acta Endocrinol* 13 241 (1953)
- 224 Vilanova, V and Canadell J M *J Clin Endocrinol* 9 883 (1949)
- 225 Wallach J B and Scharfman W B *J Am Med Assoc* 148 729 (1952)
- 226 Warming Larsen A Hamburger C and Sprechler, M *Acta Endocrinol* 11 400 (1952)
- 227 Watson E M and Pearce R H *Am J Clin Pathol* 19 442 (1949)
- 228 Weissman I *J Clin Endocrinol and Metabolism* 12 1223 (1952)
- 229 Wells L J *Proc Soc Exptl Biol Med* 63 417 (1947)
- 230 Werner S C *J Clin Endocrinol* 13 1332 (1953)
- 231 White F P and Sutton L F *J Clin Endocrinol* 11 1395 (1951)
- 232 Wilkins L *The Diagnosis and Treatment of Endocrine Disorders in Childhood and Adolescence* Charles C Thomas Springfield Ill 1950
- 233 Wilkins I Crigler J F Jr Silverman S H Gardner I I and Migeon C J *J Clin Endocrinol and Metabolism* 12 277 (1952)
- 234 Wilkins I Fleischman W and Howard J E *Endocrinology* 26, 385 (1940)
- 235 Wilkins I Crigler J F Jr Silverman S H Gardner I I, and Migeon C J *J Clin Endocrinol and Metabolism* 12 1015 (1952)
- 236 Wilkins L *J Pediat* 41 860 (1952)
- 237 Wilson H *J Clin Endocrinol and Metabolism* 13 1465 (1953)
- 238 Wislocki G B Dempsey L and Fawcett D W *In* *The Normal and Pathological Physiology of Pregnancy* Williams & Wilkins Baltimore 1948
- 239 Wolfson W Q *Arch Internal Med* 92 108 (1953)
- 240 Zierler K I *Bull Johns Hopkins Hosp* 89 203 (1951)

- Ashley M M 405 (311) 810
 Ashmore J 575 (7) 578 621 655
 Ashton A 767 (30) 785
 Ashworth R B 40 (243) 54
 Askonas B A 480 (25) 500
 Asling C W 237 (8 405) 239 (??) 244
 (405) 250 (305) 297 307 436 (??4)
 437 (?? 560 660) 600 606 515 515
 Asper S P Jr 486 (50) 457 (30) 490
 (30) 500
 Assenmacher I 319 327 328 331 329
 Astwood F B 203 (79) 221 222 230
 231 233 252 (9) 253 (12) 256 (9)
 256 (10) 279 (12 315) 286 (280)
 293 (10 11 250) 294 (9) 295 (10)
 281 288) 297 304 305 360 371 378
 3 9 4?? (12) 408 440 (37) 441 (32)
 442 (33) 445 (35) 460 (148) 461 (31
 36 254) 462 (32 34 38) 464 (32 35)
 465 (35 686 687) 466 (35 36) 468
 (35) 500 503 506 516 822 (8) 820
 (3 168 200) 830 (9) 838 (6 7) 870
 872 874 875
 Aterman K 760 (31) 781 (32) 785
 Atkinson W B 359 3 9 670 (10) 678
 Atna A 488 (127) 503
 Aub J C 154 (1) 167 (5) 169 170
 459 (590) 487 (39) 500 515
 Audus L J 2 15 (12) 31 (12) 40 (11)
 43
 Auger H O 830 (10) 870
 Augustinsson K B 119 (12) 144
 Austin P R 218 (28) 231
 Austing M F 460 (85) 502
 Averill S C 372 379
 Avery G S Jr 42 (??) 48
 Awapara J 673 (11) 678 728 (33) 785
 Avelra B J 290 (13) 292 (13) 297
 666 (12) 678
 Axelrod L R 625 655 667 (13) 678
 Axelrod Heller D 436 (74) 506
 Ayer D F 221 (57) 222 (5 9 87) 226
 (9 81) 231 233 286 (20) 297
 Ayres G B 254 302
 Azimov G I 345 3 9
- B**
- Babcock J C 556 (100) 581
 Babcock V I 779 (1061) 813
 Babel J 154 (103) 172
 Bacchus H 741 (31 37 38 39) 765 (35
 36) 785 786
 Bachmann W L 540 (8 10 12) 542 (9
 11) 552 (8) 554 (8 11) 556 (8 11)
 559 (9 11 12) 578
 Backus G L 20 (260) 54
 Baer Z M 58 (24) 59 (24) 78 (25 27
 28) 79 (24 26) 81 (24) 84 101 (13)
 135 (14) 144
 Badenoch J 732 (41) 786
 Bader M F 09 (693) 711 (693) 804
 Badinez O 751 (10) 785
 Badrick F I 288 (507) 289 (707) 305
 Baeder D H 761 (991) 764 (998 991)
 812
 Baggenstoss A H 766 (1018) 812
 Bagnall A W 749 (42) 786
 Bahn R C 337 (9-12) 359 860 (11)
 860
 Bailey C C 189 (4) 195
 Bailey R I 314 3 9
 Baillarge M 9 (137a) 10 (137b 137d)
 12 (137a-f) 14 (137e f) 51
 Bainborough A R 456 (40) 500
 Baker B L 241 (14) 258 (1) 259 (1)
 260 27 346 381 671 (14 119) 678
 680 719 (138) 744 752 (563) 754
 (48 137) 755 (564) 762 (483) 763
 (482 483) 766 (47) 786 788 798 800
 Baker C 712 (1124) 714 (1126) 815
 Baker A 188 195
 Baker R 156 170
 Baker R S 28 (91 93) 23 (93) 43
 (92) 50
 Balant C P 547 (13) 578
 Balch H E 271 (206) 273 (205) 30
 327 (114) 8 3
 Baldin J T 665 (271) 671 (271) 684
 742 (1218) 819
 Baldridge G D 447 (49) 748 (49) 760
 (49) 778 (643) 773 (643) 786 802
 Balesdent Marquet M I 66 (23) 84
 Balfour W 106 (15) 144
 Ball C D 42 (213a) 53
 Ball E G 185 (23) 195
 Ball S M 161 174
 Ballabio C B 709 (954) 711 (954) 811
 Balmann J H 3 3 (15 16 17) 354 355
 356 357 378 379 704 (50) 734 (51)
 780

- Allcroft W M 695 (1600) 814
 Allgra G 670 (4) 677
 Allegretti N 496 (23) 500
 Allen B M 491 (24) 500
 Allen H C Jr 825 (3) 870
 Allen M J 692 (506-507) 793
 Allen W M 604 664
 Allende M F 740 (314) 793
 Almeida C R M de 18 (7) 48
 Alpert M 290 (4) 297 671 (5) 677 725
 (1255 1256) 742 (19) 785 819
 Alrich L M 766 (16 747) 769 (17) 785
 805
 Althausen T L 484 (25) 500
 Altman A D 345 379
 Altmann G 70 (8 9) 77 (8) 83
 Altschul A M 647 (100) 680
 Altszuler N 741 (37 29) 785 786
 Amar R 60 (10 11) 85
 Ambrus C 479 (26) 500
 Ambrus J 479 (26) 500
 Amelung D 647 655
 Amendolla C 500 (311) 552 (312) 587
 Ames R G 396 (7 203) 495 451
 Amin A H 100 (4) 104 (8.1) 105 137
 (8) 144
 Amoroso I C 169 (51) 171
 Anromin G D 763 (1131) 816
 Ancowitz A 243 (210, 257 280) 261
 (210) 265 (50) 299 303
 Anderson A B 474 (27) 500
 Anderson E 192 (2) 195 256 (178) 258
 (5) 265 (147 178) 297 301 374 389
 455 (477) 511 703 (940) 810
 Anderson L M 250 (258) 304
 Anderson G F 763 (21) 785
 Anderson G W 410 (10) 495
 Anderson H L Jr 195 (32) 196
 Anderson H V 642 (105) 658
 Anderson J A 713 (748) 805
 Anderson W 695 (1030) 814
 Andersson B 400 417 (9) 495
 Andreai S R 24 (9) 48
 Andreie W A 24 (8 9) 29 (8) 30 (8)
 43 741 (22) 785
 Andrewartha H G 74 (12) 83
 Andrews F M 765 (460) 791
 Andrews F N 318 (1a) 379
 Andrews H L 437 (87) 50
 Andrus M 610 (6) 678
 August A A 700 (8.0) 808
 Anker H S 578 (1) 578
 Anliker R 554 (141 144) 556 (144, 150)
 558 (145) 582
 Anner G 537 539 (339) 541 (3 4) 548
 588
 Anselmino H J 251 (6 170) 257 (6
 170) 260 (170) 297 301
 Anslow W I Jr 395 495
 Anthony A 673 (7) 678
 Antonchuk N 278 (127 128) 281 (121)
 300 665 (94) 680 698 (376 377) 701
 (378) 794
 Antonucci R 549 (5) 550 (5 6) 552 (6)
 553 (5 6) 554 (5) 555 (5) 560 (6) 578
 Antopol W 609 (399) 701 (23) 747 (556)
 785 795 809
 Aptigot S 597 (57) 606
 Aposhian H V 618 648 (108) 659
 Appel S B 696 (24) 714 (24) 745 (24)
 785
 Apter N S 718 (246) 791
 Arigoni D 536 (61 2641) 578 585
 Arndt U W 181 (3) 195
 Arner B 848 (4) 810
 Arnstein I H 718 (78) 787
 Aron C 669 (8 9) 678
 Aron H 748 (905) 801
 Aronoff S 17 (314) 56
 Arons W L 688 (1151) 696 (1151) 697
 (1151) 701 (1151) 721 (1151) 727
 (1151) 745 (1148) 765 (1151) 766
 (1151) 768 (1151) 777 (1151) 810
 842 (215) 875
 Aronson S M 731 (25) 749 (1072) 785
 812
 Arreguin B 592 (20) 655
 Arrillaga F C 871
 Arroyave G 655
 Arth G E 539 (258 290) 586
 Artress J I 674 (203) 682
 Arty I 58 (13) 65 (14 15-17 18 19 20
 21 22 23 63 64) 83 150 (9 10) 144
 Ashoe Hansen G 743 (26 27 29 29a)
 753 763 (29) 785 795 834 (2) 870
 Asdell S A 332 (2) 379
 Asero B 78 (203 204) 87 104 (11) 144
 146
 Ash I 221 (57) 237
 Ashenbrucker H 748 (463) 797

- Baylis R I S 288 (19) 292 (19) 293
 (19) 294 (19) 297
 Bazeley P I 206 (76 77) 253
 Bazemore A W 291 (87) 292 (5 9 87)
 231 235 286 (20) 297
 Beach F A 673 (17 18 126) 6 8 680
 Beach R G 22 (14) 48
 Beall D 543 (120) 591 604 58* 655
 Beare J I 489 (57) 501
 Beary D F 29 (572) 800
 Baton G H 489 (57) 501
 Beaton J P 489 (57) 501
 Bechgaard P 479 (58) 501
 Beck E M 737 (968) 797
 Beck J C 867 (135) 873
 Beck I A 751 (211) 752 (67 63) 786
 791
 Becker D 451 (51) 451 (512) 512
 Becker D A 456 (59) 501
 Becett S 494 (313) 508
 Becks H 293 (72) 210 (362) 241 (71
 294 361) 432 (79) 227 307 500
 Beer C T 609 (97) 610 (99) 611 (99)
 616 (99) 633 (99) 635 (99) 636 (99)
 657
 Beevers H 39 (88) 40 (15) 44 (88) 48
 50
 Begany A J 764 (988 990) 771 (990)
 113 (990) 812
 Behr W T 674 (19) 678
 Behrens M 2 (16) 3 (76) 49
 Behrens O K 190 (135) 191 (135 136)
 198
 Beierwaltes W H 452 (329) 507 781
 (61 1295) 786 835 (15) 870
 Bekaert J 448 (145) 503
 Beldes E J 490 (80) 501
 Belehradsk J 426
 Belisle J 549 (295) 586
 Belko J S 749 (610) 803
 Bell D M 187 (6) 195
 Bell F T 197 (86) 197
 Bell G I 699 (65) 786
 Bell G M 497 (304) 507
 Bell J H 740 (1100) 814
 Bell P H 2 0
 Bell P E 699 (65) 786
 Bellamy A W 497 (256) 506
 Bellanca J 723 (117) 788
 Belleau B 519 (21) 579
 Ben M 708 (1123) 712 (1124) 714
 (1123 1126) 815
 Benacerraff B 757 (460) 772 (416) 796
 797
 Benati O 100 (90 91) 146
 Benbow J T 498
 Benditt I P 674 (77) 683 763 (66) 786
 Benedict L M 447 (270) 473 (270)
 496 (417) 507 509
 Benedict F G 490 (60) 501
 Benedict J D 732 (68) 786
 Benedict W H 767 (67) 786
 Benfey B G 137 (19) 144 473 496
 Bengtson L P 669 (71) 678
 Benirschke K 610 (14) 615 (14) 655
 Benjamin F B 754 (69) 779 (69) 786
 Bennet Clark T A 2 3 (17 18) 6 (17
 18) 7 (17) 45 (16) 46 48
 Bennett E I 26 (19) 48
 Bennett H S 690 786
 Bennett J H 770 (314) 793
 Bennett L L 175 (7) 195 238 (139)
 243 (54) 247 (139) 248 (74) 249 (137)
 251 (23 137 375 413) 253 (24) 271
 (75) 290 (137) 297 299 300 301 306
 303 305 (223) 431 607 (74 341) 705
 (73) 707 (73) 708 718 (341) 732
 (341) 733 (74 341) 735 (75) 736
 (75) 786 794 818
 Bennett M F 61 (520) 62 (577) 66 (73)
 85 94
 Bennett R 413 (729) 517
 Bennett V 443 (779) 517
 Bennett W A 337 (9-12) 379 691 (76
 77) 751 (511) 771 (910 917) 869
 (11) 787 799 809 870
 Benoit J 319 327 328 331 379
 Bensley S H 164 (12) 170
 Benson J A 756 (470 421 472) 796
 Bentinck R C 705 (73) 707 (73) 718
 (78 414) 720 (414) 786 787 796
 851 (16) 870
 Bentley J A 6 (21) 9 (20) 15 (21) 48
 Benua R S 452 (61) 453 (61) 501
 Berdisco G 319 (295) 431
 Berengo A 674 (22) 678
 Berg J L 715 (647) 716 (647) 809
 Berg O 435 501
 Bergen J R 719 (518) 799

- Balomey A A 395 (11) 495
 Bancroft R W 187 (163) 190
 Bandurski R S 35 (31) 39 (32), 40 (32)
 41 (31) 44 (32) 45 (32) 48
 Banerjee D K 537 (172) 583
 Banerjee S 740 (52) 53 54) 786
 Bines D 542 543 (15) 550 (14) 578 582
 Bingham A D 762 (55) 766 (55) 786
 Bannister B 539 (173) 583
 Bannister J 100 (16) 144
 Barber A 768 (56), 786
 Barberio J R 775 (57) 786
 Barbieri E 485 (521) 512
 Barclay W R 761 (278) 763 (278) 792
 Barger G 474 (283) 506
 Bargmann W 105 109 144 391 496
 Bargon N 676 (15) 678
 Barnaby M 496 (41) 500
 Barker M H 462 (43) 464 (42) 500
 Barker S B 187 (123) 198 452 (402)
 474 (49) 475 (45) 356) 476 (45) 46
 47 49 401) 490 (50) 51) 491 (51) 500
 501 508 509, 782 (458) 797
 Barker-Jorgensen C 397 338 399 (15)
 4 6
 Barkley L B 539 579
 Barlow G 698 (1120) 708 (1122 1123)
 712 (1124) 714 (1122 1193 1126) 727
 (1125) 815
 Barlow H W B 2 (114) 23 (114) 50
 Barnabé L 496 477
 Barnes A R 765 (497) 798
 Barnes C S 540 (17) 562 (17) 569 (17)
 579
 Barnes J M 128 144
 Barnett H L 154 (97) 172 851 (12) 870
 Barnicot A 158 164 170
 Barott G W 705 (177) 707 (177) 721
 (177) 732 (177) 753 (177) 789
 Barrett G R 332 (172) 354
 Barrett H W 254 (15) 297 467 (52) 501
 Barrett T F 830 (68) 872
 Barnett R J 315 324 329 331 445
 579 581 583
 Barron E S Guzman 729 (639) 802
 Barry A G 270 (78) 233
 Barry M C 609 616 (100) 618 (100)
 621 (100) 655 657
 Barry S R 442 (164) 504
 Barsantini J C 361 379 480 (140) 503
 Bartels, L C 827 (13) 870
 Barthel M 497 (359) 608
 Barthmaalt R 863 (160) 874
 Bartlett M V 479 (357) 480 (357) 608
 Bartlett I D 243 (122) 248 (16 1)
 122) 249 (16) 297 300
 Bartlett R G 674 (203) 682
 Barton A D 729 (58) 786
 Barton D H R 523 521 (18 19) 525
 529 (217) 540 (17) 544 (20) 567 (17)
 567 (20) 569 (17) 579 584
 Burton M N 739 (806) 807
 Bartter F C 153 154 (11) 170 707
 (343) 718 (59) 721 (743) 753 (343)
 783 (59) 786 794 850 (14) 870
 Ba chiere L 763 (60) 766
 Baserga R 751 (604) 753 (604) 788
 Bass A C 706 (705) 708 (705) 710
 (205) 790
 Bass A D 249 (90) 299
 Bassett G 747 (181) 789
 Bastien P A 751 (654) 803
 Bastian J W 331 (766) 587
 Bateman J H 703 (1232) 719 (1252) 818
 Bates R W 211 231 246 (312) 297 (18)
 297 305 373 374 375 385 667 (16)
 678 703 (940) 810
 Bather R 497 (53) 501
 Battle F F 493 (424) 509
 Batres E 552 (270) 556 (69) 580 586
 Batts A A 251 (375) 306
 Bauchau A G 63 (31) 66 (30) 84
 Bauer F K 830 (68 69 195) 872 875
 Bauer W 151 (1) 169 487 (39) 500 715
 (153) 789
 Bauld W S 509 (124) 606 (121) 631
 (121) 658 667 (163) 681
 Bauman K L 237 (326) 305
 Baumann C A 187 (134) 198 488 (350)
 389 (173) 504 507
 Baumann E J 436 (54 55 56) 461
 (432) 501 510 781 (61) 786
 Baxter B 447 (270) 473 (250) 506
 Bay R 759 (108) 787
 Bayle J 718 (1158) 817
 Bayles T B 747 (319) 749 (1174) 793
 817
 Bayliss M J 758 (322) 289 (321 322)
 305 623 (260 760a) 662

- Blakeslee A F 27 (40) 35 (215) 54
 Blalock A 715 (89) 787
 Blanch B 751 (211) 791
 Bland M U 731 (468) 797
 Blanquet I 438 (69 70) 601
 Blaschko H 58 (33 34) 78 (35) 84 107
 107 144
 Blatt O 460 (465) 510
 Blattacharya P 665 (270) 611 (270) 685
 Bleifer D 293 (114) 294 (114) 300
 Bliss C I 166 167 (18) 170
 Bliss D E 60 (36 39-41) 63 (36 37
 41) 66 (36-40) 84 98 (26 27) 109
 (26 27) 113 (27) 138 (26) 139 (26
 27) 140 (21 25) 144
 Bliss E I 256 (32) 258 (31) 291 (31)
 297 694 (1119) 696 (1119) 815
 Bloch E 610 611 (74) 612 (15) 613 615
 655 657
 Bloch K 188 (8) 195 475 (86) 50 578
 (1 33) 578 579 591 (16 18 174) 592
 (17 160 161 326) 593 (226) 602 (16)
 655 659 670 674
 Block M M 716 (1263) 819
 Block R J 416 431
 Blodinger J 410 (10) 425
 Blom P S 830 (19) 870
 Blondeau R 15 (223) 55
 Blood J P 705 (177) 707 (177) 721
 (177) 732 (177) 753 (177) 769
 Bloodworth J M B 710 (90) 787
 Bloom B 716 (92) 732 (92) 787
 Bloom B M 59 (173) 585
 Bloom F 743 (91) 752 (91) 787
 Bloom F C 156 (70) 171
 Bloom W L 777 (766) 806
 Blum T 445 (71) 601
 Blumenthal H T 246 (181) 249 (406)
 301 307
 Blumgart H I 496 (72) 501
 Blunt J W 766 (537 669 680 681 888)
 767 (93 537) 779 (761) 787 799 808
 805 809
 Bo W J 670 (31) 678
 Boak R A 777 (1180) 817
 Boas F 28 (25) 48
 Boas N F 290 (193) 302 742 (583) 801
 833 (127) 873
 Boatman J B 439 (73) 501
 Bobin G 50 (42 43) 84
 Bocchi A 22 (26 183) 43 (26) 48 50
 Bocklage B C 576 (34) 579 598 (19
 215) 655 661
 Boda J M 344 380
 Bode N R 26 (27) 48
 Bodenchuk J M 607 (98) 680
 Bodenheimer F S 71 (41) 84
 Bodenstein D 71 (50 52) 72 (46 47 49)
 73 (46) 74 (46 48) 75 (53) 77 (51) 84
 Bolian D 95 (28) 109 144
 Boehme C C 489 (98) 502
 Bogart R 674 (47) 678
 Bogdanove I M 495 (22) 506 553
 (49) 797
 Bogen I 777 (1159) 817
 Bogogna G 164 (132) 173
 Bogoraze D 78 (124) 79 (54 55 122
 123 124) 84 86
 Bogoroch R 460 (24) 501
 Bohnel F 816
 Borsson C J 68 (56) 84
 Boland I W 696 (95) 718 (94) 766
 (95) 787
 Bolene C 489 (75) 501
 Boling I 725 (636) 753 (635) 802
 Bollman J I 153 (73) 154 171 485
 (200) 504
 Bologna U 489 (16) 501
 Bolony F 164 (54) 171
 Bolton W 674 (30) 678
 Bondy P K 239 (33) 253 (33) 297 452
 (78) 481 (77) 501 753 (96) 779 (97)
 736 (99) 782 (98) 787 845 (66) 871
 Bone A D 102 (284) 150
 Bonelli M 164 170 172
 Bongiovanni A M 784 (100) 787 844
 (21) 850 (20) 870
 Bonhomme P 440 (507) 511
 Boniface J 741 (285) 792
 Bonner C D 669 (85) 679 693 (101)
 701 (534) 787 799
 Bonner D 36 (135) 51
 Bonner J 3 5 (332) 15 (191 192) 26
 (19 33 101) 28 (93 285 3 3) 29
 (93 285) 30 (31) 31 (29 33 181) 36
 (125) 37 (53) 39 40 (201 331) 41
 (31 35) 43 (31) 44 (32) 45 (32) 46
 (28) 48 50 51 52 53 55 56 57 (20)
 655
 Bonner W D Jr 15 (297) 24 (298) 25
 (297) 39 (296 297) 43 (296 297a)
 55

- Bergenstal D M, 696 (542 543) 697
 (542) 718 (216) 750 (543 545) 752
 (545) 791 799 800
 Berger J 42 (22) 48
 Berger M, 597 655 656
 Berger S 192 (36 34) 196 261 (112) 300
 Bergh A 276 (43) 292
 Bergman A J 362 379
 Bergmann F 121 (20 311 312) 144 161
 Bergmann W 535, 536 587
 Bergmeyer H U 676 (62) 679
 Bergstroem S 534 (22) 579
 Birksen J 439 (339) 440 (339) 485
 (700) 490 (81) 501 504 508
 Berlin B S 776 (79) 779 (708) 787, 804
 Berlin J 552 (271) 568 (269) 569 (269)
 586
 Berliner R W 155 170
 Berman J 725 (674) 803
 Berman R A 667 (247) 683
 Bern H A 673 (23) 678
 Bernard B de 41 (23a 337a) 48 50
 Bernheim F 716 (406) 797
 Bernini G 397 496
 Bernstein D F 693 (80 81) 787
 Bernstein S 522 549 (5) 550 (5 6) 552
 (6) 553 (5 6) 551 (5) 558 (5) 559
 (24) 560 (6) 578 579
 Bernston H S 708 (94) 233
 Berrut R 571 (324) 587
 Berry C M 321 (71) 381
 Berry J W 763 (397) 778 (396) 795
 Berson S A 781 (82) 787 879 (71) 870
 Berthrong M 771 (910) 809
 Bertrand I 494 (63) 501
 Bessesen D H 499 (64) 501
 Best C H 238 (26 212 324 325) 239
 (212 325) 251 (24) 254 (15 77) 297
 302 305 712 (1032) 755 (902) 812
 813
 Best W R 744 (84) 747 (83) 787
 Bethel J J 489 (173) 504
 Bevans M 486 (692) 516
 Beyer C A 670 (24 25) 678
 Beyer R F 539 (288 290) 586
 Beythien A 47 50
 Bezzer A E 762 (603) 801
 Bhaskar S N 179 (14 15) 170
 Bhattacharya P 341 (127) 382
 Biale J B 40 (201) 53
 Biasotti A 324 585 496
 Biwatt F 366 379
 Bible S W 703 (85) 787
 Bickle A S 6 (21) 15 (21) 48
 Bide A L 552 (25 26) 519
 Bidmead D S 721 (1187) 817
 Bieller H V 441 (65) 501
 Bielschowsky F 163 (99) 172 498 (66
 67) 501
 Bien I J 740 (85a) 787
 Bieri J G 235 (158) 270 (158) 301
 Bieri A 167 (16 17) 170
 Biggers J D 494 (68) 501 668 (96 27
 28) 678
 Bijvoet J M 533 (27) 5 9
 Billensten D C 60 (32) 84
 Biller H 427
 Billingham R I 75 (85) 78
 Billows J A 759 (1197) 817
 Binhammer R T 287 (150) 301
 Biozzi G 772 (416) 796
 Birch A J 557 543 579
 Birke G 611 633 (12 176) 675 (174)
 655 634 (35) 795
 Birmingham M H 289 (78) 297
 Birnie J H 394 (74) 496 477 710 (105)
 712 (375) 787 795
 Bischoff F 600 656
 Bishop M A 272 (8) 281
 Bishop I M F 744 (753) 790
 Biskind C R 691 (400) 795
 Bissegger A 128 151
 Bissell A 442 (746) 445 (35) 462 (35)
 464 (35) 465 (35) 466 (35 746) 468
 (35) 500 517
 Bisso A 745 (77) 787
 Bissonnette T H 317 319 (27) 3 8
 Bitancourt A A 56 (74) 48
 Bitman J 667 (79) 678
 Bittner J J 751 (731) 805
 Bjering T 394 (21) 426
 Bjørneboe M 774 (88 325) 787 793
 Black H M 673 (97) 680
 Black K O 260 (27) 212 (29) 97
 Black P T 752 (30) 297
 Blackburn C M 490 (446) 510
 Blackman S S Jr 851 (18) 8 0
 Blackmore K F 393 (99) 498
 Black Schaffer B 714 (8 0) 808
 Bladon I 553 (30) 554 (31 32) 556 (31
 32) 579
 Blanch W 463 (775) 505

- Britten L J 17 (43) 49
 Brobeck J R 281 (131) 283 (131) 241
 719 300 303 792 (740) 747 (741)
 805 814 (76) 870
 Brock H 341 (31) 342 (31) 3 9
 Brodie L C 727 (111 1099) 733 (104)
 87 814
 Brodtkin E 106 145
 Brody S 416 (625) 488 (628) 514 718
 (119) 788
 Broh Kuhn R H 185 195 451 (464)
 510 777 (113) 782 (113) 788
 Brohn S 163 1 0
 Brooks P V 529 (36) 563 (36) 579 603
 655
 Brooks V B 117 (32) 145
 Brophy D 490 (99) 507
 Brougher J C 166 170
 Brown C Y 697 (27 525) 766 (527
 5 8) 799
 Brown D H 243 (275) 260 (2 8) 304
 Brown D M 234 (92) 233 293 (135) 00
 Brown F F 451 (93) 507
 Brown I M 697 (577 595) 166 (597
 993) 799
 Brown F A Jr 58 (66 67 68 10) 60
 (10) 61 (65 66 70 75 84 88 90)
 595 61 (69 72 76 77 78 19 80 81
 82 83 84 85 86 87 89 422 423 4 4
 591 593 594) 66 (71 73) 67 (70)
 78 (74) 85 92 94 116 (35) 135 (33
 34) 145
 Brown F C 435 (666) 515
 Brown G C 719 (12) 785
 Brown G L 117 (36 37) 133 145
 Brown H 126 (115) 788 843 (708A) 875
 Brown J B 4 (45) 49 345 383
 Brown J H U 691 (114) 788 845 (27
 25) 870
 Brown J W 17 (119 324) 87 56
 Brown M 496
 Brown M B 674 (125) 680 757 (605)
 801
 Brown R G Jr 195 (195) 148
 Brown R W 315 316 (32 33) 380
 Brown W E 342 371 (99) 3 9 380 671
 (37) 678
 Browne J S L 712 (55) 273 (55) 230
 (55) 298 605 606 637 (143) 649
 (143) 659 663 741 (22) 784 (1175)
 785 817 850 (291) 876
 Brownell C 598 (198) 837 (198) 838
 (198) 875
 Brownell G I 449 (684) 516
 Browne A C 645 (24) 646 655
 Brownlee G 765 (116) 788
 Bruce H M 290 291 295 (41) 298 660
 (39) 6 8
 Brucke H V 101 145
 Bruckmann G 189 (17) 195
 Bruger A 488 (643) 515
 Bruger M 441 (333) 507 717 733 (835)
 807
 Brumby I J 366 380
 Brumfield L T 24 (46) 49
 Brummel L 288 (301) 289 (307) 305
 616 (40) 678
 Bruner J A 337 (35) 380
 Bruning K J 552 (900) 556 (901) 584
 Brunn F 396 496
 Brunner M I 642 (53 105) 656 658
 Bruns H J 567 (95) 872
 Bryan W H 42 (47) 44 (47) 49
 Bryans F I 0 380
 Buchanan G D 165 (140) 173
 Bucher A L I 592 655
 Buchholz C 496 (753) 517
 Buchtal F 190 (41) 126 (41) 145
 Buchwald K W 173 (117) 788
 Bucklin D H 70 (90) 85
 Budziarek R 556 (31 38) 579
 Buckmann D 73 (92) 76 (91 92) 85
 Bueding I 51 (94) 85
 Buehler H J 667 (135) 659 681
 Bulbring F 79 (95) 80 (95) 85 101
 (46) 119 (44) 128 (43) 133 145
 Bugbee E I 236 (42) 2 0 298 401
 (116) 403 (116) 428
 Bullough W S 421 496 671 (41 42 43
 44) 678 754 (118) 788
 Bullwinkle B 72 (153) 51
 Bunding I 566 (739a) 386
 Bunding I M 219 221 (51) 232 233
 Burchenal J H 752 (119) 753 (120) 788
 Burgen A S V 117 (47) 145
 Burger H 611 (45) 617 (45) 678
 Burger M 191 (18) 193 194 195
 Burke D C 500 (59) 579
 Burkell C C 748 (121) 783
 Burkhardt J 319 (37) 380
 Burn D 514 (332a) 587
 Burn G P 396 (29) 496

- Bonomo I C96 (350) 766 (350) 794
 Bonanes R W 236 (34) 297
 Boon C J 295 (173) 301
 Boon H L 9 (311) 56
 Booker W M 741 (102) 787
 Bookman J J 721 (103) 725 (103) 787
 Boone M A 449 (79) 501
 Boot I M 61 (518) 94
 Boothby W M 449 (575) 481 (52), 486
 (82) 490 (80 81) 501 502 512
 Bowman J I 491 (700) 516
 Borden A L 727 (111 1099) 755 (104)
 787 814
 Borek I 591 (18) 655
 Borell U 208 991 416 (53) 502 670 (32
 33 34) 678
 Borgatti G 373 (28a) 579
 Borgstrom F 557 (35) 579
 Borman A 698 (300b) 794
 Bornstein I P 838 (22) 850 (63) 870
 871
 Bornstein J 186 (9 10) 193 195 257
 (35 36) 258 (37) 997 843 (23) 870
 Boroughs H 41 (35) 48
 Borst J G G 843 (74) 870
 Borth R 345 579
 Borthwick H A 27 48
 Bose J C 19 49
 Boss W R 395 426 710 (105 106) 787
 Bosshard W 528 (218) 586
 Bostrom U 669 (35) 670 (35) 678
 Boswell T 693 (1233) 818
 Botkin A I 453 (84) 460 (85) 500
 Bouckaert J P 175 (12) 195 417 (159)
 430
 Boucot N 848 (51) 871
 Bounhiol J J 68 (14 56 59 63 64) 71
 (57) 72 (58 60 61) 73 (62) 83 (58)
 83 84 139 (9) 144
 Bourcet P 450 (234) 505
 Bourne G H 691 787
 Bourne W 395 (192) 430
 Bovarnick M 475 (86) 502
 Boyet D 119 (29) 145
 Boyet Nitti F 119 (29) 145
 Bowden C H 453 (779) 454 (779) 518
 Bowen H F 157 165 173
 Bower L F 838 (22) 870
 Bowers J Z 759 (108) 787
 Bowman R H 240 (38) 297 674 (36)
 678
 Boyer G E 188 198 224 (7) 276 (7)
 231 249 (40) 294 (40) 998
 Boyd E M 490
 Boyle D 782 (852) 808
 Boyle F P 3 (128) 51
 Boylston G A 426
 Braccun C 743 (139) 788
 Bradbury J T 342 371 (79) 579 580
 671 (37) 672 (768) 678 684
 Bradley H T 492 (493) 511
 Bradley T R 358 (30) 399 421 426
 Bradlow H L 576 (46 61) 579 580
 609 (99) 610 (99) 611 (99) 616 (99)
 633 (99) 635 (99) 636 (99) 657
 Brady F J 437 (87 135) 502 503
 Brady R O 188 (13 14) 195 475 (509)
 513 610 612 615 623 655
 Branche G 701 (534) 799
 Brand F 177 195 406 426
 Brand F B 448 (88) 502
 Brandt W 191 (18) 195
 Brandt W L 711 (958) 811
 Branham S I 758 (805) 807
 Brannon F S 491 (485 641) 493 (485
 641) 511 515
 Brant F 841 (25) 842 (25) 870
 Branzi G 674 (164) 681
 Braun Menandez E 714 (109) 787
 Brauner I 28 (38 40) 29 (39 40) 43
 44 46 (41) 47 (42) 49
 Brauner M 28 (40) 29 (40) 49
 Braunsteiner H 435 (89) 436 (89) 502
 Braver R W 451 (91) 452 (91) 502
 Bray P F 745 (624) 802
 Brayer F T 438 (735) 517
 Brayne M K 476 (90) 502
 Brazeau P 674 (88) 680
 Brendler H 673 (38) 678
 Brennan M J 749 (784 1044) 806 813
 Brenner J 598 (235) 585
 Brenner S 121 (30) 145
 Bridges W L 705 (426) 796
 Briggs F N 784 (30) 298 449 (705)
 451 (91 104 705) 452 (91) 502 516
 Brill B A 243 (209) 245 (209) 302
 Brink N G 221 (87) 222 (5 8 9 87)
 224 (7 41) 226 (7 9 87) 231 232 233
 270 (40 415) 286 (20) 294 (40) 297
 298 308
 Briot M 757 (460) 774 (461) 787
 Bristol W R 712 (110) 787

- Cardesa A F 374 (148a) 333 60 (103) 63²
 Cardona Lynch I 777 (716) 804
 Cardwell H M I 537 (44) 338 (44) 579
 Caré F 68 (711) 88
 Caren R 199 (161) 199
 Carey I J 111 112 145
 Carey N 750 (1005) 812
 Cargill W H 155 170
 Carlisle D B 60 (102 105 105 113 114) 61 (7 101 108 110 112) 62 63 (102 103 106 107 108) 65 (102 103 104 108) 66 (104 108) 67 (7) 79 (98 99 100) 81 (98 99 100 101 109) 82 (100) 83 85 86 109 (7) 111 (67) 138 (65) 139 (6 65) 141 144 145
 Carlo J 745 (876) 808
 Carlson S P 100 (68) 145
 Carmel W J 833 (77) 834 (77) 872
 Carmichael J 736 (128) 788
 Carnes W H 156 161 161 170 173
 Carol J 542 (14 15) 543 (15) 559 (14) 578 667 (50) 679
 Carpenter C M, 777 (1189) 817
 Carpenter F H 221 (41 42) 231 232
 Carpenter R K 288 (56) 298
 Carr F H Jr 454 (102) 502
 Carr J E 226 (14) 231
 Carroll E 707 (343) 718 (59) 721 (343) 753 (343) 783 (59) 788 794 850 (14) 879
 Carroll K K 692 (132 133) 735 (132 133 134) 788 845 (30) 870
 Carruthers F P 260 (243) 303
 Carryer H M 757 (136) 765 (136) 788
 Carson D A 483 (508) 512
 Carson H L 74 (115) 86
 Carstam S P 61 (116) 62 (117 119) 81 (118) 86
 Carstensen L 221 (68) 232
 Carter A C 206 (90) 233 271 (355) 273 (355) 306 827 (184) 876
 Carter G S 476 (103) 502
 Carter J P 766 (16 74) 769 (17) 785 805
 Carter P 594 (87) 635 (85) 65 667 (78) 679
 Cartwright G E 748 (463) 797
 Casanova R 558 (45) 559 (45) 579
 Cash J R 766 (747) 805
 Casada I F 312 (43) 380 671 (251 251a) 63 (55) 679 683
 Casper son T 436 (775) 437 (228) 505
 Caspi L 638 639 (41) 647 (48) 648 (48) 619 (46-48) 656
 Cassen B 333 (3) 830 (68 69) 870 872
 Cassidy J W 743 (864) 808
 Castellani I 672 (51) 679
 Castor C W 719 (138) 754 (137) 788
 Catchpole H R 156 (41) 171 312 380
 Cater D B 277 (57 89) 278 (57) 298 299
 Cates J I 666 (12) 678
 Cavallero C 258 (58) 298 724 (141) 743 (137) 788
 Catz B 459 (101) 460 (101) 467 (105) 502
 Cavallito C J 24 49
 Cavanaugh L A 271 (350 351) 306
 Cavett J W 446 (106) 502
 Cawley E P 784 (1702) 817 833 (38) 871
 Cayala J 763 (182) 789
 Cazal P 78 (124) 79 (54 55 122 123 124) 82 (120 121) 84 86
 Cerceo E 604 605 661 667 (191) 682
 Chabrier P 465 (113) 502
 Chadwick L F 77 (175) 86
 Chaikoff I I 187 (34) 188 (5 20 22 68 85 164) 195 196 197 199 311 (113) 312 (742) 382 436 436 (511) 438 (796) 441 (709 714 792 735) 442 (211 476 638 709 710 714) 443 (107 711 713 732 791 792 794 795) 445 (107) 446 (637 713) 447 (754 790) 449 (405 707 712) 450 (706) 451 (91 704 705) 452 (91 708) 454 (238) 455 (471) 462 (211) 464 (657 797) 469 (210 637) 473 (793) 485 (184) 502 504 505 511 512 515 516 517 518 519 576 (46) 579 833 (65) 871
 Chakravarti R N 552 (94) 581
 Chalmers T M 305 (34) 496 711 (142) 788
 Chamberlain R 849 (51) 871
 Chamberlin E M 552 (275) 554 (47) 563 (47) 579 586
 Chamberlin J M 554 (294) 586
 Chambon Y 669 (52) 679
 Chamorro A 459 (108) 502 672 (53) 679
 Champagnat P 20 22 (61) 49

Burn J H 79 (9a) 80 (9a) 85 98 119
 (44) 128 (43 52 57) 121 (40 51)
 131 (49) 132 133 134 145 212 (30)
 231 241 (43) 252 (43) 293 391 (30)
 392 (30) 393 (30) 496 497 (CS2) 516

Burnes W W 613 (46) 678

Burnett C H 709 (507) 800

Burns B D 125 (58) 146

Burns C M 160 (22) 170

Burns F J 449 (94) 502

Burns T W 187 (6) 195 286 (44) 287
 (370) 293 (13a) 298 300 305 730
 (962) 811

Burris M J 674 (41) 678

Burris R H 29 (318) 42 (200) 53 56

Burrows B A 709 (507) 800

Burrows H 317 334 380 66a (48) 678

Burstein S 611 617 619 (768) 622 (31)
 631 (31-33) 63a (31-33) 637 (2) 30)
 647 (30) 618 (31) 66a (274) 671 (274)
 656 656 662 684 784 (1259) 819

Burstrom H 2 3 (53) 13 (137) 15 (48)
 49) 16 35 (53) 40 (51) 42 43 (50 51)
 44 45 46 49 51

Burt A S 315 380

Burton R B 634 (330) 63a (3a) 656 664

Burwell R I Jr 514 579

Buschle W 420 496

Bush I I 288 (47) 289 (45-48) 291
 (41) 292 (46) 298 694 788 707
 (1027) 818

Bussanello F 674 (22) 678

Buta S 496 (23) 500

Butcher E O 693 (124) 788

Butenandt A 82 514 (42) 560 (41) 579
 604 615 610 656

Butler G C 60a 656

Butler H M 396 498

Butler L C 739 (17a) 788

Butt A J 761 (901) 761 (901) 812

Butt W R 217 231 667 (49) 679

Buttaro C A 164 (23) 170

Butterly J M 840 (23) 870

Buu Hoi N P 597 (11 57 58) 655 656

Buxton C L 606 656

Buxton P A 496

Byers S O 43a (615 616) 514 672 (218)
 692 732 (36) 794

Byrne W W 336 380

Byrom F B 481 (95) 486 502

Bzwatner W G 462 (91 449) 502 510

C

Cabanis P J G 98 145

Caccianza I 164 170

Cagan R N 479 (97) 509 727 (121)
 730 (126) 788

Calhoun G I 783 (589 590) 801

Call C A 217 (4a) 232

Calhoun D 708 (9a9) 811

Calkins I 837 (172) 873

Callan H G 79 (96) 85

Callow N H 615 616 656

Callow R A 501 (43) 579 615 656

Calvin L D 717 (179) 733 (179) 789

Calvo J M 490 (98) 502

Camarata S J 840 (84) 876

Cameron A 670 (56) 679

Cameron A T 736 (128) 788

Cameron G R 761 788

Cameron M I 76 (97) 77 (97) 85

Campbell J 203 (12) 231 236 243 (a1)
 52 54) 249 (54) 251 (27) 253 (49)
 251 (27 53) 257 (54) 267 (51) 26a
 267 (52) 274 (50) 297 298

Campbell W W 161 (149) 174

Canus G 21 24 (5a) 49

Canadell J M 833 (224) 87a

Campy F G 449 (326) 507

Canivene R 362 369 370 312 380

Cannon B 775 (1193) 817

Cannon I J 768 (689) 803

Cannon J A 715 (130) 788

Cannon W B 94 115 130 (61) 131 (60)
 145 43a (214) 604

Cantraw A 773 (18) 799 457 (131)
 481 (625) 482 (676) 503 514 594
 (230) 601 736 (917) 782 (851 8a7)
 808 810 824 (42 151) 828 (41) 834
 (42) 871 874

Cape R D T 747 (131) 788

Caplin S M 73 (278) 55

Capot I 438 (6) 70) 501

Cappiello M 243 (319) 249 (319) 305

Capra P 483 (100) 484 (99) 100) 502

Capraro A 397 496

Capretta G 489 (101) 502

Carballeira A 272 (55) 2 3 (55) 700
 (5a 402) 298 307

Carbone J 733 (637) 802

Card L L 374 384

- Chisby K H 462 (463) 513
 Cloud L P 815 (73) 818 (73) 8 0
 Clough H D 190 (89) 197
 Cluxton H F 694 (107) 775 (10 9) 814
 Cobb S 718 (153) 789
 Cochran T H 733 (911) 771 (912) 809
 Cochrane G C 696 (158) 789
 Cockett A T 719 (37) 793
 Code C I 512 (76) 574 (7 6) 584 704
 (737) 746 (274) 756 (1194) 757 (136)
 765 (136) 7 0 (233) 771 (233) 772
 (737 2 4) 774 (733) 775 (233) 789
 791 792 805 817
 Coleridge R A 561 (97) 576 (207) 686
 Coggeshall C 274 (65) 998
 Coggeshall H C 484 (119) 503
 Coggins W 266 (99) 299
 Cohen A G 484 (170) 503
 Cohen H 295 (66) 998
 Cohen M R 556 (173 201) 873 875
 Cohen R A 494 (171) 503
 Cohen R S 780 (159) 789
 Cohen S L 545 (48) 580 667 (96 186)
 669 (107) 640 (107) 680 687 745
 (450) 794
 Cohn C 458 (646) 516 777 (160) 789
 Cohn M 405 496
 Cohn D V 153 (63 65) 154 (64 66 67
 69) 165 (61 67) 166 (67) 171
 Cole D F 711 (161 167 163) 789
 Cole H H 338 (49) 344 380
 Cole J W 764 (164) 766 (165) 789
 Cole P D 240 (235) 303 349 383
 Cole W 549 (177 179) 583
 Collens W S 193 195
 Collingsworth D R 642 (108) 656 658
 Collins D A 715 (166) 789
 Collins L 697 (1121) 698 (1170 1121)
 708 (1172 1123) 714 (1122 1123
 1176) 722 (1176) 815
 Collings M C 195 (32) 196
 Collip J B 157 173 251 (2 0) 252 (30)
 253 (86 170) 295 (769) 297 299 304
 471 (37 3) 476 487 704 (825) 807
 Collonge A 700 (183) 704 (183) 789
 Colb V 104 (111) 144
 Colowick S I 184 (101) 185 (75) 195
 197 257 (755) 304
 Colpoys F L 766 (318) 793
 Colton F B 569 (49 50) 580
 Comar C I 158 (96 145) 172 173
 Common R H 642 (195) 674 (195) 682
 Commoner B 33 (69) 45 46 (68) 49
 Comsa J 483 (173 121 125 126) 497
 (122) 503
 Concha F 488 (127) 503
 Condliffe P G 204 271 (31 32) 931
 Conn J W 276 (14) 231 260 (68) 271
 273 (67) 291 (69) 298 696 (172) 697
 (170) 698 (170) 705 (170 172 177)
 707 (177) 708 (170) 713 (167 169)
 171 174) 721 (172 177) 724 (173 174
 175 176) 775 732 736 (168 313) 753
 (177) 781 (168) 789 793 841 (35)
 812 (34) 8 0 871
 Connor M H 333 332
 Connor T B 153 (77 78) 160 (77 78)
 162 (77 78) 171 172
 Conroy L 745 (1141) 816
 Consolazio C F 740 (1100) 742 (608)
 801 814
 Constable B J 734 (468) 797
 Constantin J M 539 (288) 549 (51)
 550 (51) 558 (57) 580 586
 Constantine A G 766 (878) 807
 Conston A S 866 (31) 870
 Contoulis J 540 (8) 552 (8) 554 (8)
 556 (8) 578
 Cool B 46 (69a) 49
 Cook C 767 (30) 768 (148) 785 789
 Cook D L 714 (149) 733 (179) 789
 Cook J W 616 (75) 65
 Cook R P 120 145
 Cooke R L 153 (86) 154 (86) 155 172
 Coon J M 333 426
 Coon W W 740 (1165) 741 (1168) 816
 Coons A H 445 (480) 511
 Cooper W D 400 (200) 430
 Coover M O 484 (600) 514
 Cope C L 634 (54) 635 (54) 656
 Cope O 463 (553) 499 (551) 512 515
 Copp D H 250 (395) 307
 Coppedge R L 211 231 371 375 380
 Corbett B D 439 (481) 511
 Corcoran A C 493 (178) 503 700 (735)
 714 (734 736) 740 (963) 805 811
 Corcoran J W 563 (160) 582
 Corey F J 574 (54) 548 (53 54) 549
 (55) 580
 Corn A T 481 (310) 507

- Chandra R 479 (110) 487 (110) 498 (109) 502
 Chang C 212 (63) 214 (63) 298
 Chang H 158 159 170
 Chang H C 101 (60) 146
 Chang M C 671 (199 231) 682 683
 Chang T S 158 170 171
 Channon H J 502 656
 Chipman D W 708 (143) 788
 Chapman F M 478 (111) 502
 Chipman R I 742 (608) 801
 Chipman W H 759 (190) 790
 Charalampous F C 189 (21) 195
 Charipper H A 245 (336 337) 282 (336) 306 476 (246) 491 (219) 505 736 (200) 790
 Charniaux H 64 (126-128) 86
 Charniaux Cotton H 64 (130) 65 (129) 86
 Charney C W 866 (31) 870
 Chart J J 870
 Chase H B 755 788
 Chase J H 791
 Chash B W 427
 Chauchard B 672 (54) 674 (54) 679
 Chauchard P 672 (54 150) 674 (54 150) 679 681
 Chaudonneret J 68 (131) 86
 Chauvet J 404 414 495
 Chavre V J 756 (201) 790
 Chefurka W 75 (132) 86
 Chemerda J M 552 (65 270) 554 (47 294) 563 (47) 579 580 586
 Chen G 212 (63) 244 (63) 298 757 (145) 788
 Chen J 422 (35) 46
 Chen R W 576 (46) 579
 Cheng C P 281 (59 60) 282 (61 62) 298 719 (1231) 720 (146) 788 818
 Cheng D W 737 (760) 781 (760) 805
 Cheng P 673 (55) 679
 Chernick S S 188 (22 80) 195 197
 Chesney A M 461 (112) 507
 Chesterman F C 703 (147) 788
 Cheymol J 465 (113) 467 (114) 507
 Chlucwicz Z 62 (133) 86
 Chilcote M E 154 (150) 161 (150) 174
 Childs D S 440 (115) 441 (115) 502
 Chinn L T 537 (172) 683
 Chiu C Y 722 788
 Chornack F W 484 (478) 511
 Chou C 242 (63) 244 (63) 298
 Chow B F 210 215, 218 238 (61) 231 238 (64) 278 (153) 298 301 416 (201) 431
 Christen R 701 (10) 785
 Christensen W R 185 (23) 195
 Christian R I 673 (55) 679
 Christiansen G S 40 (63 64) 41 (63-60) 43 49
 Christiansen R G 537 583
 Chu J I, 494 (116) 502
 Chung A C 365 (240) 366 (239a) 380 386
 Cianficconi, F 68 (307) 91
 Cierszko I S 206 231
 Cipra A 461 (432) 510
 Claes H 31 (66) 49
 Clapham H M 488 (109) 502
 Claringbold P J 494 (68) 501 668 (78) 678
 Clark A J 131 (70) 146
 Clark C T, 101 (277) 180
 Clark D E 452 (324) 507 787 (150) 788 830 (183) 875
 Clark H E 30 (67) 49
 Clark I 728 (151) 729 (152) 789
 Clark I C Jr 616 618 (50 1) 656
 Clark L D 718 (153) 789
 Clark R F 497 (670) 516
 Clark W I le Gros 320 380
 Clarke A P W 716 (157) 789
 Clarke F H 816
 Clarke J C 243 (365) 245 (365) 306
 Clarke P M 375 376 380
 Clary M L 670 (06) 679
 Clausen K H 408 (117) 507
 Clauser H 403 (77 141) 414 (72) 427 499
 Clawson T A 461 (112) 507
 Claxton B F 740 (154) 741 (150) 789
 Claxton J C 452 (118) 507
 Clayton R B 503 (30) 579
 Clayton T 558 (241) 585
 Clegg M T 344 380
 Cleghorn R A 716 (157) 717 (156) 789
 Clemons H 467 (487) 511
 Cleroux A P 290 (281) 304
 Cleveland A S 750 (545) 752 (545) 800
 Cleveland L R 68 (134) 82 (134) 86

Curtis J M 591 (68) 656
 Curtis P H 700 (134) 790
 Cushing H 246 (35) 200 (387) 397
 874 (33) 871
 Cuthbertson D I 217 (76) 769 (74)
 270 (75) 799
 Cutkomp L K 77 (416) 9
 Cutler H H 841 (40) 871
 Cutuly I 278 (71) 799
 Cutuly E C 278 (77) 799
 Czarski P 161 (33 34) 170

D

Daane T A 349 380
 Dacanay J G 719 (493) 720 (493) 798
 Daellenbach H R 502 (146) 554 (146)
 556 (146) 580
 Daft F S 737 (190) 790
 Dailey M E 482 (720) 617
 Dalberg L Y 42 (94) 50
 Dale H H 98 131 146 407
 Dalghesh C L 136 (78) 146
 Dalton A J 498 (136 4 0) 503 511
 781 (196) 790
 Dameron J T 693 (197) 790
 Damgaard E 498 (741) 517 749 (1166)
 816
 d'Amour F E 759 (1038) 815
 Danford H G 706 (198) 790
 Danford P A 706 (198) 790
 D'Angelo S A 208 731 273 (78) 297
 407 (137) 503 706 (200) 790 824
 (42) 808 (41) 834 (42) 871
 Daniel P M 375 (762b) 586
 Danielli J F 176 127 146
 Daniels R 339 (342) 588
 Dannenbaum H 615 620 (39) 666
 Dannenburg A M Jr 777 (716 117)
 804
 Danowski T S 400 (130) 503 705 (701)
 790
 Danto J L 740 (707) 790
 Dao T I T 696 (544) 750 (044) 784
 (203) 790 799
 Darling P C 441 (454) 797
 Darrach M 775 (817) 807
 Dart I M 701 (534) 790
 Das N 427
 Dauben H J Jr 550 (60) 580
 Dauben W C 534 (67) 549 (63 64)
 502 (60) 576 (46 61) 579 580
 Dauber D 486 (133) 503
 Dandel I 537 656
 Dandel R 507 656
 Dauphinee J A 749 (378) 793
 Daus M A 579 (107 161) 561 (161)
 512 (161) 563 (161 167) 580 583
 Dauter I 339 380
 Davenport H W 287 (370) 305 790
 (062) 706 (704) 790 811
 Davenport V D 703 (178) 719 (1228)
 818
 David K 610 610 656
 Davidoff I M 824 (39) 871
 Davidson C S 741 (1106) 816
 Davidson I W F 703 (17) 231 76
 713 (01 02 54) 719 (04) 703 (49)
 007 (04) 762 (51) 260 (00 51 52 54)
 767 (02) 774 (50) 798
 Davidson J A 449 (19) 501
 Davies B M A 154 (06 31) 160 167
 (36 37 8) 168 170
 Davies R I 40 (231) 53
 D'Avignon M 877 (43) 871
 Davis A K 706 (200) 708 (200) 710
 (200) 711 (377) 790 793
 Davis C 691 (815) 807
 Davis F A 22 (213) 53
 Davis J O 700 (706) 709 (06) 710
 (706) 790
 Davis J P 175 (772) 150
 Davis J S 376 (150) 377 (189) 584
 Davis J W 540 (318) 587 624 (303)
 675 (03) 633 (003) 675
 Davis M E 691 (707) 00 (708) 790
 805 (44) 811
 Davis R F 76 (13) 49
 Davidson C 289 (171) 301
 Davison I 717 (119) 733 (179) 799
 Davoll H 406 (147) 407 409
 Dawson A B 313 346 380 380
 Day B F 17 (74) 49
 Day H G 488 (341) 503
 Day M F 77 (145) 8 (144) 86
 De Andino A M 742 (709) 790
 Deane H W 186 (145) 198 278 (154)
 290 301 690 691 (210 220) 737 (711)
 750 (210) 781 (210) 777 (1138) 790
 791 796 816

- Cori C F 181 (101) 180 (26), 191 (142)
 194 195 197 198 202 (110) 203
 (110) 207 (280) 300 304
 Cori G T 180 (25) 195
 Cori O 481 (310) 507
 Cornblath M 213 (278) 260 (278) 304
 Cornbleet T 704 (60) 779 (60) 780
 Cornelis G 192 (153) 199
 Corner G W 670 (58) 679
 Corner G W Jr 438 (111) 507
 Cornfield J 703 (940) 810
 Cornforth J W 336 (57a) 537 (44
 57) 538 (41) 542 (56) 557 (56) 579
 580 656
 Corns W G 42 (70) 49
 Cornubert G 63 (137) 64 (136 138) 60
 (139 140) 66 (136 137 140 142 143
 514) 86 94
 Corrigan K E 938 (170) 874
 Corsden M 463 (221) 467 (221) 505
 Corson D R 436 (129 650) 503 515
 Cortegiani I 101 (72) 106 146
 Cortell R E 440 (637) 475 (130) 503
 515
 Cortis Jones B 221 231
 Cosgriff S W 749 (180) 789
 Cosgrove J B R 458 (131) 503
 Coste F 671 (59) 679 740 (181) 663
 (182) 789
 Costerousse O 597 (57) 600
 Cotes P M 243 (70 71) 262 (71) 260
 (71) 210 (70) 238 360 (51) 366 380
 Cournot L 674 (103) 680
 Courrier R 452 (132 133) 503 700 (183)
 704 (183) 789
 Courvoisier B 151 (103) 170
 Coussens R 417 (109) 430
 Covian M R 714 (109) 780
 Coward K H 370 (95) 381
 Cowgill G R 458 (134 300) 503 507
 Cowie A T 169 (31) 170 338 (53) 34
 (54) 361 363 380 400 (39) 420 610
 (185) 754 (184) 789
 Cowie D B 437 (87 130) 507 503
 Cowie L M 42 (730) 500
 Cox H R 207 (204) 304
 Cox P I 302 380
 Crabb F D 743 (623) 703 (150) 789
 802
 Craig J M 774 (187) 780
 Craig L C 176 (59) 177 (58 60) 178
 190 401 406
 Craig R I 714 (427) 796
 Crain D 535 539
 Crandon, J H 741 (188) 789
 Crane R K 477 (491) 480 (491) 511
 Craver B A 670 (56) 679
 Crawford J D 104 (32) 150 162 10
 173
 Crawford T B B 100 (8) 102 (73) 104
 (80) 100 137 (8) 144 146
 Creaser C W 434 (242) 505
 Crepea S B 761 (189) 776 (189) 790
 Crevier M 496 (283) 515
 Crichton J A 365 (51) 366 (51) 380
 Crightler J F Jr 603 (321) 664 714
 (1212) 782 (772) 783 (1213) 806 818
 801 (36 230) 802 (36) 871 80
 Crile G 839 (360) 871
 Criscitello M G 696 (412) 714 (472)
 797
 Crispell K H 271 (70) 272 (3) 213
 (72) 298 299
 Crispin D J 704 (826) 807
 Crochton J A 243 (70) 270 (70) 298
 Crockett C I 747 (319) 793
 Cronkite I I 492 (334) 507 709 (190)
 790
 Crooke A C 208 217 221 (20) 231
 Crooks H M Jr 528 (214 219) 509
 (219) 561 (219) 567 (214) 584 604
 (197) 600 (185) 600
 Crooks B A 363 (50) 380 400 (39) 400
 (40) 417 406
 Crossman J 305 (219) 430
 Crowdy S H 37 (71 72) 49
 Crowfoot D 176 106 512 (59) 580
 Crovatto H 420 400
 Cruik S 164 (100 101) 172
 Cruickshank I A 501 (297) 576 (297)
 586
 Crapo A C O (58 60) 60 60
 Culiner A 871
 Cumming C R 737 (268) 808
 Cummings M M 770 (192) 772 (191)
 777 (113 766 932) 790 806 810
 Cunningham O 78 (74) 85
 Curtis A C 781 (1702) 817 833 (38) 871
 Curtis G M 442 (701) 449 (545) 810
 516 827 (172) 838 (54) 871 874

- Devitt J F 743 (230) 791
 De Vries J A 791
 de Vries I A 843 (24) 870
 de Walleville H 345 (28b) 579
 de Wit J C 661 (26a) 684
 Des B P B 770 (73) 771 (233) 772
 (237) 774 (237) 775 (233) 791
 Dey F I 371 381
 Devsach I J 436 (161) 603
 Dhy e F G 672 (112) 680 737 (234)
 791
 Diamond M C 712 (23a) 791
 Di Bella S 676 (15) 678
 Dickel D F 534 (67) 580
 Dickel H 754 (736) 791
 Dickens F 117 (47) 145 158 170 351
 391
 Dieker S E 393 (54) 416 47
 Dickie I F 449 (774) 517
 Diezfulasy E 347 381 59a 656 733
 (237) 791
 Dilecock K A II 670 (716) 687
 Diefenbach A F 743 (180) 789
 Diermeier H F 249 (90) 299
 Di George A 837 (47a) 871
 Dijkmans J 497 (160) 503
 Dill D B 741 (188) 789
 Dillaha C J 7a4 (239) 7a5 (238) 239
 948) 491 811
 Dillaha J 727 (48) 797
 Diller I C 7a1 (241) 791
 DiMarco A 774 (141) 788
 Di Mauro S 845 (90) 872
 Dimond A E 36 (319 320) 56
 Dine R F 488 (167) 504
 Dingemanse E 239 (711) 271 (87) 799
 307 419 (52) 477 544 545 580 594
 610 (59) 615 (59) 656
 Dingman J F 688 (1151) 696 (1151)
 697 (1151) 704 (1151) 721 (1151) 727
 (1151) 745 (1149) 76a (1151) 766
 (1151) 768 (1151) 777 (1151) 816
 D'lorio A 496 (583) 513
 Dirks H B Jr 474 (48) 475 (356) 476
 (46 47) 501 508
 Dirnagl K 439 (163) 504
 Dirscherl W 191 (43) 196 676 677 (63
 64) 679
 Di Stefano H S 249 (90) 799
 Dittman W A 214 (99) 216 (99) 218
 (99) 793
 Dittmer K 467 (52) 501
 Dixon H B F 221 272 224 791 277
 (89) 279 299
 Dixon J S 279 (229) 303
 Dixon T F 179 170 173
 Djeras C 791 (68) 529 (212) 740 (76
 77 184) 541 (74 284) 749 (67 78
 766 268) 770 (313) 557 (7a 763 264
 76a 270) 553 (347) 5a4 (10 313)
 5a6 (69 72 73 767) 5a7 (7a) 5a8
 (313) 568 (269) 579 (769) 570 (71)
 580 583 584 585 586 588 608 (179)
 660
 Dobija M 777 (686) 803
 Dobriner K 522 546 (706) 549 (17a)
 561 (708) 565 (705) 568 (323) 580
 583 584 587 604 (170) 609 (6 99) 610
 (99 316) 611 (99) 612 (134) 613 (134)
 615 (170 316) 616 (99 100 316) 618
 (100) 621 (100) 632 (170 760b) 633
 (99) 63a (99 171) 636 (99) 655 656
 657 658 659 660 667 664 667 (65)
 679 751 (1113) 752 (1113) 815 861
 (48) 871
 Dobson M M 673 (746) 683 784 (1132)
 816
 Dotyns B M 211 231 2a4 (91) 799
 412 (164) 457 (165) 499 (a51) 504
 512 833 (49) 871
 Dockrell A W 278 (253) 303
 Doe R P 786 (92) 291 (92) 799
 Doerner A A 700 (247) 749 (951) 775
 (742) 791 811
 Dohan F C 76a (93) 799 634 (748) 637
 (218) 636 (218) 667 849 (51) 859
 (70) 871
 Dohrn P F R 62 86
 Doisy E A 217 (45) 232 575 (7) 576
 (34) 578 579 594 (148 176 790 318)
 598 (19 215) 621 (3) 655 656 659
 660 681 683
 Doisy E A Jr 57a (7) 576 (34) 578
 579 598 (19 215) 621 (3) 637 (147)
 65a 659 681
 Dole J C 768 (471) 797
 Dolin G 246 (94) 768 (94) 299
 Domini G 128 (755) 150
 Donahue J K 81 (161) 87

- Deanesly R 491 (141) 603
 de Angelis W 480 (140) 603
 Deb C 740 (52 53 54) 786
 De Bastiani G 257 (79) 799
 De Beruchamps I M 50 (146) 86
 De Bodo R C 243 (82 210) 256 (363)
 257 (80) 261 (210) 265 266 (82) 270
 (81) 299 300 306 373 386 395 (22)
 426 726 (213 214 215 216) 790
 De Bourgraaf J F 419 (52) 400
 Debre, R 718 (217) 790
 de Bruin H 4 (145) 51
 De Buen A M 68 (147 148) 86
 De Candia S 485 (142) 503
 de Castillo J 117 (80) 146
 De Costa I J 700 (218 219) 790
 Dedman M L 221 231 276 (85) 256
 (85) 293 (65) 299
 de Duve C 175 (12) 190 (144 157) 191
 195 198 199
 Defretin R 59 (149) 86
 De Gasperis A 437 (143) 503
 De Groot J 252 (84) 283 (84) 299 321
 380
 Deiss W P 450 (144 381) 503 508
 De Jongh S I 249 (401) 200 (121) 266
 (400) 300 307 594 650
 de la Balze F A 811
 Delabarre F 671 (59) 679
 De Lamater I D 780 (220) 790
 de Lattin G 60 (321) 89 90
 de Lawder A M 190 196
 Delav J 494 (63) 501
 Delbarre F 747 (181) 789
 del Castillo E 449 (684) 516 828 (198)
 837 (198) 838 (198) 875
 De Lema B 68 (151 153 154 155 156)
 72 (150) 78 (155) 86 87
 Delf E 862 (104) 872
 Deltour G H 442 (595) 446 (603) 447
 (556) 448 (145 604) 453 (585) 503
 513 514
 Deluca H A 735 (915) 810
 de Lustig F S 804
 Demajo S 251 (215) 302
 De Maria W J A 154 (66 68) 111
 De Martini F 753 (221) 754 (221) 790
 De Meio R H 600 661
 Démeusy N 65 (158 159 160) 66 (14-
 143 160 514) 86 87 94
 Deming J M 45 (186) 50
 Deming Q B 702 (222) 711 (409) 790
 796
 De Moor I 243 (256) 246 (256) 249
 (256) 257 (256) 258 (256) 262 (255)
 265 (255) 303 790
 De Mowbray R R 744 (223) 790
 Dempsey L 707 (343) 718 (59) 721
 (343) 753 (343) 783 (59) 786 794
 800 (14) 861 (235) 870 876
 Dempsey I W 437 (146 147 149) 442
 (147) 460 (148) 462 (146) 503 691
 (225) 791
 Dempster G 767 (840) 779 (840) 807
 Den H 265 (83) 299 725 (216) 790
 Denney D B 524 (240) 585
 Denffer D von 2 (76 77) 3 (76 77) 23
 (77) 30 (75) 31 (78) 49
 Denny F W 778 (446) 797
 Denny Brown D 835 (46) 871
 Dent C F 154 170
 Dent F M 741 (107) 787
 Dent J H 770 (227) 791
 Depraeter M 701 (1158) 816
 Derbes V J 770 (227) 771 (226) 791
 de Robertis I 437 (151 152 154 155)
 442 (154) 447 (150 151 152 153)
 448 (150) 456 (152) 462 (151) 503
 De Roeth A 106 (81) 146
 de Ropp R S 32 (104) 3, (252 253)
 36 (252 254) 50 54
 Derriek J B 253 (86) 299 421 (53) 427
 Derrien Y 446 (157 158) 448 (158 58)
 503 513
 Derwans A 764 (228) 791
 Desaulles I 704 (754) 707 (229 913)
 791 805 811
 Desclaux L 672 (239) 683
 Desclin L 313 332 348 351 360 361
 362 368 369 370 380 381 609 (61)
 679
 Deuel H J Jr 476 (437) 485 (251) 488
 (252) 506 510 518
 Deulofeu V 531 656
 Deulch G 452 (155) 503
 Deulch A S 509 (309) 58
 Devis R 863 (47) 871
 Devis Vandon I 863 (47) 871
 de Visscher M 440 (634) 497 (110) 503
 515

Durfee H K 3 (125) 51
 Durgin M L 74 (73) 79?
 Dury A 705 (71 72 23) 71 (71)
 747 (270) 79
 Duthie J J R 716 (70 931) 810
 Du Toit C H 451 (900) 519
 Dutt R H 334 335 336 (76) 381 671
 (70) 679
 du Vigneaud V 99 (79 272 290) 149
 150 401 402 403 404 (176 161 164
 209 211) 405 (36 176 176) 406 (51
 147 171) 497 498 (171 210) 499 41?
 (148) 413 426 427 428 429 430 431
 Dvoskin S 443 (174) 445 (174) 504
 Dworetzky M 746 (274) 77 (274) 79?
 Dye J A 478 (175 176 177) 504
 Dyer F J 375 (95) 381
 Dyer H M 401 498
 Dyk horn S W 375 385
 Dyrenfurth I 290 (402) 50
 Dziemian A J 417 (178) 504
 Dzieniatkowski D 492 (174) 504

E

Eakin R M 100 (84) 146
 Earle D P Jr 210 (81) 239 305 (22)
 426
 Eastcott H H G 756 (75) 79
 Easterling L 489 (698) 516
 Eastham J F 549 (63 64) 55 (65)
 56 (46) 579 580
 Eberhard T 789 (852) 808
 Ebersole C M 613 (246) 683 781 (1132)
 815
 Ebert R H 761 763 (748) 79?
 Ebling P S 611 (71) 679
 Echallier C 63 (180) 87 141 (85) 146
 Edelmann A 692 (279) 79
 Edman I 408 (60) 497
 Edsall J T 406 496
 Edwards G A 66 (181) 87
 Edwards W B 700 (242) 775 (247) 721
 Egg Larsen N 165 (114) 173
 Egami A 79 (182) 87
 Egerton B W 740 (504) 798
 Eggleston L A 187 197
 Egusa M 698 (191) 680
 Ehnobom K 68 (183) 87
 Ehrenstein M 543 (85) 546 (137) 547
 (13) 571 (131) 578 581 583
 Ehrich W I 761 (991) 764 (928 990
 931) 771 (990) 773 (990) 812
 Eichenberger K 52 (136) 554 (144
 146) 556 (141 146) 58?
 Eichenman I I 773 (780) 791
 Eichna I W 493 (180) 504
 Eickner L 669 (72) 69
 Eichorn K B 837 (81) 872
 Eidinoff M L 577 (86) 578 (114) 581
 599 609 (69 99) 633 (99) 635 (983
 99) 655 657
 Eiknes K 256 (32) 297 694 (281) 79?
 Eiman J W 851 (50) 871
 Eiseheid A M 732 (752) 79
 Eiden H A 79?
 Eisenberg F 668 (70) 616 (73) 679
 683 718 (414) 720 (284 414) 79? 796
 Eisenberg H 153 (74) 171
 Eisenmenger W J 844 (21) 870
 Eisenstein A B 740 (286) 741 (285
 986) 79?
 Eisler A 205 (66) 998
 Elberg S S 205 991
 Elhagez F A A 535 580
 Elhel I I 05 (555) 707 (857) 751 (287
 288) 753 (857) 754 (857) 79? 808
 Elks G F 554 (31) 556 (31) 579
 Ellnbogen L 176 (93) 19?
 Ellinger F 750 (289 291) 760 (700)
 79
 Elliott B B 27 50
 Elliott F H 289 (28) 997
 Elliott G C 153 (43) 171
 Elliott H A 485 (736) 505
 Elliott H W 616 (73) 679 720 (51
 413) 79 796
 Elliott K A C 106 145 704 (292) 79?
 Elliott T R 97 146
 Elliott W H 575 (7) 576 (34) 578 59
 598 (19 215) 621 (3) 637 (147) 655
 659 661
 Ellis B 546 (87) 581
 Ellis S 195 (37) 250 (96) 251 (96) 166
 299
 Ellsworth R 154 (2) 155 170 171
 Elmadjian F 692 (875) 694 696 (875)
 744 (293) 745 (793) 79? 808
 Elmer A W 439 (181) 504
 Elminger P J 795 (126) 60
 El Rawi I 459 (104) 460 (104) 509

- Donald T C 752 (243) ~91
 Donaldson D M 771 (726) 804
 Donard T 669 (136) 681
 Done A K 694 (214) 701
 Donsach I 470 (106) 604
 Douin M N 608 (98) 642 (98) 657
 Douini P 217 ~31
 Donker J D 306 381
 Donohue W I 751 (1017) 815
 Donovan B T 330 381
 Doonan P D 724 (508) ~98
 Dorfman A 674 (227) 683 718 (246)
 761 (245) 763 (66) 786 791
 Dorfman L 522 580
 Dorfman R I 154 (122) 167 (122) 168
 173 259 (163) 301 500 (80) 591 (305)
 599 604 (275) 605 606 610 (14 15
 269) 611 612 (15 134 209) 613 (~0.)
 615 (14 15 211 212) 616 617 619
 619 622 (34) 623 (15 117 121 122)
 624 (303 311) 625 (303) 626 632 (80)
 633 (72 80 303) 634 (31-33) 635
 (31-33) 636 (144) 637 (29 30 144)
 638 (96 306) 644 (80) 645 (114 117)
 646 647 648 (31 96 112 255) 649
 (72 80 144) 650 (71 72) 651 (71)
 6 2 (~3 73a) 654 (254a) 656 656
 657 658 ~59 661 662 663 667 (221)
 668 (66) 673 (140) 675 676 (108)
 679 680 681 683 701 702 (219 251)
 703 (252) 704 (247 249 250) 720
 (480) 731 (478 479) 791 797 798
 ~0 (140) 875
 Dorrance W R 459 (6~7) 516 ~82
 (1049) 783 (~09) 795 813
 Dorsey E A 665 (135) ~31
 Doty P 176 (28) 196
 Doughaday W H 243 (~74) 260 (278)
 304
 Dougherty J 449 (167) 504
 Dougherty T F 243 (209) 245 (209)
 302 671 (68) ~79 704 (257 9~1) 744
 (256) 745 (254) ~61 (253-~55) 762
 791 803 811
 Douglas W W 137 (82) 146
 Dow C 4~7
 Dowling D F 341 (75) 381
 Downs W G Jr 245 (95) 299
 Doyle I M 667 (134) 681
 Doyle L P ~65 (~60) 791
 Drabkin D L 477 (168 169), 478 (168)
 504
 Drach I 61 (16. 163 164 165) 63 (162-
 164) ~7 (162 164) 87
 Drachman S R 721 (103) 725 (103)
 732 (5 6 9) 785 787
 Dragstedt I R 494 (4~6) 511 756
 (1260) 819
 Drake M P 227 (48 49) ~3~
 Drake T G 161 (61) 171
 Draper H H 737 (261) 739 (261) ~91
 Dratz A F 154 (67) 165 (67) 166 (67)
 171
 Drechsler K 479 (760) 518
 Drending A S ~40(10 805 81 82 83
 84) 342 (~) 559 (9) 578 581
 Drill V A 4~6 (170), 488 (170 172) 4~9
 (171) 504 712 (110) 78~
 Dritch R 732 (7 8 9) 785
 Drummond J C 488 (407) 509
 Drury D R 182 187 (163) 189 196
 199 714 (958) 722 (1206) 811 818
 Drury M I 766 (262) ~91
 Drusch H E 777 (1189) 81~
 Dry dale G R 4~9 (173) 504
 Dubnik C S 408 (136) 503 781 (196)
 790
 Dubois E F 494 (570) 513
 du Buy H G 39 (265) 54
 Ducommun P 674 (69) 6~9
 Ducommun Lehmann b 674 (69) 678
 Dudley H W 401 45~
 Dufast P P 121 (202) 148
 Duff I F 727 (929) 732 (9~9) 765 (929)
 766 (263) 781 (1225) 79~ 810 818
 Duff J I 123 144
 Duff S R 537 (44) ~38 (44) 5~9
 Duff B J 750 (~64) 792
 Dugal L P 741 (~65 266) 792
 Dugas M 384
 Duggar W M 1~ (202) 53
 Dumm M E 737 738 755 79 809
 Duncan G G 193 (30) 196
 Dunn J S 189 196
 Dupont Raabe M 68 (168 170 172) 76
 (167 169 171 173 174) 80 (169) 81
 (166 171) 87
 Durand J B 60 (39 40) 66 (39 40) 84
 98 (26) 109 (26) 138 (26) 1 ~ 144 146
 Durehon M 59 (42 43 175-1~9) 84 8

Evans I D 770 (1015) 780 (1016) 812
 Evans R D 438 (111) 456 (340) 502
 508
 Evans R M 551 (31) 556 (31) 579
 Evans V J 769 (310) 793
 Evan W F 493 (691 695) 516
 Evens M 25 27 50 53
 Everall J 751 (311) 793
 Everett C M 492 (667) 516
 Everett J W 330 333 (82-81) 334 (273)
 368 371 381 384 385 671 (162 276)
 681 683 853 (130) 869 (130) 873
 Everett N B 346 381
 Eversole W J 391 (74) 496 4 7 12
 (375) 795
 Evertzen A 1,7 (79) 197
 Fwer R F 598 399 (621 63) 427
 Fva M 781 (175) 818
 Fyer H 191 (43) 196

F

Fager J 458 (468) 459 (468 469) 471
 (469) 488 (466) 510 511 742 (781)
 782 (780 783) 806
 Fahey J I 749 (317) 793
 Fajans S S 271 (61) 273 (67) 291 (61)
 998 691 (831 837) 696 (172) 677
 (170) 698 (170) 705 (170 172 177)
 707 (171) 708 (170) 721 (170 177)
 725 732 736 (168 313) 753 (111) 781
 (168) 782 793 807 742 (34) 870
 Falk M S 7,0 (314) 793
 Farah A 706 (425) 712 (425) 796
 Farber R K 243 (722) 754 (722) 503
 6 4 (152) 691 733 734 (692) 804
 Farber S J 770 (81) 299 395 (22) 427
 Farkas A 577 (88) 581
 Farkas L 577 (88) 581
 Farm r T H 271 (23) 281 776 (85)
 786 (85) 293 (85) 929
 Farnsworth E B 704 793
 Farquhar M G 314 381
 Farquharson R F 857 (53) 8 1
 Farrar B 729 (17) 787
 Farrar M W 599 (16) 579
 Farrell G I 254 (108) 288 (377) 360
 563 (163) 583 816
 Fatt P 114 115 178 147 147
 Faustini R 78 (205) 87 104 (92) 146

Fauvet F 361 381
 Favour C B 749 (1174) 770 (707) 804
 817
 Fawcett C H 14 50
 Fawcett D M 441 (190) 442 (191) 467
 (190) 473 (190) 504
 Fawcett D W 861 (238) 876
 Fawcett J K 756 (775) 797
 Featherstone R M 122 (218) 160
 Feder N 73 (794) 75 (455) 89 93
 Fedor I J 698 (1120) 708 (1122 1123)
 717 (1171) 714 (1122 1123 1176) 722
 (1175) 815
 Feil M S 702 (751) 791
 Feinberg S M 773 (1103) 815
 Feldberg W 101 104 (120) 106 (113
 114 115) 115 119 (114 116) 136 137
 (82) 146 147
 Feldberg W J 117 (36 37) 145
 Feldman J D 703 (316) 744 (316) 748
 (317) 793
 Feldman W H 467 (197) 504
 Feldott G 477 (376) 482 (193) 504 508
 Feldstein I M 717 (179) 733 (179) 789
 Feller D D 187 (34) 196 441 (709) 447
 (709) 517
 Fellinger K 435 (89) 436 (89) 507
 Fels G 209 210 931 604 (289) 678
 Fettermann G H 455 (138) 503
 Felts J M 188 (85) 197
 Feng Y S L 739 (79) 805
 Fenwick M 288 (307) 289 (307) 505
 Ferguson D J 673 (81) 674 (81) 679
 Ferguson J K W 490 (530) 516
 Ferguson K A 289 (48) 998
 Fern J 671 (87) 679
 Ferner H 192 (47) 196
 Fernholz F 544 (337) 587
 Ferrari J 794 656
 Fern M G 28 (82 333) 50 56
 Ferstl A F 288 (109) 300
 Fertman M B 838 871
 Fesard A 119 (121) 147
 Feurer M 546 (251) 558 (259) 585 586
 Fevold H L 369 579
 Fickett W 333 (343) 588
 Fieber M H 8,0 (186) 875
 Fielsing U 324 335
 Fienberg R 766 (318) 793
 Fierce W I 771 222 (112) 223 233
 Fieer L F 522 527 (92 96) 5,0 538

- Flück H 239 (97) 272 (55) 273 (55)
 290 (55) 298 299
 Flson R 506 (201) 8,5
 Flter S K 763 (291) 792
 Flvehjem C A 497 (675) 516
 Fly F 400 (61) 427
 Flk R S 285 (380) 307 694 (214) 745
 (621) 791 802 800 (109 208) 872
 870
 Immcus C W 490 (665) 515 665 679
 Imerson G A 489 (182) 504
 Imerson K 841 (214) 875
 Imerson W J 672 (74) 674 (4) 670
 Immes I F 490 (60) 501
 Immelt J W 703 (1232) 719 (1232) 818
 Inami M 60 (188 191 191) 61 (186
 192) 62 (184 189) 63 (191) 78 (519)
 81 (187) 87 94
 Inbick H C 722 (1137) 816
 Enders I 59 (195) 87
 Ingbaek I 117 (80) 146
 Engel C R 566 (147) 567 (147 148) 587
 Engel F I 256 (98) 260 (339) 266 (99)
 299 306 495 (716) 517 718 (878) 720
 (97 296 297 301) 730 (217 298 299
 982 983) 732 (982) 734 (295 300)
 735 736 742 (984) 787 797 808 811
 841 (52) 871
 Engl G I 718 (302 303) 727 793
 Engel L L 545 (318) 587 510 (84) 514
 600 601 605 615 635 (85) 647 (257
 258) 666 (250) 667 (78) 668 657 667
 679 683 706 (1146) 816
 Engel M B 156 160 171 177
 Engel M G 256 (98) 299
 Ingelbach W 241 (100) 299
 Ingelbreth Holm J 753 783
 Ingelman I I 470 (250) 506
 Ingelman M 597 (301) 515 (301) 609
 Ingfeldt B 158 163 171 174
 Ingle I T 693 (591) 801
 Ingstrom A 436 (183) 504
 Ingstrom W W 672 (71) 679 712 (304)
 793 841 (131) 845 (131) 873
 Innor A H 206 (76 77) 233
 Forich S 164 (101) 170
 Intemann C 452 (708) 485 (184) 504
 516 753 (1011) 812
 Ipstein S H 608 (201a 235 236) 609
 (235) 611 (201 234) 647 (201a) 613
 (235) 657 660 661
 Ipstein D 782 (851) 808 838 (85) 81
 Ipstein M 591 (177) 670
 Ipstein I R 721 (410) 725 (410) 796
 Iranko O 417 (62) 427
 Ireach A 673 (80) 679
 Irickson A L 554 (47 294) 563 (47)
 579 586
 Irickson R I 552 (275) 554 (47) 563
 (47) 579 586
 Irshoff B H 238 (101) 299 337 381
 453 (188) 159 (185 186) 504 739
 (305 306) 793
 Irwiner V 78 (197-199 200 201 202-
 204 205 207 516) 79 (196) 87 (207)
 87 88 94 100 (90 91) 103 104 (11)
 108 135 136 144 146 150
 Irulkar S D 187 (52) 196
 Irwan W F 478 (452) 510
 Isamilla R 271 (25) 297
 Isamilla R F 271 299
 Isar S 192 (33) 196
 Isselher A I 746 (307) 747 (304 308)
 793
 Ivers A D 747 (945) 810
 Itheridge I M 716 (309) 793
 Itkin W 100 (94 95) 146
 Ittlinger M C 461 (36) 466 (36) 500
 535 (71) 870
 Ettorre R 518 (91) 581
 Eufinger H 485 (189) 504
 Lusman F I 581 (464) 797
 Ivans H M 175 (7) 195 202 (37 60
 71) 204 (60) 205 (29) 210 (70) 214
 (67) 216 (67) 217 (67) 218 (28) 219
 220 221 (31) 231 237 256 257 (405)
 238 (139 250) 239 (22 107 117 151
 152 247) 240 (151 367) 241 (71 104
 204 249 361) 242 (103 151 152 248
 398) 243 (24 151 152 277 267) 245
 (261 267) 246 (103) 247 (139 237
 247) 248 (24 117 247) 249 (261) 250
 (276 278) 251 (23) 252 (167) 253
 (24) 254 (232) 270 (103) 271 (25 277
 350) 272 (117) 275 (103) 276 (350)
 285 (130) 286 (130 150) 287 (398)
 289 (227) 297 299 300 301 307 308
 304 305 306 307 317 369 381 383
 385 434 (213) 492 (29 500 660) 500
 505 515 516 670 (153) 680 692
 (750) 731 (314) 735 (75) 754 (49)
 786 794 795 867 (187) 876

- Frans I D 770 (1015) 780 (1016) 81°
 Frans R D 438 (111) 456 (340) 50°
 508
 Evans P M 551 (31) 556 (31) 579
 Evans V J 769 (310) 793
 Evan W F 493 (691 695) 516
 Evenari M 25 27 50 53
 Fverall J 551 (311) 793
 Fverett G M 49° (667) 516
 Fverett J W 330 333 (82-84) 334 (233)
 368 371 381 384 385 671 (162 776)
 681 683 853 (130) 869 (130) 873
 Everett V B 316 331
 Fversole W J 391 (74) 4°6 4°7 712
 (375) 795
 Fvertzen A 177 (79) 197
 Fwer P F 398 399 (671 63) 42°
 Eja M 581 (1275) 818
 Ever H 191 (43) 196
- F
- Fager J 458 (468) 459 (468 469) 471
 (470) 488 (466) 510 511 742 (781)
 48° (180 783) 806
 Fahey J L 749 (312) 793
 Fajana S S 771 (61) 273 (67) 291 (65)
 298 691 (831 832) 696 (172) 697
 (110) 698 (170) 705 (170 172 177)
 01 (177) 708 (170) 721 (11° 17)
 725 732 736 (168 313) 753 (177) 781
 (168) 782 793 807 742 (34) 870
 Falk M S 770 (314) 793
 Farah A 706 (475) 712 (475) 796
 Farber R K 243 (772) 254 (772) 03
 614 (157) 681 733 734 (692) 804
 Farber S J 770 (81) 229 305 (22) 476
 Farkas A 517 (58) 581
 Farkas L 577 (88) 581
 Farmer T H 221 (73) 231 216 (85)
 286 (85) 233 (85) 299
 Farnsworth F B 704 793
 Farquhar M G 314 381
 Farquharson R F 857 (53) 871
 Farrar B 729 (97) 787
 Farrar M W 539 (16) 59
 Farrell G I 254 (108) 288 (319) 300
 563 (16) 583 846
 Fatt P 114 115 176 146 147
 Faustini R 78 (705) 87 104 (97) 146
 Fauvet E 361 381
 Favour C B 749 (1174) 770 (707) 804
 817
 Fawcett C H 14 50
 Fawcett D M 441 (190) 442 (191) 462
 (190) 443 (190) 504
 Fawcett D W 861 (238) 876
 Fawcett J K 756 (275) 792
 Feather tone R M 122 (248) 150
 Feder V 73 (294) 75 (455) 89 93
 Fedor I J 698 (1170) 708 (1122 1173)
 712 (1174) 714 (1122 1123 1176) 722
 (1175) 815
 Feil M S 702 (251) 791
 Feinberg S M 773 (1103) 815
 Feldberg W 101 104 (170) 106 (113
 114 115) 115 119 (114 116) 136 17
 (82) 146 4
 Feldberg W J 117 (36 3) 145
 Feldman J D 703 (316) 744 (316) 748
 (317) 793
 Feldman W H 467 (197) 504
 Feldott G 477 (376) 452 (193) 504 508
 Feldstein I M 717 (179) 733 (173) 789
 Feller D D 187 (34) 196 441 (109) 442
 (109) 51
 Fellinger K 435 (89) 46 (89) 50°
 Fels G 209 210 231 604 (289) 663
 Fettermann G H 455 (138) 503
 Felts J M 188 (85) 197
 Feng V S I 739 (759) 805
 Fenwick M 798 (307) 289 (307) 305
 Ferguson D J 673 (81) 674 (81) 679
 Ferguson J K W 490 (530) 51°
 Ferguson K A 259 (48) 298
 Fern J 671 (87) 679
 Ferner H 192 (47) 196
 Fernholz F 544 (332) 58°
 Ferrari J 594 656
 Ferri M G 78 (82 333) 50 56
 Ferstl A E 288 (109) 300
 Fertman M B 838 871
 Fessard A 119 (121) 147
 Feuer M 546 (751) 558 (280) 585 586
 Fevold H L 369 379
 Fickett W 533 (343) 588
 Fieber M H 830 (186) 875
 Fielding U 324 385
 Fienberg R 766 (318) 793
 Fierce W I 221 277 (112) 223 233
 Fieser L F 522 527 (92 96) 550 538

- Irick H 239 (97) 272 (55) 273 (55)
 290 (55) 298 299
 Iron R 856 (201) 875
 Flster S K 763 (231) 792
 Ilvehjem C A 497 (675) 516
 Ili F 400 (61) 427
 Ili R S 285 (380) 307 631 (211) 745
 (621) 791 807 850 (108 208) 872
 875
 Immens C W 490 (665) 515 (68 679
 Imerson C A 489 (182) 504
 Imerson K 841 (211) 875
 Imerson W J 672 (71) 674 (71) 679
 Immes I I 490 (60) 501
 Immett J W 703 (1232) 719 (1232) 818
 Inami M 60 (188 191 194) 61 (186
 192) 62 (184 185) 63 (191) 68 (519)
 81 (187) 87 94
 Inbaek H C 722 (1137) 816
 Lnders J 59 (195) 87
 Ingbaek L 117 (50) 146
 Engel C R 566 (147) 567 (147 148) 562
 Engel F L 256 (98) 260 (339) 266 (99)
 299 306 495 (716) 517 718 (878) 29
 (97 296 297 301) 730 (717 738 299
 982 983) 732 (982) 734 (95 300)
 735 756 762 (984) 787 797 808 811
 841 (52) 841
 Engel G I 718 (302 303) 79 73
 Engel I I 545 (318) 587 590 (541) 594
 630 601 605 615 635 (55) 647 (247
 258) 666 (250) 667 (78) 668 657 66
 679 685 706 (1146) 816
 Engel M B 156 160 1 1 1 2
 Engel M G 256 (98) 299
 Engelbich W 271 (100) 299
 Engelbreth Holm J 753 793
 Engelmann I P 410 (250) 506
 Engelmann M 597 (501) 598 (301) 663
 Engfeldt B 154 163 171 174
 Engle F T 693 (591) 801
 Engstrom A 436 (183) 504
 Engstrom W W 672 (79) 679 712 (104)
 793 841 (131) 845 (131) 873
 Ennor A H 206 (16 74) 233
 Enrich S 164 (101) 170
 Entenman C 452 (708) 485 (184) 504
 516 753 (1011) 810
 Epstein S H 608 (201 235 236) 607
 (235) 641 (201 231) 647 (201 643
 (235) 657 660 661
 Epstein D 782 (851) 808 838 (55) 841
 Epstein M 501 (177) 660
 Epstein I R 724 (410) 725 (410) 796
 Iranko O 417 (62) 427
 Irech A 673 (80) 679
 Erickson A L 554 (47 291) 563 (41)
 579 586
 Erickson R L 552 (275) 554 (47) 563
 (47) 579 586
 Ershoff B H 235 (101) 299 331 381
 453 (188) 489 (185 187) 504 733
 (305 306) 793
 Erssimer A 78 (197-199 200 201 202-
 204 205 207 216) 79 (196) 82 (202)
 87 88 94 100 (90 91) 103 104 (11)
 108 135 136 144 146 150
 Erulkar S D 187 (57) 196
 Erway W F 478 (452) 510
 Escamilla R 271 (25) 297
 Escamilla R F 271 299
 Esser S 192 (33) 196
 Essellier A F 746 (307) 747 (304 308)
 793
 Etess A D 747 (945) 810
 Etheridge F M 716 (309) 793
 Etkin W 100 (94 95) 146
 Ettlinger M G 461 (36) 466 (36) 500
 538 (7) 80
 Ettorre R 548 (91) 581
 Eufinger H 488 (189) 504
 Lusman I P 781 (467) 797
 Evans H M 175 (7) 195 202 (34 60
 71) 204 (60) 209 (23) 210 (7) 214
 (67) 216 (67) 217 (64) 218 (78) 219
 220 221 (31) 231 232 236 237 (405)
 238 (159 250) 239 (77 107 117 151
 152 247) 240 (151 307) 241 (21 104
 204 219 361) 242 (103 151 152 248
 398) 243 (24 151 157 277 267) 245
 (261 262) 246 (103) 247 (139 277
 247) 248 (24 117 247) 249 (276) 250
 (276 228) 251 (27) 252 (167) 253
 (24) 254 (232) 270 (103) 271 (25 271
 350) 272 (117) 275 (103) 276 (350)
 285 (130) 286 (130 150) 287 (398)
 289 (277) 297 299 300 301 307 308
 304 305 306 307 312 363 331 333
 385 434 (243) 492 (29 500 660) 500
 505 518 515 570 (183) 582 602
 (380) 731 (344) 735 (75) 754 (48)
 786 794 795 867 (187) 876

- Foglia V G 671 (86) 679
 Foldes F F 490 (701) 504
 Foley E J 703 (331 332) 703
 Folkers K 221 (81) 222 (8 8 9 87)
 21 (7 47) 226 (7 9 87) 231 232 233
 279 (40 415) 286 (20) 294 (40) 297
 298 308 558 (289) 586
 Folkow B 147
 Folley S J 169 170 171 243 (70) 270
 (10) 298 3 8 339 340 341 347 (54)
 351 (91 107) 352 353 354 355 356
 357 358 (30) 359 361 362 (93 94)
 363 364 365 366 (51) 367 369 372
 375 376 378 379 380 381 382 383
 400 (39 65) 421 (25) 426 427 704
 (50) 431 (334 335) 732 (335) 734
 (51) 784 786 799
 Follis R H 701 (336 337 338) 754 (336
 337 338) 793
 Follis R H Jr 487 (707) 504
 Fondal L 676 (736) 683
 Fonken G S 560 (703) 584
 Fontaine M 398 (65a) 497
 Fontant B J 457 (1067) 814
 Foote F W 499 (196) 504
 Forattini C 488 (703) 504
 Forbes A I 154 (3) 170 703 (339) 718
 (59) 483 (59) 788 794 850 (14) 870
 Forbes T R 668 (81) 679 680
 Forchielli I 638 (96) 647 (96) 648 (96)
 657
 Ford E 451 (15) 454 (15) 500 482
 (15) 785
 Ford W 81 (247) 88
 Ford Z W Jr 491 (699) 516
 Foreman N 606 (158) 481 (634) 789 807
 Forester R H 117 (261) 150
 Forker R F 551 (331) 587
 Forman C W 154 (97) 172
 Forsey R R 833 (142) 873
 Forsham P H 283 (296 307) 291 (233
 309 391) 292 (115) 293 (114 115 308
 309) 294 (114 309) 300 303 305 307
 4 9 (704) 495 (301) 504 507 697
 (341 1150) 698 (345) 709 (885) 718
 (341 1150) 721 (1150) 732 (68 341)
 733 (341) 744 (907) 745 (1150) 748
 (514) 781 (346 513) 686 794 799
 809 816 822 (125) 841 (214) 873 875
 Forsyth B T 706 (342) 794
 Fortier C 844 (56) 871
 Fortune I Y 434 (205) 504
 Foster C G 499 (510) 519
 Foster G I 239 306 443 (706) 475 (56)
 507 504
 Foster R J 37 (53) 50
 Foster W C 437 (445) 510
 Foulks J 674 (58) 680
 Fourman P 707 (343) 721 (343) 753
 (343) 794
 Fowler J L A 716 (157) 789
 Fox C I 772 807
 Fox F 492 (207) 505
 Fraenkel Conrat H 239 (117) 248 (117)
 272 (117) 300 446 (708) 505 731
 (341) 794
 Fraenkel Conrat J 239 (117) 248 (117)
 272 (117) 300 485 (200) 505
 Frame E G 248 (118) 249 (118) 500
 Francis T 749 (12) 785
 Frank A H 338 340 385
 Frankel S 727 (919 937) 728 (918) 810
 Franklin A E 443 (760) 449 (760) 506
 Franklin A L 442 (711 638 710) 446
 (637) 462 (211) 464 (637) 469 (210
 637) 505 515 517
 Franks W R 497 (53) 501
 Franksson C 102 (104) 146 694 (381)
 795
 Frantz R 733 (637) 802
 Fraps P M 333 338 340 382 385
 671 (182) 680
 Fraser A M 393 (66) 401 427 430
 Fraser R 208 232 470 (166) 504
 Frawley T F 291 (391) 292 (115) 293
 (115) 300 307 697 (1150) 698 (345)
 713 718 (1150) 721 (1150) 745 (11 0)
 794 816 841 (57) 842 (215) 871 875
 Fredga A 11 (84 85) 50 536 (104) 581
 Fredericq E 176 (42) 177 (42) 196
 Frédéricq H 131 (174) 147
 Fredrickson D S 697 (11 0) 718 (11 0)
 745 (1150) 781 (346) 794 816
 Free A A 452 (118) 502
 Freed S C 715 (944) 794 810
 Freed V H 5 (86) 50
 Freedberg A S 451 (217) 486 (72) 501
 506 714 (660) 803
 Freedman H H 295 (66) 298 783 (48)
 794

- (152) 546 (89) 548 (91 99) 549 (90
 99) 552 (91) 556 (91 98 100) 558
 (97 152) 566 (96) 581 582 604 (67
 170) 615 (170) 632 (260b) 656 660
 Fieser M 523 527 (92) 570 552 (94) 581
 Fignolo W 480 (140) 503
 Finch C A 748 (514 1141) 799 816
 Finch S C 747 (319) 793
 Findlay C W 767 (320) 793
 Finerty J C 287 (159), 301 671 (208)
 682 758 (321) 793
 Finestone A J 763 (1019) 812
 Fingal F 719 (322) 793
 Fingerman M 61 (75 525) 62 (76 77 78
 79 80 82 83 421) 85 9* 94 116 (35)
 117 (261) 145 160
 Fink K 443 (194) 504
 Finl R M 443 (194) 504
 Finkel A J 768 (1204) 818
 Finland M 270 (200 202, 281 (100) 291
 (202) 302 724 (616) 725 (616) 743
 (617) 761 (617) 779 (614) 780 (612
 613 615) 801 802
 Finnerty J 222 (5) 231 256 (10) 257
 Finney D J 212 231 391 (30) 392 (30)
 393 (30) 426
 Fiori C 366 379
 Fiori W 373 379 381
 Fischel I F 761 (324 376) 774 (88 323
 324 325 376) 780 (901) 787 793 809
 Fischer A 2 (76 77) 3 (76 77) 28 (77)
 49
 Fischer E 533 581
 Fischer F I 104 (149) 147
 Fischer R 531 (102) 581
 Fischer P 78 (25) 84 135 (14) 144
 Fischer W H 698 (173) 806
 Fish C A 265 (93) 299 600 647 (89)
 657 659
 Fish W A 449 (94) 802
 Fish W R 599 605 617 636 (144) 637
 (144) 649 (141) 657 659
 Fisher A 779 (1023) 812
 Fisher C 321 (71) 381
 Fisher D A 492 (414 518 519 520) 509
 512
 Fisher I R 437 (195) 504
 Fisher C 838 (55) 871
 Fisher C R 692 (977) 735 (917) 737
 (234) 791 811
 Fisher J D 270 221 (51) 222 (102) 232
 233 277 (390), 292 (390) 294 (390)
 307 740 (1101) 815
 Fisher L, 764 (311) 793
 Fisher R J 252 (110) 253 (110) 300
 Fishman, J B 202 (115) 203 (115) 204
 (92) 233 234 236 252 (414) 265
 (414) 271 (414) 308
 Fishman I 840 (29) 870
 Fishman L W 669 (85) 679
 Fishman W H 669 (85) 670 (83) 616
 (84) 693 (101) 701 (534) 679 787 99
 Fitch D R 764 (989) 812
 Fitts W T, 715 (1214) 818
 Fitzgerald J D I 771 (746) 803
 Fitzgerald P 499 (196) 504
 Fitzgerald P J 352 (244) 586
 Flanagan C K 729 (728) 804
 Flanagan J B 711 (327) 793
 Flataker L 763 (1916) 818
 Fleischman W 434 (197) 493 (713) 458
 (198) 492 (197) 504 518 653 (370)
 664 851 (234) 876
 Fleisher G A 569 (103) 581
 Fletcher A A 749 (328) 793
 Flink C B 786 (92) 291 (92) 299 691
 (341) 718 (341) 732 (341) 733 (341)
 745 (450) 793 794 797
 Flink M G 286 (92) 291 (92) 299
 Flint A H 461 504
 Flock F V 455 (200) 504
 Florentin A A 286 (391) 307
 Flores I 67 (209 211 212 213) 78
 (212) 79 (212) 81 (210) 88 100 (129)
 101 (129) 105 (122) 135 141
 Florey Elisabeth 67 (212) 78 (211) 73
 (212) 88 100 (122) 104 (129) 105
 (122) 135 147
 Florkin M 79 (26) 84
 Florschutz P A 79 (316) 56
 Flowe B H 756 (949) 811
 Fluckiger F 81 (914) 88
 Fluharty R C 440 (443) 499 (501 557)
 510 512 513
 Fluhmann C F 751 (330) 793
 Flux D S 365 (88) 381
 Fox I P 187 (41) 192 (35 36 38 39
 40) 193 (37) 196 260 (111 113) 261
 (119) 300
 Fogel S 45 (68) 46 (68) 49

- (1050) 741 (1052) 782 (1019 1051)
 783 (367 368 369) 794 795 815
 Gaburda G J 741 (1156) 816
 Gaidum J H 100 (S) 101 (S₁) 105 137
 (S) 136 137 144 147 474 (217) 491
 (211) 505
 Gadeke R 497 (218) 505
 Gaebler O H 243 (122) 248 (16 17 17)
 123) 249 (16 123) 250 (201) 251
 (125) 258 (173) 297 300 303 480
 (439) 510 674 (19) 678
 Gade K 192 (41) 196
 Gainer J H 747 (609) 801
 Gaines W I 399 (73) 400
 Galbraith H W 248 (123) 249 (123)
 258 (123) 300
 Gallagher T F 401 430 528 (116) 509
 (309) 548 (191) 549 (21) 557 (35)
 563 (117) 567 (116) 568 (194 220
 323) 569 (194) 577 (86 113 191 197
 194) 578 (113 114) 579 581 583 584
 587 603 610 611 612 (134) 613 (134)
 616 618 621 633 (99) 635 (98a 99)
 636 (99) 655 657 658 667 (176) 681
 Gallagher U M 597 (217) 661
 Gallone L 494 (219) 505
 Gallup A H 39 (212) 53
 Galston A W 15 (50) 26 (33) 28 (90
 91 93 96) 29 (93) 31 (89) 39 40
 (66) 42 (94) 43 (97) 48 50 54
 Ganong W F 246 (707) 307
 Garcia J 47 (185) 59
 Garcia J F 295 (126) 300 756 (70) 786
 Garcia Lopez G 751 (1075) 814
 Gardiner J E 100 (135) 147
 Gardner L I 653 (301) 664 691 (819)
 714 (1212) 782 (112) 783 (310 771
 1213) 784 (370 771) 795 806 807
 818 848 (61 146) 851 (235) 852 (233)
 871 873 876
 Garlick W R 476 (46) 501
 Garrard A 40 (11) 48
 Garrison W M 436 (274) 506
 Gasie G 751 (10) 785
 Gaskell J F 103 (136 137) 113 147
 Gaskell W H 58 (240) 88
 Gaskill C J 342 371 385 856 (704) 875
 Gassner F X 463 (270) 467 (52) 501
 505 669 (97) 610 (92) 613 (93) 680
 Gastineau C F 784 (71) 795
 Gatti T 253 (409) 307
 Gatica J 549 (268) 586
 Catt egen R 718 (436) 796
 Gauch H G 17 (702) 53
 Caudefroy C 658
 Caudino N M 710 (373) 715 (310) 795
 Gaunt R 246 (94) 268 (94) 278 (127
 128) 281 (127) 299 300 394 (74)
 426 497 463 (221) 487 (221) 505
 665 (94) 680 698 (376 377) 701 (378)
 710 (105) 712 (808) 787 795 807 848
 Gautheret R J 36 (95) 50
 Gav I N 757 (664) 712 (664) 803
 Gav Y 465 (113) 502
 Gebert W H 533 576 (280) 586
 Gejer K 70 (242) 88
 Geiger E 459 (104) 460 (104) 502
 Geigv R 72 (241) 88
 Geilung E M K 190 196 390 (76) 422
 (35) 400 407
 German Q M 780 (613) 801
 Gerst S H 605 (262) 660
 Geller H M 672 (179) 682
 Geller H O 779 (633 1152) 80 816
 Gellert M 176 (28) 196
 Gelluzzi W 494 (219) 505
 Gemmull C I 186 196 490 (200) 223
 224) 505
 Gemzell C A 284 (120) 285 (130) 286
 (130) 300 607 (380) 694 (81) 735
 (319) 795
 Genazzini E 472 (474) 511
 Genell S 848 (67) 871
 Genitis V E 850 (63) 871
 Gephart F T 544 (332) 587
 Gersh I 391 427 436 (208) 437 (276
 277) 448 (227) 505
 Gershberg H 281 (131) 283 (131) 300
 485 (223) 505 737 (268) 799
 Gersich R 758 (505) 759 (504 595) 769
 (594) 801
 Gerke J R 608 (98) 640 (98) 657
 Gerlach U 469 (705) 505
 Germuth F G 763 (385) 711 (382) 772
 (383 386) 773 (384 386 387) 774
 (384) 776 (384) 795
 Gerold C 558 (742) 585
 Geschwind I I 271 (31-33) 231 230
 250 (208) 277 (303) 279 (209) 287

- Freedman R I 751 (503) 798
 Freeman I B, 772 807
 Freeman M 497
 Freeman S 158 (30 52 53) 170 171
 Freiberg I K 252 (141) 257 301
 Freiburg S R 42 (87) 50
 French R C 31 (88) 44 (88) 50
 French T H 353 (218) 354 (18) 379
 381 385
 Frenk S 765 (349) 794
 Frenkel H 697 (431) 796
 Frenkel M 843 (71) 872
 Frentz J 64 (215) 88
 Freud J 238 (119) 239 (211) 271 (87)
 299 300 307 610 (59) 615 (50) 656
 Freudenberg K 191 (43) 196 427 536
 (106) 581
 Frey I 370 382
 Frey J R 674 (242) 683
 Freyberg R H 696 (350) 766 (350), 794
 Freytag L 758 (698) 804
 Friberg O 417 (62) 427
 Frick W 554 (70) 580
 Fridenson A 594 (102) 658
 Fried J 559 (107) 563 (249) 581 585
 608 (232) 609 620 (232) 612 (732)
 657 661 698 (30a 30b) 794
 Fried I 371 383
 Fried P H 347 351 387 460 (58 59)
 871
 Friedberg F 251 (120) 300 798 (352
 729) 794 804
 Frieden E 474 (212) 475 (213) 476
 (213) 505
 Frieden F H 672 680
 Friedgood C F 758 (353) 794
 Friedgood H B 313 346 382 435 (214)
 505 823 (60) 871
 Friedlaender A S 772 (354) 794
 Friedlaender S 772 (354) 794
 Friedlander W J 718 (355) 794
 Friedler G 782 (501) 808
 Friedman B K 104 (317) 151
 Friedman C I 710 (360) 714 (359 361)
 715 (357 358) 794
 Friedman J 753 (1199) 760 (1127) 779
 (418) 796 815 817
 Friedman M 485 (615 616) 514 672
 (218) 682 715 (347 944) 732 (356)
 794 810
 Friedman, M H 345, 387
 Friedman S M 710 (360) 714 (359 361)
 715 (357 358) 794
 Froelich A L 192 (44) 196
 Frohman C I 187 (40) 196
 Fromageot C 177 196
 Fromageot I 403 414 477 429
 Frost R 66 (216) 88
 Frost R C 157 (143) 175, 710 711 (367)
 794
 Fruhman G J 747 (363 364) 794
 Fruton J S 730 (530) 799
 Fry F C 278 (274) 281 (131) 783 (131
 241 242) 300 303 304, 697 (740)
 747 (741) 805
 Fuenzalida, F 670 (90) 697 (365) 680
 794
 Fuerst A 546 (106) 558 560 (149) 581
 582
 Fuhrman F A 397 427
 Fujimaki Y 487 (215) 505
 Fujimoto G I 574 (110 111) 581
 Fukuda S 70 71 (22-224) 80 (219) 88
 141 (125-128) 142 (125-128) 144
 Fukushima D K 544 (707) 546 (206)
 563 (112) 571 (310) 577 (86 113 115)
 578 (113 114) 581 584 587 609
 (99) 612 (134) 613 (134) 616 (99
 100) 618 (100) 621 (100) 633 (99)
 635 (99 99) 657 658
 Fuller H J 32 (102) 50
 Fuller M 771 (788) 806
 Fullerton C W 493 (216) 505
 Fulton G P 713 (1215) 716 (1244 1245)
 743 (366) 794 819
 Funnel J W 671 (91) 680
 Fuoco L 721 (141) 788
 Furter M 576 (276) 680
 Futterman P 190 (87) 197

G

- Garenstroom J H 760 (121) 300
 Cibe M 58 (234 239) 59 (234, 60 63
 (279 233) 65 (14 15-17 18 19 20
 21 22 23 63 64 235 236) 78 (230
 737 235) 79 (225 276 231 232 233)
 82 (239) 83 (239) 88 84 88 138 (130)
 139 (9 10) 141 (129) 144 147
 Gabrilove J L 453 (677) 516 505

- Goodall McC 103 (143) 147
 Goodgal S H 3_u (284) 55
 Goodman I D 479 (439) 510
 Goodman L 457
 Goodman I S 73^o (61 6^o) 998 703
 (173^o) 719 (3^o 1232) 795 818
 Goodwin L G 212 (30) 231 391 (30)
 39^o (30) 393 (30) 4^o
 Goodwin M I 601 (107) 658
 Goodwin R H 74 (97) 50
 Goodwin W F 830 (68 69) 8^o
 Goranson F S 187 (52) 196
 Gorbman A 434 (742) 43_u 438 (243
 244) 446 (597) 501 505 514
 Gordon G S 238 (139) 217 (139) 501
 663 (230) 685 718 (78) 719 (13) 720
 (284 413 414) 785 787 79^o 796
 Gordon A H 154 (36 37) 16_u 166 167
 (36 37 38) 168 170 171 450 (24_u)
 505
 Gordon A S 708 231 273 (48) 299 457
 (137) 4 6 (246) 491 (739) 503 505
 736 (200) 743 (411) 747 (363 364 412
 844) 748 (412 75^o 842) 761 (411)
 783 (348) 790 794 796 805 808 824
 (4^o) 834 (4^o) 871
 Gordon C 418 498
 Gordon F S 293 (140) 501 870
 Gordon H T 126 (206) 147 151
 Gordon M 491 (739) 505
 Gordon M B 669 (72) 679
 Gordon O S 676 (73) 679
 Gordon S 40^o (161) 404 (161 211) 409
 (211) 430 431
 Gordon S A 2 3 5 27 (99) 30 (99) 50
 Gottlieb J 488 (189) 504
 Gottschalk R G 751 (415) 75^o (415)
 796
 Gough J 468 (720) 804
 Gough N 667 (237) 685
 Govaerts J 153 171 440 (507) 511
 Gow R C 154 (87) 172
 Grabar P 772 (416) 796
 Graber R P 558 (119) 581
 Grad B 288 (222) 289 (32^o) 505 475
 (396) 509 623 (200a) 662 713 (417)
 796
 Grady A B 272 (2) 230 233 (11) 997
 422 (12) 4^o
 Graef I 738 (803) 755 809
 Graff J 481 (306) 482 (306) 507
 Graff S 699 (399) 795
 Graffeo A J 759 (419) 796
 Grafflin A I 437 (747 218) 505
 Graham A B 760 (1127) 779 (418) 796
 815
 Graham A R 179 132 (145 146) 147
 Graham G 760 (141) 272 (141) 501
 Graham J B 759 (419) 796
 Griham R M 458 (550) 512 759 (419)
 796
 Granata L 257 (79) 999
 Grandori L 68 (744) 88
 Grant G A 577 (745) 543 (120) 58^o 585
 Grant J K 645 (74) 646 648 (104) 655
 658
 Grassé P P 77 (746) 88
 Grasso R 434 (154) 442 (154) 503
 Graves R C 62 (86 572) 85 94
 Gray B J 501
 Gray C H 843 (70) 87^o
 Gray C L 600 (13) 655
 Gray F 705 (476) 796
 Gray J L 479 (97) 50^o 730 (176) 731
 788 801
 Gray L A 371 (29) 379 671 (34) 678
 Gray R 26 (101) 50
 Gray S J 756 (420 471 422) 796
 Gray S W 81 (247) 88
 Gray W D 248 (142) 283 501 692 (423)
 771 (424) 773 (424) 796
 Green A A 626 (270) 645 (270) 66^o
 Green C D 498 (475) 511
 Green D M 661 705 (426) 706 (425)
 712 (475) 714 (477) 717 (179) 733
 (179) 789 796
 Green H N 761 (478) 796
 Green J 487 (249) 505 843 (71) 872
 Green J A 468 (702) 516
 Green J D 323 324 325 377 (114) 331
 33^o
 Green M 32 (102) 50
 Green M H 455 (138) 503
 Green S 741 (102) 787
 Green T W 470 (250) 506
 Green Armytage V B 595 (105) 658
 Greenbaum A L 243 (146) 248 (148)
 752 (145) 253 (147) 254 (146) 501
 339 (98) 340 341 354 381 382 731
 (334) 793

- (132) 291 (132) 295 (305) 300 303
 305 374 393 421 (170) 450
 Geurrier Y 161 (54) 171
 Geyelin H R 481 (230) 505
 Ghirelli F 78 (25 27 29 206) 84 88
 135 (14 93) 144 146
 Ghosh A 673 (131) 680
 Ghosh B N 207 221 (34 93) 231 233
 233 (135 136) 300 782 (1230) 818
 Ginchetti A 128 (138) 147
 Gibbs C B F 190 (89) 197
 Giffey J W 208 210 (98) 233
 Gifford H 711 (429) 796
 Gilbert C 494 (231) 505
 Gilbert C W 507 (217) 601
 Gilbert J 365 (240) 386
 Gill W M 365 (240) 366 (405) 380 386
 Gillespie J O 770 (502) 798
 Gilliland I C 208 232
 Gillman J 494 (231) 505
 Gilman A 119 (174) 148 427 674 (88)
 680
 Ginnoulhac E 492 (232) 506
 Ginsburg M 393 497
 Girard A 594 658
 Girard R J 270 (259) 303
 Giroud A 691 (389) 739 (388 390 301
 392 393 897) 795 809
 Gish D T 414 (208) 431
 Guttlin D 176 (93) 197
 Gitman L 479 (233) 505
 Gladyshev B N 437 (368) 508
 Glafkides C M 249 (137) 251 (137) 290
 (137) 300
 Glascock R F 353 (15 16 17) 354 357
 (16) 379 734 (51) 788
 Glaser G H 717 (395) 718 (394) 763
 (397) 778 (396) 821 795 799
 Glaser R W 81 (243) 88
 Glassman M D 193 (37) 196 260 (111
 113) 300
 Glaubach S 393 (81) 427 699 (398 396)
 756 (398) 795
 Glaz E 283 (409) 307
 Glendenning M B 670 (188) 689 700
 (815) 808
 Glenn E M 845 (64 178) 871 874
 Gley E 450 (234) 505
 Glick D 691 (400) 743 (973) 795 811
 Glock G E 354 332 781 (401) 795
 Godfrid M 217 (45) 939
 Godley A F 462 (235) 505
 Godlowski Z Z 747 (402 403) 795
 Godwin J T 352 (244) 386
 Coemine J 489 (98) 502
 Goering H I 524 (117) 661
 Goetz F C 683 (1151) 696 (1151) 697
 (1151) 704 (1151) 721 (1151) 727
 (1151) 745 (1148) 765 (1151) 766
 (1151) 768 (1151) 777 (1151) 816
 Goffart M 131 133 (38) 134 (139 141)
 145 147
 Gofman J W 485 (236) 505
 Gold A 824 (156) 875
 Gold N I 670 (95) 680
 Goldacre P L 23 (96) 50
 Goldberg R C 311 (113) 389 441 (795)
 443 (795) 454 (237 238) 505 518 533
 (65) 871
 Goldberg M W 523 (277) 576 (246) 577
 (116) 528 (278) 531 (799) 681 686
 Goldblatt H 715 (404) 795
 Golden A 845 (68) 871
 Golden W R C 721 (405) 796
 Goldfarb W 488 (302) 507
 Goldfine M M 667 (96) 680
 Goldfischer M 481 (609) 516
 Goldie E A G 440 (483) 511
 Goldman L 754 (406) 770 (407 408) 796
 Goldner M G 190 (77 156) 192 (77)
 197 199
 Goldsmith E D 476 (246) 491 (799)
 505 673 (97) 680
 Goldstein M S 182 (81) 183 (50 51 80
 81) 184 196 197 477 (240) 505 726
 (409) 796
 Goldzieher J W 594 658 667 (98) 680
 724 (410) 725 (410) 796 838 (67) 871
 Goldzieher M A 724 (410) 725 (410)
 796 858 (67) 871
 Goler G G 669 (72) 679
 Golla Y M L 278 (138) 300
 Gollan F 497 (241) 505
 Goltz H L 759 (775) 760 (775) 806
 Golub O J 453 (188) 504
 Gomez E T 278 (426) 508
 Gómez Mont J 337 387 796 819
 Gomeche P 763 (182) 789
 Gomori C 98 (142) 109 147
 Good R A 758 (1143) 816

Gut M 549 (90) 570 (192) 58* 586 692
 (34) 656
 Gutfreund H 1 6 (54 55 56) 190
 Gutman A B 447 (270) 473 (270) 506
 Gutsche C D 537 (172) 593
 Gut ell T S 560 (203) 594
 Guttenberg H von 47 50
 Guy elman J B 66 (200) 88
 Gysel H 292 (157) 301 695 (753) 707
 (140) 796 805

H

Habbe I 642 (101) 680
 Hader A 743 (749 1065 1066) 763
 (1067) 766 (1065 1066) 800 814
 Hackel D B 771 (445) 797
 Hackett D I 39 (111 112 112a 113)
 43 (109 110 111 112a) 44 (113 301)
 50 55
 Hackett J W 449 (94) 502
 Haifield C J 767 (1033) 813
 Haifon I 74 (251) 88
 Haenni E O 42 (15) 543 (15) 518
 Hafez F S L 318 382
 Hafkenschie J H 696 (1258) 714
 (1758) 819
 Haft D 183 190
 Hagdahl I 221 (68) 232
 Hagen P 104 (148) 147
 Hagerman D D 669 (102) 680
 Hagerwood M A 787 (98) 787
 Hahn E O 418 (446) 79
 Hahn L 153 (43) 171
 Hahn P 491 (271) 506
 Haines A J 707 (645) 802
 Haines S F 439 (339) 440 (339) 508
 535 (141) 836 (141) 878
 Haines W J 248 (266) 304 604 626
 642 (53 105) 646 (106) 656 658
 Hair L Q 371 383
 Haist P E 674 (130) 681
 Halberg F 745 (447-450) 797
 Haley T J 216 232
 Hahkis M N 357 384
 Halkerston I D K 221 233 288 (304)
 289 (307) 305 644 (40) 6 8
 Halkerston J M 735 (907) 809
 Hall C E 238 (108) 240 (158) 287 (159)
 301 698 (453) 700 (452) 701 (452)
 714 (451 1006) 797 812

Hall M 130 (253) 150
 Hall M G 741 (454) 79
 Hall O 287 (159) 301 698 (453) 700
 (452) 701 (452) 714 (451 1007) 797
 812
 Hall S R 315 382
 Hall T C 421 (84) 42 754 (739) 755
 (455) 797 805
 Hall W H 163 (99) 172
 Hallett J W 779 (456) 797
 Halma N S 314 382 495 (247) 506 782
 (457 458) 753 (459) 797 823 (78
 79) 872
 Halpern B N 674 (103) 680 757 (460)
 774 (461) 797
 Ham A W 161 171
 Ham G C 416 (85) 497
 Hamasaki S 74 (252) 88
 Hamberg U 102 (105) 146
 Hambre P J 610 (132) 675 (132) 680
 Hamburger C 671 (104) 680 742 (62)
 797 841 (226) 8 6
 Hammed K A 137 (134) 14
 Hamilton H C 392 (86) 40
 Hamilton J 837 (81) 869 (80) 872
 Hamilton J B 617 657
 Hamilton J D 741 (746) 805
 Hamilton J G 436 (273 274 275) 500
 Hamilton L D 748 (63) 797
 Hamilton M B 450 (514) 612
 Hamlin A 763 (397) 795
 Hamlin E 104 (149) 147
 Hummett F S 492 (276) 500
 Hammond J Jr 317 341 (126 127)
 382 669 (105) 680
 Hamner C L 42 (2131) 53
 Hamolsky M W 451 (277) 506
 Hanan P 774 (842) 797 808
 Hanaoka K I 65 (370) 82 (371) 91
 Hanaoka T 4 0 (87 88 89) 427 428
 Hancock C P 2 (114) 23 (114) 50
 Hancock J 366 380
 Hancock T J 766 (165) 789
 Handler P 153 (63 65) 154 (64 66 67
 68) 185 166 1 1 716 (466) 797
 Hanenson I B 335 (219) 431
 Hanngren A 468 (278) 506
 Hansch C L 9 (210 211) 10 (211)
 127 (115 209-212) 13 39 (212) 50 53

- Greenberg D M 187 (72) 197 251 (120)
300 502 (215) 602 728 (352) 794
- Greenberg J 270 (81) 999 390 (22) 470
- Greenberg M 187 (41) 196
- Greenberg S M 485 (251) 506 781 (32)
785
- Greenblatt I J 479 (233) 505
- Greenblatt R B 856 (72) 872
- Greene C R 755 (1173) 817
- Greene R R 669 (192) 682
- Greene J A 484 (119) 489 (202) 503
506 708 (143) 788 870
- Greenman I 705 (201) 790
- Greenman F S 202 (37) 230 239 (151)
152, 241 (140) 242 (151 152) 243
(151 152) 286 (150) 301 711 (429)
796
- Greenwald I 154 171
- Greenwood C F 239 306
- Greenwood F C 354 390
- Greenwood J I A 716 (157) 789
- Greaves D P 767 (702) 804
- Greaves J D 202 (144) 253 301
- Greggerman R I 77 (248) 88 103 147
- Gregory F G 32 80
- Gregory F J 777 (1050) 813
- Greenhalgh C W 553 (30) 579
- Greep R O 160 (108 111) 169 (14 15
59) 170 171 172 210 215 (16) 218
231 237 (348) 238 (64) 246 (207) 278
(103 154 155) 282 (102a) 290 998
301 302 306 324 329 332 371 379
382 416 (201) 431 666 (142) 681
690 691 695 (431) 701 (645a) 790
(210) 781 (210) 790 791 796 802
- Greer M A 322 333 (122) 382 441
(747) 455 (747) 457 (253) 461 (36
204) 466 (36) 500 506 517 822 (73)
829 (74) 835 (7 75) 839 (76) 870 870
- Grenell R G 718 (432) 796
- Griesbach W E 163 (99) 172 312 314
315 325 386 391 (163) 430 457 (572)
506 510 573 (160 166) 834 (164) 874
- Griffith P C 771 (912) 809
- Griffith M 169 (53) 196 241 (156) 301
- Grimes H M 236 (47) 270 299
- Grimson H S 756 (949) 811
- Griwald C C 703 (39) 724 857 866
(116) 873
- Grob D 717 (433) 796
- Grobstein C 492 (206) 506
- Groen H 697 (434 435) 796
- Grokoest A W, 718 (436 1169), 753
(221) 754 (221) 790 790 816
- Grollman A, 156 157 171, 509 (793)
647 (293) 665 715 (437) 716 (1109)
751 (415) 752 (415) 796 816
- Grolnick M 751 (438) 796
- Gross F 292 (157) 301 707 (440) 76,
(433) 698 (441), 796 797
- Gross I J 60 (721), 86, 90
- Gross J 151 171 436 (397) 443 (208
200) 445 (208-262 264) 449 (167)
450 (245) 451 (2 7 209) 452 (201
207) 453 (207) 454 (250 260) 453
(263 265) 490 (261 263 264 265
266) 504 505 506 509 761 (442) 797
- Gross R 747 (443) 797
- Gross S 863 (212) 878
- Grossman L B 485 (676) 516
- Grossman M I 704 (67) 779 (69) 786
- Grote I W 401 (116) 403 (116) 478 481
- Cruder H 31 (78) 49
- Grundt H M 572 682 701 (444 1131)
797 815
- Grueter F 347 351 356
- Gummer R H 671 (251 251a) 687
- Gruner P 278 (166) 301
- Grun J A 479 (263) 506 673 (99) 680
- Gryniewicz S F 803 (77) 834 (77) 877
- Gubler C J 748 (463) 797
- Gudernatsch J F 491 (209) 506
- Gudmundson H R 293 (244) 303 696
(725) 755 (720) 804
- Guénal J 496 (583) 513
- Guernsey F S 3 (164) 18 19 (163) 31
(163) 33 (163 164 166) 34 (162) 39
40 (165) 41 (167) 5
- Guha B C 427
- Gudry M A 677 (100) 680
- Guillain J 494 (63) 501
- Guillemin R 246 (760) 303
- Gulland F N 47
- Gunkel J E 21 (100 106) 60
- Gunn D I 75 (249) 88
- Gupta P S 163 (128) 173
- Gur J R N 176 (93) 197
- Gurin S 198 (13 14 121) 195 198 507
619 (170) 660 662
- Gustin S T 480 (436) 510

- 658 660 616 (108) 680 720 (140)
 431 (418 479) 797 798
 Hayashida T 795 (167) 301
 Hayden H S 838 (175) 874
 Hayes D R 211 (184) 302
 Hayes M A 767 (483) 763 (482 483)
 764 (481) 798
 Haynes R 250 (163) 301 623 (121 172)
 676 (48) 657 658
 Hays F J 770 221 (51) 773 (39) 276
 (39) 237 240 (161) 249 (165) 250
 (165) 301 366 (137) 332
 Hays R I 451 741 (107) 787
 Hazard J B 839 (364) 871
 Healdley N I 696 (95) 719 (94) 66
 (95) 787
 Heard R D H 574 (175) 515 (174) 587
 535 596 (1) 599 606 609 631 (174)
 634 (233) 655 658 661
 Heath C 487 (39) 500
 Hebb C O 106 (15 153) 144 147
 Hebbich L 288 (109) 300
 Hébert S 672 (74) 683
 Hechter O 770 (81) 733 257 (743) 785
 (739 777) 290 (201) 294 (783) 702
 303 304 301 (331) 606 673 624 676
 (792) 628 (759 292) 638 (48) 646
 (163) 647 (48) 648 (48 151) 649 (46
 48 259) 650 658 659 660 663 664
 677 680 744 (484) 798
 Heck F J 751 (1102) 815
 Heckel N J 866 (82) 872
 Hedge A N 837 (199) 875
 He J 541 (124 128) 542 567 (130) 510
 (131) 587 698 (1707) 818
 Heftmann E 667 (109) 680
 Hegri E S 100 (154 155) 147
 Hegsted M D 189 (21) 195
 Heideberger M 446 (294) 448 (91
 235 296) 506 507 798 801
 Heideman L M 209 210 37
 Heier A 226 (43) 232
 Heiffer M H 741 (39) 785 786
 Heilman D N 763 (486) 798
 Heilman F R 751 (487) 798
 Heims A 72 (392) 80 (392) 91
 Heinbecker P 270 (411) 30 394 (270
 271 222) 395 (222) 431 459 (501
 507) 511 783 (837) 807 844 (83) 872
 Heinemann M 459 (476) 510
 Heine R W 735 (799 1190 1191) 806
 817
 Heinemann M 831 (179) 863 (159) 83
 874
 Heinzelman R A 104 (756) 150
 Hellauer H I 101 (40) 145
 Hellbaum A 671 (91) 680
 Heller C C 865 (86) 866 (85 87) 877
 Heller F I 840 (84) 877
 Heller G G 864 (144) 868 (144) 873
 Heller H 81 (768 769) 82 (764) 89
 313 798 (964 994) 427 498 709 743
 (489 490) 798
 Heller M 156 171 449 (5) 550 (5) 553
 (5) 554 (5) 558 (5) 558
 Heller R 46 (1197) 51
 Hellman L 592 (83) 609 (99) 610 (99)
 611 (99) 612 (134) 613 616 (99) 633
 (99) 635 (99) 636 (99) 657 658 732
 (491) 798
 Hellstrom J 102 (104) 146 694 (381)
 795
 Helmreich M I 606 (765) 617 (766)
 648 (765 766) 659 660
 Hemberg T 5 (120) 6 (171) 25 (123)
 76 51
 Henning A F 507 703 (532) 719 (497
 493) 735 (493) 798 799
 Hemphill R E 717 (494) 798
 Henbest H B 4 (45) 6 (124) 49 51
 52 (25 26) 553 (0) 554 (31 37) 56
 (31 32) 579
 Hench P S 705 (1080) 707 (1080) 708
 (1080) 709 (1080) 721 (1080) 724
 (1080) 752 (1080) 753 (1080) 765
 (883 1080 1186) 766 (883) 798 809
 814 817
 Henderson J B 764 (509) 798
 Henderson J H M 36 (125) 51
 Henderson J W 768 (498) 798
 Henderson N 160 (22) 170
 Hendricks S B 77 (36) 48
 Heneage P 724 (1059) 1060) 813
 Heni F 748 (166) 301
 Henly A A 221 (20) 731
 Hennemann I H 154 (3) 10
 Henriques O B 750 (499 500) 798
 Henriques S B 750 (499 500) 798
 Henry J A 153 (71 82) 171 172
 Henry M L 183 (50) 196

- Hansel W 331 334 (129) 335 (129) 38°
671 (106) 680
- Hansen B A M 2 (116) 50
- Hanser G 72 (290 291) 89
- Hanson D A 669 (279) 684
- Hanson F R 642 658
- Hanson J 603 (641) 802
- Hansler H 164 (69) 171
- Hanstrom B 58 (253 255) 60 (254 255) 67 (255) 82 (253 255) 88 89
138 (150 151) 147
- Hanze A R 560 (203) 584
- Hara J 67 (256 257) 77 (256) 80 (256) 89
- Hardegger E 579 (279) 586
- Harder R 31 (117 118) 51
- Harding L W 719 (372) 793
- Hardy J D 781 (467) 797
- Hare K 391 (101) 498
- Hare R S 391 (101) 498
- Harfenist F J 176 (59) 177 (58 60) 178 196
- Hargis B J 774 (723) 804
- Hargreaves H P 726 (115) 788
- Harrington C R 443 (282 283) 287
445 (281 286) 447 (289) 448 (287 288 289) 474 (27 279 283 396 397) 475 (280 396 397) 500 506 509
- Harrton L B 544 (207) 584
- Harker J E 77 (259) 89
- Harmon A 248 (17) 237
- Harms J W 579 (260 261) 89
- Harms W 60 (259) 89
- Harnik M 539 (342) 588
- Harper E O 452 (290) 506
- Harris C W 147 282 (81 160) 283 (84) 299 301 370 321 322 324 325 376
327 329 330 363 (55) 380 391 38°
396 (91) 400 (39 92) 417 (93) 476 428
- Harris J I 180 (61) 178 219 232 279 (229) 303
- Harris K D 480 (291) 506
- Harris L J 734 (468) 740 (469) 797
- Harris R 705 (744) 806
- Harris R S 669 (107) 670 (107) 680
- Harris S 770 (470) 771 (470) 772 (470) 773 (470) 797
- Harris T N 713 (3) 770 (470) 771 (470) 772 (470) 773 (470) 785 797
- Harris W 741 (102) 787
- Harrison, C U 768 (471) 797
- Harrison H C, 247 (161) 260 (161) 301
- Harrison J H 696 (472 473) 714 (472) 750 (473) 797
- Harrop G A 493 (216), 505
- Hart D S 317 (135) 318 (136) 38°
- Hart E R 131 (200 201) 148
- Hart P D 777 (474) 797
- Hartman F W 779 (1058) 813
- Hartmann M 604 605 658
- Hartop W L Jr 609 (749) 66°
- Hartroft P M 706 (475) 797
- Hartroft W S 706 (475) 797
- Harvey A M 717 (433) 796
- Harvey W R 75 (762) 89
- Hasegawa K 70 71 (265) 89, 141 147
- Haskell T H 24 49
- Haskin D 754 (476) 797
- Hasman M 43 44 46 (41) 47 (42) 49
- Hassel O 523 (173) 524 579 582
- Hastings A B 186 (104 145 155) 188 (175) 198 199 439 (761) 518 777 (1138 1177) 816 817
- Hatch R N 670 (169) 681
- Haterius H O 245 (336) 252 (336) 306
- Hathway D E 554 (31) 556 (31) 579
- Hatzioles B C 365 (240) 596
- Haugaard N 181 (63) 155 (129 130) 186 (131 132 133) 196 198 257 (3,2 373) 306 727 (1084 1085 1086 1087) 814
- Hauptmann K H 677 (63) 679
- Hauser E A 761 (991) 764 (991) 81°
- Hausler H R 243 (57 54) 249 (54) 257 (54) 265 (52 54) 767 (52) 298
- Hausler E P 594 658
- Havel V J 63 (766) 66 (766), 89
- Haven A C Jr 549 (51) 550 (51) 558 (119) 580 581
- Havens W P Jr 773 (280 4,7) 791 797
- Havinga F 8 (119) 51
- Hawk W D 776 (79) 787
- Hawn C van Z 772 (778) 806
- Hawrylewicz F J 708 210 (98) 214 (99) 216 (99) 218 (99) 233
- Hawkins R D 479 (292 293) 506
- Hawkins W W 737 (868) 808
- Hayano M 606 623 624 (704) 676 (78 204) 631 (204) 632 (701) 645 646 (204) 647 (89 120) 648 (112) 657

- Hirschmann H 57 (158) 57 (156 15
 158 161) 532 (155 159) 540 (164)
 543 (164 165) 561 (159 161) 567
 (161) 563 (160 161 167 163) 587
 588 594 604 605 615 659 664
 Hirschmann R 57 (166) 583
 Hirshfeld A I 774 (14) 785
 Hiscoe H B 111 150
 Hisaw F L 334 338 370 555 555 669
 (114 115 253) 641 (113) 672 680 693
 Hitchcock A F 9 (338 339) 57
 Hjertquist S O 163 (50) 171
 Hoigland H 717 (515 516 517) 719
 (517 518) 720 (1036) 799 815
 Hoir W S 434 (305) 492 (303 304) 507
 Hoberman H D 481 (306) 482 (306)
 507 725 (519 1203) 797 818
 Hobson B M 419 (129) 428 429
 Hoch F L 477 (307) 507
 Hochhauser M 584
 Hoch Ligeti C 716 (309) 753 (570) 793
 799
 Hochster R M 646 (116) 680
 Hodges J P 281 (168 169) 301
 Hodgkin A L 126 (161 167) 148
 Hoher R 176 148
 Hofer P F A 718 (571) 799
 Hofelt B 102 107 (158) 108 (158) 147
 Hoff H C 781 (677) 803
 Hoffman M M 539 (124 176) 606 (124)
 631 (124) 637 (143) 649 (143) 650
 (143) 658 659
 Hoffman W C 718 (572) 799
 Hoffmann E 419 (308) 460 (309) 507
 Hoffmann F J 178 (163) 148 493 (311)
 507
 Hoffmann F 128 (163 204) 148 149
 251 (6 140) 257 (6 170) 260 (170)
 297 301 481 (310) 493 (311) 50
 Hofmann A 136 150
 Hofmann F G 289 (171) 301 666 (142)
 681 695 (431) 701 (645a) 711 (573)
 796 799 807
 Hogben C A M 153 (73) 154 171
 Hogben L 392 (105) 418 419 (106) 428
 Hogberg B 473 (117) 428
 Hogg L W 705 (177) 707 (177) 721
 (177) 737 (177) 753 (177) 789
 Hogness J R 469 (312) 507 735 (574
 525) 799
 Hohmann W 536 (106) 581
 Holbrook W I 777 (111 1099) 755
 (104) 787 814
 Holec H G O 196 (313) 507
 Holden M 779 (576) 799
 Holden W D 764 (164) 766 (165) 769
 (538) 789 799
 Holdsworth M 30 51
 Holland B C 491 (485) 493 (485) 511
 Hollander D H 770 (1160) 816
 Hollander J I 677 (527 578) 766 (527
 528) 797
 Hollander V I 610 (316) 664 676 (120
 560) 705 (557) 706 (560) 707 (557)
 711 (560) 751 (557) 753 (557) 754
 (55) 808 817 845 (901) 877
 Hollander W 154 (77) 172
 Hollenhorst R W 767 (67) 768 (498)
 786 798
 Holley A D 3 (178) 51
 Holley H L 742 (579) 799
 Holley R W 3 (128) 51
 Hollifield C 272 (73) 299
 Hollings H I 278 (416) 251 (416) 309
 665 (764) 684 701 (1718) 818
 Hollinhead W H 322 340 584
 Holm N 440 (140) 503
 Holman H R 730 (530) 799
 Holmes C B 701 (745) 805
 Holmes J H 722 (531) 799
 Holmgren H 208 231 476 (83) 507
 Holmstrom F G 371 583
 Holt L F 751 (950) 811
 Holtermann H 776 (43) 737 256 (177)
 795 (368) 301 306 537 (44) 538 (44)
 579
 Holtkamp D E 507 703 (532) 719 (492
 493) 735 (493) 798 799
 Holtman D F 494 (314) 507
 Holton P 392 (107) 428
 Holtzer A 774 (461) 797
 Holvasz R P 518 (167) 583
 Holzbauer M 419 473 450
 Homan J D H 795 (173) 301
 Homburger F 669 (85) 679 693 (101)
 701 (534) 787 799
 Homiller R I 178 (149) 189
 Honour A J 439 (315 481) 507 511
 Hooker C W 668 680
 Hooten I N 145

- Henry W I 746 (501) 798
Henschler D 77 (270) 89
Hensler \ H 770 (502) 798
Hentstell H H 701 (503) 798
Herdan G 744 (540 1217) 747 (540)
748 (540) 799 819
Herlant M 311, 316 310 382 383
Herling F 549 (175) 583
Herold F T 833 (77) 834 (77) 87
Herrald I 201 (6) 207 (6) 297
Herr M I 549 (132 151) 582
Herrick F H 740 (504) 798
Herring V V 202 (167) 301
Herrold F A 463 (220) 505
Herschberg A D 674 (110) 680
Hershberg F B 529 (213) 539 500
(139) 518 (21 138 242) 575 (135)
576 (135) 582 585 620 (136) 559
Hershberger I G 544 (136) 582 668
(111) 680
Hertig A F 670 (238) 683
Hertz R 672 (112) 680 692 (506 507
977) 735 (977) 737 (234 978) 774
(505) 791 798 811
Hertz S 440 (298 299) 607
Herz J F 556 (90) 581 698 (350b) 794
Herz J F 608 (98) 612 (98) 607
Herzig I T 516 (137) 565 (149 200)
566 (147) 567 (147) 571 (131) 582
586
Herzog H L 550 (139) 558 (138) 582
670 659
Hess B 477 (434 435) 510
Hess G P 221 (13 31 32 41 42) 931
930
Hess M 287 (159) 301 671 (208) 682
Hess S M 542 (140) 582
Hess W C 724 (508) 798
Hesse A 5 (126) 51
Hestrin S 670 (275) 684
Hetzel B S 453 (300) 507
Heusler K 537 (346) 538 (346) 552
(149) 504 (141 143) 582 588
Heusser H 546 (251 202) 552 (140) 554
(141 144 146) 556 (144 146 150)
518 (145) 565 (149 200) 566 (147)
567 (141 148) 582 585 586
Hevesy G 153 (43) 171
Hewer T F 806 (88) 827 (88) 872
Hewl F W 549 (131 151) 582
Heyman A 770 (1015) 780 (1016) 810
Heymann H 538 (152) 552 558 (159)
582
Hickey F C 449 (94) 502
Hickey M D 766 (262) 791
Hickey R C 391 (101) 408
Hidalgo J 764 (509) 798
Hidy P H 189 (64) 196
Higgins G M 698 (512) 746 (274) 741
(510) 701 (511) 702 (213) 772 (274)
782 (512) 791 792 799
Hild W 81 (271) 89 100 113 147 391
(101a) 408
Hildebrandt F 487 (215) 605
Hill A V 408
Hill B R 604 (67 170) 615 (170) 632
(260b) 656 660
Hill D F 27 (111 1099) 750 (104) 787
814
Hill D L 77 (125) 88
Hill F R 760 (1149) 816
Hill C C 455 (138) 503
Hill M 318 (142) 383
Hill S R 490 (301) 507 581 (51) 589
Hill S R Jr 901 (391) 507
Hillarp A A 102 107 (158 109) 108
147 148 321 585
Hillman R W 763 (21) 785
Hills A C 185 (130) 180 (60) 197
198 207 (312) 293 (392) 306 30
459 (204) 504 721 (1080) 748 (514)
794 799 814
Hills O W 846 (89) 870
Hillward I W 496 (313) 501
Himm J M 58 (34) 78 (30) 84
Himwich H F 188 (302) 507
Hinckley G V 703 (1212) 19 (1732)
818
Hindebro H 614 (110) 680
Hine D C 453 (300) 50
Hines M M 116 (35) 143
Hines M N 61 (500) 61 (76) 78 79
81 82) 85 94
Hinman F 851 (16) 870
Hinslaw H C 461 (190) 504
Hirschmann F 604 610 659
Hirschmann F B 577 (158) 29 (157
158 161) 572 (159) 561 (159 161)
562 (161) 563 (160 161 162 163)
560 (154) 58 583

Hunter O B 715 (553) 800
 Hunter S F 732 (551) 800
 Hurloch B 631 (54) 635 (54) 656
 Hursh J B 460 (370) 807
 Hurtz V 335 (77) 470
 Hurvthal I M 837 (91 96) 867 (91)
 872
 Huston T M 332 (151) 385
 Hutelangs I M 765 (200) 791
 Hutchinson J H 837 (137) 873
 Hutchison R 447 (371) 807
 Hutter O F 117 (166) 148
 Huxley A F 176 (167) 148
 Huxley J S 470
 Hyde J F 101 106 (707) 151
 Hyman G A 740 (555 556) 800
 Hysom G 670 (132) 675 (137) 680

I

Ichikawa M 72 (273-277) 80 (770) 89
 Ift J D 371 383
 Ikonen M V 496 (377) 807
 Illingworth B A 243 (183) 757 (183)
 807
 Infant A 470
 Ing H R 177 (167) 123 (168) 124 (168)
 148
 Ing K B 755 (550) 800
 Ingalls T H 241 (184) 807
 Ingbar S H 439 (323) 807 (77) (557)
 774 (616) 775 (558 616) 780 (615)
 724 (616) 775 (558 616) 780 (615)
 800 801 807
 Ingestad T 38 (131) 51
 Inhoffen H H 540 (344) 545 (171) 683
 588
 Ingle D J 187 (139) 198 241 (14) 758
 (1) 259 (1) 266 (185 187 188) 793
 (187) 795 (186 188 189) 796 797
 807 351 374 (161a) 383 671 (110)
 677 (118) 680 700 (560 566) 701
 (560 567) 708 (571 584) 717 (577)
 710 (138) 770 (567 574 576 577 578
 579) 721 (559 573) 727 (577 573)
 778 (581) 779 (570) 736 (580 582)
 757 (563 568 553) 754 (48) 755 (564
 566) 756 (561) 757 (561 569) 758
 (575) 780 (565) 786 788 800 801
 Ingram J M A 14 (81) 80
 Ingram W R 478

Irazu J 871
 Irving G W Jr 401 402 (76) 476 478
 Irving J T 169 (79) 174 439 (707) 805
 Irwin J O 418 (111) 478
 Isaac J P 669 (8) 678
 Isaacson J F Jr 669 (170) 680
 Isacoff I 188 (72) 195
 Iseri O A 154 (110) 160 (108 111) 167
 (110) 168 (110) 177
 Isselbacher K J 631 639 (779) 647
 (298) 648 (778) 649 (778) 663
 Itatani M K 498 (766) 518
 Ito T 75 (778) 89
 Ito V 668 (171) 673 (177) 680
 Itz J 419 (684) 516 828 (108) 837
 (108) 838 (108) 875
 Iversen K 834 (5) 870
 Ivon Sallman I 767 (585) 768 (585) 801
 Ivy A C 478
 Ivy H B 68 (1095) 814
 Iwakura C 66 (279) 89

J

Jackson C M 736 (586) 801
 Jackson N R 703 (1737) 719 (1737) 818
 Jackson S H 751 (1047) 818
 Jackson W P U 865 (104) 875
 Jacob J 720 (587) 801
 Jacob M 483 (477 478 479) 510
 Jacob S 776 (869) 808
 Jacob W 497 (218) 808
 Jacobs E 153 (71 87) 154 (81) 155 160
 171 172
 Jacobs M 488 (676) 516 846 (107) 875
 Jacobs M D 705 (1050) 818
 Jacobs R 595 (127) 603 (127) 658
 Jacobs W P 17 (137) 77 (133) 24 (137)
 51
 Jacobsen P P 606 (133) 673 (133) 674
 (133 150 151 167 168, 675 (133
 167) 616 (168) 618 (151) 658 659
 Jacobsen D 376 377 329 363 (55)
 367 368 380 387 388 400 (39 77)
 476 478
 Jacobson D 787 (160) 801
 Jacot B 705 (101) 807
 Jarvinen K A J 754 (592) 801
 Jaffe H 436 (783) 467 (784) 518
 Jagendorf A T 36 (134 135) 81
 Jager B V 787 (379) 505 730 (967) 811

- Hopke C J 773 (477) 797
 Hopkins T F 230 (81) 233 287 (283)
 294 (283) 304
 Hopkins T R 153 (74 77 78) 160 (77
 78) 162 (77 78) 171, 70
 Hopper A I 492 (316) 507
 Hopwood M L 463 (220) 505 673 (93)
 680
 Horeau A 452 (132 133) 505
 Horecker B J 182 (66 67) 190
 Horlick L 486 (139) 503
 Horne, V 15 (223) 53
 Horsfall F L 779 (630 631) 800
 Horsley V A 492 (317) 507
 Horst W 450 (318) 507
 Horwitt B N 608 617 631 636 (144)
 637 (144) 647 (146) 648 (146) 649
 (144) 659 722 (986) 812
 Horstmann P M 271 (174) 301 648
 (250) 663
 Horwitz J P 512 (11) 554 (11) 506
 (11) 509 (11) 508
 Hotchkiss R D 311 383
 Hotop H 420 (115) 408
 Hotta S 188 (65) 197 576 (46) 579
 Houser H B 778 (446) 797
 Houssay A B 781 (535) 799
 Houssay B A 243 (180) 248 201 (175)
 203 (175) 256 (175 179) 207 260
 (175) 265 (175 177 178 180) 266
 (176 177) 271 301 324 374 383 484
 (319) 507
 Howard J E 153 (74 76 77 78) 154
 160 162 171 172 603 (320) 664 784
 (508) 801 851 (234) 860 (105) 872
 876
 Howard R P 156 (129) 160 173 865
 (91) 872
 Howell C 698 (376 377) 795
 Howell D S 705 (206) 709 (206) 710
 (206) 790
 Howell R W 23 (129) 51
 Howes F I 753 (536) 766 (537 880
 881 888) 767 (93 320 537) 787 795
 799 804
 Hsia R S L 637 659
 Hsiang T T 33 (130) 51
 Hsieh K M 246 (141) 249 (406) 301
 307
 Hsieh W M 101 (69) 146
 Hsu Y T, 753 (570) 799
 Huang R I 664
 Huang Minlon 537 (96) 558 (334) 566
 (96) 571 (168) 581 583 587
 Hubay C A 769 (538) 799
 Hubble D 783 (539) 799
 Huber P 474 (6) 500
 Hubl H, 59 (272) 89
 Hubble J 200 (96) 251 (96), 260 (121)
 299 300
 Huddleston B 182 (81) 183 (50 51 81)
 194 (161) 196 197 199 722 (160) 789
 Hudgins P C 770 (192) 770 (191), 717
 (193 932) 790 810
 Hudson G 741 (510), 747 (540) 48
 (540) 799
 Hudson I 723 (117) 788
 Hudson P B 697 (541) 799
 Hubener H J 647 (2) 651
 Huter F 497 (365) 508
 Huff R I 290 (126) 300
 Huff S J 455 (139) 503
 Huffman M N 560 561 (169 170) 683
 594 599 (293) 647 (293) 659 663
 Hufnagel C A 775 (571) 786
 Huggins C 670 (246) 683 696 (512 743
 544) 697 (542) 700 (543-546) 700
 (545) 784 (1132) 799 800 815 803
 (92) 872
 Hughes J F 829 (157) 874
 Hughes J S 740 (504) 798
 Hughes R E 734 (468) 797
 Huis in t Veld L G 544 545 580
 Huisman T H J 702 (546) 800
 Hull R B 719 (547 708) 800 804
 Hume D M 282 (182) 283 (296) 288
 (265) 307 304 305 330 383 744
 (902) 809 844 (93) 872
 Humel E J Jr 194 198
 Humphrey J H 104 (164) 148 771
 (548) 773 (548) 800
 Humphreys R J 149 740 (549) 800
 Humphreys S R 747 (1101) 815
 Humphries I 561 (208) 684
 Hundley J M 750 (550) 800
 Hungerford G F 744 (551) 746 (502)
 800
 Hunt C 719 (518) 799
 Hunter C D 656
 Hunter J H 550 (236) 585

Hunter O B 775 (553) 800
 Hunter S F 730 (551) 800
 Hurloch B 634 (54) 63a (54) 650
 Hursh J B 460 (700) 800
 Hurst N 305 (20) 400
 Hursthal I M 837 (94 96) 867 (94)
 870
 Huston T M 332 (151) 505
 Hutchings I M 785 (700) 91
 Hutchison J H 837 (137) 873
 Hutchison R 417 (301) 807
 Hutter O F 117 (166) 148
 Huxley A F 126 (167) 148
 Huxley J S 400
 Hyde J E 101 106 (707) 161
 Hyman G A 740 (55a 55b) 800
 Hysom G 670 (130) 67a (130) 680

I

Ichikawa M 72 (273-277) 80 (776) 89
 Ifft J D 301 383
 Ikonen M V 496 (320) 807
 Illingworth R A 213 (183) 200 (183)
 303
 Infante A 400
 Ing H R 100 (167) 123 (168) 124 (168)
 148
 Ing K B 700 (500) 800
 Ingalls T H 241 (184) 300
 Ingbar S H 459 (323) 807 09 (507)
 701 (616) 72a (558 616) 780 (615)
 721 (616) 72a (508 616) 780 (615)
 800 801 800
 Ingestad T 38 (131) 51
 Inhoffen H H 540 (344) 545 (171) 585
 588
 Ingle D J 187 (139) 198 241 (14) 246
 (1) 250 (1) 266 (18a 187 190) 704
 (187) 705 (186 188 189) 206 217
 302 301 374 (183a) 383 671 (119)
 672 (118) 680 700 (500 501) 01
 (500 500) 706 (571 554) 717 (517)
 719 (135) 720 (507 574 576 577 5 8
 579) 721 (509 573) 700 (72 73)
 723 (531) 709 (570) 736 (580 582)
 700 (563 565 583) 754 (48) 700 (564
 566) 700 (561) 703 (561 560) 708
 (565) 740 (56) 786 788 800 801
 Ingram J M A 14 (81) 80
 Ingram W P 423

Irazu J 871
 Irving C W Jr 401 402 (6) 408 408
 Irving J T 183 (49) 170 492 (701) 605
 Irwin J O 419 (111) 408
 Isaac J P 609 (8) 678
 Isaacson J I Jr 609 (120) 680
 Isaacs I 188 (22) 196
 Isert O A 151 (110) 160 (108 111) 167
 (110) 168 (110) 172
 Isselbacher K J 631 639 (705) 647
 (705) 648 (705) 649 (705) 665
 Itatons M K 498 (706) 518
 Ito T 100 (778) 80
 Ito N 605 (101) 673 (100) 680
 Itz J 419 (681) 516 808 (195) 837
 (195) 838 (198) 875
 Iversen K 834 (a) 870
 Ivon Sallman I 767 (58a) 68 (585) 801
 Ivy A C 420
 Ivv H B 788 (109a) 814
 Iwakura C 66 (279) 88

J

Jackson C M 736 (506) 801
 Jackson N R 03 (1230) 719 (1230) 818
 Jackson S H 751 (1047) 815
 Jackson W P U 58a (194) 875
 Jacob J 720 (587) 801
 Jacob M 483 (427 428 409) 610
 Jacob S 776 (800) 808
 Jacob W 497 (218) 608
 Jacobs F 103 (71 80) 154 (81) 105 160
 171 172
 Jacobs M 488 (676) 516 846 (107) 876
 Jacobs M D 0a (1000) 815
 Jacobs R 50a (127) 609 (127) 658
 Jacobs W P I (130) 22 (130) 24 (130)
 51
 Jacobson I I 606 (103) 623 (133) 674
 (133 100 151 167 168) 62a (133
 167) 648 (168) 649 (151) 658 659
 Jacobson D 326 377 379 363 (5a)
 367 368 380 380 383 400 (70 00)
 420 423
 Jacobson D 282 (160) 301
 Jacot B 20a (191) 300
 Jarvinen K A J 751 (500) 801
 Jaffe H 406 (783) 407 (784) 518
 Jagerdorf A T 36 (134 13a) 51
 Jager B V 707 (700) 700 730 (70) 811

- Jahan I 154 172
 Jahn J I 271 (206) 302 696 (158) 789
 Jaler J W 290 (192 193) 302 653 654
 659 693 (511) 742 (588) 782 (1051)
 783 (89 590) 801 813 842 (99)
 847 (97) 851 (101) 850 (98) 861 (100
 101) 872
 Jaques I B 771 (788) 800
 Jaques R 104 (164 169) 148
 Jarnik Y R 665 (220) 671 (220) 683
 Jarpa A 751 (10) 785
 Jarmjarm Y 701 (1158) 810
 Jeanloz R W 606 (133) 623 (133) 624
 (133 167 168 205) 625 (133 167) 646
 (168) 648 (151) 658 659 660
 Jefferies W McK 210 232 254 (191)
 284 (196) 302 694 (1119) 696 (1119)
 815 826 (102) 872
 Jeffers W A 696 (1258) 714 (1255) 819
 Jefferson J M 320 383
 Jeffries J W 720 (576) 800
 Jeger O 531 (102) 534 (62) 536 (62
 264) 554 (144) 556 (144 150) 558
 (145) 581 582
 Jelinek J C 224 (7) 226 (7) 231
 Jelinek V C 279 (40) 294 (40) 298
 Jencsel N D 462 (96) 502
 Jenkins D 284 (106) 291 (309) 293
 (309) 294 (309) 302 306 688 (1151)
 696 (473 1151) 697 (1150 1151) 704
 (1151) 718 (1150) 721 (1150 1151)
 727 (1151) 744 (1147) 750 (473) 745
 (1150) 765 (1151) 766 (1151) 768
 (1151) 777 (1151) 797 816 842 (215)
 845 (173) 874 876
 Jenkins G N 476 (103) 502
 Jenkins R 496
 Jenneret P 747 (308) 793
 Jensen H 175 (69 70) 197 453 (84)
 460 (85) 479 (97) 502 730 (126)
 731 788 801
 Jensen J M 452 (324) 507
 Jerchel D 2 51
 Jessar R A 697 (527 528) 766 (527
 528 669) 799 803
 Jevnik M A 550 (139) 558 (138) 582
 620 (136) 659
 Jewell P A 295 (111a) 428
 Jimenez F V 484 (325) 507
 Joel N 449 (326) 507
 Johnson A S 69 (280) 76 (281) 89
 John J I 227 (114) 873
 Johns H I 252 (144) 253 301
 Johns W I 539 (288 290) 588
 Johnson A D 705 (426) 796
 Johnson A S 758 (255) 759 (291 595)
 769 (594) 801
 Johnson B 697 (170) 698 (170) 708
 (170) 789 842 (34) 870
 Johnson B B 290 (13) 291 (13) 297
 666 (12) 678 702 (596) 801
 Johnson B C 737 (261) 739 (261)
 1181) 791 817
 Johnson B J 705 (177) 707 (177) 721
 (177) 732 (177) 753 (177) 789
 Johnson C 716 (79) 787
 Johnson C F 459 (426) 510 831 (179)
 873
 Johnson G I 436 (324) 507
 Johnson H W 451 (328) 452 (328) 454
 (328) 507
 Johnson J A Jr 553 (740) 588
 Johnson P 770 (933) 810
 Johnson I C 452 (329) 507
 Johnson R B 755 (104) 787
 Johnson R I 270 (238) 303 349 383
 384
 Johnson R H 604 (107) 658
 Johnson R M 488 (330) 507
 Johnson S 423 (112) 428 744 (484) 798
 Johnson S G 848 (103) 872
 Johnson T B 443 (331) 507
 Johnson W S 537 539 588
 Johnston M W 713 (171 174) 724 (173
 174 175) 789
 Johnston S F 753 (170) 788
 Johnston T V 708 (272) 797
 Johnwick E B 833 (38) 871
 Joiner C I 724 (597) 804
 Joly I 76 (287) 82 (287) 89
 Joly P 68 (282) 70 (283) 76 (284 286
 287) 82 (285) 89
 Jones A M 428
 Jones B M 71 (288) 89
 Jones D M 670 (262) 684
 Jones E R H 4 (45) 6 (124) 49 51
 552 (25 26) 553 (30) 554 (31 32)
 556 (31 32) 579
 Jones G F S 784 (598) 801 860 862
 (104) 872

- Jones H B 483 (236) 505
 Jones H S 713 (703) 804 841 (120) 8 J
 Jones H W 837 (122) 8 J
 Jones H W Jr 600 659 647 (123) 680
 Jones I C 239 (197) 2 S (133 197) 301
 302 332 382 393 4 S
 Jones I S 767 (333 599) 768 (583) 801
 Jones J K 667 (124) 680
 Jones J R 781 (64 122) 786 818
 Jones M E 66 (146) 681
 Jones R 522 (49) 549 (175) 580 583
 Jones T I 695 (431) 9
 Jones W J 341 383
 Jongbloed A 672 (236) 674 (236) 683
 Jon on A 13 (137) 51
 Jordan M 471 (664) 515
 Jordan P H 746 (668) 803
 Jordan R M 366 383
 Jores A 470 478
 Jorgensen H E 528 (247) 585
 Joseph N R 156 (47) 171
 Joseph S 246 (91) 268 (91) 299
 Josephson I S 671 (148) 682
 Jo ey A I 447 (600) 801
 Jost A 673 (123) 680 693 801 833 872
 Julah J D 47 (332) 507
 Juhn M 246 (198) 302
 Julia M 9 (137a) 10 (137b 137d) 12
 (137a-f) 14 (137e f) 51
 Julian P I 343 (177 149) 553 (1 6)
 565 (148) 567 (148) 569 (146 180 181
 182) 583
 Jutisz M 443 (338) 447 (338) 513
- K**
- Kabat E A 461 (324) 762 (603) 774
 (324) 793 801
 Kadin M 834 (104) 8 J
 Kadota I 194 (71) 197
 Kaegi K 544 (297) 586
 Kagan J 673 (126) 680
 Kagawa C M 707 (604) 801
 Kahnt F W 641 642 645 (153) 646
 (207) 659 661
 Kaiser P 72 (233) 89
 Kaji S 72 (274 245) 80 (246) 89
 Kalman M 677 (124) 680
 Kamings C E 697 (434) 796
 Hamm O 401 403 428 431 528 (215)
 529 (215) 584 604 605 (185) 660
 Kammlade W G Jr 335 383
 Kammer M I 444 (333) 507
 Kamm S 488 (198) 504
 Kaplan A 560 (38) 871
 Kaplan H S 674 (128) 680 732 (603)
 801
 Kaplan N O 187 (2) 197 737 (606)
 801
 Kaplan S J 719 (944) 811
 Kappas A 698 (1900) 817
 Kapp S A 830 (183) 875
 Kar A B 671 (129 130) 673 (131) 680
 783 (940) 811
 Kark A L 736 (1200) 759 (810) 807 819
 Kark R H 744 (84) 781
 Kark R M 234 (199) 302 742 (608) 745
 (607 798) 801 806
 Karlson A G 777 (609) 801
 Karlson P 72 (900 201) 82 89
 Karnemaat J N 642 (108) 658
 Karnofsky D 497 (334) 507 670 (132)
 673 (132) 680 699 (610 611) 798
 (918) 801 810
 Karp A 482 (333) 507
 Karpel W J 549 (149) 566 (176) 567
 (148) 569 (180 181) 583
 Karvonen M J 417 (67) 477
 Kasdon S C 669 (83) 679 701 (334)
 799
 Kas F H 200 (200) 281 (900) 290
 (201) 291 (702) 302 709 (557) 774
 (616) 723 (538) 743 (617) 761 (617)
 749 (614) 780 (612 613 615) 800 801
 802
 Kassell B 448 (88) 50
 Kassenhar A A H 207 (87) 233 674
 (133) 676 (90) 68 681
 Kastrup H 197 (47) 196
 Katherman R E 600 (13) 655
 Kato J 13 (138) 14 (138) 22 (138) 51
 Katonach F 700 (850) 808
 Katsch G F 743 (411) 761 (411) 796
 Kat oyannis P G 99 (250) 150 404
 (211) 409 (211 211a) 431
 Katz A 531 (183) 583
 Katz B 114 115 126 146 147 722
 (160) 731 (788) 789 792
 Katz L 486 (139) 503 716 (1089) 814
 Katzenellenbogen E R 522 (79) 568
 (323) 580 587 635 (171) 636 660

- Katzman I A 217 23^o 667 (134 135)
 659 681
 Kaufman I H 691 (1053) 813
 Kaufman N, 769 (619) 802
 Kaufmann S 540 (76 77 191) 568
 (269) 569 (209) 580 583 586
 Kavanagh I 21 (97) 50
 Kawakami A 81 (292) 89
 Kawamoto K 497 (336) 50^o
 Kay J H 776 (620) 802
 Kayakan S 741 (6^o1) 80^o
 Kazmin V I 637 (143) 649 (143) 659
 Kersling H H 122 (248) 160
 Keating F R 439 (339) 440 (339 447)
 451 (337 338) 456 (340) 490 (446)
 608 610
 Keating F R Jr 440 (115) 441 (115)
 450 (19) 451 (16 17) 452 (16 17 61)
 453 (61) 500 501 50^o
 Keaty C 671 (91) 680
 Kefford V P 2 (17 18) 3 (17 18) 6 (17
 18) 7 (17) 23 (139) 48 51
 Kehl R 669 (136) 681
 Kelley B 485 (341) 508
 Kelley L W 284 (195) 50^o
 Kelley R M 751 807
 Kelley V C 385 (380) 507 634 (244)
 745 (621) 791 80^o 800 (108 708) 87^o
 876
 Kellgren J H 740 (622) 80^o
 Kellner A W 492 (670) 516
 Kelly F J 870
 Kelly I W 526 (102) 872
 Kelsall M A 743 (623) 752 (186) 789
 802
 Kemmerer A R 727 (111 1099) 700
 (104) 787 814
 Kemp A D 539 (173) 583
 Kemp T 239 (703) 502
 Kempf R 671 (137) 681
 Kendall F C 448 (343 744) 474 (342)
 508 545 (747) 560 (316) 569 (49 50
 103) 571 580 581 584 585 587 618
 (512) 700 (1195) 705 (1050) 707
 (1050) 708 (1050) 709 (1050) 721
 (1050) 724 (1050) 73^o (1050) 751
 (457) 753 (1050) 765 (495 496 497
 655 1050) 780 (565) 78^o (512) 798
 800 802 814 817
 Kendall F E 486 (692) 516
 Hendrick M I, 743 (617) 761 (617) 802
 Henk R 58 (213), 89
 Kennedy B J 206 (90) 233 271 (355)
 273 (355) 506 627 (181) 875
 Kennedy I P 477 (491) 480 (491) 511
 Kennedy G C 669 (39) 678
 Kennedy M H 572 (27) 655
 Kennedy T H 461 (345) 508 602 (6^o6)
 80^o
 Kenny A D 153 (112) 154 (55 110)
 165 166 167 168 (84 110) 169 172
 173 667 (177) 68^o
 Kent C C Jr 669 (135) 681
 Kent L J 797 (111 1099) 787 814
 Kepler I J 601 605 615 616 618 660
 844 845 (109) 848 (132) 850 (13^o)
 873
 Kerby G I 776 (730) 804
 Kerkin M 285 (330) 505
 Kerns K R 30 (67) 49
 Kerpel Fromus E 395 428
 Kerppola W 723 (627) 80^o 846 (111)
 873
 Kerr F H 672 (139) 691
 Kerr H D 875 (112) 873
 Kersley G D 724 (1157) 817
 Kerwin J F 719 (493) 735 (493) 798
 Kestor A S 352 (244) 388 442 (346)
 456 (691) 503 516
 Ketchel M 73 (791 296) 75 (455 456)
 80 (295) 89 93
 Ketellapper H 47 51
 Ketterer B 390 (718) 418 472 475 475
 (218) 428 431
 Keutmann I H 631 (330) 635 (35) 656
 664
 Key J A 767 (628) 769 (628) 802
 Keyes P H 695 (699) 802
 Khalil A 25 (5) 48
 Kheim T 721 (653) 803
 Kiang S I 243 (210) 257 (50) 261
 (210) 261 (80 83) 299 30 726 (714
 215 216) 790
 Kubrick I A 741 (701) 244 (106) 299
 502
 Kuehly C I Jr 474 (48) 476 (47 49)
 501
 Kilbourne I D 779 (630 631 637) 80^o
 Kilby R A 691 (77) 787
 Kilvington T B 396 428

- Kimball C I 100 (73) 19^o
 Kimball R F 80 (707) 89
 Kimura S J 79 (633) 80^o
 Kincaid F D 66 (708) 89
 Kincaid R C 713 (978) 810
 Kinde \ \ 489 (314) 508
 King F 211 219 23^o
 King F H 768 (411) 797
 King J T 751 (731) 805
 King J W 43 (92) 80
 King I S 374 386
 Kingsbury B F 438 (318) 508
 Kingsland \ 437 (798) 519
 Kingston R 592 (27) 655
 Kinnev T D 769 (619) 771 (445) 797
 80^o
 Kinsell I W 271 243 (705) 50^o 696
 (158) 783 725 (636) 779 (428) 733
 (635 634) 781 (634) 80^o 804 874
 (115) 875 (115) 877 (114) 867 (113)
 873
 Kinsella R A Jr 667 (134) 681
 Kipfer H 476 (7) 800
 Kirk R L 473 478
 Kirkwood J G 533 (343) 588
 Kirkwood S 441 (190) 442 (191) 46^o
 (190) 473 (190) 504
 Kirschner I 458 (468) 459 (467 468)
 511 78^o (782 783) 806
 Kit S 729 (639) 802
 Kitchell R I 693 80^o
 Kittinger G W 609 (249) 662
 Kjems E 498 (349) 508
 Kjerulf Jensen K 468 (117 350) 50-
 508
 Kleeman C R 153 (86) 154 (86) 17^o
 Kleigman A M 779 (643) 779 (643) 802
 Klein B V 188 (138) 198
 Klein D T 35 (140) 51
 Klein G 479 (351) 508
 Klein J P 191 (97) 197 479 (357 353)
 508
 Klein M 347 351 (184) 359 360 363
 365 367 369 (185 186) 370 (157) 383
 84
 Klein R 154 (87) 172 271 (219) 273
 (719) 30^o 303 653 (321) 664 693
 (641) 706 (640) 719 (642) 783 (707)
 1211 1213) 802 818
 Klein R L 727 (127) 788
 Klein R M 35 (140 141 147) 51
 Klein S P 182 (81) 183 (81) 197
 Klein W H 75 (170) 36 (168) 5^o
 Kleinberg W 245 (337) 795 (66) 298
 306
 Kleinholtz L H 61 (299) 62 (300) 63
 (266) 66 (716 766 301) 88 89 116
 (170) 135 (171) 148
 Kleinman A 480 (400) 509
 Klemperer W W 706 (424) 712 (475)
 796
 Kligman A M 747 (49) 748 (49) 769
 (49) 786
 Kline D L 482 (354) 508 778 (644) 80^o
 Kline I T 643 (140) 681
 Klinefelter H F Jr 857 866 (116) 873
 Klingsberg A 608 (974) 609 (974) 654
 Klitgaard H M 452 (355) 453 (355)
 474 (48) 475 (356) 476 (46 47) 490
 (50) 501 508
 Klyne W 579 (36 188) 532 536 540
 (185) 544 (70) 563 565 (36 186 189)
 567 (70) 579 583 605 (23) 655
 Knauff R F 707 (645) 80
 Knobil E 246 (707) 30 666 (142) 640
 (141) 681 695 (431) 701 (645a) 776
 802
 Knoepfelmacher A A 478 (750) 517
 Knoll J I 577 (86) 581 592 (83) 65
 Knorpp C T 781 (64 1275) 786 818
 Knowles F G W 60 (113) 61 (112 07
 304 305 66-608) 67 (03) 86 89 90
 111 139 (142) 145 148
 Knowles W S 559 (16) 579
 Knowlton A I 715 (646 647) 716 (644)
 771 (646) 80^o 847 878 844
 Knowlton H I 842 (99) 87
 Knox W E 28 (143 144) 51
 Knuchel W 677 (61) 679
 Kobayashi H 59 (09) 90
 Kober S 594 (60) 656
 Kobernick C D 733 (648) 80
 Kobernick S D 772 (789) 773 (789)
 806
 Koch F C 707 720 (48) 238
 Koch G 548 (171) 583
 Koch H J 61 (310) 80 (310) 90
 Kochakian C D 154 (90) 164 (89 90)
 17 240 (208) 302 479 (357) 480

- (357) 508 616 618 619 618 (30) 51
 156 159) 656 659 674 (269) 676
 (143, 236) 677 681 683 684 729
 (651) 731 (652) 732 733 803 803
 Kocsis J J 221 (51) 333 377 383
 Kocchlin B A 518 (191) 577 (191) 102)
 583
 Kofod Johnsen V 134 (173) 148 160
 Kogl F 4 (145) 10 (146) 61
 Kohler V 497 (303) 503
 Koelle G B 119 (174) 143
 Koelling G 518 (171) 583
 Koenig V 211 219 232
 Koeppe H 457 (738) 503
 Koester H 537 (193) 519 (296) 584 586
 Koff A K 863 (118) 879
 Kogure M 142 (175) 148
 Kohler D 436 (41) 500
 Kohn R R 81 (518) 95
 Koidsumi K 75 (311) 90
 Kottle L S 674 (88) 680
 Kouvusalo M 457 (360) 508
 Kohnsky M 722 (160) 789
 Koller F 546 (105) 581
 Koller G 58 (312) 59 (313) 60 (312) ---
 (314 316) 81 (315) 90 138 148
 Koller M 673 (80) 679
 Kollros J J 77 (504) 94
 Konnerth A 715 (437) 796
 Konsuloff S 419 428
 Konvuchenko T U 496 (322) 507
 Kooistra G 77 (317) 90
 Kopeć S 71 (318) 90
 Kopp L 717 (708) 793
 Kordik P 119 (177) 173 (168) 121 (168)
 148
 Koskowski W 107 (118) 148
 Kostial K 117 (166) 148
 Kountz W B 727 (607) 803
 Kowalewski K 781 (551) 803
 Kracht J 442 (361) 508
 Kraht U 442 (361) 508
 Kraft K 476 (367) 508
 Krah M F 146 188 197 243 (278) 260
 (278) 304 27 (655) 803
 Kraitz F W 154 (141) 155 (142) 157
 (143) 165 (140) 178 751 (285) 797
 Kraitz L 157 (143) 173 302 383 458
 (468) 459 (467 468) 468 (702) 511
 516 782 (782 783) 806
 Kramar J 716 (606) 803
 Kramer H 193 194 195
 Kramer L B 437 (363) 508
 Kramer M 5 (147) 7 10 (147) 51
 Kramer W 188 (8) 195
 Krantz S C Jr 670 (141) 681
 Kraych S 539 (342) 588
 Krebs H A, 187 197
 Krehl W A 737 (991 1773) 738 (1271
 1272) 811 818
 Kreiss R E 213 (24) 248 (24) 753 (74)
 707 735 (75) 786
 Kribben F J 32 (153), 52
 Krieger V I 396 428
 Kringsman B J 131 (179) 148
 Kretchevsky T H 598 (116) 518 (191)
 567 (116 191) 598 (191 770) 569
 (191) 577 (191 192 194) 578 (114)
 581 583 584 609 (99) 610 (316) 615
 (316) 616 (316) 633 (99) 635 (98)
 99) 607 664
 Kroc R L 459 (646) 516
 Krochert G 492 (364) 508
 Krog V 3 (172) 57
 Krohn P L 671 (145) 681 775 (56 607)
 787 803
 Kroneberg G 497 (366) 508
 Kroon D B 708 (119) 300 413 (366)
 480 (366) 508
 Kroukh A 428
 Krueger H 674 (47) 68
 Krusenkemper W 676 (62) 679
 Krupp M A 470 (250) 506
 Kuang M H 397 (124b) 499
 Kuehl I A Jr 222 (8 9) 224 226 (7
 9) 201 239 2 9 (40) 794 (40) 998
 Kuffer S W 120 (181) 126 (180 182)
 148
 Kuhl W J 781 (608) 803
 Kuhl W J Jr 455 (229) 505
 Kuhn W 533 (190) 584
 Kuizeng M H 604 (107) 658 70,
 (843) 706 (94) 801 808
 Kulescha Z 28 (1494) 36 (148 149 150)
 51
 Kulikarni A B 546 (202) 585
 Kulonch I 671 (146) 681
 Kumagi I F 243 (209) 745 (709) 307
 744 (706) 791 803
 Kunderman P J 768 (912 913) 810

Kunkel H C 404 405 (125 126 196)
498 499
Kuntz A 111 (183) 148
Kuntz F 66 (320) 67 (319) 90
Kupperman H S 371 383
Kurith P 522 (146) 554 (146) 556 (146)
582
Kurland G S 451 (774) 486 (72) 501
506 714 (660) 803
Kurtz M 243 (710) 257 (50) 261 (710)
265 (89) 2 0 (81) 222 302 726 (214)
790
Kuseu T 696 (350) 766 (350) 794
Kue G 19 (151) 22 (151) 51
Kutsky R J 37 (152) 51
Kuuisto A 491 (567) 508 782 (661)
803
Kuzell W C 765 (662) 803
Kuzin A M 434 (368) 508
Kuznetzova S M 406 (52) 507
Kvanne F 592 (83) 657
Kyle L H 724 (508) 798
Kyle T I 605 659

L

Lacasseagne A 459 (369) 508 597 (57
55) 656
Lachance J I 734 (663) 803
Lachare A 454 (370 371) 508 517
Lacour F 346 383
Ladl L 428
Lalman A J 315 335 336 345 383
Lafon M 443 (608) 445 (462) 446 (462)
448 (601) 510 514
Lahr E L 344 (722a) 385
Laibach F 32 (153) 50
Laidlaw J C 284 (196) 302 449 (372)
508 688 (1151) 696 (1151) 694 (1151)
704 (1151) 721 (1151) 724 (1151) 744
(1147) 765 (1151) 766 (1151) 768
(1151) 777 (1151) 816
Lair J A 12 (163) 383
Lamaver L B F 207 (52) 233
Lambert E H 494 (343) 508
Lamont W A 214 (99) 216 (99) 218
(99) 233
Lance M 735 (525) 799
Landau D 748 (412 872) 796 808
Landau R L 673 (147) 681 867 (119
120) 873

Landau W W 757 (664) 742 (664) 803
Landauer W 69 (665) 803
Landgrebe F W 102 (754) 150 358 (30)
379 392 (130) 393 (178 217) 396
(217) 417 (178 217) 418 (717) 419
(173 217) 421 (25 134 136 217)
418 (717) 419 (129 217) 421 (25 134
136 217) 422 (131 134) 423 (178 132
217) 424 (131 137 134a 135) 425
(178) 426 429 431 492 (374) 508
Landmann W A 227 (48 49 113) 237
234
Lands A M 134 148
Lane M 762 (916) 810
Lanfranski J 669 (136) 681
Lang A 0 (154) 31 (182) 52
Lang W 546 (233) 585
Langdon R G 592 (160 161) 659
Langemann H 107 (187) 148
Langford H 784 (508) 801 860 (105)
872
Langley J 121 148
Lanman J T 693 (666) 751 (950) 803
811
Laqueur L 239 (211) 302 594 (60 63)
610 (59) 615 (59) 656
Laqueur G I 719 (667) 803 845 (121)
873
Lardella G 531 (102) 581
Lardon A 529 534 (72 76) 540 (195a)
551 (198 199) 579 584
Lardy H A 477 (345 346 473) 482
(193) 489 (173) 504 508 509
LaRoche G 492 (377 348) 508
Larsen P 5 27 (158) 52
Larsen W E 824 (115) 825 (115) 873
Larsh J F Jr 497 (379) 508
Larson F C 450 (144 381) 484 (380)
503 508
Larson H O 540 (344) 588
Lasagna G 343 (87b) 381
Lasater M B 606 (135 265) 648 (265)
659 662
Lasater T F 708 (1045) 813
Lasher N 754 (476) 797
Laski B 751 (1047) 813
Laskin D M 160 172
Last J H 746 (668) 803
Lattes R 766 (669 881) 767 (93) 787
803 809

- Laubach G D 552 (200) 5.6 584
 Laughlin R M 833 (77) 834 (77), 872
 Laurent F 671 (3) 679
 Lauridsen J 749 (670) 803
 Laville G 35 (251) 55
 Lavietes P H 488 (162) 504
 Law L W 752 (671) 803
 Lawler H C 99 (279) 150 402 (164)
 404 (164 203) 414 430 431
 Lawley T B 670 (238) 683
 Lawrence A G 776 (79) 787
 Lawrence J S 747 (600) 801
 Lawrence R T B 235 (212) 239 (212)
 302
 Laws G F 549 (17) 553 (202) 562 (17)
 569 (17) 579 584
 Lawson A 465 (352 619), 466 (649) 508
 515
 Lawson L J 528 (215) 529 (215) 584
 604 (187) 605 (185) 600
 Lawton A H 437 (87 135) 502 503
 Lawdell O E 751 (950) 811
 Layton L L 766 (672) 767 (672) 769
 (672) 770 (672) 803
 Lizaurescu D 164 (155) 174
 Lazarow A 725 (673 674) 803
 Lazarus S S 190 (77 156) 192 (77) 197
 199 243 (404) 248 (404) 256 (213)
 302 307 746 (1180) 817
 La o Wasem E A 395 (137a) 429
 Leach R B 218 (70) 232
 Leaf A 271 (293) 304 712 (675) 718
 (59) 783 (59) 786 803 850 (14) 870
 Leake T B 718 (78) 787
 Leary D C 459 (426) 510 831 (129) 873
 Leath M J 478 (728) 517
 Leatham J H 218 237 778 (127) 281
 (127) 300 359 379 479 (268) 481
 (353) 482 (384) 485 (383) 506 509
 665 (91) 672 680 681 698 (376) 377
 676) 699 (956) 701 (378) 715 803
 811
 Leblond C P 436 (387) 438 (388) 443
 (258 260) 445 (385) 449 (167 258-
 263) 451 (257 259) 452 (257) 453
 (257) 454 (259) 475 (386) 504 506
 509 596 (1) 655 691 (399) 741 (265)
 750, 781 (677) 792 795 803
 Lecoq R 672 (150) 674 (149 150) 681
 Lederer J 459 (300) 494 (399) 609
 Ledogar J A 600 659
 Lee M 238 (335) 247 (335) 248 (335)
 272 (335) 305
 Lee M O 242 (216) 254 278 (214) 307
 Lee N D 187, 197 469 (312) 507 728
 (679 680) 735 (524) 799 803
 Lee R 243 (365) 245 (365) 306
 Lee R F, 718 (681) 803
 Lee L Y 101 (69), 146
 Leech F B 497 (392) 498 (391) 509
 Lees A D 45 (159) 52
 Leffell E C 365 (240) 386
 Legay J M 68 (322) 90
 Legendre R 78 (323 324) 90
 Leger J 772 (652) 803
 Legrand J J 65 (325) 90
 Lehman F P 766 (16 747) 769 (17)
 785 805
 Lehmann J N 705 (426) 796
 Lehniger A 477 (393) 509
 Lei H P 243 (51) 262 (51) 265 (51) 298
 Leiffer E 154 (92) 152
 Leigh C 605 (235 236) 609 (235) 641
 (234) 643 (235) 661
 Leigh H M 641 (59) 657
 Lein A 450 (394) 490 (406) 503
 Leininger R F 436 (327) 507
 Leith W 772 (682) 803
 Leloup J 435 (395) 509
 Le Maistre C 776 (684 685) 747 (683
 685) 803
 LeMay M 761 (324) 774 (324) 795
 Lemon H M 618 618 (163 298) 659
 664
 Lemonde P 777 (685) 803
 Lender T 51 (326) 90
 Lenel M R 65 (159) 87
 Lenel R 67 (327) 90
 Lenhard R 522 549 (5) 550 (5) 553 (5)
 551 (5) 555 (5) 555 (24) 578 579
 Lenk C T 570 (71) 580
 Lennon B E 500
 Lens J 177 (79) 197
 Lenz A L 602 (291) 628 (291) 663
 Leonard S L 278 (217) 303 674 (151)
 681
 Leonhardt K 671 (45) 677 (45) 678
 Leopold A C 2 (261) 3 (164) 18 19
 (163) 22 (14) 25 (1 0) 27 31 (160
 163 171) 33 (160 163 164 166 169)

- 31 (167) 36 (168) 39 40 (165) 41
(167) 48 (161) 48 50 52 54
Leopold I II 734 (1178) 767 (687 688
747) 768 (687 688 689 742) 749
(456) 797 805 805 817
Lepovskij S 672 (185) 674 (185) 682
Lerman J 474 (396 397 635) 415 (396
397) 495 (398 690) 509 515 516 837
(172) 875
Lerner S R 442 (211) 462 (211) 805
Leroy E P 767 (916) 810
Lesh J B 271 222 (117) 232 233 300
Lesnick G 846 (192) 875
Le Tourneau D 3 (172) 52
Levedahl B H 617 (766) 618 (766) 662
Leventhal M L S.6 (123 202) 857 (702)
873 875
Leveque T F 60 (32 328) 84 90 391
429
Lever J D 601 804
Levi C 485 (430) 510
Levi H 397 (15b) 496
Levie L H 238 (119) 241 (273) 300 303
Leviel F 34 (178) 52
Levin L 243 (272) 254 (271 272) 303
504 656 659 674 (157) 681 733 (691)
734 (602) 804
Levin R H 540 (210) 560 584
Levine B 748 (1192) 817
Levine H 672 (179) 682
Levine H D 697 (1057) 713 (1057) 813
Levine M D 437 (399) 509
Levine R 175 (177) 182 (81) 183 184
194 (161) 196 197 198 199 772 (160)
796 (409) 789 798
Levine S Z 454 (540) 513
Levinsky N G 309 429
Levinson G I 673 (17) 678
Levitt J 45 (175 176) 46 (175) 54
Levitt M F 709 (693) 710 (373) 711
(693) 795 804
Levitz M 575 (204) 576 (704) 584
Levy A L 406 429
Levy H 600 606 (133) 623 (133) 624
625 638 (48) 646 647 (48) 648 (48
151) 649 (48 166) 656 658 659
Levy I I 779 (79) 303
Levy L 600 638 (166) 659
Levy R P 284 (195) 304 876 (107) 873
Levy S E 715 (89) 787
Lewis I 718 (804) 804
Lewis A A G 395 (34) 496 711 (142)
788
Lewis I M 737 (491) 806
Lewis J I 744 (497) 806
Lewis I A 676 (770) 682 700 (435) 704
(694 696) 715 (844) 756 (847) 804
805 808
Lewis R A 271 (719) 273 (719) 781
(719) 302 303 604 (56) 615 (86) 653
(371) 657 664 701 (697) 706 (1146)
718 (577) 735 (697) 746 (939) 783
(1700 1211) 799 804 810 816 850
(124) 841 (124) 843
Lewis R B 758 (698) 804
Lewis S L 77 (39) 90 101 (190) 148
Lexander K 7 (177) 55
I Hélias C 68 (330 332 333) 72 (331)
74 (334 335) 90
I Heureux M A 155 (151) 166 168
(146) 172 173 174
Ihoste J 68 (336 337 338 339) 90
I C H 137 (191 192) 148 202 203 204
205 211 214 216 217 219 220 221
225 231 232 236 237 (405) 238 (139)
250) 239 (22 151 152) 240 (362) 241
(21 104 249) 242 (151 152 305) 243
(24 151 152 247 262 365) 244 (106
151 152) 245 (761 262 365) 247 (139
272) 248 (74) 249 (726) 250 (226 228
395 396) 251 (375 413) 253 (24) 254
(232) 258 (1) 259 (1) 265 266 (187)
270 (238) 271 (75 27) 275 (230) 276
(359) 277 (775 303 304) 279 (229)
286 (150) 287 (132) 289 (227) 290
(413) 294 (132 187 225) 295 (162
305) 296 297 299 300 301 304 303
304 306 307 308 349 374 38 388
384 395 (233) 424 (170) 430 431 435
(209) 503 719 (138) 735 (75) 736 (72
580) 744 (551) 746 (552) 748 (905)
754 (48) 765 (904) 786 788 800 809
824 (115) 875 (115) 873
Li T H 101 (69) 146
Libby R L 878 (3) 870
Liberman M J 669 (138) 681
Libretti A 672 (2) 674 (2) 677
Liddle G W 291 (233) 293 (114) 294
(114) 300 303 705 (73) 707 (73) 786
822 (125) 873

- Lieberman S 544 510 (318) 516 (206)
 561 (705) 560 (705) 568 (270) 577
 (115) 581 584 587 590 (172 173)
 592 (173) 602 (231) 601 (244) 615
 616 678 (291) 632 (170) 633 (173)
 630 (171) 656 659 660 667 668 667
 (65 154) 679 681 861 (48) 871
 Lieberman A 672 (79) 679 712 (301) 703
 Light A I 671 (154) 681
 Lightbody H D 480 (400) 509
 Lightfoot I N 556 (201) 584
 Lihenthal J I Jr 700 (1203) 819
 Lihing M 463 (721) 457 (221) 505
 Liljestrand A 153 (30) 172
 Lim R K S 101 (69) 146
 Limasset I 37 (178) 52
 Lin R C Y 147
 Lindberg H A 462 (43) 500
 Lindberg M C 606 (119) 624 (204) 626
 (204) 631 (204) 632 (204) 646 (120
 204) 647 (120) 658 660
 Linder P J 17 (179) 52
 Lindgren F T 485 (236) 505
 Lindhard J 120 (41) 126 (41) 145
 Landsky J F 766 (160) 789
 Ling H W 201 (13) 202 (43) 998
 Linke O 79 (340) 90
 Linnemann H 489 (645) 515
 Linser H 3 (180) 6 (180) 52
 Lapmann F 477 (307 491) 480 (491)
 507 511 737 (606) 801
 Lipner H J 452 (402) 476 (49) 478
 (401) 501 509
 Lipnik M J 747 (49) 748 (49) 769 (49)
 786
 Lipp P G 46 (306) 55
 Ippman R W 772 (699) 804
 Lipsett M N 736 (700 701) 804
 Lipschutz A 672 (150) 681 697 (365)
 794
 Iisak L 101 (193) 148
 Lissner H 851 (16) 870
 Lissitzky S 438 (244) 443 (588) 438
 (744) 445 (589 591 593) 446 (500
 598 602 603 605 606) 447 (586 588
 592) 448 (604) 454 490 (592) 505
 515 514
 Lister A 767 (702) 804
 Lattell R 549 (5) 550 (5 6) 552 (6)
 553 (5 6) 554 (5) 558 (5) 560 (6) 578
 Little C 819
 Little H N 591 (174) 660
 Littner N 161 (61) 171
 Litwans J 747 (94, 916) 810
 Iitzka C 416 (403), 509
 Liverman J I 31 (181 182) 52
 Livermore A H 402 409
 Livingston S 719 (642), 802
 Iltamas R 447 (404 405) 609
 Iobotsky J 616 (52) 656
 Locher F 600 658
 Locke B D 767 (585) 768 (585) 801
 Locke W 713 (703) 804 841 (126) 875
 Lockett M F 202 (231) 503 710 (704)
 804
 Lockie L M 766 (828) 807
 Ioeb E N 715 (646 647) 716 (647) 771
 (646) 802
 Ioeb L H 763 (397) 778 (390) 775
 Ioecken B 500 (60) 580
 Loken F 660 (260) 670 (259) 684
 Loewe L 727 (127) 788
 Loewe S 762 (972) 811
 Ioeu O 98 133 148
 Logan G B 754 (371) 795
 Logan J B 277 (764) 304
 Logan J E 735 (915) 810
 Logan R E 490 (406) 509
 Logaras G 488 (407) 509
 Logemann W 537 (193) 584
 Lojkin M E 710 (802) 802
 Lona F 22 (183) 52
 London F 766 (1140) 816
 Long C N H 247 (161) 252 (353) 760
 (161) 278 (273 274) 281 (131) 283
 300 301 303 306 495 (716) 517 692
 (706 740) 721 (405) 723 (1220) 740
 (833) 741 (700) 747 (741) 798 804
 805 807 818
 Iong D A 498 (408 409) 509 761
 (706a) 804
 Long J A 192 (2) 195 258 (5) 297
 Long J B 770 (707) 804
 Long M F 252 (338) 306
 Longino F H 756 (949) 811
 Longmire W P 775 (130) 788
 Lopez Toca R 751 (1075) 814
 Loomis T A 670 (762) 684
 Loomis W E 52
 Loosli C G 779 (547 708) 800 804

- Lorune J A 286 (38) 307 312 31,
 381 383
 Lorenz H 869 (11) 8 0
 Lorenz N 337 (9-12) 379 410 (18) 449
 (544) 451 (15) 451 (15) 450 (18) 500
 512
 Lotapiech W D 213 (36) 253 (236)
 63 710 (09) 804
 Lott M H 569 561 (169 170) 583
 Lotz W I 158 (96 14) 172 173
 Louchart J 783 (599 590) 801
 Loughhead R 867 (170) 8 3
 Lughlad R 673 (147) 681
 Louis I H 260 (68) 271 (67) 273 (67)
 231 (6) 298 696 (172) 637 (170)
 638 (10) 705 (10 172 177) 707 (107)
 708 (10) 713 (169 171 174) 721
 (172 177) 724 (173 174 175 176)
 732 (172 177) 736 (313) 753 (1 1)
 789 793 841 (35) 842 (34) 870 871
 Lourie E M 101 (46) 145
 Loustalot A J 42 (18) 52
 Love W D 693 (710) 746 (710) 804
 Lovell B J 553 (30) 554 (31 32) 556
 (31 32) 579
 Low B W 181 (82) 19
 Low I F 45 (186) 52
 Lowe C 413 (771) 518
 Lowe C U 724 (711) 748 804
 Lowe J 804
 Lowenthal J 771 (88) 806
 Lowman E W 765 (1186) 817
 Lowry E I 763 (294) 792
 Lowy E 499 (410) 509
 Lozner L L 494 (411) 509
 Lucas C C 254 (53) 298
 Lucas V 458 (469) 471 (469) 488 (466)
 510 511 782 (790) 802
 Luckhardt A B 161 173
 Luckwill L C 26 (187 188) 34 52
 Lulena F P 113 (263) 125 (195) 148
 180
 Ludwig A W 833 (127) 8 3
 Ludwig D 75 (341) 90
 Ludwig D J 613 (156) 681
 Luscher M 77 (343-345) 90
 Luetscher J A Jr 270 (13) 232 (13)
 297 666 (12) 678 702 (222) 790 846
 Luft R 705 (714) 707 (714) 708 (713
 714) 711 (714) 804 818 (4 128) 870
 873
 Lugovoy J H 211 (34) 304 754 (877)
 808
 Lukens I D W 189 (83) 192 243 (256)
 246 (256) 248 (37) 243 (256) 257
 (256) 258 (36) 265 (93) 299 303
 606 (1258) 714 (1258) 819
 Lukes R M 539 (288 290) 586
 Lukoschus F 77 (312) 90
 Lumry R 600 (236) 616 (296) 648 (296)
 663
 Lund C C 496 (41) 509 741 (188) 789
 Lundberg A 131 (196) 133 148
 Lundegårdh H 45 (190) 46 (190) 52
 Lundgren H I 418 (413) 509
 Lundgren M M 210 (202) 291 (202)
 302 419 (614) 801 802
 Lunenfeld B 345 (256) 379
 Luongo M A 790 (715) 804
 Lurie M B 717 (716 717) 804
 Lusted I 492 (414) 509
 Lutwak Mann C 666 (202) 613 (155
 159) 681 68
 Lyall A 481 (249) 505
 Lynn W G 434 (415) 435 (415) 509
 Lynn W S Jr 266 (99) 299 649 (145)
 660
 Lyon D M 414 (21) 500
 Lyon R A 542 383
 Lyon R I 847 (203) 852 (203) 875
 Lyon T P 485 (236) 505
 Lyons C K 756 (949) 811
 Lyons W R 214 32 243 (365) 245
 (365) 248 250 (235) 303 304 318
 349 350 352 361 372 3 3 378 379
 380 381 383 384 385 610 (183) 682
 Lytle D A 641 (88) 652
- M
- McAdams G B 449 (634) 515
 McAlpine H T 788 843 (30) 856 (211)
 864 (211) 870 875
 McArthur J W 155 (138) 165 (135) 173
 213 282 440 (443) 499 (551) 510 512
 Macaulay M H I 39 (130) 429
 McCall F 184 (1175) 817
 McCull P J 667 (146) 681
 McCann S M 283 (240) 284 (108) 300
 303 322 (181) 384 844 (134) 873

- McCarrison R 459 (414) 510
 McCartney W 475 (254) 506
 Macchi I A, 288 (259) 290 (201) 502, 503
 McClendon J I 437 (445) 446 (106) 502 510
 McClintock L 597 (301) 598 (301) 663
 McClosky W T 430
 McClure C D 764 (509) 798
 McCollum I V 462 509
 McConthey W M 440 (447) 490 (446) 510
 MacCorquodale D W 594 (290 318), 660 663
 McCracken B H 421 (181) 427 688 (1151) 696 (1151) 697 (1151) 704 (1151) 721 (1151) 727 (1151) 754 (739) 755 (455) 765 (1151) 766 (1151) 768 (1151) 777 (1151) 797 805 816
 McCrawley E I 718 (432) 796
 McCreery I 673 (140) 681
 McCrory W W 154 (97) 170
 McCubbin J W 497 (506) 511
 McCullagh E P 595 (199) 604 (275) 615 (275) 660 662 624 (136) 867 (136) 873
 McCullagh R D 278 (77) 299
 McCune D J 440 (533) 512
 McDermott W 253 (241 242) 503 602 (749) 747 (741) 776 (685) 777 (685) 803 805
 MacDonald A D 392 (105) 428
 McDonald H J 221 232
 McDonald P R 707 (742) 769 (689 742) 803 805
 MacDonald R H 768 (178) 789
 Macdougall I 260 (29) 272 (29) 297 672 (139) 681
 MacDowell I C 703 (1195) 817 827 (188 189) 875
 MacDowell R J S 128 (203) 149
 McLachern D 490 (92) 62 721 805
 McGavack T T 700 (744) 805
 McGinty D A 441 (450) 442 (448 450) 462 (96 449) 460 (788) 507 510 518 626 645 (200) 660
 McGurr E M 837 (137) 873
 McGoon D C 733 (911) 771 (912) 809
 McCovern N H 592 (27) 655
 McGraw A B 781 (1130) 815 836 (209) 875
 MacGregor A G 489 (759) 518
 McGrew R V 601 (186) 660
 McGuckin W F 318 (227) 685
 McGuire T F 714 (1203) 818
 McHenry L W 489 (57) 501
 McIntosh A V Jr 560 (203) 584
 McIntosh F C 101 148
 McIntosh, H W 260 (243) 503 01 (745) 805
 MeJunkin F A 157 (152 153) 174
 McKay E A 445 (633) 515
 McKay, E M 187 (163) 199 370 (184) 430 722 (1206) 818
 McKergue G 484 (486) 511
 McKendry J B R 824 (136) 873
 McKenzie B F 560 (316) 587
 Mackenzie C G 462 464 (416) 469 (416) 509
 Mackenzie J B 462 464 (416) 469 (416) 509
 Mackenzie H R 272 (50) 273 (50) 290 (55) 298 436 (129 650) 503 510 710 (360) 794
 McKeown T 320 330
 McKibbin J M 737 (211) 790
 MacKinnon D 434 (305) 507
 MacLachlan E A 667 (247) 683
 MacLagan V F 453 (419 710 719) 454 (779) 470 (418 604 683 710 780) 476 (90 418 419 774 777 715) 502 509 516 518
 McLamore W M 537 (316) 538 (346) 688
 McLean C R 771 (746) 805
 McLean F C 156 (70) 171
 McLern I 243 (146) 253 (147) 254 (146) 301 354 33
 McLellan I J 449 (24) 517
 MacLeod I D 288 (36) 298
 McLester J S 742 (529) 799
 McLetchie N G B 189 (31) 190
 McManus J F A 311 384 766 (747) 805
 McMillin G C 486 (40) 500
 MacMullen J 188 (120) 198
 McNamara H 154 (97) 172 501 (12) 870
 McNaught M I 488 (109) 502
 Macpherson H T 177 (84) 19
 MacPhillamy H B 516 (209) 584

- McQuarrie I 192 (88) 197 719 749 805
 McQuillan M T 456 (451) 510 81
 (749) 805
 McRae D H 15 (191 192) 37 (83) 50 52
 MacRae J 46 (306) 55
 McShan W H 316 317 (43 254) 309
 (702c) 316 377 330 334 335 336
 418 (457) 510 647 (170) 631
 MacVicar R 489 (75) 501
 Madden R J 756 (718) 804
 Maddin S 745 (207) 790
 Maddock C I 740 741 (1187) 817
 Maddock W O 218 237 591 660
 Madison I 718 (1009 1010) 812
 Madison I J 745 (719) 804
 Magallon D T 671 (166 167 168) 631
 Magarej F R 768 (720) 804
 Magerlein B J 550 (210) 560 (203)
 584 611 (234) 661
 Magil F B 193 (37) 196 260 (111
 113) 500
 Magnani A 549 (177) 583
 Magnani B 459 (101) 502
 Magnin G L 761 (189) 776 (189) 90
 Magnus Levy A 490 509
 Magrassi B 479 (421) 509
 Magy D B 239 (247) 247 (247) 248
 (247) 503
 Maher A 68 (346) 90
 Maier Huser H 403 (72) 414 (72) 477
 499
 Maigre F 107 (178) 148
 Maisterrena J 293 (308) 305
 Majno G 164 172 173
 Majumdar A C 496 (427) 509
 Malamud B 493 (424) 509
 Malan P 14 (771) 37 (271) 38 (271) 54
 Malcolm J 163 182
 Malev G F 417 (423) 509
 Malkiel S 748 (722) 750 (777) 773
 (1103) 774 (723) 804 815
 Malinow M R 493 (424) 509
 Malm O J 165 (114) 173
 Malmjac J 164 172
 Malmros H 486 (425) 509
 Malone J P 766 (267) 791
 Malpress F H 338 339 341 (99) 347
 (101) 351 (107) 359 761 362 363
 (102) 380 381 387 384
 Mamby A R 712 (675) 803
 Mamoli L 620 660
 Man L B 459 (476) 485 (516 517) 510
 512 831 (179 139) 813
 Mancera O 579 (217) 519 (268) 550
 (313), 554 (313) 556 (2 73) 555
 (313) 580 584 587 587 608 660
 Manchester R C 595 (147) 49
 Mancini R L 417 (751) 517 804 871
 Mandel I 483 (427 428 479) 510
 Mandel P 483 (427 478 479) 510
 Mandel W 293 (714) 303 606 (725)
 755 (775) 804
 Mandell L 552 (65) 530
 Mandl A M 669 (157) 631
 Manicardi I 674 (72) 678
 Mankle E A 765 (667) 803
 Mann C P 863 (158) 874
 Mann F C 467 (19) 504
 Mann J M 642 (108) 658
 Mann P 697 (541) 799
 Mann P J F 106 (193) 148
 Mann T 666 (202) 668 (160) 673 (158
 159) 631 682
 Mansor L F 703 (535) 719 (492 493)
 735 (493) 798 799
 Mansur M 483 (515) 51
 Manunta G 169 17
 Many A S 736 812
 Maqsood M 449 (430) 510 751 (721) 804
 Marano B J 592 (53) 657
 Marasco J P 112 (615) 803
 Maraud R 438 (70) 501
 Marbarek F P 271 272
 Marchetti L 717 231
 Marcus R 440 (619) 514
 Marcus S 702 771 (776) 804
 Marden W G R 347 (171) 384
 Marder S N 752 (605) 801
 Maren T H 437 (431) 510
 Marescaux J 669 (8 9) 678
 Margen S 271 (25 206) 29 30 725
 (6 6) 729 (778) 733 (635 637) 804
 827 (114) 833
 Margolin S 718 (72) 797
 Margulis R R 749 (784) 768 (1006) 806
 814
 Marine D 461 (432) 510 781 (61) 786
 Marinelli L D 499 (557) 513
 Marion G B 37 (172) 384
 Markee J F 322 330 333 (84) 334
 (733) 381 382 385 61 (161 162
 276) 631 633 853 (1 0) 869 (130) 873

- Marker R L 525, 529 (215 219) 510
 561 (217 218 219) 565 (216) 567
 (214) 584 594 601 605 615 660
 Markhardt B 672 (79) 679
 Markoff G \ 476 (433) 478 (433) 510
 Marks H I 258 (216) 262 (215 216)
 263 (216) 303
 Marois M 443 (599) 452 (132 133) 503,
 514
 Maroney S P 80 (347) 90
 Marrazzi A S 151 (199 200 201) 148
 Marré E 24 (233) 33 (197 198) 41
 (131a) 42 (196) 46 (299) 53 55
 Marrian G F 594 655 656 659 660 667
 (163 257) 681 683
 Marsh J B 181 (63) 186 (131 132)
 196 198 257 (372 373) 306 727
 (1086) 814
 Marshall C W 568 (220) 584 624 (168)
 646 (168) 658 659
 Marshall F H \ 317 318 384
 Marshall I M 728 (729) 804
 Marshall W A 321 (1) 379
 Marsili G 674 (164) 681
 Marti H \ 772 (699) 804
 Martin C 745 (1098) 747 (1098) 814
 Martin G J 295 (66) 998
 Martin I G 154 172
 Martin S P 776 (730) 804
 Martinet M 755
 Martinez C 751 (731) 805
 Martius C 477 (434 435) 510
 Marvin H M 728 (33) 785
 Marx L 239 (203) 302 485 (767) 518
 Marx W 202 (71) 237 238 (250) 233
 (247) 240 (362) 241 (21 219 301)
 242 (248) 244 (106 248) 247 (247)
 248 (247) 297 299 303 306 476 (437)
 485 (436 767) 510 518
 Mason A B 783 (732) 805
 Mason F J 769 (619) 807
 Mason H L 569 (50) 572 (225 226) 574
 (226) 580 584 590 (195) 604 605
 615 616 618 634 (198) 635 (194)
 646 (251) 647 (282) 649 (283) 680
 683 697 (1079) 701 (1077 1078) 705
 (1080) 707 (737 1080) 708 (1080) 709
 (1080) 721 (1080) 724 (1080) 725
 (1079) 727 (1077) 732 (1080) 755
 (1080), 765 (882 1080) 766 (882) 784
 (371) 795 805 809 814 841 (131) 844
 (110 197) 84, (131) 848 (132, 850
 (132), 878 875
 Mason R C 670 (165) 681 778 (971)
 810
 Masoro F J 168 (22 85) 195, 197
 Massart L, 121 (202) 148
 Masson G 361 379
 Masson G M C 361 384 700 (735) 714
 (734 736) 715 (733) 740 (963) 805
 811
 Masson I 867 (133), 873
 Mast H 278 (166) 301
 Masters W H 671 (166 167 168) 681
 Mateer F M, 705 (201) 790
 Matelli M 11 (84 85 197) 50 53
 Mathies J C 300 479 (438) 480 (439)
 810
 Mathieson D R 705 (1080) 707 (1080)
 709 (1080) 709 (1080) 721 (1080) 724
 (1080) 732 (1080) 733 (1080) 765
 (1080) 814
 Mathies J C 250 (251) 303
 Matthews J D 208 231
 Matthews J I 361 386
 Matthews S A 455 (668) 516
 Matsumoto F 65 (348) 80
 Matsumoto K 10 (350) 64 (341 350) 90
 Mattis I A 452 (290) 506
 Mattos L 750 (500) 798
 Mattox V R 548 571 572 (225 226)
 74 (226) 584 707 (731) 805
 Maughan G H 478 (175 176) 504
 Maumenee A I 775 (738) 805
 Maurer W 450 (440 441) 810
 Maurice G 774 (461) 797
 Maw W A 672 (195) 674 (195) 687
 Maxwell R 708 (1123) 712 (1124) 714
 (1123 1126) 815
 May C L 423 429
 May R 476 (442) 510
 Mayer A M 27 53
 Mayer G 347 351 (184) 359 360 362
 363 365 367 369 (185 186) 372 380
 383 384
 Mayer J 202 232
 Mayer M M 797
 Maylath F R 767 (687 688) 788 (687
 688) 803
 Maynard F L 743 (366) 794

- Mazoné H 672 (150) 674 (150) 681
 Mead E R 740 (504) 798
 Mead J F 474 (489) 511
 Meadows E C 766 (450) 767 (750) 805
 Means J H 440 (443) 449 (453) 463
 (503) 474 (397) 475 (397) 490 (454)
 495 (398) 509 510 515 780 (751) 805
 838 (138) 8 3
 Medawar P B 75 (86) 787
 Mederos L O 494 (373) 508
 Mednicoff I 749 (1068) 814
 Meeks R C 700 (566) 722 (572) 755
 (566) 755 (575) 800
 Megel H 748 (752) 805
 Méhes G 490 (455) 510
 Mehl J W 488 (462) 518
 Mehler A H 28 (143 144) 51
 Meier A L 479 (456) 480 (456) 510
 Meier H L 534 (256) 585
 Meier J R 441 (795) 443 (795) 518 576
 (46) 579
 Meier R 608 (753) 704 (754) 707 (979)
 805 811
 Meier Burkhardt R E 479 (457) 480
 (457) 510
 Micklejohn H P 725 (755) 740 741
 (755) 805
 Meisinger M A P 222 (8 9) 274 (47)
 531 2, 2
 Meister L 293 (744) 303 696 (725) 755
 (725) 804
 Meister P 604 682
 Meister P D 608 609 (225) 641 642
 643 (225) 657 660 661
 Mites J 35 34 35 35 35 360 361
 36 36 371 375 378 384 386 400
 (531) 480 (158 479 460) 510 512
 670 (169) 681 739 (77 779) 805
 Melampy P M 673 (201) 682 737 (760)
 781 (760) 805
 Melcher G W 779 (761) 805
 Melin H A 837 (43) 871
 Mellinger P C 61 763 (500) 776 (900)
 869
 Melchior J B 352 354
 Mendel B 479 (237) 506 726 (409) 756
 Mendelsohn M L 706 (850) 711 (850)
 860) 868
 Mendenhall P 221 (75) 2, 2
 Mendes E G 61 (351) 66
 Mendes M V 68 (352) 60
 Menkin V 761 (762) 805
 Mentzer C 24 (199) 53
 Menzel K H 540 (344) 588
 Meranze D S 866 (31) 8 0
 Merckenschlager, F 28 (25) 48
 Merkin M 286 (41) 293 (135) 298 300
 Mermelstein H 493 (515) 514
 Merrill P 479 (467) 509 511 782 (782)
 806
 Merritt H H 717 (795 764) 795 805
 Meserve F P 476 (437) 510
 Metcalf W K 744 (1247), 819
 Metuzals J 323 384
 Metznerberg R C 12 (115) 50
 Metzger N 4 6 (51) 501
 Meudt W 7 (315 315a) 56
 Meyer A S 618 624 625 676 631 (204)
 632 638 (203) 646 (204) 660
 Meyer E W 549 (177 179) 569 (180 181)
 182) 583
 Meyer H 72 (93) 91
 Meyer K 236 (103) 242 (103) 246 (103)
 270 (103) 275 (103) 299 501 (228)
 585 766 (880 881 888) 767 (73 599)
 782 801 803
 Meyer P J 744 (765) 805
 Meyer P K 316 326 342 (43 257)
 369 (202c) 376 (189) 380 384 385
 386 478 (452) 510 544 (137) 582
 668 (111) 673 (240) 677 (170) 680
 681 683 702 (604) 707 (1072) 746
 (1070-1072) 747 (260 1071) 752 801
 814
 Meyerheim G 647 (2) 655
 Meyers T H 831 (139) 873
 Messire C 559 (22) 230 326) 70 (221)
 585 589 641 (152) 642 (152) 643
 659 661
 Mishle Veloss C 276 (22) 293 417
 (144) 424 429
 Michael M 77 (76) 806
 Michael M Jr 761 (77) 763 (76) 806
 Michaels G D 271 (276) 273 (275) 502
 725 (636) 807 824 (115) 825 (115)
 877 (114) 873
 Michaelson I C 6 3 (765) 767 (765)
 769) 866
 Michaux A 740 (75) 873

- Michel O , 438 (214) 412 (595) 443 (599)
446 (596 597, 602, 603 605 606) 448
(604) 505 514
- Michel R 436 (803) 438 (214) 412 (599)
413 (588 599) 415 (461 462 589 591
593) 416 (157 158 461 462 590 596
597 598 602 603 605 606) 417 (211
461 586 588 592 600) 448 (158 587
601 604) 453 (585) 451 490 (592)
503 505 510 513 514 519
- Micheli R A 552 (60) 580
- Michie A J 154 (101) 172
- Michon J 58 (353) 91
- Miesha M 164 (155) 174
- Middleton H H , 128 (201) 149
- Middleton S 128 (163 294) 148 149
- Midonkawa O 192 (71) 197
- Miescher K 532 (337) 533 (338) 537
539 (339) 541 (2 3 4 127 128) 542
567 (130 232) 578 582 585 588 641
646 661 698 (773 1207) 806 818
- Miettinen J K 536 (104) 581
- Migeon C 286 (394) 307
- Migeon C J 623 (208) 661 667 (171)
681 714 (1212) 733 (770) 782 (772)
783 (370 771 1213) 784 (370 771) 795
806 818 851 (235) 852 (233) 876
- Migeon C L 653 (321) 684
- Migone A 672 (1) 677
- Milanes F 751 (1075) 814
- Milani L 672 (172) 681
- Miles A A 498 (408 409) 509
- Milkovic S 496 (23) 500
- Miller A M 605 615 616 617 (311)
618 624 (311) 661 668 850 (140) 875
- Miller D C 371 384
- Miller E 529 (36) 565 (36) 579 605 (23)
655
- Miller F R 826 (159) 847 (203) 852
(203) 874 875
- Miller H 489 (759) 518
- Miller I H 42 (900) 53
- Miller J N 777 (1159) 817
- Miller L 625 635
- Miller L L 443 (729) 517
- Miller M 604 (275) 615 (211 275) 616
(274) 618 (211) 661 664
- Miller O B 439 (463) 510
- Miller R A 278 (253 254) 303
- Miller R W 773 (250) 792
- Miller Z 741 (774) 806
- Milnerd A 39 (32) 40 (32 201) 44 (32)
45 (32) 48, 53
- Milnikan C H 835 (141) 836 (141) 875
- Mills F S 833 (142) 875
- Mills J A 536 (233 234) 585
- Mills L C 284 (196) 302
- Milman A F 243 (756 257) 246 (256)
249 (756) 252 (257) 256 (257) 256
(257) 257 (256 257) 258 (256 751)
260 (257) 262 (255) 265 (750) 303
- Milne M D 153 154 160 172
- Minnini G 488 (655) 515
- Minz B 484 (120) 503
- Mira O J 760 (1127) 815
- Miramontes L 541 (74 284) 580 586
- Mirand I A 759 (775) 760 (775) 806
- Mirick G S 770 (776) 773 (776) 779
(1040) 806 818
- Mirsky I A 183 (57) 185 190 (87) 195
196 197 249 (258) 303 331 334 484
(461) 510 727 (113) 752 (113) 788
- Mislow K 528 (235) 585
- Mitchell C A 717 (853) 808
- Mitchell G M 358 (30) 379 421 (20)
422 (131) 423 (132) 424 (131 137
134a) 496 499
- Mitchell H H 483 (479) 511
- Mitchell J W 17 (179 909) 52 53
- Mitchell M I 246 (260) 250 (259) 303
- Mitchell N 479 (233) 505
- Mitolo M 79 (354) 91
- Mitsui T 10 (903 204) 53
- Mittleman A 697 (541) 799
- Mobile A 620 (136) 659
- Modern F W S 293 (244) 303 696
(725) 755 (725) 804
- Moewus F 3 (205) 53
- Moffett R B 556 (236) 568 (237) 569
(237) 585
- Mogabgab W J 771 (1144) 774 (177)
778 (777) 806 816
- Mohney J B 460 (390) 507
- Moinardi C 674 (259) 685
- Mokia G G 70 (355) 91
- Mole R H 460 (465) 510
- Molhuysen J A 843 (74) 870
- Molitor H 393 (81) 427
- Moll F C 772 (778) 806
- Molomut N 250 (369) 306 743 (779)

- 1065 1066) 763 (1067) 766 (1065
1066) 777 (1063) 806 814
Mond B 590 (172) 600 667 (153) 681
Morey W I 352 (214) 380 434 (440)
456 (651) 458 (468 552) 459 (467 468
469) 463 (507) 471 (469) 473 (552)
488 (466) 498 (471) 510 511 516
740 (781) 781 (1749) 782 (780 782
783) 806 819
Monroe B G 435 (472) 511
Monroe R A 450 (473) 452 (473) 453
(473) 511
Montagnani M 489 (610) 514
Montalbetti J A 756 (1969) 759 (910)
807 819
Montalenti G 79 (356) 91
Monte R W 749 (784 1044) 806 815
Mooly L S 156 (88) 197
Moolten S E 763 (942 943) 810
Moon H D 243 (762) 245 (761 262) 304
Moon V H 561 (785) 763 (785) 806
Moore B 695 (786) 806
Moore C R 615 (173) 681 784 806
Moore D F 771 (788) 806
Moore D H 396 (7) 405 797
Moore F D 463 (553) 513
Moore F J 736 (700 701) 804
Moore J A 551 (331) 567 (530) 587
Moore L A 495 (771) 517
Moore M C 830 (183) 875
Moore S 771 (24) 772 (24) 774 (24) 731
779 (88) 799
Moore W W 352 384
Mor M A 672 (174) 681
Morandi L 747 (308) 793
More R H 733 (648) 772 (789) 713
(789) 807 806
Morel C J L 756 (1260) 819
Morel F 452 (132 133) 503 711 (790)
806
Morel G 3 (776 330) 22 (276) 29 (330)
36 (206) 53 55 56
Morelli A 472 (474) 511
Moretti G F 419 (145) 429
Moretti G P 69 (357) 91
Morgan A F 481 (93) 507 737 (791
792) 739 (125) 775 (793) 788 806
Morgan H R 780 (264) 797
Morgan J A 775 (793) 806
Morgulis R R 749 (1044) 813
Morley E H 187 (139) 193 295 (189)
397 717 (777) 720 (574 577 578 579)
771 (573) 722 (573) 800
Morley H U 465 (649) 466 (649) 516
Morrill M F 154 (32) 155 (37) 170
Morris C J O R 221 (70 23) 231 276
(85) 256 (85) 273 (85) 299 423 424
429 653 667 (49) 661 679 783 (732)
805
Morris G M 745 (862) 808
Morris H P 498 (136 475) 503 511
781 (196) 790
Morris M P 42 (185) 57
Morris P 771 (20 23) 731 276 (85)
286 (85) 773 (85) 799 667 (49) 679
Morrison J F 206 733 254 (263) 304
Morton J B 704 (1135) 762 (1135) 816
Morton M L 436 (511) 447 (476) 455
(477) 511 512
Mosca L 258 (58) 298
Mosceley V 484 (478) 511
Moses C 439 (73) 501
Mosher H A 768 (794) 806
Moshin J R 717 (1189) 817
Moskowitz M 704 (62) 732
Moss L D 708 (273) 792
Mote J R 688 (795 796) 806
Mou T W 290 (701) 302
Moulder J W 498 (766) 518
Moulder P V 863 (97) 87
Moyer A W 277 (764) 304
Mozziconacci P 718 (217) 790
Muhlboch O 594 (63) 656
Muehrke R C 284 (199) 307 744 (84)
745 (607 798) 747 (797) 787 801 806
Mueller G C 670 (175) 681
Mueller G P 563 (238) 585
Mueller J F 738 (799) 806
Mueller J M 406 412 429
Mufson M 745 (865) 808
Muhler J C 674 (279) 683
Muir R M 9 (210 211) 10 (211) 12
(115 209-212) 13 33 (707 208) 39
50 53
Mukherjee R 483 (479) 511
Mulay A S 671 (178) 687
Mulberger R D 767 (742) 768 (742) 805
Muhnos M G 710 (802) 736 (800 801)
806 807
Muller R 2 51 698 (753) 805

- Muller W H 40 (68) 46 (68) 49
 Mullick V 183 (51) 196
 Mulvaney T H, 834 (143) 878
 Munday K 421 (134) 422 (131) 409
 Munroe J S 213 (52 54) 249 (54) 257 (54) 265 (52 54) 267 (52) 298
 Munson P L 153 151 (82) 160 165 166 167 (85 106 107) 168 169 172 175 220 231 243 284 (30) 298 301 667 (176 177) 681 683 692 (423) 756
 Mureddu F 169 172
 Murlin J R 190 (73) 193 195 197
 Murneek A F 33 (195 247) 42 (257) 53 55
 Murphy A J 485 (201) 504
 Murphy J 838 (22) 870
 Murphy J B 752 (803 801) 807
 Murphy J S 779 (1040) 813
 Murphy R F J 395 (149) 406 (149) 429
 Murray H 608 (235 236) 609 (235) 641 (234) 643 (235) 661
 Murray H C 608 (201a) 641 (88 201) 612 (201a) 657 660
 Murray I L 605 690 641 642 643 661
 Murray R 758 (805) 807
 Murrill N M 576 (282) 586
 Mushett C W 739 (806) 807
 Mussett M V 154 (37) 167 (37 38) 168 170
 Mu ulun N 837 (96) 867 (95) 872
 Muus J 445 (480) 511
 Myant N B 43 (315 481) 440 (483) 449 (482) 451 (482) 452 (482) 507 511
 Myers J D 484 (484) 491 (485) 493 (485) 511
- N
- Naber G H 668 (272) 684
 Nace H R 225 (239) 585
 Nachmison H 468 (702) 516
 Nachmansohn D 59 (358) 91 106 (205 206) 119 (205) 121 (206) 149
 Nadel I 637 (30) 647 (30) 655 671 (178) 682 740 (807) 807
 Naegle C F 700 (242) 775 (242) 791
 Nagano T 62 (309 360) 66 (359) 91
 Nagareda C S 719 (808) 807
 Nagayama T 425
 Nogle F O 768 (1005) 814
 Nahum I H 672 (179) 682
 Nairn V 744 (1247) 819
 Najjar V A 175 (90) 197
 Nakamura K 3 (337) 5 (337) 7 28 (337) 66
 Nakamura N 64 (486) 66 (486) 83
 Nakashima M 714 (359) 715 (358) 794
 Nalbandov A V 312, 333 (249c) 334 335 336 369 374 383 384 385 386
 Nall D M 616 (156) 618 (151) 619 (157) 648 (156) 659
 Nasset L S 495 (756 757) 518
 Nastuk W I 126 149
 Natelson S 291 (781), 304 754 (871) 803
 Nathanson I T 501 (87) 657 667 (15) 672 (74) 674 (74) 679 751 807
 Naurais I 164 (100) 172
 Nava G 486 (142) 503 672 (180 181) 674 (180 181), 682
 Nayar K K 68 (361-363) 91
 Naylor A W 22 (213) 36 (135) 51 53
 Neal W B Jr 484 (486) 511 759 (510) 807
 Nebel I 548 (171) 583
 Nedzel G A 773 (381) 795
 Needham D M 722 788
 Neely W B 42 (213a) 53
 Nefores M N 483 (155) 519
 Neghme A 751 (10) 785
 Nether R 572 (308 308a) 513 (308a) 514 (308a) 587 672 (286 287) 641 (152) 642 (152) 659 663 671 (182) 682 687 (1079) 707 813
 Nekhorocheff I 715 (217) 790
 Nelson A A 691 (811) 807
 Nelson C T 754 (1136) 772 807 816
 Nelson D H 284 (327) 286 (32 261) 288 (31 265) 291 (31) 297 304 305 674 (281 814 815) 676 (815) 745 (957) 759 (108) 787 792 807 811 845 (64 156 178) 871 874
 Nelson J F 186 (91) 197
 Nelson J W 278 (966) 304
 Nelson M M 385 610 (153) 682
 Nelson N A 543 588
 Nelson R J 771 (816) 807
 Nelson T S 782 (150) 788
 Nelson W A 757 (664) 772 (664) 807
 Nelson W O 319 351 361 385 559 (117) 660 865 (86) 866 (85 87) 867 (145) 872 874

- Nelson W I 311 (150), 429
 Nelson W V 861 (141) 868 (141) 8 3
 Nemir R I 467 (487) 511
 Nes W R 569 (10) 590
 Nettett F B 186 (104) 198
 Nesha C 742 (608) 801
 Nets A T 437 (57 135) 502 602
 Netten C 21 (197) 53
 Neulerger A 411 (488) 511
 Neufeld A H 157 173
 Neurath H 176 (47 147 148) 177 (42)
 190 199 414 429
 Neutlings J I J 295 (173) 301
 Nerverre C 104 (100) 172
 Newbold G T 518 (3) 579
 Newcomb I H 41 (714) 42 (4 14)
 41 (4) 49 53
 Newman H J 672 (743) 683
 Newsom S W 775 (917) 807
 Newton J D 103 (1232) 719 (1732) 818
 Newton W H 256 (768) 304
 Nezamus J I 266 (190) 295 (188 189)
 302 314 (163a) 333 317 (517) 320
 (36) 314 576 507 518 500) 21 (573)
 728 (581) 736 (582) 752 (584) 757
 (569) 800
 Nguyen van Thoi 443 (608) 514
 Nicholas H J 576 (34) 579 598 (19
 715) 635
 Nicholas N J 598 (215) 661
 Nichols C W Jr 312 (217) 315 (242)
 380 438 (796) 519
 Nichols J 691 (818 819 820) 692 (1014)
 807 810 818 (146 147) 875 8 4
 Nicol J A C 59 (364) 91
 Nicolayson R 161 173
 Niergang F 18 53
 Nielson F D 642 (108) 658 107 (642)
 802
 Niemeyer H 417 (491) 480 (491) 511
 Nieva F S 5 30 (99) 50
 Niemann C 474 (489 490) 511
 Nigrelli R F 491 (739) 505 671 (97)
 680
 Nihet Y 482 (492) 511
 Nilson B 107 (150) 108 (159) 148
 Nilsson C 13 (131) 51
 Nishimatsuji J 72 (275 2 6 777) 89
 Nishikawa M 419 (292 293) 568
 Nissim J A 102 (821 822) 807
 Nitsch J P 17 (18) 33 (718) 34 53
 Nittono Y 75 (363) 91
 Nivard R J F 4 (119) 51
 Nixon J N 08 (936) 712 (935) 810
 Nixon W I 478 (728) 517
 Noall M W 604 (761) 662
 Nobel C K 492 (493) 511
 Noble R I 116 (708) 149 292 (69)
 304 687 (823) 704 (823 826) 712
 (824) 732 (134) 756 (82) 788 807
 815 (80) 8 0
 Nolan I 667 (96) 680
 Nontder J F 435 (494) 511
 Norcross B M 766 (828) 677
 Norris E H 438 (115) 511
 Northrop I C 737 1160) 781 (710) 865
 Norton H W 333 (749c) 335 (156) 353
 358
 Nothacker W C 768 (86) 80
 Novack A 128 (829) 807
 Novak V J A 72 (391 36) 80 (366
 367) 91
 Novikov L H K 83 (830) 807
 Nowikoff M 82 (308) 91
 Nowinski W W 137 (155) 503
 Novot D S 591 (740) 585
 Nuttle C F 20 53
 Nutman P 0 23 (720) 30 61 53
 Nyden S I C 3 (184) 682

O

- Oakley W G 260 (141) 772 (141) 307
 Oakwood T S 578 (215) 579 (215) 584
 615 (185) 680
 Oerle I A 06 (584) 521
 Oberman R R 394 (152) 499
 Oberrauch H 552 (744) 585
 Oerti C 128 (704) 119
 Ochs C 41 53
 Ochs I 718 (1010) 812
 O'Connell I W 778 (766) 304
 O'Connor D 450 (745) 505
 O'Connor W J 499
 Oda J M 295 (196) 200
 Odeblad I 431 (496) 511 609 (100) 6 8
 Oell C A 166 (127) 173
 Oell R T 767 (625) 802
 Oell T T Jr 449 (497) 511
 O'Donnell A 102 (127 128) 609 (121)
 653

- O'Donnell W M 691 (831 832) 807
 O'Donovan D K, 261 (270) 253 (170) 503
 Oesterling M J 740 (833) 807
 Östland E 77 (363) 81 103 131 (210) 149
 Ogden G F 437 (87 140) 502 503
 Ogilvie A I 461 (498) 474 (498) 511
 Ogilvie R F 257 (271) 266 (271 272) 304
 Ogryzlo M A, 749 (378) 793
 Ojers G W 775 (834) 807
 Okey R 672 (185) 674 (185) 682
 Oldham F K 390 (76) 427 429
 Oleson J J 277 (261) 304 721 (835) 807
 Olhagen B 750 (836) 807
 Olivecrona H 848 (4 128) 870 873
 Oliver I 746 (601) 798
 Oliveureau M 435 (395 500) 492 (499) 509 511
 Oliveto I P 520 (213) 558 (241 242) 565
 Ollajos R W 154 173
 Olmsted P C 594 (87) 657
 Olsen L J 719 (322) 793
 Olsen N S 194 (92) 197
 Olson J A 768 (1096) 814
 Olson K J 673 (46) 678
 Oneley J L 176 (93) 497
 Ondarza R 447 (405) 509
 O'Neal I W 459 (501 502) 511 783 (837) 807
 Oneson I B 545 (48) 580 667 (186) 682
 Opdyke D F 246 (312) 305
 Opienska Blauth J 166 173
 Oppenauer R 615 681
 Oppenauer R V 552 (244) 585
 Oppenheim E 717 733 (838) 807
 Oppenheimer I 278 (128) 300
 Oppermann A 31 (117) 51
 Opsahl J C 278 (273 274) 304 763 (839 1039) 807 813
 Orbison J I 766 (165) 789
 Ord W M 481 (503) 511
 Ormsby H L 767 (840) 770 (840) 807
 Orr K D 696 (24) 714 (21) 745 (24) 785
 Ortavant R 339 (576) 380
 Orten J M 187 (45) 196
 Osato R L 409 431
 Osbond J M 552 (56) 557 (56) 580
 Osborn C M 305 406 710 (106) 787
 Osborn, M M I 608 (201a) 642 (201a) 660
 Osborne D J 25 (221) 53
 Osborne M M Jr 154 (32) 15, (32) 162 (32) 170
 Oshry I 436 (56) 501
 Osnos M 796 (841) 808
 Osterberger A I 448 (344) 508
 Oswald A 447 (504) 511
 Ott I 399 (155) 429
 Ott W H 780 (901) 809
 Ottaway J H 256 (275) 260 (275) 304
 Ottinger B 763 (385) 712 (386) 713 (384 386 387) 774 (381) 776 (384) 795
 Otto G F 437 (431) 510
 Otto H 464 (505) 511
 Otto J F 252 (276) 289 (276) 304 725 (558) 800
 Ottolenghi A 78 (207) 88
 Otsu T 65 (310) 82 (371) 91
 Oughton D I 504 (31) 556 (31) 579
 Outshoorn A J 102 (73) 146
 Overbeel G A 295 (173) 301
 Overbeek J A 304
 Overholser M D 278 (426) 308
 Overman J R 774 (812) 797 808
 Overman R R 489 (171) 504 706 (205) 708 (205) 710 (205) 711 (327) 733 (1254) 790 793 819
 Overton E 429
 Owen E C 479 (110) 487 (110) 488 (109) 504
 Oyaert W 417 (159) 430
 Oyama J 763 (385) 772 (386) 773 (386 387) 774 (465) 795 797
 O'Leary K 69 (372) 91
- P**
- Pabst M I 703 (843) 808
 Pace M 670 (4) 677
 Pacheco H 2 63
 Padawer J 747 (844) 808
 Paesi F S A 671 (187) 682
 Pagan C 42 (185) 52
 Page E 375 384 734 (663) 803
 Page I W 670 (188) 682 700 (845) 808 864 (147) 874
 Page I H 100 (276) 104 (276) 105 135 136 149 150 493 (128) 497 (506)

- 503 511 700 (735) 704 (695 696)
 14 (131 736) 715 (816 817) 750
 (814) 756 (817) 804 805 808
 Page J I 452 (118) 502
 Lahaut J 410 (507) 511
 Paine V I 111 (115) 149
 Pakesch I 435 (51) 436 (59) 502
 Palade G I 107 149
 Palay S L 107 (112) 149
 Palm L 479 (438) 480 (439) 510
 Palmer A 663 (113) 684
 Palmer J G 254 (327) 305 45 (957)
 811
 Palmer R A 747 (131) 788
 Palmer W G 876 (102) 872
 Palmer W W 416 (294) 447 (270) 415
 (94) 473 (10) 483 (108) 506 510
 Palmieri M L 488 (134) 505
 Pan S Y 215 (118) 218 (118) 34
 Panisset M 777 (686) 803
 Panos T C 758 (321) 793
 Panouse J B 61 (373) 65 (314) 91
 Panzenhagen H 747 (848) 808
 Papineau Couture G 524 (215) 585
 Papkoff H 203 (63) 221 (57) 232 270
 (230) 303
 Papper S 705 (941) 810
 Pappenheimer A M 75 (315) 91 161
 (97) 170
 Pappo R 539 (173) 583
 Pardoe U 101 (46) 145
 Parente N 616 (156) 618 (156) 659
 Parhon C I Jr 416 (8) 500
 Parhon M 487 (503) 510
 Park C R 183 186 195 197 213 (178)
 257 (36) 700 (218) 997 304
 Parker G H 100 111 (115) 131 (113)
 149 419 (151) 429
 Parker L W 160 (108) 175
 Parker M W 27 (10) 48
 Parkes A S 270 (10) 231 270 (41) 998
 318 (142) 383 470 (665) 491 (141)
 503 515 673 (849) 803
 Parkinson D 670 (188) 682
 Parkinson J I 4 8
 Parmer L C 700 (850) 808
 Parrish A E 725 (1255 1256) 819
 Parrott D M 843 (70) 872
 Parry D W 25 (225) 53
 Parson W 271 (72) 212 (73) 273 (72)
 998 299
 Larson U 668 (160) 681
 Partridge J W 271 (106) 302 775 (636)
 733 (135) 151 (631) 802 897 (114)
 875
 Larvonen S 863 (148) 874
 Laschke K I 273 (78) 299 457 (137)
 451 (625) 482 (676) 503 514 594
 (130) 661 674 (255) 683 136 (917)
 742 (209) 765 (313) 782 (151) 852
 790 794 808 810 811 (42 152) 875
 (151) 896 (152) 898 (41) 934 (42
 152) 937 (475) 838 (51) 841 (152)
 842 (150) 945 (33) 848 (33) 857 (153)
 861 (150) 868 (149) 870 871 874
 Lasqualetti A T 491 (510) 510
 Pasqualini R Q 313 385 479 867 (154)
 874
 Passano I M 60 (114 376 378 379)
 63 (376 377 379 380) 66 (319) 86
 91 108 113 (118) 115 (218) 138 (214)
 139 (116 117) 149
 Pásztor 101 (193) 148
 Pataki J 540 (76 17 184) 568 (109)
 569 (269) 580 583 586
 Pat J W 110 (51) 786
 Patel I R 271 (351) 306
 Paterson A B 498 (391) 509
 Paterson J C 717 (153) 308
 Paterson J A I 579 (36) 565 (10) 579
 Paton W D M 155 (58 219 221 223)
 131 (277) 137 (82) 145 146 149
 Patt H M 161 173 612 (189) 685 759
 (1111) 815
 Pattee C J 744 (1175) 817
 Patterson E 597 661
 Patterson C N 713 (1245) 716 (1245)
 819
 Patterson J W 190 (96 97 98) 197
 Patterson P A 699 (610) 801
 Patton H 458 (688) 516
 Patton H D 782 (384) 307
 Patras M C 154 (150) 161 (150) 174
 Pauker R S 673 (18) 678
 Paulish G 331 (200a) 384
 Pautsch F 62 (382) 81 (381) 82 (382)
 91
 Pawan G L S 305 (34) 416
 Payne F 347 383
 Payne P W 203 221 (3 79) 272 (2 3)
 279 281 283 243 (179) 254 (279)
 276 (10) 219 (12) 286 (280) 293 (10)

- 11 280) 290 (10 281) 297 301 422
 (12) 426 731 (854) 774 (804) 808
 422 (8) 870
 Irazin M 467 (114) 607
 Isaaccock W C 406 (59 340) 463 (003)
 499 (540) 501 508 515
 Iearce, R H, 533 (227) 876
 Pearlman M R J 540 (417) 012 (248)
 577 (246) 585 099 (226) 681 867
 (150) 874
 Iearlman W H 577 (246) 585 594 59)
 600 604 600 615 616 661 667 (100
 191) 687 863 (155) 874
 Pearce A G I 311 312 313 314 385
 Iearson J A 40 (276) 53
 Iearson O H 484 (500) 486 (000) 487
 (050) 490 (500) 515 696 (860 1201)
 638 (1200) 700 (807) 706 (859 860)
 707 (807) 711 (859 860) 701 (287
 288) 703 (807) 753 (857) 808 817
 845 (90) 872
 Iearsons J 461 (603) 516
 Ieckham B M 669 (192) 682
 Peezenik O 690 (861) 808
 Pedersen K O 204 (80) 217 221 (57
 69) 232 446 (157) 448 (735) 503 507
 Iedrick L 771 (424) 773 (424) 786
 Ieerdeman A I 533 (27) 579
 Ieters G 417 430
 Peevers R W 002 (20) 579
 Pekkarinen A 487 (300) 508 742 (1167)
 816
 Iellegrino P C 745 (862) 808
 Iellerin J 496 (503) 515
 Pelser H 607 (434) 746
 Iencharz R 672 (145) 614 (180) 687
 Pencharz R I 731 (107) 300
 Penew I 49 (309) 508
 Ienhos J C 670 (193) 687
 Pentz I I 721 (301) 797
 Perinio A 447 (333) 507
 Percefull S C 722 (531) 99
 Ierera G A 714 (863) 808
 Pérea J M 79 (383 384) 91
 Iernetti H 449 (684) 516 888 (198)
 837 (198) 838 (198) 875
 Ierisutti G 183 (57) 190 (87) 196 197
 Perkins F B 138 149
 Perkins H R 151 170 175
 Perkoff G T 845 (156) 874
 Ierlman D 563 (249) 585 608 (98) 670
 642 (98) 657 661, 698 (300b) 794
 Ierlman I 436 (011) 400 (417) 511 517
 Perlman I I 620 (130) 659 743 (864)
 808
 Ierlmutt J 637 (1121) 698 (1121) 722
 (1120) 815
 Perlmutter M 740 (860) 808
 Perman J M 476 (669) 516
 Iernow B 100 137 (220) 149
 Ieron F 500 (127) 609 (127) 637 (233)
 658 661
 Ierrault M 411 (512) 517
 Ierry W F 488 (131) 503 737 (868)
 781 (1207) 782 (866) 783 (867) 808
 819 829 (107) 874
 Ierry W I M 170 (221) 134 (141)
 147 149 220 (10) 921 290 (41) 928
 498 (409) 509
 Iersike F C 481 (513) 517
 Ierska H 107 (226) 149
 Ierska I 76 (869) 808
 Ieruzzi P 128 (138) 147
 Ieschel I 714 (870) 808
 Ie ez M 071 (374) 887
 Ietermann M 400 (514 576) 517 515
 Ieters B A 707 (12401) 819
 Ieters J B 831 (179) 813
 Ieters J H 483 (515) 517 700 (201)
 790
 Ieters J I 459 (426) 483 (776) 494
 (411) 480 (516 517) 509 510 517
 517 863 (158) 874
 Ietersen A P 443 (736) 03 (236) 503
 Ietersen W F 306 381 400 (61 180)
 407 450
 Ieterson D H 608 (701a) 609 670
 641 (88 201) 642 (701a) 643 637 660
 661
 Ietit A 571 (324) 587
 Ietit D W 467 (100) 502
 Ietrovitch A 60 (9) 678
 Ietrow A A 546 (87 200) 581 580
 Ietrziska T 546 (254) 585
 Ieyser I 107 (237) 134 149
 Pfeifferberger M Jr 844 (87) 870
 Pfeiffer C A 671 (194) 687
 Pfeiffer C C 122 (227) 127 149 115
 (081) 803
 Pfeiffer I W 74 (380) 91

- Pfaffner J J 591 (324) 664
 Pflugfelder O 68 (380) 71 (386) 91
 Pfoertner W 612 (101) 680
 Phillips R W 338 340 355
 Phillips W I J, 672 (190) 674 (195) 682
 Photopoulos C 761 (376) 774 (386) 793
 Piana A 495 (740) 517
 Piana G 674 (202) 683
 Piccione V 489 (76) 501
 Picha G M 681
 Pick R 716 (1089) 717 (871) 808 814
 Pickering D I 492 (414 518 519 520)
 509 512 876 (159) 874
 Pickford G I 79 (387) 91
 Pickford M 115 (228) 117 (228) 149
 425
 Pickles V R 417 (93) 428
 Piepho H 72 (389 391 392 393) 73
 (390) 80 (388 392) 91
 Pierce F T 716 (92) 732 (92) 787
 Pierce J G 402 (161 162 163 201)
 403 404 (161 201) 406 (51 147) 412
 (145) 427 429 430
 Pierrotti L 164 (95) 172
 Pigraud H 863 (160) 874
 Piliero S J 747 (412) 748 (412 812)
 796 808
 Pilet F F 3 (277) 18 (227) 19 55
 Pillsbury D M 747 (49) 748 (49) 769
 (49) 778 (643) 779 (643) 786 80
 Pilser H 843 (71) 82
 Pincus G 230 233 257 (253) 258 (230
 252) 291 (284) 294 (253) 303 304
 591 (331) 594 (230) 599 (230) 604
 (229) 600 606 (133) 610 (15) 612
 (15) 610 (15) 618 (206) 623 (151)
 624 (133 150 151 167 168 200) 620
 (133 167 202) 646 (168) 647 (89 238)
 648 (151) 654 (254a) 655 658 659
 660 661 662 664 667 (196 197 221)
 668 (198) 671 (199 234) 682 683 692
 (875) 693 (874) 694 696 (875) 702
 (727) 744 (293) 745 (293 873 876)
 792 804 808
 Pincus I J 192 (99) 193 (99 100) 194
 (99) 197 848 (161) 874
 Pincus J B 754 (877) 808
 Pine I 718 (878) 808
 Pintar I 490 (455) 510
 Pinto F M 671 (96) 679
 Piotrowski L J 207 233
 Piquet B, 763 (182) 789
 Pirani C I 740 (1109 1101) 766 (879)
 809 814 815
 Pirozynski W J 743 (230) 791
 Iitesky I, 746 (668) 803
 Littendrieh C S 71 (394) 91
 Pittoni A 180 (521) 512
 Pitt Rivers R 443 (280) 445 (264 286
 522 524) 447 (522) 449 (261 262)
 450 (245) 454 (260) 475 (263 260
 523) 490 (261 263 264 260 266) 505
 506 512
 Pitts R F 154 172 706 (920) 708 (959)
 709 (920 921) 710 (921 934) 754
 (921) 810 811
 Iitzer H S 524 (19) 579
 Pizarro O 701 (10) 785
 Place C H J 670 (200) 682
 Plager J 667 (171) 681
 Plager J E 626 628 (208) 646 (240
 241) 661
 Plamondon C A 486 (30) 497 (30)
 499 (30) 500
 I lantin I O 611 633 (12) 630 (12a)
 655 694 (381) 795
 Plate W P 856 (162) 874
 Platt R M 28 (229) 29 (229) 36 (229)
 58
 Plattner P A 527 (118) 546 (201 202
 203 254) 560 (149) 566 (147) 567
 (147 148) 581 582 585
 Plimpton C H 185 (23) 195
 Plongeron R 403 (141) 429
 Plotz C M 766 (537 880 881 888)
 767 (93 537) 787 799 809 847 874
 Plotz E 691 (207) 790
 Plotz E J 691 (207) 700 (208) 790
 Plummer A J 463 (220) 505
 Plummer H S 449 (520) 512
 Plummer W A 430 (81) 501
 Pochin E E 439 (481) 440 (483) 449
 (482) 451 (482) 452 (482) 459 (506)
 511 512
 Pohl R 44 53
 Pohley F M 212 216 218 233
 Polley H F 705 (1080) 707 (1080) 708
 (1080) 709 (1080) 721 (1080) 724
 (1080) 732 (1080) 753 (1080) 765
 (495 496 497 882 883 1186) 766

- (582 683 1018 1186) 798 800 812
 814 817
 Polley J R 714 (61) 794
 Pomerantz L 736 (800 801) 806
 Pon V G 279 (229) 303
 Ponz F 479 (527) 612
 Poos G I 539 (288 290) 586
 Pops G T 324 385
 Popenoe I A 99 (229 279) 149 160
 402 (164) 493 (165) 494 (164 209)
 413 414 450 451
 Popjak G 353 (218) 586 536 (57a) 580
 592 666 667
 Poppell H F 484 (550) 486 (555) 487
 (555) 490 (550) 515
 Porath J O 221 (33) 231 279 (229)
 290 (300) 503 506 424 (131) 170)
 429 430
 Poratti G 436 (9) 600
 Porter C C 724 (1025) 723 (1020)
 731 (1075 1109) 815 815
 Porter J C 673 (701) 687
 Portfolio A G 771 (415) 797
 Possompès B 68 (306 307 401) 71 (401)
 72 (795 398-402) 91 92
 Posternak J 119 (121) 147
 Potick D 256 (179) 391
 Potter D D 109 140 149
 Pottner M A 617 (266) 618 (266) 662
 Potts A M 401 450 702 (201) 791
 Poupa O 491 (271) 506
 Poutasse F 610 (769) 612 (269) 662
 Power M H 439 (330 541) 440 (115
 339 417) 441 (115) 450 (19) 500 500
 508 510 512 697 (1079) 701 (1077
 1078) 700 (1080) 707 (1080) 708
 (1080) 709 (1080) 721 (1080) 724
 (1080) 720 (1079) 727 (1077) 732
 (1080) 753 (1080) 765 (1080) 814
 841 (40) 844 (110 197) 516 (89) 871
 872 873 875
 Powning R F 77 (145) 86
 Praetz B 561 (208) 577 (115) 581 584
 Prager J 574 (111) 581
 Prager R 178 (204) 149
 Praglin J 770 (1036) 813
 Praymorsky M 106 (299) 151
 Prato G S 461 (653) 515
 Praytor F H 460 (85) 692
 Prelog V 524 (19) 531 (67 207 256 205)
 546 (207) 577 580 585 601 610
 (797) 615 (297) 660 663
 Presch H P 473 (163) 504
 Preston R 770 (407) 796
 Prestrud M C 260 (190) 300 344
 (163a) 353 728 (541) 729 (570) 736
 (540 582) 752 (583) 900
 Price C A 45 (231) 53
 Price D 666 (202) 687
 Price M 166 (127) 173
 Price W H 194 197 201 (120) 257
 (280) 300 304
 Prichard M M I 320 (202b) 386
 Priestley J T 696 (884) 714 (884) 809
 Prima C 488 (578) 512
 Prins D A 576 (303) 560 (700) 585 587
 618 (509) 673 (117) 645 (117) 653
 673
 Prinzie A 489 (520) 512
 Prisament M 677 (207) 68
 Procopio J 270 (347) 306
 Profumo P 47 (196) 53
 Prosser C I 101 (232) 120 (232) 122
 (232) 149
 Trout F S 576 (262) 585
 Ruddick J F 766 (700) 767 (700) 805
 Rueter R D 490 (570) 512
 Runtz F T G 283 (392) 307 409 (704)
 504 709 (880) 710 (154) 741 (105)
 783 794 809 846
 Puhlsley L I 200 (286) 304
 Pummer W J 540 (51 52) 581
 Puntriano G 460 (531) 512
 Purinton C O 695 (746) 806
 Purnell J F 768 (689) 803
 Purves H B 312 314 315 585
 Purves H D 391 (169) 430 457 (537)
 461 (715) 606 508 510 612 (676)
 807 823 (165 166) 534 (164) 84
 Purvis O 32 33 (234) 50 68
 Puschel I 783 (530) 807

Q

- Quarton G C 718 (153) 789
 Quashnock J M 708 (906) 810
 Quastel J H 106 (198) 148 443 (269)
 449 (260) 506 676 (116) 680
 Querido A 207 253 750 (1172) 817
 Quimby F H 430 (573 763) 512 518
 Quimby F H 674 (703) 687

Quinby W C 775 (1193) 817
 Quinn E L 835 (167) 856 (167) 874
 Quittner H 747 (855) 879

R

Riab W 149 476 (534) 497 (535) 512
 745 (549) 809
 Raadts F 5 7 54
 Rabbi A 455 (536) 512
 Raben M S 202 203 204 221 (3 79)
 222 (2 3) 239 231 233 236 253 (12)
 256 (410) 257 (92) 260 (410) 267
 (791) 263 (791) 264 (257 791 292)
 255 (297) 270 (290) 271 272 273 275
 (790 292) 276 (10) 279 (12) 286
 (250) 233 (10 11 250 259) 295 (101)
 231 258 259) 236 (410) 297 304 307
 422 (12) 455 822 (8 169) 829 (165)
 870 874
 Rabinovitch B 176 (28) 196
 Rabinovitch M 592 (215) 602 674 (204)
 682
 Rabinowicz M 740 (887) 809
 Rabinowitz J L 592 662
 Rachmilewitz M 748 (317) 793
 Padeliffe C F 487 (537) 51
 Radike A W 164 (147) 173
 Raeside I J 670 (705) 682
 Raffelson H 539 (16) 5 3
 Raffi A 398 (65) 177
 Ragan C 718 (476 1169) 740 (555 556)
 753 (721) 754 (251) 766 (537 669 880)
 891 889) 767 (93 537) 779 (761)
 787 780 786 799 800 803 805 809
 816 847 (163) 874
 Pahn U 72 (241 404 405) 88 92
 Raile R B 285 (380) 307 604 (244)
 745 (624) 791 80 850 (105 208)
 672 675
 Rauman R J 782 (150) 788
 Rajagopalan S 558 (97) 581
 Rakoff A E 347 371 385 594 (730)
 539 (776 731) 601 857 (153 171)
 858 (150 171) 860 (58 59) 863 (155)
 871 874
 Rall J E 434 (470) 438 (539) 439 (541)
 440 (115) 441 (115) 449 (538) 450
 (19 476 477 478) 451 (542) 452
 (538) 454 (542) 4 6 (55) 484 (555)
 496 (555) 487 (555) 490 (555) 498

(554) 499 (549 556) 500 591 59
 511 512 513 839 (176) 874
 Ralli F P 737 (268) 738 755 77 809
 Ralls J W 539 (342) 588
 Ramasarma G B 727 (937) 810
 Rambo J H T 767 (894) 809
 Ramoy F R 746 (501) 798
 Ramirez F 540 (12) 559 (12) 578
 Rammelkamp C H 775 (416) 797
 Pamminga F F 843 (71) 87
 Rampona J M 708 (1125) 714 (1122)
 815
 Ramsay D A 549 (175) 583
 Ramsburg H H 756 (718) 894
 Pamsey C 756 (421) 796
 Ramstad E 25 (170) 52
 Randall H T 696 (860 1201) 706 (860)
 711 (860) 750 (1201) 868 817
 Randall L M 342 (243) 343 388
 Randall R A 449 (544) 455 (543) 512
 Randall S S 427 443 (777) 445 (524)
 448 (287 288) 506 512
 Randle P J 261 (794) 304
 Randoia L 740 (895) 809
 Panson S W 321 (71) 381
 Rapaport E 729 (728) 804
 Rapp B 462 (501) 519
 Rappaport J 25 (739) 27 (740) 44 54
 Rapport M M 103 104 (317) 149 151
 Rapport R L 449 (545) 512 672 (219)
 682 807 (172) 874
 Rasmusen A F Jr 497 (675) 516
 Rasmusen H 493 (546) 512 712 (675)
 803
 Rasquin P 492 (547) 512
 Ratsumamanga A R 739 (790 391 897)
 740 (887 896) 795 809
 Rausser R 555 (247) 585
 Rawlins A G 761 (895) 809
 Rawlins T E 37 (152) 51
 Rawls W B 774 (410) 775 (410) 796
 Pawson R W 434 (470) 435 (558) 440
 (443) 451 (542) 454 (542) 455 (59)
 340 545 631) 457 (548) 458 (468 548)
 550 552 550) 459 (467 468 469 549)
 463 (552 553) 471 (469) 473 (548)
 552) 484 (555) 486 (555) 487 (555)
 488 (466) 489 (763) 490 (555 763)
 498 (471 554) 499 (551 556 557)
 500 501 508 510 511 512 513 516

- 618 742 (781) 782 (780 782 783) 806
 839 (176) 874
 Ray L W 372 379
 Ray P M 29 (279) 55
 Ray R 717 (179) 733 (179) 78
 Ray R D 492 (560) 513
 Ray S N 740 (469) 797
 Ray T W 476 (161) 503
 Raybuck H F 161 (119) 173
 Raymond J 674 (706) 682
 Reid C H 693 (893) 738 (893) 809
 Reaven G 156 (6 7) 170
 Rebell G 778 (613) 779 (613) 802
 Reber F 501 (326) 587
 Reboredo A 751 (1075) 814
 Rebuck J W 761 763 (900) 809
 Recant L 233 (295) 283 (296) 304 305
 744 (902) 750 (901) 809
 Recht P 361 380
 Reddy Wm J 281 (196) 302 845 (173)
 874
 Redemann C F 474 (490) 511
 Redemann C T 5 (241) 54
 Redlich A 434 (305) 607
 Reece R I 351 (221) 385
 Reed B A 164 (89) 172
 Reed I 437 (445) 510
 Reed J 660 (72) 679
 Reed T G 762 (483) 763 (483) 778
 Rees M W 177 (102) 197
 Rees R J W 777 (474) 797
 Rigan F D 700 (242) 775 (242) 791
 Rehm M 68 (406 407) 76 (408) 92
 Reich H 537 (760) 585 606 (765) 648
 (260) 662
 Reich M 437 (561) 513
 Reichenbach I 400 (440) 510
 Reichert F L 485 (184) 504
 Reichstein T 570 526 527 (328), 529
 534 (22 196) 501 (199 199 326) 508
 (328) 509 (329) 563 565 (327) 571
 572 (308 308a) 573 (308a) 574 (308a)
 579 584 585 587 604 610 615 622
 (247 256 257) 655 662 664 687
 (1029) 707 815
 Reid D H 780 (715) 804
 Reid F 185 (103) 193 (11) 195 198 202
 204 233 243 (71) 257 (301) 258 (37)
 262 (71 234) 264 265 (71 207) 275
 (297 299) 297 298 303 305 306 330
 421 (135) 424 (135) 499, 727 (903) 809
 Reid I 260 (29) 272 (29) 297
 Reifenstein F C 156 (123) 160 170 173
 Reifenstein R W 756 (420 421 422) 796
 Reilly J 677 (707) 682
 Reilly W A 471 (562) 513 877 (81) 877
 Reinders D I 43 (242) 46 (242) 54
 Reinecke R M 237 (326) 305
 Reinecke I I 443 (563) 410 (563) 564
 565 449 (79) 501 513 750 (1155),
 781 (1185) 817
 Reinecke L M 608 (235 236) 609 (215)
 641 (88 234) 643 (235) 657 661
 Reiner M 450 (658) 515
 Reinhard I G 64 (409) 60 (409) 92
 Reinhard M C 759 (775) 760 (775) 806
 Reinhardt W O 278 (200) 200 (395)
 396 277 (303 304) 290 (500) 303
 305 307 421 430 744 (551), 746 (552)
 749 (900) 760 (704) 800 809
 Reinhold D 504 (294) 586
 Reiss M 221 233 278 (138) 268 (56)
 307 289 (307) 298 300 305 374 385
 674 (40) 678 717 (906) 735 (907) 809
 Reiss R 733 (637) 802
 Reiss R S 430 (301) 507 697 (341) 715
 (341) 732 (341) 733 (341) 781 (513)
 794 799
 Reisser O 195 (100) 198
 Reisman A S 709 (507) 800
 Remilton L 418 422 428
 Remington J W 434 (566) 513
 Remington R F 414 (566) 513
 Remy C 479 (567) 513
 Renault J 467 (114) 502
 Rennels F G 671 (208) 682
 Renner M 70 (410) 92
 Renold A I 186 (104) 186 291 (309)
 293 (115) 293 (115 308 309) 294
 (309) 300 305 697 (1150) 718 (1150)
 744 (1150) 816
 Renzi A A 390 496 710 (106) 707
 Resende F 15 (241a) 54
 Reessler C 99 (280) 150 404 (211) 406
 (171) 407 408 (210) 409 (211 211a)
 430 431
 Reverberi G 64 (411-413) 9
 Rev R 430
 Reyes A 496
 Reynier M 61 (515) 94

- Reynold S R M 862 (174) 874
 Reynolds L 838 (175) 874
 Rhamy R K 713 (928) 810
 Rhein W v 77 (414) 92
 Rhoads C P 604 (67 1 0) 632 (260b)
 667 (6a) 656 660 6 9 7a1 (908 1113)
 7a (119 1113) 788 809 815 861
 (48) 871
 Rhodes A 40 (243) 54
 Rice C O 434 (568) 446 (106) 502 513
 Rice F I 17 (244 2a1) 54
 Rice K K 710 (909) 809
 Rice K L 7a2 (583) 800
 Rich A P 733 (911) 771 (910 912) 809
 Richard J 291 (233) 303 822 (12a) 8 3
 Richards A G 71 (415) 77 (416) 92
 Richards C E 475 (459) 513
 Richards J B 257 (310) 305 809
 Richards J H 293 (135) 300
 Richardson F M 522 (24a) 585 634
 63a (248) 636 (248) 662 859 (a0) 871
 Richardson H B 483 (5 1) 484 (570)
 513
 Richardson K C 2a2 (311) 305 361
 (56) 366 (55) 380 400 (39) 426
 Richert D A 479 (567) 513
 Richfield D F 754 (406) 710 (408) 796
 Richter C P 462 464 (513) 513 710
 (909) 809
 Richter J 221 223
 Richter J W 222 (5 9 87) 224 (7) 226
 (7 9 87) 231 729 (40) 286 (20) 294
 (40) 297 298
 Richter R B 718 (246) 791
 Riddell C B 4a8 (550) 463 (5a3) 500
 512 513
 Riddle O 246 (312) 278 (2a4) 303 305
 346 373 374 375 385
 Ridgway L P 619 (610 611) 801
 Ridout J H 254 (15) 297 758 (992) 812
 Riedel B E 73a (914 915) 809 810
 Piegel B 516 (262) 685 609 662
 Riegel C 781 (467) 797
 Rienhoff W F 448 (274) 513
 Rietsma J 3a (245) 64
 Riggs D S 439 (475) 449 (575 634)
 450 (575) 4a4 (102) 475 (569) 50
 513 516 828 (198) 837 (198) 838
 (198) 825
 Rigler R 419 423 430
 Rus W 673 (209) 682
 Riker A J 37 (302) 55
 Riker W J Jr 119 (236) 122 (236) 125
 (236) 149
 Riley D P 181 (3) 195
 Rimington C 465 (3a2) 508
 Rinehart J F 315 381
 Rinfret A P 291 (233) 303 222 (12a)
 873
 Ring J 669 (210) 610 (210) 682
 Ringold H J 550 (60) 552 (7a 263 264
 26a) 5a (7a) 580 585
 Riniker B 536 (6a 264a) 518 585
 Ripstein M P 60a 663 693 (509) 738
 (599) 809
 Rislér H 68 (417) 9
 Ritchie A C 762 (916) 810
 Rittenberg D 518 (33) 519 591 (18
 2a1) 655 662
 Ritter H 227 (264) 304
 Rivero Fontan J I 482 (152) 517 236
 (917) 742 (209) 790 810 828 (41) 871
 Pives M 419 (145) 429
 Rob C G 756 (27a) 792
 Robb P D 480 (639) 515
 Robbins J 450 (514 576 577 578) 451
 (242) 4a4 (542) 484 (55a) 486 (555)
 487 (a5) 490 (555) 499 (5a6) 512
 513 839 (126) 824
 Robert H 527 (118) 581
 Roberts A 440 (298 299) 507
 Roberts C W 90 (280) 150 404 (211)
 431
 Roberts E 727 (919 937) 728 (918) 810
 Roberts E H 23 55
 Roberts I S 594 658
 Roberts K 39a (119) 230
 Roberts K F 706 (920) 709 (920 921)
 710 (921) 754 (921) 810
 Roberts M 670 (211) 682
 Roberts R H 20 (247) 54
 Roberts S 669 (245) 670 (24a) 676
 (24a) 683 729 (922) 761 (923) 810
 Robertson E 729 (651) 732 (650) 802
 803
 Robertson E C 161 (61) 171
 Robertson G G 669 (212) 682
 Robertson H A 286 (355) 307
 Robertson J D 107 (237) 149 457 (579
 580 581) 515

- Robertson O H 192 (582) 513
 Robertson R N 40 (296 218) 65 54
 Robertson W B 134 149
 Robillard F 496 (583) 513
 Robinson A M 667 (213) 687
 Robinson C 181 (106) 198
 Robinson D 438 (111) 503
 Robinson G F Jr 335 585
 Robinson H J 270 (313) 505 736 (926)
 776 778 (924 926 927) 810
 Robinson J 132 133 134 (53) 146
 Robinson J R 45 (250) 64
 Robinson R 337 (44 57) 538 579 592
 (252) 662
 Robinson S 713 (928) 810
 Robinson T J 335 339 (224 226) 585
 671 (214) 687
 Robinson W D 727 (929) 732 (929) 765
 (929) 766 (263) 781 (1225) 792 810
 818
 Robitschen T 618 (163) 648 (163) 659
 Robson H N 716 (930 931) 810
 Robson J B 670 (211) 687
 Robson J M 668 669 (105) 6 0 (161)
 680 687
 Roca J 447 (405) 509
 Roche J 436 (803) 438 (244) 447 (515)
 443 (588 599 608) 445 (587) 591 593
 416 (157 158 590 596 597 598 602
 603 605 606) 447 (586 588 512 600)
 448 (158 584 587 601 604) 453
 (585) 454 490 (502) 505 505 518
 514 519
 Roche M 201 (391) 307 697 (341) 718
 (341) 732 (69 341) 733 (341) 765
 (1140) 786 794 816
 Roche P 777 (932) 810
 Rockenbach J 780 (1008) 817
 Rockwell F M 770 (407 933) 796 810
 Rodgers O G 618 (206) 624 (204) 625
 (202) 626 (201) 631 (204) 632 (204)
 646 (204) 660
 Rodriguez R D 672 (217) 68
 Rodriguez R R 243 (190) 265 301 374
 (1432) 585
 Roe J H 484 (609) 514
 Roemmelt J C 710 (934) 810
 Rosler H 450 (318) 507
 Rogers G R 484 (486) 511
 Rogers J 545 (318) 587
 Rogers W F Jr 467 (784) 618
 Rogier I R 539 (173) 683
 Rogoff J M 708 (936) 715 (935) 810
 Rohrbaugh I M 17 (251) 64
 Rohmann L 598 (216) 565 (216) 585
 604 605 660
 Rojas G 427 497
 Rokk L 647 (2) 655
 Rolf D 270 (411) 307 394 (220 221
 222) 395 (272) 431
 Romani J 361 585
 Romano J 718 (703) 793
 Romanoff L B 610 661
 Romanoff L P 703 (727) 745 (876) 804
 808
 Romeis B 311 535
 Romero M 549 (206) 585
 Romo J 540 (76 77 154) 549 (266) 552
 (765 270 271) 556 (72 967) 550 555
 585 586
 Rondoni P 489 (610) 514
 Ronkin R R 80 (347) 90
 Ranzoni L 177 199 77 (910 917) 810
 Rondenburg J W M 31 (317) 66
 Roos A 424 (1341) 499
 Root H F 274 (65) 298
 Ropes M 154 (1) 169 487 (31) 500
 Ro e B 757 772 (692) 605 810
 Ro e C I 166 167 (18) 170
 Rose L 848 (51) 859 (50) 871
 Ro e F R 493 499
 Rose H M 406 (661) 803
 Rosenberg F 781 (218) 507 653 (321)
 664 701 (697) 703 (947) 735 (697)
 46 (799) 793 (1709 1711) 804 810
 818
 Rosenbaum J D 60 (941) 810
 Rosenberg A W 708 (936) 810
 Rosenberg I N 753 (17 714) 254 (314)
 255 (314) 256 (314) 259 (15) 260
 (315) 273 (289) 295 (281 285 289)
 296 (314 315) 304 505 447 (617) 443
 (611) 449 (612) 514
 Rosenberg N 768 (919 913) 810
 Rosenblueth A 128 130 (61) 131 (60)
 145 149
 Ro enblum I 449 (614 611) 514 515
 Rosenfeld M 416 (85 174) 4 7 450
 Rosenfeld R R 612 (134) 613 (134) 658

- Rosenfell R S 577 (83) 657
 Rosenheim O 46 (200) 585
 Rosenkrantz H 615 (217) 619 644 (303)
 675 (303) 633 (303) 638 (96) 47
 (96) 648 (96) 657 658 661 662
 Rosenkranz G 371 (272) 579 (212) 540
 (76 77 184) 44 (74 284) 546 (314)
 549 (66 768) 540 (311 313) 552 (70
 263 264 265 270 241 312) 543 (347)
 544 (40 313) 546 (6) 72 73 267) 50
 (45) 558 (313) 568 (269) 569 (469)
 580 583 585 586 587 588 608 (179)
 670
 Rosenman P H 485 (615 616) 514
 672 (218) 687 715 (347 914) 794 810
 Rosenthal M C 747 (1094) 814
 Rosenthal R L 747 (945 946) 810
 Roskelley R 409 (649) 516
 Ross F 603 (79) 618 (49) 657 703 (707)
 791
 Ross E J 183 184 (108) 198
 Ross J B 710 (503) 800
 Ross J F 747 (319) 793
 Ross O B 489 (75) 501
 Ross S 743 (407) 204 (407) 307
 Ross W F 166 173
 Rossi A 763 (60) 786
 Rossi C A 485 (536) 517
 Rosster R J 448 (618) 479 (614) 514
 735 (914 915) 809 810
 Roston J 102 (226) 149
 Rosvold H E 719 (944) 811
 Rowatt B 879 (17) 80
 Rotblat J 440 (619) 514
 Roth A A 865 (86) 87
 Roth F J 760 (1127) 815
 Roth G J 168 172
 Roth P 46 (670) 490 (60) 491 (621)
 501 514 642 (735) 683
 Rotheild I 677 (219) 687
 Rothen A 215 (16) 31
 Rothen V 416 (204) 451
 Rothman S 754 (239 446) 755 (208 239
 948 1173) 791 797 811 817
 Rotondaro F A 667 (50) 679
 Rottger E 718 (355) 794
 Rowan W 317 385
 Rowe C R 756 (949) 811
 Rowe I W 397 (86 175) 401 (116) 403
 (116) 428 477 479
 Rowland S J 365 (88) 381
 Rowlands I N 439 (315) 507
 Rowlands I W 344 (731) 385
 Rowley F M 714 (1006 1007) 817
 Rowson I I 341 (31) 341 (31) 349
 Roy A 333 (98) 340 341 381 665 (270)
 671 (220) 683
 Roy A C 783 (900) 811
 Roy W R 218 (70) 737
 Rozich R 316 384
 Rozin S 609 (76) 684
 Rubin B A 579 (212) 594
 Rubin B L 154 (124) 167 (122) 168
 173 608 (179) 611 (74) 648 (705) 654
 (404) 657 660 667 667 (221) 683
 Rubin L 464 (797) 519
 Rubin M 539 (134) 587
 Rubino F 489 (672) 514
 Rubinstein I N 673 (240) 683
 Rucart G 164 (123 124) 173
 Ruffeldt O 595 (250) 667 673 (223) 683
 Rufelt H 74 (200) 54
 Ruff A 508 (773) 586
 Rugh R 439 (673 624) 514
 Runner M N 315 315 336 383
 Rupp J 491 (625) 492 (676) 514 674
 (222) 683 874 (157) 826 (152) 834
 (152) 841 (152) 847 (150) 861 (150)
 874
 Ruschug H 563 (744) 586 604 (789) 663
 Russek A S 749 (901) 811
 Russek H I 747 (951) 811
 Russell J A 207 (115) 203 (115) 734
 236 243 (183 257 319 370) 248 (118)
 249 (118 318 319) 251 (319) 202 203
 (110) 257 (707) 258 (207) 260 (754)
 766 (318) 271 (414) 296 300 30 303
 305 308 728 (95) 729 (952) 731 811
 Rutgers J J 594 (102) 683
 Rutishauser F 164 (98 175) 172 173
 Ruyle W V 502 (275) 554 (47 204) 563
 (47) 579 586
 Ruzick L 573 (277) 546 (276) 5 8
 (2 8) 531 (279) 565 (270) 586 604
 (244) 610 (797) 615 (794) 607 663
 Ryan K J 600 601 647 (257 258) 663
 Rybak B 80 (418) 97
 Ryby I 469 (576) 799
 Ryden I 569 (180 187) 583
 Ryer A I 533 546 (287) 586

S

- Sabi A 623 619 662
 Sabih D 488 (127) 503
 Saho I F 509 (107) 581 698 (350a) 794
 Sacccone A 705 (744) 805
 Sacks J 187 (109 110) 108
 Sacktor B 70 (53 419 490) 84 92
 Sadhu D I 46 (628) 488 (127 628) 814
 Saffran J 595 (127) 596 (1) 609 (127) 655 658
 Saffran M 288 (322) 289 (321 322) 305 623 662
 Sakamoto H 673 (123) 680
 Sala G 161 170 172 709 (951) 711 (904) 721 (141) 788 811
 Salamon I 27 586
 Salamon I I 612 (200b) 662
 Salassa R M 696 (834) 714 (884) 782 (905) 809 811
 Salcedo J Jr 204 (376) 307
 Salgado I 270 (317 306 674 (69, 679
 Salhanick H A 334 385 601 662 671 (224) 688
 Salmon A A 600 (262) 662
 Salmon T N 239 305
 Salmon U J 60, 662
 Saloun R 66 (216) 88
 Salter J M 238 (212 324 325) 239 (212 325) 302 305
 Salter W T 439 (630) 440 (293) 440 (480 663) 447 (289 629) 448 (259) 449 (634) 450 (632) 474 (631 630) 450 (639) 506 507 511 515 838 874
 Saltstein H C 758 (590) 759 (591 595) 766 (964) 769 (594 964) 801 811
 Simes G L 699 (906) 811
 Siammartino R 221 383
 Samter M 787 (53) 787
 Samuels L T 237 (326) 236 (267 391) 287 (329) 288 (31) 291 (31) 297 304 305 307 461 (636) 415 600 (296) 604 (261) 606 (135) 613 (218 288) 616 (296) 617 626 646 (241) 648 649 (320) 659 661 662 663 694 (281 814 815) 696 (815) 730 (962) 735 (379) 740 (907) 759 (108) 787 792 795 807 811 845 (178) 874
 Samuels P B 743 (230) 791
 Sanborn R C 75 (421) 92
 Sandberg A A 281 (327) 286 (32) 297 305 745 (907) 811 840 (106 178) 844
 Sandern M I 61 (81 88) 62 (80 83 87 89 422 423, 424) 85 92
 Sandford I 481 (82) 486 (82) 602
 Sandford K 481 (82) 486 (82) 602
 Sandoval A 544 (251) 586
 Sandule co G 594 (102) 658
 Sanger F 177 178 (111) 179 180 198 227 283 406 (176) 430
 Santa N 739 (392 393) 795
 Santamaria I 192 (36 38) 196 261 (112) 300
 Sapirstein I A 714 (958) 811
 Sarcar U 163 (125) 173
 Sarett L H 529 530 (286) 531 (257) 532 (256 257) 539 (288 290) 549 (51) 550 (51) 553 (52 259) 567 (250) 580 586
 Sartorius O W 395 (179) 430 708 (959) 710 (934) 810 811
 Satina S 27 (240) 35 (210) 84
 Sato H 497 (336) 607
 Sauer G 506 (150) 582
 Saunders F J 661 714 (427) 796
 Saurino V R 780 (220) 790
 Seward K 289 (163) 301 525 (291) 586 610 611 (31-33) 612 615 619 (269) 623 (121 122) 626 634 (31-33) 635 (31-33) 645 (270) 645 (31) 655 656 657 658 (62 667 (22) 683
 Savit J 77 (504) 94
 Savaya I 61 (420) 92
 Sawyer C H 322 330 333 (82-84) 334 (233) 331 334 335 671 (162 276) 681 683 853 (130) 869 (130) 883
 Sawyer J I 6 (6 7) 110
 Sawyer M K 137 (240) 149 430
 Sawyer P N 770 (57) 780
 Sawyer W H 137 (240) 149 397 399 398 (178) 429
 Sav H J 550 (6) 552 (6) 553 (6) 560 (6) 578
 Sayers D 818
 Sayers G 166 174 207 (30) 220 221 (24 93) 231 233 236 (331 332) 281 (328 331) 282 (61 62) 283 (328 330 331 380 381) 280 (44) 287 (310 329 382) 290 (28 331) 293 (135 136) 298 300 305 307 375 376 (32 33) 380 391

- (180) 450 6.0 (772) 653 (21) 662
672 (760) 701 703 (1227) 1237 713
(1279) 1231 1237 713 (116 760) 730
(967) 745 (960) 757 752 (1230) 789
829 811 818 843 (2784) 850 (708) 855
Sijers M A 270 (78) 933 946 (131
317) 281 (131) 285 (170) 286 (41) 290
(131) 298 305 630 (772) 677 701 811
Scala F 450
Schachner H 447 (638) 446 (637) 464
(634) 469 (637) 515
Schachter M 104 (109) 137 (8) 146 148
Schaefer A F 755 (5-9) 785
Schaefer I I 721 (107) 725 (103) 787
Schaeffer R L 271 (133 334) 305
Schaffenburg C A 275 (199) 660 740
(363) 811 867 (175) 853
Schaffer N K 738 (335) 247 (331) 248
(335) 272 (335) 285 (14) 30 305
Schairer M A 755 (45) 786
Schallek W 170 (300) 151
Schaller F 68 (176) 77 (42, 4 8) 92
Scharfman W B 810 (275) 856
Scharles F H 480 (639) 515
Scharrer B 58 (431 437 434 440 445-
449) 59 (479 445 449) 60 (418 419)
61 (410) 67 (437) 68 (431 43 433
434 435 436 434 440) 69 (434 437)
70 (434 438 439) 71 (437 434 440
448 419) 73 (437) 74 (442) 6 (434
447 449) 77 (449) 78 (470 431) 79
(418 449) 81 (431) 82 (436 443 444)
9 93 98 105 107 113 (247 244) 138
139 149 150 331 385
Scharrer T 58 (445-449) 59 (445 449)
60 (448 449) 71 (445 449) 6 (449)
77 (449) 79 (448 449) 92 93 98 105
107 113 (247) 138 (246 247) 131 147
149 150 331 385 391 (14) 476 479
Scheer B T 66 (98 450 451 461) 64
(451) 80 (450) 89 93
Scheer M A R 66 (450 451 461) 67
(451) 80 (450) 93
Scheie H G 725 (834) 807
Scheinberg I 491 (640 641) 493 (640
641) 494 (641) 515 718 (764) 769
(364) 811
Scheinberg S R 758 (595) 759 (594
595) 766 (964) 769 (594) 801 811
Schenker V 606 (133) 673 (133) 674
(133 150 151 167 168) 625 (133
167) 646 (168) 648 (151) 658 659
704 (765) 811
Scherr M A 560 (203) 884
Scherr G H 701 (966) 811
Scheuing M 150 (253) 150
Scheurmann R 25 (76) 84
Schieve J F 718 (967) 811
Schiffer F 734 (968) 811
Schuld K T 450 (441) 510
Schuller L 768 (969) 811
Schiller S 604 615 616 618 667 674
(277) 646 (108) 680 683 770 (480)
770 (301) 63 (66) 786 792 798
Schuller O 572 (3084) 573 (3084) 574
(3084) 587 588 (286 287) 663 687
(1079) 707 813
Schlapp W 392 (104) 478
Schlaver C 714 (870) 808
Schmid H 544 (792) 586
Schmid J 758 (109) 500
Schmid K 450 515
Schmidlin J 555 (293) 557 (793 563
(325) 567 (237) 571 (325) 585 586
587 698 (1707) 818
Schmidt C 410 (69) 518
Schmidt I I 73 (454) 80 (452-454) 81
452) 93
Schmidt I H 750 (940) 811
Schmidt Thomé J 560 (41) 579
Schneebein G I 704 (257 941) 67 791
811
Schneider C I 14 (271) 37 (771) 38
(771) 54
Schneider E 488 (613) 515
Schneider J J 616 618 634 (276 277)
635 (276 277) 637 638 (778) 646
(771) 647 (778 779) 648 (779 780
25) 663 740 (807) 804 842 (150) 845
(33 140) 848 (33) 861 (150) 840 844
Schneider R 164 (176) 175 635 (171)
656 670
Schneider W P 537 (172) 556 (98) 581
585
Schneiderman H A 43 (109 110) 0
73 (794 460) 75 (455 156 454 458
459 460) 89 93
Schneerson S S 771 (1052) 813
Schoch F P 743 (973) 811
Schocken V 28 (257) 54

- Schoenheimer R 301 (251) 662
 Schoenewaldt, E F 511 (291) 586 598
 (302) 663
 Scholz C R 549 (75) 576 (209) 580 584
 Schooley J I 211 231 316 385
 Schoplich R R 560 (55) 871
 Schour I 169 (14 15 79) 170 172
 Schrink A R 19 (259) 20 (255 259) 260
 (323) 54 56
 Schreiber I C 556 (201) 584
 Schrek R 703 (974 976) 741 (974-976)
 811
 Schrieker J A 692 (917) 730 (977) 737
 (978) 811
 Schubert J 768 (1204) 818
 Schueler F W 122 (245) 150
 Schuler W 704 (751) 707 (229 979) 791
 805 811
 Schulman I 467 (457) 511 751 (950) 811
 Schulman J Jr 437 (611) 442 (611) 515
 Schultz R B 737 (951 1223) 775 (1221
 1222) 811 818
 Schulze I 459 (615) 515
 Schupbock S 161 (95) 172
 Schusdek A 188 (5) 195
 Schwabe C W 66 (451 461) 61 (451) 93
 Schwartz A 773 (1152) 810
 Schwartz H S 490 (51) 491 (51) 501
 Schwartz I I 270 (81) 299
 Schwartz R 453 (515) 510
 Schwartz S 452 (261) 506
 Schwartz T B 257 (329) 305 718 (575)
 730 (299 962 952 953) 732 (952) 799
 808 811
 Schwartzman G 711 (1051) 813
 Schwarz H 166 173
 Schwarz I I 315 (22) 78
 Schwarzer K 826 (150) 827 (150) 874
 Schweizer M 215 (336 337) 282 (330
 335) 306
 Schwenk L 579 (134) 549 (295) 575
 (135) 576 (135) 582 586 591 (253)
 594 (324) 663 664
 Schwert C W 414 499
 Schwinck I 68 (467) 75 (463) 93
 Scotom H Jr 558 581
 Scott C C 459 (616) 515
 Scott F I 25 (170) 33 (169) 52
 Scott J C 309 (155) 400
 Scott J L Jr 260 (319) 306 734 (300)
 735 736 762 (951) 792 811 911 (52)
 871
 Scott, H G 456 (24) 471 (562) 492
 (520) 506 510 513
 Scott Watson H M 163 (51) 111
 Seow R O 239 (22) 29 482 (61, 618)
 515 674 (275) 783
 Seudamore H H 66 (464) 93
 Seacock R R 430 431
 Seaman G R 80 (465) 93
 Searey A M 560 (203) 584
 Searle C I 465 (619) 466 (619) 515
 Searle G I 157 (34) 196
 Searle N Z 436 (55) 501
 Seastone C V 761 (159) 716 (153) 790
 Seay P 697 (1121) 698 (1121) 792
 (1123) 815
 Sebrell W H 732 (195) 790
 Secht M 37 (15) 52
 Secker J 716 (955) 812
 Secklen J 540 (29) 870
 Seegal B C 715 (616 617) 716 (617)
 769 (576) 771 (616) 799 802
 Seeley R D 482 (351) 509
 Seegal G 752 803
 Segaloff A 211 231 340 (75) 312 571
 375 385 555 606 631 647 (146) 645
 (146) 659 677 (100) 680 722 (956)
 736 810 856 (704) 873
 Segre F 436 (159 327 650) 503 50 515
 Seidlin S M 436 (52) 501
 Seifter J 761 (991) 764 (995 998 980
 990 991) 771 (990) 783 (990) 791 810
 Selenkow H A 456 (30) 487 (30) 490
 (30) 500
 Seligman A M 690 691 (112) 790
 Sell H M 5 (241) 42 (2132) 53 54 56
 Sells W A 439 (463) 510
 Sellers I A 481 (651 804) 515 519 755
 (992) 812
 Selmer R 72 (466) 93
 Selmi W 458 (203) 504
 Seltzer H S 71 (67) 213 (67) 293
 Selve H 213 (312) 216 (760) 268 (311)
 269 (310 311 312 315) 2 0 (311 314)
 317) 381 (311 316) 295 (101 314)
 302 303 308 361 362 379 386 704
 (995) 1091 714 (1006 1007) 720 750
 (499 1005) 756 (997 1004) 757 761

- 760 776 (998 1000) 777 (686) 798
803 812
- Sen S C 163 173
- Sen S P 2 (961) 54
- Seneca H 75 (1008) 810
- Senger G 674 (88) 680
- Senenbach W 718 (1009 1010) 812
- Sequeira L 29 (963) 54
- Serchi C 594 603
- Serini A 549 (296) 586
- Serini I 77 (468) 79 (467) 93
- Serini F 709 (954) 711 (954) 811
- Serres C Jr 574 (114) 581
- Severinghaus A F 310 386 509 (181)
874
- Severinghaus F H 604 (2 5) 615 (275)
662
- Sgouris J T 358 359 (193) 360 361 362
384 386
- Shacter B 753 (1011) 810
- Shafer W G 674 (229) 683
- Shaffer J M 773 (411) 797
- Shaffer P A 483 (652) 515
- Shaffhausen D D 366 383
- Shaffner C S 460 (309) 507
- Shanaphy J F 700 (249) 770 (249)
101) 791 812
- Shank R E 740 (986) 741 (986) 797
- Shannon J A 390 (183) 430
- Shapiro R 767 (1013) 812
- Shappiro D G 75 (463 460) 93
- Sharaf A A 668 680
- Sharp E A 441 (450) 442 (450) 510
- Sharp J T 691 (1053) 813
- Sharpless G R 461 (653) 515
- Shaw E 136 151
- Shaw G B 209 (74) 270 (75) 209
- Shaw J C 367 (240) 366 380 386
- Shaw J H 237 (348) 306
- Shaw W 188 (121) 198
- Shay J C 489 (460) 510
- Shedlovsky T 215 (16 114) 218 (118)
331 334
- Sheehan M M 475 (418 654 776) 466
(418 777 778) 509 515 518
- Sheehan H L 189 (31) 190 691 (870)
692 (1014) 807 812 848 (182) 874
- Sheehan J C 561 576 586
- Shelberg W E 537 (172) 583
- Sheldon W H 110 (1015) 777 (193) 790
(1016) 790 812 845 (66) 871
- Sheline G F 530 (183) 875
- Shelley H 19 (90) 80 (95) 85 119 (44)
145
- Shelton E K 271 (349 350 351) 300
- Shemano I 668 (930) 683
- Shepanek I A 767 (1017) 810
- Shepherd D M 104 (219) 103 150
- Sheppard R 703 (813) 706 (584) 801
808
- Sherwood S I 136 147
- Shetlar M R 156 173
- Shick R M 766 (1018) 810
- Shimkin M B 243 (365) 245 (360)
306 703 (1011) 810
- Shipley I G 544 (136) 582 608 (111)
680 702 (604) 801 870
- Shipley R A 194 198 252 (353) 253
(92 203) 275 (352) 306 605 (79) 614
(81) 618 (9) 606 (144) 637 (144)
649 (144) 657 659 703 (252) 791
- Shipner L B 960 (354) 306
- Shippel S 871
- Shipton M C 15 (12) 31 (12) 48
- Shock N W 603 (1054) 813
- Shoji K 23 (264) 54
- Shoppee C W 571 (301) 523 576 (303)
527 (299) 532 540 (00) 544 (304
305) 558 (45 273) 553 (45 298) 564
(303) 565 (189 259) 579 583 585
586 587 610 677 (247) 667
- Shorey J M 154 (104) 172
- Shorr E 187 (123) 198 206 233 271
273 (355) 306 483 (571) 513 716
(1263) 819 827 (184) 875
- Showacre J L 39 (965) 54
- Shubik P 751 (60a) 753 (60a) 762 (916)
788 810
- Shulman M H 716 (1944) 819
- Shuman C R 763 (1019) 810
- Shure N 773 (1184) 817
- Shwartzman G 734 (25) 779 (1070)
1021 1077 1023) 785 812
- Sicé J 597 (118) 581
- Sichel R 768 (411) 797
- Sicuteni F 488 (655) 515
- Sidorikina M Ya 497 (656 657) 498 (656
657) 515
- Siebenmann R 765 (1024) 812

- Siecke V 747 (443) 797
 Siegel A N 828 (41) 871
 Siegel B M 746 (608) 803
 Siegel I 436 (55) 501
 Siegel S 521 (306), 525 (306) 587
 Siegel S M 39 40 (206) 54
 Sieminski J S 791 (61) 786
 Sifford R H 431
 Sikand R S 672 (179) 687
 Silber R H 154 (136) 173 700 (1219)
 723 (1219) 724 (1025) 729 (1025)
 734 (1025) 735 (1219) 750 (1219) 752
 (1210) 813 818
 Silberberg M 216 (356 357) 270 (358)
 306
 Silberberg R 270 (358) 306
 Silberberger J 22 35 (267) 54
 Silberstein F 505 (105) 658
 Silides D N 202 237
 Silprandi D 187 (124) 198
 Silprandi N 187 (124) 198
 Silver H K 830 (186) 869 (186) 876
 Silver M 376 (241) 377 386 669 (231)
 683
 Silver S 450 (658) 515 830 (186) 875
 Silverman S H 714 (1212) 818 851 (36
 235) 852 (36 233) 871 876
 Silverstein R 753 (332) 793
 Silverstone G 553 (30) 579
 Silveus E 482 (725) 517
 Simay Kramar M 716 (656) 803
 Simeox S J 827 (172) 874
 Simkins S 476 (659) 515
 Simmons F A 865 (91) 872
 Simms H D 737 (792) 806
 Simond A C 236 (42) 270 298
 Simonsen M 722 (1137) 816
 Simpson M F 702 (37 60 71) 204 (60)
 209 (29) 210 (29) 214 (67) 216 (67)
 217 (67) 218 (28) 231 232 236 237
 (405) 238 (250) 239 (22 107 151 152
 247) 240 (151 303) 241 (21 104 204
 249 361) 242 (103 151 152 248 308)
 243 (24 151 152 277 262) 244 (105
 106 151 152 248) 245 (261 262) 246
 (103) 247 (232 247) 248 (247) 250
 (96) 251 (96) 254 (237) 270 (103) 271
 (227) 275 (103) 276 (259) 287 (338)
 289 (227) 297 299 300 301 302 303
 304 305 306 307 312 369 381 483
 492 (75 560 660) 500 513 515 731
 (314) 794 867 (187) 875
 Simpson S 492 (661) 515
 Simpson S A 572 (121 507 309 509a)
 573 (308a) 574 (307 309a) 587 587
 622 (286 287) 663 669 (737) 683
 687 (1029) 702 (1026) 707 (444 1131)
 797 813 815
 Simpson S I 673 (233) 685
 Sinex F M 186 (145) 188 198 727
 (1135) 728 (1030) 813 816
 Singer B 702 813
 Singer M 130 (259 253) 150 437 (149)
 503
 Singer M J 293 (744) 303 696 (725)
 755 (725) 804
 Singh G R 396 (29) 426
 Singh T 539 588
 Sinkoff M W 213 (89) 256 (363) 265
 (82 83) 266 (82) 290 306 373 386
 726 (213 214 215 216) 790
 Sinn L 190 (135) 191 (135 136) 198
 Siperstein E 317 315 386
 Siperstein M D 576 (46) 579
 Sirek O V 712 (1032) 813
 Sirota J H 155 173
 Si son J H 709 (557) 800
 Si sons H A 767 (1033) 813
 Sita G E 552 (275) 554 (47) 563 (47)
 579 588
 Sjogren B 705 (714 1034) 707 (714) 708
 (713 714) 709 (1031) 711 (714 1034)
 804 813 813 (4) 870
 Skaggs R H 758 (143) 789
 Skanse B 440 (663) 441 (662) 499 (557)
 513 515
 Skoog F 14 (271) 18 19 (269) 22 25
 (269) 35 (267) 37 (271) 38 (271) 53
 54
 Skowran S 471 (664) 515
 Sladek S A 492 (670) 516
 Slater R J 405 (175) 428
 Slater R R 45 (300) 46 (300) 55
 Slattery P A 243 (565) 245 (365) 306
 Slaunwhite W R Jr 594 (87) 613
 (788) 657 663 667 (78) 679
 Slechta R F 671 (234) 683
 Slein M W 185 (75) 195
 Sles or A 697 (74 311) 709 (1035) 715

- (311) 732 (311) 733 (741 741) 786
794 815
- Sloven L W 483 (408) 512
- Slocombe A G 172 (301) 151 719 (416)
720 (1036) 799 815
- Slocumb C H 705 (1050) 707 (1050)
708 (1050) 709 (1050) 721 (1050) 724
(1050) 732 (1050) 753 (1050) 765
(435 496 497 883 1186) 766 (883
1186) 798 809 814 817
- Slosser J 481 (87) 486 (82) 507
- Slotka K H 604 665
- Sluczewski A 672 (235) 683
- Smelser G K 254 (364) 306
- Smith A H 187 (45) 196
- Smith A L 763 (926) 776 (926) 778
(976 927) 810
- Smith A U 490 (665) 515 693 (819) 808
- Smith B 81 (268 909) 89 398 (909) 498
- Smith C C 164 173 751 (115) 815
- Smith D C 435 (666 668) 446 (669)
497 (667 670) 515 516
- Smith D E 672 (189) 687 759 (1111)
815
- Smith D J 757 (1037) 815
- Smith D I 759 (1038) 815
- Smith D R 817 (903) 857 (203) 875
- Smith F 755 (91) 933
- Smith F A 489 (291) 506
- Smith E L 204 291 290 231 233 287
(7 379) 293 (7 135 136) 296 300
325 750 (967) 811
- Smith F M 766 (763) 797
- Smith I 595 (184) 459 759 (1046) 815
- Smith G F 6 (124) 51
- Smith G H 763 (1039) 815
- Smith G N 676 (700) 615 (200) 660
- Smith G V 861 (190) 863 875
- Smith G V S 594 663
- Smith H L 765 (497) 798
- Smith H M 490 (60) 501
- Smith J A 187 (41) 192 (36 38 40)
196 260 (113) 261 (112) 300
- Smith J M 779 (1040) 815
- Smith M C 243 (365) 245 (365) 306
- Smith M I 430
- Smith M S 11 54
- Smith O W 594 (290) 663 861 (190)
863 875
- Smith P F 239 306 491 (671) 515 827
(188 189) 875
- Smith R A 342 (713) 343 366
- Smith R C 770 (1212) 819
- Smith R H 185 (103 176) 198 257
(301) 262 (301) 305 478 (672 673)
479 (672 673) 480 (672 673) 516 727
(903 1041) 809 815
- Smith R I 437 (671) 516
- Smith R W 749 (784 1044) 768 (1043
1096) 781 (1042) 806 815 814
- Smith R W Jr 206 (90) 253 271 (355)
306 827 (184) 875
- Smith S C 497 (675) 516
- Smith S G 708 (1045) 815
- Smith S W 150
- Smith T C 676 (236) 683
- Smith V R 332 (172) 384
- Smith W W 759 (1046) 815
- Smull K 718 (746) 791
- Smyles E 500 (172) 660 667 (153) 681
- Smyth C J 764 (509) 798
- Smyth F 492 (700) 512
- Smyth F S 492 (414) 509
- Smyth N P D 781 (1130) 815 836
(709) 875
- Snair W D 243 (51) 262 (51) 265 (51)
298
- Snelling C E 751 (1047) 815
- Sniffen R C 865 (91) 870
- Snoddy C S Jr 557 (166) 583
- Snodgrass R F 71 (471) 93
- Snyder J G 695 (1048) 815
- Sobel E H 155 (138) 162 (138) 173
- Sobotka H 488 (676) 516 846 (192) 875
- Sodig H G 5 7 54
- Soffer L J 459 (677) 488 (676) 516 705
(1050) 771 (1052) 782 (1049 1051)
783 (367 568 369) 794 795 815 828
833 (127) 846 (191 192) 875 875
- Soiva K 863 (148) 874
- Sokoloff L 691 (1053) 815
- Solandt D V 498
- Solem J H 295 (368) 306
- Soley M H 436 (775) 506 837 (81) 872
- Solganits R I 482 (678) 516
- Soloman B 436 (783) 518
- Solomon D H 693 (1054) 747 (1104)
815 815
- Solomon S 602 678 663

- Solomon S S 595 (127) 609 (127) 658
 Solotorovskiy M 761 (1110) 774 (1110)
 777 (1055) 813 851
 Soloway A H 529 (309) 571 (310) 587
 Soloway S 732 (65) 786
 Somers G F 452 (118) 507
 Somerville I F 667 (217) 683 863 (193)
 875
 Somerville W 697 (1057) 713 (1056
 1057) 813
 Sommers S C 670 (238) 685 770 (1053)
 813
 Sondheimer F 521 (272) 579 (212) 537
 (316) 538 (316) 544 (281) 46 (314)
 550 552 (312) 554 (70 313) 558 (313)
 580 584 586 587 588 608 (179) 660
 Sonneborn T M 80 (472) 93
 Sonenberg M 352 586 456 (679) 680
 631 473 (679) 816
 Sorkin M 526 587
 Sorkin S Z 846 (192) 875
 Sorrentino J 829 (17) 870
 Soskin S 175 (127) 198 260 (354) 307
 Sougin Mibashan R 565 (194) 875
 Soukop S W 376 (189) 377 (189) 384
 Soukavine A 672 (239) 683
 Sourkes T I 724 (105) 1060 819
 Southam C M 79 (1061) 813
 Spadolone I 128 (138 255) 147 160
 Spain D C 743 (1065 1066) 747 (1064)
 757 (1062) 763 (1067) 766 (1065
 1066) 814
 Spain D M 20 (363) 306 743 (779)
 777 (1063) 806 814
 Spait T H 749 (1065) 814
 Sparrow F M 775 (1069) 814
 Spaul E A 417 (186) 480
 Specht N W 830 (195) 875
 Speer M I 104 (756) 160
 Speirs R S 701 (1070) 703 (1072 1741)
 746 (1070-1072 1741) 747 (848 1071)
 752 (671) 803 808 814 819
 Spence A W 461 (437) 510
 Spence T 497 (392) 509
 Spencer A G 702 (1073) 814
 Spero C B 641 (201) 660
 Sperti I 257 (79) 299
 Spies J R 708 233
 Spies T D 745 (1074) 751 (1075) 814
 Spingarn C I 710 (803) 807
 Spink W W 779 (2) 784
 Spinks A 497 (682) 516
 Spiro H M 756 (122) 790
 Spisni D 164 (132) 173
 Sprague R G 634 (195) 660 696 (884)
 697 (1079) 701 (1077 1078) 704 705
 (1080) 707 (1078 1080) 708 (1080)
 709 (1080) 714 (884) 721 (1080) 724
 (1080) 725 (1079) 727 (1079) 732
 (1080) 753 (1080) 765 (1080) 809
 814 843 (2094) 844 (110 194) 848
 (196) 873 875
 Sprechler M 671 (104) 680 741 (1081)
 814 841 (796) 876
 Spring F S 556 (37 38) 579
 Springer J J 635 (85) 657
 Sprott W I 453 (419 759) 454 (759)
 475 (683 780) 609 516 518
 Squires W I 180 (970) 811
 Srere P A 576 (46) 579
 Stack Dunne M P 137 (754) 150 219
 221 (24) 222 (24) 224 (24) 231 233
 247 (57 89 370 741) 278 (57 340)
 249 (88 370) 291 298 299 306
 Stacie W C 181 (128) 185 (179 180)
 186 (65 125 131 132 133) 197 198
 257 (372 373) 306 76 (1053) 27
 814
 Staehelin D 791 (391) 307 765 (1149)
 816
 Stafford R O 613 (46 240) 678 683
 Stahl I 575 (135) 516 (135) 589
 Stalker H D 74 (115) 86
 Stamler J 716 (1089) 717 (871) 808 814
 Stanbury J B 411 (800) 449 (684) 469
 (235 801) 505 516 519 828 (198)
 837 (198 199) 838 (198) 875
 Stand R 695 (1090) 814
 Stanley M M 440 (37) 441 (685) 449
 (685) 465 (686 687) 500 516 579
 (9 200) 830 (9) 870 875
 Staple I 640 (175) 660
 Staple T W 745 (798) 806
 Stare F J 187 (134) 198
 Starling W W 546 (250) 585
 Starr I 435 (558) 457 (558) 458 (688)
 467 (105) 50 513 516
 Stasili N R 459 (640) 516
 Statland H 459 (689) 495 (690) 516
 Staub A 190 191 193 239 (134) 300

- Stavray G W 129 132 (145 146) 147
 Stead E A Jr 305 (149) 406 (149) 409
 491 (641) 493 (641) 515
 Stears J C 619 (139) 681
 Stebbins R B 278 (416) 281 (416) 215
 (374) 308 308 665 (961) 684 701
 (1002 1215) 739 (906) 807 814 818
 Stecher R M 605 (310) 606 (310) 663
 Steelman S I 207 208 210 211 214
 216 218 219 231 235 457 (165) 504
 833 (49) 871
 Steen A S 760 (1033) 814
 Steeves F A 29 (216) 55
 Steeves T 3 (276) 29 (263) 54 55
 Stefanini M 742 (1094) 814
 Steffensen I H 768 (1043 1095 1096)
 815 814
 Steger K 479 (691) 516
 Stegwee D 77 (473) 83
 Stehle R I 305 (197) 401 491 499
 (189 191) 423 430
 Sterdle H 81 (44) 93
 Stern I F 856 (201 202) 507 (202) 875
 Stein J D Jr 251 (315) 306
 Stein K F 745 (1078) 747 (1097 1098)
 814
 Stein M 331 (200a) 384
 Stein O 743 (317) 793
 Stein R H 766 (598) 807
 Stein Y 743 (317) 793
 Steiner A 486 (692) 516
 Steinback H L 847 (203) 852 (203) 8 5
 Steinbeck A W 288 (19) 292 (19) 293
 (19) 294 (19) 297
 Steinberg H 840 (29) 870
 Steinmetz C G 768 (689) 779 (456)
 797 803
 Stephens C A L 727 (111 1099) 255
 (104) 787 814
 Stephens G C 62 (85 522) 63 (475) 66
 (476) 85 93 94
 Stephens G J 64 (477) 66 (477) 93
 Stephenson F F M 540 (10) 578
 Stepto R C 740 (1100 1101) 766 (819)
 809 814 815
 Sterling C 29 (277) 55
 Stern T N 733 (1254) 819
 Sternberg W H 342 371 385 856 (204)
 875
 Sterne G D 458 (559) 513
 Sternheimer R 483 (693) 516
 Stetten D Jr 188 198 254 (316) 307
 482 (335) 507 732 (68) 786
 Stevens A R 748 (1141) 816
 Stevens H 208 (94) 233
 Stevenson J A F 719 (947) 811
 Stevenson R 556 (371) 579
 Steward F C 23 (278) 55
 Stewart G N 715 (9 5) 810
 Stewart G S 157 16 173
 Stewart H J 493 (694 695) 516
 Stewart J 695 (185) 789
 Stewart L H 861 (205) 875
 Stewart W C 397 450
 Stickney J M 751 (1102) 815
 Studworthy G 618 (157) 619 (157) 659
 Stier R A 713 (1103) 815
 Stimmel B F 599 647 (293) 663 607
 (241) 683
 Stitch S R 667 (124) 680
 Stobaugh R E 563 (238) 585
 Stock C C 609 (611) 751 (1113) 752
 (119 1113) 788 801 815
 Stodola F H 560 (316) 581
 Stoerk H C 154 (136) 161 (27 135) 170
 173 700 (1219) 715 (646 647) 716
 (647) 723 (1219) 734 (1109) 735
 (1219) 743 (1105) 750 (1219) 751
 (1107) 753 (1219) 753 761 (1110)
 710 (1106) 771 (646) 774 (58 325
 1110) 777 (1055) 787 792 793 809
 813 815 818
 Stokem M 609 (99) 610 (99) 611 (99)
 616 (99) 633 (99) 635 (99) 636 (99)
 657
 Stokes A M 190 (89) 197
 Stokes W M 535 536 583 587
 Stoll R 438 (69 70) 501
 Stoll W A 136 150
 Stone D 695 626 (292) 628 (259 292)
 640 (200) 662 663
 Stone R E 745 (1074) 751 (1015) 814
 Storaasli J P 826 (102) 872
 Stork G 556 (73 267) 580 586
 Stormont R T 293 (317) 307
 Stover J H Jr 117 (261) 150
 Stowe B B 2 (281) 6 (280 281) 7 (281)
 29 (279) 55
 Strandh J R E 163 (50) 171

- Straube R I 672 (189) 682 709 (1111) 815
 Strauss F 415 (71) 501
 Strauss M B 757 (1062) 814
 Strauss R 863 (212) 875
 Straw, R F 659 667 (175) 681
 Street H I 23 55
 Streeten D H P 698 (1151) 696 (1151) 697 (1151) 704 (1151) 721 (1151) 727 (1151) 745 (1145) 760 (1151) 766 (1151) 768 (1151) 777 (1151) 816
 Stricker P 317 251 586
 Strickroot F L 271 (334) 305
 Striganova A R 164 173
 Strisower B 480 (276) 605
 Strisower I II 157 (34) 196
 Stroud S W 599 663
 Studer A 674 (242) 683
 Studer P I 630 (171) 680
 Stumpe W M 763 (21) 785
 Sturgis S H 670 (95) 680 857 (206) 859 (206) 875
 Sturkie P D 482 (696) 516 672 (243) 683
 Sturm F 752 (403 404) 807
 Sturtevant F M Jr 674 (244) 683
 Stutinski F 60 (478) 68 (479) 75 (480) 81 (480) 93 179 (262) 160
 Sudak F A 713 (1215) 716 (1245) 819
 Sue P 436 (388) 509
 Sugura K 751 (1112 1113) 752 (1113) 815
 Suster I F 729 (728) 804
 Sullivan J 462 (38) 500
 Sulman F G 295 (278) 307 423 (190) 480
 Sulzberger M B 754 (1114 1115) 815
 Summers C H R 544 (304 305) 559 (45 273) 559 (45) 579 580 587
 Summer A K 692 (1014) 810 849 (182) 874
 Sunderman F W 831 (207) 875
 Suneson S 62 (119) 86
 Sung C 496 (677) 516
 Suomalainen P 81 (481 482) 33
 Sur N 783 (350) 811
 Surányi I A 544 (42) 579
 Suranyi I 620 (30) 666
 Sure B 451 (699) 489 (635) 516 751 (1116) 816
 Sussmann I A 747 (556) 809
 Sutherland L W 101 (142) 190 (144) 194 193 723 815
 Sutherland K 730 (1101) 766 (879) 809 815
 Sutton I I 851 (231) 876
 Sutton R F 503 (340) 688
 Svedberg T 448 (296) 507
 Sverdrup A 209 (79) 251
 Swahn B 486 (475) 601
 Swan J M 99 (740) 150 404 (211) 409 (211 211a) 431
 Swanberg H 670 (3) 677
 Swann H G 690 (1118) 815
 Swanson C I 30 (284) 55
 Swanson I W 494 (700) 516
 Swartz I B 777 (717) 804
 Sweet M I 288 (379) 307 600 616 617 (266) 615 (735) 648 (766 296) 662 663 694 (119) 696 (119) 815
 Suedin B 153 (90) 172
 Sweeney M J 243 (122) 248 (122) 300
 Swenson R F 442 (70) 516
 Sweet W A 29 (264) 54
 Swift M A 672 (149) 692
 Swingle W W 691 (1121) 698 (1120 1121) 708 (1122 1123) 712 (1124) 714 (1122 1123 1126) 722 (1125) 815
 Swinard C A 282 (61 62) 698
 Sidnor K I 280 (80 381) 287 (459) 327 343 (205a) 800 (705) 875
 Sykes J F 366 386 667 (29) 678
 Siverton J T 703 (1198) 760 (1127) 779 (415) 796 816 817
 Sigo C M 242 (383) 204 (383) 307 669 (215) 610 (24 25 24a) 676 (745) 678 683 729 (1203) 733 784 (1178) 815 818
 Szerb J 720 (547) 801
 Szilágyi D F 791 (1130) 815 836 (709) 876
 Szmujszkowicz J 539 (173) 663

T

- Tagmann I 546 (257) 580 604 (244) 610 615 662 663
 Tagnon R F 467 (184) 518
 Tainter M I 113 160
 Tut J F 572 (121 307 308 308a) 573 (308a) 574 (307 308a) 582 587 672

- (28 287) 663 687 (1029) 707 (1026)
 707 (441 1131) 797 813 815
 Takaoka M 72 (483) 80 (483) 93
 Takemura K H 502 (60) 580
 Takewaki K 61 (484) 61 (480 486) 66
 (486) 67 (484) 93
 Taki I 79 (487) 93
 Tala P 208 233
 Talalay I 673 (246) 683 784 (1137) 816
 Talbot J 720 (294) 79
 Talbot N B 151 (32) 150 162 (32 138)
 170 173 667 (247) 683 713 (703) 804
 841 (126) 875
 Talbot T R 701 (808) 808
 Talbott J H 766 (828) 807
 Talesnik J 128 (163) 148
 Talmage R V 154 (41) 155 157 158
 165 177 173 302 333 468 (702) 516
 710 711 (362) 794
 Tamaoki B 668 (121) 673 (122) 680
 Tambiah M S 2 (18) 3 (18) 6 (18) 48
 Tang P C 77 (394) 307
 Tang Y W 3 28 (280) 79 (280) 55
 Tanzig S 47 55
 Tarail R 00 (701) 790
 Tarr R 797
 Tata J 404 (601) 514
 Titeno I 773 (632) 80
 Tatum A I 492 (703) 516
 Taub D 537 (346) 538 (346) 588
 Taub R 177 173 124 (304) 175 (300)
 151
 Taubenhaus M 704 (1130) 761 (1133)
 767 (1013 1134 1135) 810 815 816
 Tuurog A 441 (709 714) 442 (709 710
 714) 443 (107 711 713 732) 440
 (101) 446 (713) 447 (734) 449 (700
 707 712) 450 (706) 451 (91 704 700)
 452 (91 708) 461 (797) 507 516 517
 519
 Taylor A 434 (715) 517
 Taylor A B 843 (709a) 875
 Taylor B 704 (1135) 762 (1013 1130)
 81 816
 Taylor D A H 551 (43) 579
 Taylor D B 119 (764) 122 (774) 150
 Taylor F H 494 (411) 509
 Taylor F N L 741 (454) 797
 Taylor H C 861 (48) 871
 Taylor H M 706 (949) 811
 Taylor J C 790 (11) 475
 Taylor I W 767 (628) 80
 Taylor N B G 116 (708) 149
 Taylor N R W 286 (380) 307
 Taylor S I Jr 404 (176) 400 (176)
 479 480
 Tchernoff G 12 (137 e f) 14 (137 e f)
 51
 Teel H M 736 (386) 246 (387 388) 249
 (388) 700 (387) 270 307
 Teich S 540 (318) 587 590 (173) 592
 (173) 633 (173) 660
 Teicher R 754 (1136) 816
 Teilum G 777 (1137) 816 860 (710) 875
 Telfer M E 670 (248) 683
 Telfer W H 70 (488-497) 93 94
 Telikka A 491 (367) 508 782 (661) 803
 Templeton R D 151 (102 153) 174
 Teng C 106 (145) 198 777 (1138) 816
 Teng C T 186 (104 140) 198
 ten Holt S P 843 (24) 8 0
 Tennenbaum M 106 (198) 148
 Tenney A 449 (21) 500 702 (10) 785
 Teodoru C V 704 (20) 785
 Tepperman H M 166 (93) 168 172 1 3
 735 736 816
 Tepperman J 219 (90) 288 (780) 299
 307 479 (567) 490 (716) 515 517 730
 736 816
 Terepka A R 154 (90) 164 (90) 177
 Terpstra J 830 (19) 8 0
 Terpstra W 2 (286) 5 7 (286) 55
 Terroux K G 1 6 (48) 145
 Terry L L 766 (1140) 816
 Terry M L 104 (32) 155 (37) 162 (3)
 170
 Terhakovec G A 761 (780) 763 (780)
 806
 Tesar W C 217 (264) 304
 Teubner F G 5 (288) 33 (287) 42 (287)
 55
 Tew M B 806 (11) 864 (211) 875
 Tewkesbury I B 443 (331) 507
 Teys eyre J 672 (233) 683
 Thalhimer W 747 (1064) 814
 Thatcher J S 164 (147) 173
 Thayer S A 506 (34) 579 594 (69 148
 176 318) 598 (19 715) 655 656
 659 660 661 663 664

- Thiers R M 451 (699) 516 781 (1116)
 815
 Thierion M, 741 (766) 792
 Thibault C, 339 (77b) 580
 Thibault O 451 (370 371 717-720) 508
 517
 Thiersch J B 748 (1111) 818
 Thimann K V 2 (281, 290) 4 (200 291)
 6 (280 281 294 295) 7 (281 290 291)
 9 (291 293) 10 (291 293) 12 (290
 291 293) 13 (291 293) 15 (291 293
 297) 16 (291) 20 (291a) 21 (10, 106)
 24 (298 299) 25 (293 297) 27 (295)
 28 (293) 29 (279 280) 31 (171) 39
 (63 111 112 112a 113 296 297a) 40
 (63 64) 41 (61-63) 43 44 (113 295
 301) 45 46 (292 299 300) 48 49
 50 55 58 128 150
 Thing E 137 (266) 150 418 423 430
 Thoma R W, 608 (97a 98) 609 (97a)
 642 (98) 65 698 (50b) 795
 Thomas G B 295 (374) 306
 Thomas G H 554 (31) 556 (31) 579
 Thomas J F 37 (302) 65
 Thomas J R 758 (143) 788
 Thomas J W 495 (721) 517 747 (131)
 788
 Thomas K L 755 (575) 800
 Thomas L 724 (711) 743 (1142) 753
 (1143) 761 (1142) 771 (1144) 774
 (777) 778 (777) 804 808 816
 Thomopoulos T 438 (722) 517
 Thompson E C 759 (1046) 813
 Thompson E O P 170 (116 117) 190
 193 227 233
 Thompson J C 485 (723) 517
 Thompson, J L 560 (203) 584
 Thompson J W 131 (272) 149
 Thompson L M 595 (127) 603 (127) 658
 Thompson P H 419 (721) 482 (725) 517
 Thompson R 853 (212) 875
 Thompson R E 220 222 (102) 233 277
 (390) 292 (390) 294 (390) 307
 Thompson W O 449 (724) 452 (725)
 517
 Thom en F 68 (493 495 498 502) 63
 (496) 74 (491 496) 75 94 108 139
 150
 Thom en M 68 (493-502) 94 108 139
 (269) 150
 Thomson A P D 319 321 (1) 329 330
 379 386
 Thomson D L 2, 2 (30) 29
 Thore S 79 (803) 94
 Thorborg J V 673 (219) 688
 Thora G W 287 (296 302) 291 (308
 309 391) 293 (308 309) 294 (309)
 305 307 421 (84) 427 459 (204) 495
 (301) 504 507 605 (86) 615 (86) 637
 666 (750) 683 688 (1151) 696 (472
 473 1151) 697 (74 341 1057 1150
 1151) 704 (1151) 706 (571 1146) 707
 (585) 713 (1057) 714 (472) 718 (441
 52, 1150) 721 (1150 1151) 724
 (1151) 732 (341) 733 (74 341) 744
 (902 1147) 745 (1145 1148 1150) 750
 (443) 755 (455) 765 (1149 1151) 766
 (1151) 768 (1151) 777 (1151) 781
 (316 513) 786 793 794 797 799 800
 809 815 840 (213) 841 (57 214) 842
 (215) 871 875
 Thorn J W 284 (196) 502
 Thorn S W 815 (173) 874
 Thorogood I 193 199
 Thorp F 670 (169) 681
 Thor en R K 26 (541) 808
 Threlkeld A 439 (749) 517
 Thresh R 2 (13) 48
 Thurlow J 31 (334) 48
 Thygerson I 779 (633) 807 816
 Tiengo M 397 426
 Tierney N A 483 (726) 517
 Tietze I 176 (141 145) 199
 Timeris I 460 (74) 501
 Tipton I H 478 (728) 517
 Tipton S R 476 (727 728) 517 770
 (1153) 816
 Tishkoff C H 443 (723) 517
 Tishler M 554 (47 794) 558 (331) 563
 (47) 571 (168) 579 583 586 587
 Tisdall F F 161 (61) 171
 Tisclus A 221 (68) 232 404 499
 Tissières A 478 (730) 517
 Titman L C 22 (303) 47 (303a) 55
 Titus F 104 (277) 150 563 (249) 585
 608 (232) 620 (232) 642 (232) 661
 Toaff R 669 (276) 684
 Tobnan I Jr 863 (21C) 876
 Tobias C A 285 (130) 286 (130) 300
 697 (380) 795

- Tobias J M 77 (504) 94
 Toby C G 756 (827) 807
 Toennies G 148 (149) 199
 Toh C C 101 (120) 136 (18 120) 146
 147
 Tokuyama I 218 (10) 232
 Toman J F P 119 (71) 125 (70 72)
 150
 Tomasewski A J 510 (82 83) 581
 Tomich F G 490 (731) 517
 Tomkins G M 576 (46) 59
 Tompkins G 631 639 (298) 647 (708)
 648 (298) 649 (298) 663
 Tompsett R 776 (684 685) 777 (683
 685) 803
 Tong W 441 (714) 442 (714) 443 (711
 713 732) 446 (713) 447 (134) 449
 (712) 517
 Tonhazy N E 724 (14) 731 (1162 1163)
 785 816
 Toolan H W 753 (1154) 816
 Toole E H 27 (36) 48
 Toole V K 27 (36) 48
 Toompas C A 741 (35) 785
 Topper A 837 (217) 876
 Torda C 720 (1155) 816
 Tornbean J A 671 (154) 681
 Tornblom N 154 (148) 162 174
 Torrey J G 23 55
 Tortoreto J N 235 (330) 306
 Touchstone J C 634 (248) 635 (248)
 636 (248) 662
 Toverud S U 153 (112) 169 173
 Towery T R 467 (784) 518
 Tracy A 672 (256) 674 (256) 683
 Traeger C T 696 (350) 766 (355) 794
 Traut H F 851 (16) 870
 Travis D F 66 (505) 94
 Treager H S 741 (1156) 816
 Trendelenburg U 133 (56) 145
 Trentin J J 347 386
 Treshella P 843 (23) 870
 Trethewie E R 106 150
 Treverrow V 450 (733) 517
 Trezzi F 47 55
 Trikojus V M 517 781 (749) 805
 Trumberger G W 331 354 (129) 335
 (129) 338 671 (106) 680
 Tripod J 707 (229) 291
 Trippett S 406 (171) 407 408 (171 210)
 430 431
 Tristram G R 177 (150) 199
 Trotter W R 490 (266) 508
 Trounce J R 670 (216) 682
 Trust A P 488 (172) 504
 Trubowitz S 745 (862) 808
 Trunnell J B 438 (735) 517 839 (176)
 874
 Tsatsas G 534 (258) 585
 Tschopp F 559 (229) 585 698 (441 773)
 797 806
 Tsui C 22 (22) 54
 Tubianan M 449 (736) 451 (736) 517
 Tudor Williams D P H 123 (168) 124
 (168) 148
 Tuerkischer E 186 (151) 199 756
 (1157) 816
 Tui C 746 (1180) 817
 Tullner W W 672 (112) 680 692 (506
 507 977) 735 (977) 737 (234 978)
 774 (505) 791 798 811
 Tully M E 550 (139) 582
 Tulska A S 862 (118) 875
 Tum Suden C 241 (418) 308
 Tuppy H 179 (119 120) 180 198 404
 412 430
 Turchini J 60 (506) 94 436 (737) 517
 Turfitt G E 620 663
 Turnaf J 701 (1158) 816
 Turnbull J H 556 (39) 579
 Turnbull L 554 (294) 586
 Turner C D 58 (507) 94
 Turner C W 346 347 352 359 365 367
 371 375 378 379 384 388 400 (200)
 430 445 (564 565) 449 (65) 450 (423)
 452 (473) 453 (473) 501 511 515 520
 (205) 682
 Turner D L 523 (218 219) 529 (219)
 561 (217 218 219) 584
 Turner H H 271 (393) 307 853 (218)
 876
 Turner J C 740 (555 556) 800
 Turner J F 40 (248) 54
 Turner J S 46 55
 Turner L B 716 (1159) 816
 Turner R A 402 (201) 404 406 (51)
 437 431
 Turner R B 525 (259) 526 (320) 529

- (722) 561 560 (321) 571 (321 322)
574 (319) 575 (319) 585 587
- Turner T B 780 (1160) 816
- Tusini G 672 (2) 674 (2) 677
- Tuthill C H 278 (127 128) 281 (127)
300 660 (94), 680, 701 (378) 795
- Tuttle A H 439 (749) 517
- Tverdt G 102 (44) 196
- Twarog B M 100 (276) 101 (276) 100
120 (270) 120 (274 275) 133 (270)
135 150
- Tweedy W R 151 (100) 100 157 161
(149 150) 169 (79) 172 174
- Twombly G A 868 (219) 870
- Twombly G H 597 598 (301 302) 655
608
- Tyler C 416 437
- Tyler F H 281 (327) 286 (267 304) 287
(329) 304 305 307 601 (815) 636
(815) 726 (115) 730 (962) 745 (957),
788 807 811 843 (208a) 845 (156
178) 874 875
- Tyner D A 539 (342) 588
- Tyree E B 672 (189) 682 700 (1111)
816
- Tyslowitz R 462 (38) 600
- U
- Udenfriend S 104 (277) 150 176 199
- Ueberwasser H 533 (339) 588
- Uhlenhuth E 491 (738 739) 517
- Ulberg L C 671 (251 251a) 673 (55)
679 683
- Ulrick W C 430 (740) 517
- Ulrich F 200 (390 336) 307
- Ulhafer P R 528 (218 219) 579 (219)
561 (218 219) 584
- Ulstrom R A 745 (447) 737
- Umbreit W W 724 (14 1161) 731 (1161-
1163) 785 816
- Umeya T 58 (503) 94
- Umratk A 101 (40) 145
- Underdahl L O 494 (373) 508
- Ungar F 590 (60) 591 (305) 600 606
612 (705) 613 (305) 616 617 (34) 618
622 (34) 624 625 632 (80) 633 (80
303) 638 (306) 644 (80) 647 649
(80) 606 657 660
- Ungar G 498 (741) 517 749 (1166) 761
(1165) 762 (1165) 772 (1164) 816
- Uotila U 459 (743) 460 (744) 489 (742)
517 742 (1167) 816
- Upton A C 740 (1168) 741 (1168) 816
- Urbach F 780 (220) 790
- Urban F F 498
- Usang H H 134 148 150 397 427 41
Usuehi F 495 (745) 517 674 (252) 683
- Uldert I I 211 (223) 271 (87) 299 305
- V
- Vaurel F 161 174
- Vail V V 731 (652) 803
- Vaillancourt deG 718 (436 1169) 760
(669) 796 803 816
- Valere I H L 289 (98) 297
- Valeri V 674 (204) 682
- Van Bekkum D W 755 (1172) 817
- Van Bekkum G 676 (20) 678
- van Bommel A J 533 (27) 579
- Van Campenhout E 192 (153) 199
- Vancea P 164 174
- Vandemark N L 431
- Vanderberghe G 192 (44) 196
- Vanderhaeghe H 508 (323) 587
- Vander Kloot W 73 (509 510) 94
- van der Kuy A 378 383
- Vanderlaan J C 441 (748) 442 (748)
517
- Vanderlaan W P 292 (397) 295 (397)
307 441 (747 748) 442 (746 748) 455
(747) 466 (746) 517
- Van Der Scheer J 277 (264) 204 816
- Van Der Vies J 295 (173) 301 304
- Vande Wiele R 602 (991) 628 (991) 663
- van Dorp A A 569 (50) 580
- van Dyke D C 236 (399) 242 (38)
270 285 (130) 286 (130) 287 (399)
295 (126) 300 307 692 (380) 795
- van Dyke H B 210 215 218 31 234
242 (63) 244 (63) 278 (153) 298 301
396 (7 203) 400 402 (45 164) 403
(165) 416 425 426 430 431
- Vane J R 128 (57) 145
- van Middlesworth L 439 (749) 517
- van Oordt J 671 (44) 678
- van Ordstrand H S 768 817
- van Overbeek J 15 (273) 30 (222) 53
- van Putten L M 755 (1177) 817
- van Rasste M H 20 (237 238) 54

- van Roooe Belle A G C 378 (159a) 585
 van Rooven C I 767 (840) 779 (840) 807
 Van Scott I J 755 (1173) 817
 van Senden H 31 (118 267) 51 54
 van Soest F M 318 (159) 585
 van Tienhoven A 333 586
 Van Valkenburg P A 460 (370) 507
 van Wagenen G 492 (570) 517 599 (87) 617 (145) 657 659
 Van Wieringen G 249 (401) 266 (400) 307
 Vars H M 480 (723) 517 758 (303) 794
 Vartiainen S 863 (149) 874
 Vaseszky Sz 283 (409) 307
 Vasquez Lopez F 323 586
 Vaughan J H 749 (1174) 761 (316) 774 (376) 795 817
 Vaughan J R Jr 409 431
 Vaughan M 186 (133) 198 77 (1087) 814
 Vaughn D D 770 (227) 791
 Vaughn Williams E M 126 (187) 148
 Veillet A 59 (511) 61 (515) 64 (715 512 513) 65 (160) 66 (143 160 514) 67 (327) 86 8 88 90 94
 Velardo J T 315 580 669 (114 115 203) 680
 Velasco M 506 (69) 580
 Veldstra H 7 (310) 8 (307 308 310 311 312 313) 10 (308 309 312) 12 (310) 55 56
 Velick S F 176 177 199
 Velluz L 571 (324) 587
 Venning E H 700 (407) 307 605 606 631 663 693 (899) 702 738 (899) 784 (115) 809 815 817 850 (221) 861 (720) 868 (722) 876
 Venzmer G 164 174
 Verbanck M 153 (71 87) 154 (91) 155 171 177
 Vercellone A 104 (11) 144 620 660
 Verkaaik B 10 (146) 51
 Verne J 672 (254) 683
 Verney E B 395 (111a 207) 396 (207) 428 431
 Vernon L P 17 (314) 56
 Verrotti M 164 (157) 174
 Verzár F 773 (1176) 817
 Vesterdal J 741 (1081) 814
 Vestergaard P 815 (223) 876
 Vestling C S 478 (700) 517
 Veziris C D 669 (200) 683
 Vialli M 78 (516) 94
 Vianna Dias M 133 (38) 145
 Vian A 266 (98) 299
 Vicari E M 672 (206) 674 (206) 683
 Vignos P J Jr 600 (310) 606 (310) 663
 Vilanova A 833 (224) 876
 Villamil M F 442 (751) 517
 Villar Caso J 482 (752) 517
 Villee C A 186 (155) 199 669 (102) 680 727 (1177) 817
 Vimeaux J 669 (757) 683
 Vine B G 154 (85) 166 (80) 167 (85) 168 (85) 172
 Vischer F 563 (375) 571 (325) 587 608 671 641 (157) 642 (706a) 643 (206a) 644 648 (317) 659 661 663
 Visscher M B 673 (81) 674 (81) 679 745 (448 449) 751 (731) 797 805
 Vitagliano G 79 (306) 91
 Vltos A J 7 (315 315a) 56
 Vonoczky J 395 478
 Vogel A W 734 (1118) 767 (742) 768 (742) 772 (1118) 805 817
 Vogel M 700 (144) 805
 Vogt M 102 (701 282) 131 (287) 150 285 (403) 307 694 817
 Vogt Marthe 81 (517) 94
 Vogt W Jr 740 (180) 789
 Volk B W 190 (77 156) 192 (77) 197 199 243 (404) 248 (404) 256 (213) 307 307 746 (1180) 817
 Vollmer H 496 (753) 517
 Voloshin T 752 (63) 786
 Voltman A 540 (84) 581
 von Abrams G J 21 48
 Von Buddenbrock W 58 (93) 85
 von Euler U S 78 (708) 88 101 (97) 102 103 (101) 107 (107) 113 130 131 137 (96) 146 147
 von Euw J 525 527 (328) 551 (376) 508 (328) 559 (379) 563 565 (379) 571 577 (308 308a) 573 (308a) 574 (308a) 585 587 615 622 (276 787) 663 664 687 (1079) 707 815
 von Mutzenbecher P 443 (754) 518
 Voss H E 310 369 371 372 375 386

Voulgaropoulos V 648 (328) 664
 Vroman J 768 (942 943) 810
 Vuslsteke C A 192 (157) 194 199

W

- Waard J de 25 (316) 31 (317) 50
 Wachman J 190 (87) 197
 Wachowski H F 434 (415) 435 (415) 509
 Wade R 677 (123) 680
 Wagenknecht A C 29 (318) 56
 Waggener R A 478 (177) 504
 Waggoner L L 36 (319 320) 56
 Wagner K F 716 (307) 747 (307) 793
 Wagner R B 598 (214 219) 529 (219) 551 (331) 561 (219) 567 (214 330) 584 587
 Wahba N 565 (280) 586
 Wahlgren V 714 (477) 796
 Wahlstrom R C 739 (1181) 817
 Wain R L 12 (321 322) 14 (31 321) 15 (321) 16 50 58
 Wainerdi H 606 (350) 766 (350) 794
 Walaas E 195 (159) 199 669 (960) 670 (259 261) 684
 Walaas O 195 (159) 199 669 (960) 670 (258 259 261) 683 684
 Walaszek L J 221 (51) 252
 Walbach S B 740 741 (1182) 817
 Wald G 77 (248) 88 103 147
 Wald M H 462 (43) 500
 Wald N 747 (886 946) 809 810
 Waldenstrom J 745 (1183) 817
 Walderman F B 714 (1209) 818
 Walker A M 431
 Walker D G 237 (8 405) 244 (405) 297 307
 Walker J M 336 (214) 431
 Walker S M 361 386
 Walkling A V 599 (231) 661
 Wall J W 773 (1184) 817
 Wallace L R 341 536
 Wallach D P 780 (1185) 781 (1155) 817
 Wallach J B 540 (225) 876
 Wallen Lawrence 236 (390) 240 307
 Waller I R 569 (181) 583
 Wallis E S 544 587
 Wallraff E B 727 (111 1099) 735 (104) 787 814
 Walop J N 61 (518) 94
 Walters W 696 (884) 714 (884) 807
 Wang F 493 (755) 518
 Wang K J 391 431
 Wang S C 474 (48) 475 (356) 476 (47) 501 508
 Wang T V 246 (181) 249 (406) 301 307
 Wannamaker I W 778 (446) 797
 Ward I E 765 (883 1186) 766 (883 1186) 809 817
 Waring H 116 (283) 150 390 (216 216) 392 (130) 393 (217) 396 (217) 411 (217) 418 (217) 419 (216 217) 421 (134 135 136 217) 422 (134) 423 (217) 425 (218) 429 431
 Warming Larsen A 541 (246) 816
 Warren G H 764 (990) 771 (990) 773 (990) 812
 Warren J V 491 (641) 493 (641) 515
 Warren R 749 (670) 803
 Warson M D 490 (30) 512
 Wartman T G 276 (14) 231
 Warzynski R J 542 (11) 544 (11) 556 (11) 559 (11) 578
 Wassén F 716 (611) 804
 Waterhouse C 635 (35) 656
 Waterman R I 209 234
 Waterman T V 75 (519) 84
 Watkins C H 751 (1102) 815
 Watkins O 246 (548) 247 (368) 307
 Watman R V 495 (756 757) 518
 Watson D 724 (1187) 817
 Watson E M 475 (781) 518 833 (7 7) 876
 Watson S C 732 (335) 795
 Wawrzyniak M 437 (758) 518
 Wawzonick S 474 (45) 475 (356) 476 (47) 501 508
 Way E I 496 (697) 518
 Wayne F J 499 (750) 518
 Wayne H L 718 (1188) 817
 Ways B 462 (502) 464 (502) 519
 Weaver V H 770 (227) 791
 Webb H M 61 (84 88 520 525) 62 (80 86 87 89 521 522 523 524) 66 (73) 85 94
 Webb J I 623 (960) 662
 Webber L F 753 (120) 789
 Weber G 479 (760) 518
 Weber H 68 (576) 34
 Weber L 558 (242) 585

- Weber R P 3 50
 Webster B 461 (112) 502
 Webster H D 670 (169) 681
 Webster T A 217 (76) 99
 Webster W W Jr 20 (323) 56
 Wechtel C I 595 (105) 658
 Weckesser L C 703 (238) 99
 Weeks D C 41 (919) 54
 Weidlich H A 553 (333) 58
 Weigand F A 705 (901) 790
 Weil P G 850 (291) 816
 Weil R 213 (407) 254 (407) 307
 Weiland I 698 (1907) 818
 Weil Walherbe H 102 (951) 150 18,
 199
 Weimer H W 777 (1189) 817
 Weinbaum J G 691 (832) 804
 Weinberger H 271 (95) 97
 Weinmann J P 169 (15 79) 170 1 9
 Weinstein H G 498 (741) 517
 Weinstein H R 187 (41) 192 (36 38 39
 40) 193 (37) 196 960 (111 113) 261
 (119) 300
 Weinstein R C 260 (408) 307
 Weintraub A 608 (901a 235 236) 609
 (935) 641 (88 901 234) 642 (901a)
 643 (935) 657 660 661
 Weintraub R L 17 (391) 25 (96) 50 56
 Weir D R 738 (799 1190 1191) 806
 817
 Weir E G 439 (761) 518
 Weisberg H F 194 (161) 199
 Weisberger A S 743 (1192) 817
 Weisblat D I 104 (956) 160 568 (237)
 569 (237) 585
 Weise C I 488 (762) 518
 Weisel G F 598 (1375) 499
 Weisenborn F L 574 (333a) 587
 Weisenfeld S 489 (763) 490 (763) 518
 Weisman P A 775 (1193) 817
 Weiss B 442 (764) 447 (765) 518
 Weiss F 497 498 (766) 518
 Weiss M 705 (177) 707 (177) 721 (177)
 739 (177) 753 (177) 789
 Weiss P 111 150
 Weiss S B 485 (767) 518
 Weiss T 563 (41) 579 771 (926) 791
 Weiss W W 780 (715) 804
 Weissman L 836 (228) 8 6
 Weisz P 933 (409) 307
 Wei Yuan Huang 518 (99) 549 (99) 556
 (95 100) 581
 Welbourn R B 756 (1194) 817
 Welch A D 107 144 738 (1191) 817
 Welch J A 335 (156) 583
 Welch W J 708 (1192) 714 (1122) 818
 Welcher A D 410 (10) 425
 Weller I F 56
 Wells B B 700 (1195) 798 (33) 785 817
 Wells I J 278 (96) 508 603 809 813
 Wells I H 82 (921) 94
 Welsh J H 60 (39-41) 61 (598) 63 (41)
 66 (39) 18 (599 530) 19 (530) 84 94
 98 (96 27) 100 (291 293) 101 104
 (295) 105 (999 293) 106 (997 299)
 109 (96 27) 113 (27) 116 (996 987)
 190 (00) 121 122 123 124 (304)
 195 (305) 196 198 (999) 129 (289)
 130 135 (991 292 293) 136 (293 294)
 138 (26) 139 (26 94) 144 150 151
 Welt I 754 (1196) 817
 Welt I D 187 (139) 198
 Welt L G 395 (150) 4 9
 Wendler N L 552 (166) 558 (119 334)
 571 (168) 581 583 587
 Wenk I 68 (531) 94
 Wense T 103 151
 Went F W 5 (147) 7 10 (141) 14 (329)
 19 39 (329) 51 56
 Wentworth J H 759 (1197) 817
 Werder A A 153 (1198) 760 (1199) 719
 (418) 796 815 817
 Werle M D 773 (1103) 815
 Werner S C 440 (163) 458 (768) 518
 782 (1199) 817 839 (930) 876
 Wertheimer E 186 (151) 189 (17) 195
 199 415 (1) 499 816
 Wertheimer E J 754 (968) 811
 Werthessen N T 591 (253) 594 (50)
 604 (999) 661 663
 Weslaw W 488 (770) 518
 Wesson L G Jr 395 (11) 425
 West C D 484 (555) 496 (555) 487
 (555) 490 (555) 518 610 615 616
 649 664 696 (860 1201) 698 (1200)
 706 (859 860) 711 (859 860) 808 817
 West L 736 (917) 810
 West G B 102 (249) 103 150
 West T C 670 (262) 684

- Westerfeld W W 413 (771) 473 (667)
513 518 591 664
- Westeringh C van de S (312 313) 10
(312) 56
- Westermeyer V W 202 203 204 233
236 233 (12) 236 (410) 237 (292) 260
(410) 261 (410) 262 (291) 263 (291)
264 (231 292) 265 (292) 270 (290)
271 272 273 275 (290 292) 273
(259) 293 (289) 236 (410) 29 504
507 522 (163) 874
- Westman, A 367 (134) 583 733 (237)
791
- Weston R I 305 431
- Wetphal U 601 606 606
- Wetrick M I 437 (772) 461 (772) 518
- Wettstein A 552 (142 293 305) 504
(143) 557 (303) 509 (221 231 336)
563 (325) 570 (131 220) 573 (108a)
574 (3081) 582 585 586 587 588 600
(318a) 604 608 621 622 (256 287)
630 641 (152 207) 642 (152 2061)
643 (206a) 644 645 (103) 646 (207)
648 (317) 658 659 661 663 647
(1029) 707 815
- Wetmore R H 3 (76 330) 21 (106) 22
(216) 29 (330) 50 55 56
- Weyer F 138 151
- Wharton M C 717 (103) 790
- Wheeler C F 704 (116) 784 (1902) 789
817
- Whimant C 710 (1214) 818
- Whittaker W L 671 (14) 678 755 (46)
766 (47) 786
- White A 210 (103) 211 33 236 (34)
243 (383) 254 (343) 257 307 310 345
363 375 386 725 (1203) 730 (1530)
733 740 (4) 748 (108) 761 (423) 785
741 799 810 815 818
- White E 451 (503) 511
- White F P 551 (31) 876
- White H L 270 (411) 307 391 (220 221
222) 337 (222) 431
- White J M 459 (57) 501
- White M R 769 (1201) 818
- White W F 470 (200) 471 (562) 508
513
- White, W F 220 221 222 (106 109) 2 3
224 225 (39) 236 (33 106) 227 229
(114) 237 238 254 271 (110 417)
290 (165) 301 307
- Whitehead R W 764 (509) 798
- Whitehorn W V 490 (740) 517
- Whitman B 591 (321) 604
- Whitmore W F 696 (860 1901) 106
(860) 711 (860) 750 (1201) 804 817
- Whitney J L 251 (413) 290 (413) 308
330 (293) 431 708 818
- Whittaker V I 100 (16 13) 121 (309)
144 147 151
- Whittle I I 604 (187) 605 (185) 660
- Whittlestone W G 400 402 (224) 403
431
- Whorton C M 761 (767) 763 (76) 806
- Wichman R 534 (5) 870
- Wick A N 187, 187 189 722 (1906) 818
- Wickel A 757 (145) 788
- Wiedbrauck H 73 (532) 94
- Wieland I 532 (337) 533 (305) 539
(339) 588
- Wiener M 606 (119) 646 (120) 647
(190) 658
- Wiel I I 762 (21) 785
- Wiggins P I 761 (505) 768 (555) 801
- Wiglesworth V B 71 (331 578 539
511) 12 (535 536 537 538) 73 (533
534 540) 16 (542) 77 (547) 94 95
- Wight A 770 (1193) 817
- Wightman F 54
- Wijesundera S 100 (16) 144
- Wignans M 355 586
- Wilden R M 441 (40) 846 (80) 871 872
- Wilman S C 5 (332) 28 (333) 40 (331)
56
- Wilds A L 339 543 553 (340) 588
- Wiles T F 332 (172) 584
- Wilhelm A I 166 168 (146) 177 178
174 202 203 204 (92) 283 284 230
239 (33) 243 (320) 2 2 3 (33) 260
271 277 305 308 731 734 (1196) 736
(99) 787 811 817
- Wilhelms C M 114 (1208 818
- Wilkins I 211 (219) 273 (219) 40 303
483 (773) 518 603 664 701 (697) 714
(1212 735 691 782 (772) 783 (1209
1211 1213) 804 806 818 810 (121)
851 (36 124 234-236) 852 (36 234
233 236) 853 (232) 854 855 (232)
871 875 876
- Wilkins M J 44 (249) 54
- Wilkins R W 493 (1807 504
- Wilkinson J H 453 (419 770 7 9) 454

- (419) 467 (418 419) 774 777 778)
475 (418 654 653 776 780) 509 515
516 518
- Wilkinson P A 552 (25 26) 579
- Willardson D G 256 (67) 504 604
(815) 606 (815) 807
- Willebrands A F 697 (434) 796 843
(41) 872
- Willett F L 341 312 (254) 586
- Williams B S 221 (37) 431
- Williams C M 71 (543) 72 (544-546)
73 (706 454 460 509 510) 75 (137
262 375 421 456 457 458 459 460
490-497 543 547) 80 (295 453 454)
83 (546) 86 89 91 92 93 94 95
- Williams D C 667 (49) 679
- Williams H L 449 (781) 518
- Williams J H 549 (5) 550 (5 6) 552
(6) 553 (5 6) 554 (5) 558 (5) 559
(24) 560 (6) 578 579
- Williams M 440 (115) 441 (115) 450
(19) 500 507
- Williams P C 669 (237) 683
- Williams R H 187 197 436 (453) 467
(784) 469 (312) 476 (787) 507 518
728 (649 680) 735 (574 575) 799 803
- Williams R R 708 234
- Williams W 715 (1214) 818
- Williams W F 352 386
- Williams W L 348 336 721 (711) 804
- Williams Ashman H G 478 (642 673
785 786) 479 (672 643) 480 (672
673) 515 518 673 (184 763) 687 688
- Willis G C 717 (1215) 818
- Willoughby H 595 (177) 609 (127) 658
- Wilson D L 291 (391) 307 697 (1150)
699 (65) 718 (1150) 745 (1150) 786
816
- Wilson E 551 (168) 583
- Wilson G E 435 (757) 518
- Wilson H 815 (237) 8 8
- Wilson I B 106 (706) 191 (206 310 311
312) 149 151
- Wilson J C 779 (1058) 813
- Wilson M 465 (788) 518
- Wilson M L 676 (700) 645 (200) 660
- Wilson P D 715 (194) 780
- Wilson R 504 (87) 657
- Wilson W 556 (39) 579
- Wilson W P 718 (967) 811
- Wilske C 3 (334) 56
- Winkler A W 154 173 484 (411) 509
- Winne R 771 (474) 773 (474) 796
- Winnick T 452 (407) 509
- Winniford R S 563 (238) 585
- Wintenberg S 339 (575) 580
- Winter C A 775 (416) 279 (415) 281
(416) 308 665 (764) 684 700 (1219)
701 (1218) 723 (1719) 735 (1219) 750
(1719) 755 (1219) 763 (1216) 818
- Winternitz W W 723 (1700) 818
- Winters R W 737 (981 1773) 738 (1221
1772) 811 818
- Wintersteiner O 540 (164 244) 542
(748) 543 (164 165) 583 585 594
604 659 664
- Wintrobe M M 748 (463) 797
- Winzler R J 474 (212) 475 (213) 476
(213) 505
- Wirth P M 455 (138) 503
- Wise J 509 (82) 617 (81) 657
- Wolck G B 374 536 861 (238) 876
- Wisnicky W 342 (43) 580
- Wiswell J G 649 (325) 664
- Witham A C 155 1 0
- Witschi E 675 (266 267) 684
- Witt E 480 (400) 509
- Witten C L 672 (268) 684
- Witten V H 754 (1114 1115) 815
- Wittle F L 525 (215 219) 579 (215
219) 561 (219) 584
- Wittwer S H 5 (241) 54 56
- Wolf A 762 (603) 801
- Wolf A D 162 174
- Wolfe J K 667 (747) 683
- Wolfe S 765 (349) 794
- Wolferth C C 696 (1758) 714 (1258)
819
- Wolf B 75 (547) 95
- Wolf H G 770 (1155) 816
- Wolf J 438 (706) 441 (797 795) 443
(491 797 794 795) 447 (700) 449
(789) 464 (797) 473 (793) 518 519
- Wolffbrandt G 468 (350) 508
- Wolfman M 395 (219) 431
- Wolfson W Q 770 734 278 (417) 293
(417) 294 (417) 308 727 (929) 732
(979) 744 (1224) 765 (979) 781 (64
1775) 786 810 818 873 (739) 876
- Wolsky L 451 (777) 506
- Wong A S H 693 (591) 801
- Wong H 763 (66) 786

- Westerfeld, W W 443 (771) 479 (567)
513 518 594 664
- Westeringh C van de S (312 313, 10
(312) 86
- Westermeyer, V W 202 203 204 233
236 243 (12) 246 (410) 247 (292) 249
(410) 261 (410) 262 (391) 263 (291)
264 (291 232) 265 (312) 270 (290)
271 272 273 274 (290 292) 293
(289) 235 (259) 296 (410) 297 304
307 822 (169) 874
- Westman A 367 (154) 383 733 (237)
791
- Weston R I 395 431
- Westphal U 604 606 686
- Westrick M L 437 (772) 464 (772) 518
- Wettstein A 552 (112 293 335) 551
(143) 557 (293) 559 (229 231 336)
563 (325) 570 (131 230) 573 (305)
574 (308) 582 585 586 587 588 600
(318) 604 605 621 622 (256 287)
670 641 (152 207) 642 (152 206)
643 (206) 644 645 (153) 646 (207)
648 (317) 658 659 661 662 687
(1029) 707 818
- Wetmore R H 3 (276 330) 21 (106) 22
(216) 23 (330) 50 55 56
- Weyer F, 138 151
- Wharton M C 777 (193) 790
- Wheeler C F 724 (176) 754 (1202) 789
817
- Whismint C 715 (1214) 818
- Whitaker W L 671 (14) 678 755 (16)
766 (47) 786
- White A 210 (103) 211 255 236 (34)
243 (383) 254 (353) 297 307 310 345
369 375 386 754 (1203) 730 (530)
733 743 (4) 748 (254) 761 423) 785
791 799 810 815 818
- White E. 451 (503) 611
- White F P 851 (231) 8 6
- White H I 270 (411) 307 314 (220 221
222) 395 (222) 431
- White J M 459 (57) 601
- White M R 768 (1704) 818
- White W F 470 (250) 471 (562) 506
513
- White W F 270 221 222 (106 109, 223
224 225 (39) 226 (39 106) 227 270
(114) 232 250 254 279 (165 412)
550 (165) 301 307
- Whitehead R W 764 (509) 738
- Whitehorn W V, 490 (740) 517
- Whitman B 594 (324) 664
- Whitmore W F 696 (560 1201) 106
(560) 711 (560) 750 (1701) 808 817
- Whitney J L 251 (413) 290 (413) 298
395 (223) 451 709 818
- Whittaker V I 100 (16 135) 121 (309)
144 147 151
- Whittle I L 604 (187), 605 (155) 660
- Whittlestone W C 400 402 (274) 403
431
- Whorton C M 761 (707) 763 (767) 806
- Wichman P 834 (5) 870
- Wick A V 182 157 199 722 (1706) 818
- Wickel A 757 (145) 788
- Wiedbrauck H 71 (532) 84
- Wieland J 532 (337) 533 (335) 559
(339) 688
- Wiener M 606 (119) 646 (170) 647
(120) 658
- Wieland J L 763 (21) 85
- Wiggins R I 767 (585) 768 (585) 801
- Wigglesworth A B 71 (334 538 539
541) 72 (35 336 337 539) 73 (337
538 540) 6 (342) 7 (547) 94 95
- Wight A 775 (1191) 817
- Wightman I 54
- Wigesundern S 100 (16) 144
- Wijmans M 358 886
- Wildner R M 841 (40) 846 (59) 847 847
848
- Wildman S C 5 (337) 75 (3331 40 (331)
56
- Wild A I 539 543 553 (340) 688
- Wiley T F 332 (172) 384
- Wilhelm A F 166 168 (146) 172 175
174 205 203 204 (92) 233 234 236
239 (33) 243 (340) 252 25 (33) 265
271 297 405 509 731 754 (1196) 736
(99) 787 811 817
- Wilhelm C M 714 (1205) 818
- Wilkins I 271 (219) 273 (219) 307 303
453 (7 3) 518 673 684 701 (697) 714
(1717) 735 (697) 752 (772) 753 (1707
1211 1213) 804 806 818 850 (124)
851 (36 124 734 236) 852 (6 233
233 236) 853 (232) 854 855 (232)
871 873 876
- Wilkins M J 44 (249) 84
- Wilkin R W 493 (180) 504
- Wilkinson J H 453 (410 775 79) 454

- (73) 467 (418 413 744 777 778)
475 (418 604 683 776 780) 509 515
516 518
- Wilkinson P A 502 (90 26) 579
- Willardson D G 286 (967) 304 604
(815) 696 (915) 807
- Willebrands A F 607 (434) 796 813
(71) 870
- Willett E I 341 312 (207) 386
- Williams B S 91 (32) 231
- Williams C M 71 (543) 72 (541-546)
43 (996 454 460 509 510) 70 (132
762 370 421 406 407 453 409 460
400-492 543 547) 80 (990 453 454)
83 (546) 86 89 91 97 93 94 95
- Williams D C 667 (49) 679
- Williams H L 479 (781) 518
- Williams J H 549 (5) 550 (5 6) 552
(6) 503 (5 6) 504 (5) 558 (5) 509
(24) 500 (6) 578 579
- Williams M 440 (115) 441 (115) 450
(19) 500 507
- Williams P C 669 (232) 683
- Williams R H 187 197 436 (783) 467
(784) 469 (312) 406 (480) 507 518
423 (679 680) 435 (574 575) 799 803
- Williams R R 208 234
- Williams W 715 (1214) 818
- Williams W F 352 386
- Williams W L 348 386 724 (711) 804
- Williams Ashman H G 478 (672 6 3
785 786) 479 (672 673) 480 (672
673) 515 518 673 (184 263) 682 684
- Willis G C 717 (1215) 818
- Willoughby H 590 (127) 609 (127) 658
- Wilson D L 901 (391) 307 697 (1150)
699 (60) 718 (1150) 745 (1150) 786
816
- Wilson E 571 (168) 583
- Wilson G E 405 (787) 518
- Wilson H 845 (237) 876
- Wilson I B 106 (206) 121 (206 310 311
319) 149 151
- Wilson J C 779 (1008) 813
- Wilson M 465 (788) 518
- Wilson M L 676 (900) 645 (200) 660
- Wilson P D 775 (194) 790
- Wilson R 594 (81) 657
- Wilson W 556 (39) 579
- Wilson W P 718 (967) 811
- Wilske C 3 (334) 56
- Winkler A W 154 173 484 (411) 509
- Winne R 771 (424) 773 (424) 796
- Winnick T 450 (402) 509
- Winniford R S 563 (938) 585
- Wintemberger S 339 (57b) 580
- Winter C A 248 (416) 249 (415) 281
(416) 303 665 (264) 684 700 (1919)
701 (1918) 773 (1219) 735 (1219) 750
(1219) 755 (1219) 763 (1216) 818
- Winternitz W W 723 (1270) 818
- Winters R W 737 (981 1223) 738 (1221
1222) 811 818
- Wintersteiner O 540 (164 247) 542
(248) 543 (164 165) 583 585 504
604 659 664
- Wintrobe M M 748 (463) 797
- Winzler R J 474 (912) 475 (213) 476
(213) 505
- Wirth P M 455 (138) 503
- Wise J 599 (82) 617 (81) 657
- Wislocki G B 374 386 861 (238) 876
- Wisnicky W 340 (43) 380
- Wiswell J G 649 (325) 664
- Witham A C 155 1 0
- Witschi E 670 (966 267) 684
- Witt E 480 (400) 509
- Witten C L 642 (268) 684
- Witten V H 754 (1114 1115) 815
- Wittle E L 528 (215 219) 579 (215
219) 561 (719) 584
- Wittwer S H 5 (241) 54 56
- Wolf A 762 (603) 801
- Wolf A D 167 174
- Wolfe J K 667 (247) 683
- Wolfe S 765 (349) 794
- Wolferth C C 696 (1208) 714 (1258)
819
- Wolff B 70 (541) 95
- Wolff H G 1 0 (1155) 816
- Wolff J 438 (96) 441 (797 790) 443
(91 797 794 795) 447 (790) 449
(789) 464 (791) 443 (793) 518 519
- Wolffbrandt G 468 (350) 508
- Wolfman M 395 (919) 431
- Wolfson W Q 220 234 248 (417) 293
(417) 294 (417) 308 727 (979) 732
(979) 744 (1224) 765 (979) 781 (64
1270) 786 810 818 873 (939) 876
- Wolsky L 451 (777) 506
- Wong A S H 693 (591) 801
- Wong H 763 (66) 786

- Wood F H 715 (166) 789
 Wood G W 553 (30) 551 (31) 556 (31)
 579
 Wood J I 437 (709) 519
 Wood J W 759 (1276) 818
 Wood R M 777 (1236) 818
 Wood T R 166 175
 Wood W B 763 (397) 778 (398) 795
 Wood W W 533 (343) 588
 Woodbury D M 207 (35) 221 (93) 231
 233 375 376 (32 33) 580 703 (1228
 1229 1232) 719 (1227 1228 1229
 1231 1232) 782 (1230) 818
 Woodbury I A 270 (89) 233 276 (332),
 305
 Woodruff M F A 693 (1233) 818
 Woods A C 768 (1234 1235) 777 (1236)
 818
 Woods G F 553 (30) 554 (31 32) 556
 (31 32) 579
 Woods K A 698 (512) 751 (511) 782
 (512) 799
 Woodward C 691 (811) 807
 Woodward E R 759 (810) 807
 Woodward R B 537 (346) 538 539 540
 (344) 588 592 (326) 593 (376) 664
 Woollett F A 452 (118) 502 490 (731)
 517
 Woolley D W 136 151 475 (709) 519
 Woolley G W 751 (1238 1239) 752
 (1238 1240a) 764 (1237) 810
 Worcester J 713 (703) 804 841 (126)
 875
 Worcester R L 835 (167) 856 (167) 874
 Work T S 136 (78) 146
 Worrel C S 626 (200) 645 (200) 660
 Wort D J 42 (335) 56
 Worton P J 764 (989) 812
 Wotiz H H 618 (163) 648 (163) 659
 664
 Wragg L E 703 (1241) 746 (1241) 819
 Wrenn T R 365 386
 Wright, B M 462 (802) 464 (802) 519
 Wright, P A 81 (548) 95
 Wright P M 674 (269) 684
 Wright W S 192 (80) 197
 Wroblewski B 488 (710) 518
 Wrong N M 770 (1242) 819
 Wuerch J 684
 Wurl O A 770 (502) 798
 Wurzel M, 121 (20) 144
 Wickoff R W C 816
 Wyman G J 765 (1213) 819
 Wyman L C 241 (418), 308 695 (1048)
 713 (1215) 716 (1244 1215) 815 819
 Wingarden J B 441 (800) 462 (801
 802) 464 (802) 519
 Wyshak G H 188 (164) 199

X

 Xuereb C P 320 386

Y

 Yager A 747 (946) 810
 Yagi Y 436 (803) 519
 Yalow A A 436 (50) 501
 Yalow R S 781 (82) 767 829 (17) 870
 Yamada E Y 731 (479) 798
 Yamaki T 3 (337) 5 (337) 7 28 (337)
 39 (336) 40 (336) 56
 Yamamoto Y 67 (549) 95
 Yashin R 553 (347) 588
 Yasuna J M 770 (834) 807
 Yates C H 595 (127) 609 (127) 658
 Yatsushika K 72 (275) 89
 Yeates N T M 317 (263) 370 386
 387
 Yeatman J N 56
 Yeum C S 44 (301) 55
 Yoffey J M 744 (540 1247) 747 (540)
 748 (540) 799 819
 Yohalin S B 830 (186) 875
 Yoshida H 497 (336) 507
 Yoshida M 59 (309) 61 (551) 65 (550)
 90 95
 You R W 481 (651 804) 515 519 758
 (992) 812
 You S S 481 (651 804) 515 519
 Youhotsky I 536 (57a) 580
 Young F G 137 (257) 160 185 (103)
 193 (11) 195 198 219 221 (24) 272
 (24) 224 (24) 231 233 236 (422) 241
 (156) 243 (70 71) 246 (472) 247 (76
 472) 248 (147 148) 251 (419) 252 256
 (268 419 420) 257 (301 311 425)
 258 (37 246 471 475) 260 (29 470)
 262 (71 234 245 246 301 302 420)
 263 (246) 265 (71 420) 268 (472 423
 424 425) 269 (74) 270 (70 75) 272
 (29) 277 (371) 297 298 299 301 303

- 304 305 306 308 312 313 314 365
 366 399 383 387 77 (1903) 809
 Young F W 670 (169) 681
 Young J Z 78 (59) 79 (59) 95
 Young W C 60, (9a) 657 673 199
 209 680 687
 Yuzrazval S 704 (92) 792
- Z**
- Zacks S I 191 161
 Zaffaroni A 288 (92) 304 599 (12)
 684 301 (331) 606 (133) 608 (179)
 673 (133) 674 (133) 675 (133) 674
 (330) 658 670 684
 Zahler H 666 (90) 684
 Zaimis F J 195 (293) 149
 Zalesky M 278 (496) 308
 Zambotti V 41 (337a) 66
 Zamecheck N 741 (1156) 816
 Zamecnik P C 672 (74) 674 (74) 69
 Zappavodi P 77 (716 717) 804
 Zarrow I G 747 (1950) 785 (1951) 819
 Zarrow M N 331 (966) 734 335 387
 604 (961) 67 66, (771) 668 (21)
 670 (6) 671 (721 211) 678 683 684
 742 (1948 1950) 781 (1949) 83
 (1251) 819
 Zatman L J 117 (47) 146
 Zavadovski M M 40 387
 Zawadowsky B 439 (90a) 819
 Zeckwer I T 155 (12a) 819
 Zeller F A 128 161
 Zerahn K 131 148 150 337 (15b) 496
 Zerbe J W 758 (53) 794
 Zetler C 391 (101a) 48
 Zetterstrom R 158 171 174
 Ziegler M R 719 (748) 805
 Ziegler I 74 (12a) 575 (191) 687
 Zierler K L 70a (1253) 819 83, (240)
 876
 Ziff M 740 (9a) 781 (658) 787 803
 Zilli E 672 (180 181) 674 (180 181) 687
 Zilvermit D B 733 (1954) 819
 Zimmermann H J 795 (1255 1256) 819
 Zimmermann I W 9 (358 339) 66
 Zimmermann B 192 (86) 193 197 199
 Zingg W 781 (1957) 819
 Zintel H A 696 (1258) 714 (1958) 819
 859 (50) 871
 Zizine A 679 (773) 684
 Zizine L 778 (477) 308 701 (1158) 816
 Zizmer N 436 (56) 501
 Zofmann A L 482 (752) 517
 Zolman B I 749 (951) 811
 Zondek B 595 664 665 (274) 669 (276)
 670 (95) 671 (274) 684 784 (1959)
 819
 Zorn I M 155 (151) 174
 Zubiran J M 106 (1960) 759 (810) 807
 819
 Zubirán S 337 387 730 819
 Zucker M B 104 (317) 161
 Zuckerman S 317 320 329 330 331
 380 396 387 399 47 431 609 (99)
 671 (218) 675 (277) 684 784 819
 Zweifach B W 716 (1963) 819

Subject Index

A

- ACE see Adrenocortical extract
- ACTH see Adrenocorticotrophic hormone
- ACTX see Corticotropin A
- ADH see Antidiuretic hormone
- Abortion
 - hormonal imbalance and 862
 - serum protein bound iodine and 863
- Absinthin
 - as auxin inhibitor 26
- Acetate
 - as precursor of adrenocortical steroids 627-628
 - of Δ^4 androstene-3,17-dione 612
 - of cholesterol 591-592, 593
 - of estrogens 596
 - of testosterone 612
- 3 β Acetoxyallopregnan-20-one
 - labeled preparation 577
 - microbiological reactions 613-614
- 3 α Acetoxy Δ^4 cholemic acid methyl ester
 - preparation of labeled 517
- 3 α Acetoxy $\Delta^9(10)$ etanic acid methyl ester
 - synthesis 538
- 21 Acetoxy 20 ketosteroids
 - conversion to etanic acid 571
 - synthesis 569-570
- 3 α Acetoxypregnane-11,20-dione
 - labeled preparation 578
- 6 Acetoxyprogesterone 547
- 3 Acetyl-6-methoxybenzaldehyde as auxin inhibitor 26
- Acetylcholine 100-101, 106-107, 119-130, 143
 - action, calcium and 126
 - on cellular metabolism 129-130
 - concentration and 127-129
 - depolarizing 126
 - mechanism of 125-127
 - muscarinic 125
 - nicotinic 125
 - site of 119-121, 133
 - structure and 122-123
 - trophic 129-130
 - on vertebrate heart 128
 - adrenalin and 131
 - antagonists 120, 123
 - biosynthesis 99, 106
 - site of 143
 - blocking agents for 124, 125
 - chemical nature 99, 143
 - in crustaceans 61
 - distribution 98, 100-101
 - in insects 77
 - interaction with cholinesterases 121
 - intermedia and 100
 - in mollusks 79, 80
 - in nerve cells 106
 - as neurohumor 142
 - physiological role 119
 - reaction with acetylcholinesterase 128
 - receptors for 120, 121-123
 - cholinesterase and 121, 122
 - combination with drugs 123
 - linkage between acetylcholine and 123, 124
 - nature of 125, 126
 - reaction of acetylcholine with 126, 128
 - red cell cholinesterase and 128
 - relation to cell components 122
 - release 99, 113, 114
 - chemicals affecting 11
 - mechanism 114
 - storage in nervous tissue 101
 - thyroxine and 107
 - as transmitting agent 114
 - transport axonal 113
 - in worms 99
- Acetylcholinesterase reaction with acetylcholine 128
- N-Acetylhistidine, hormonal activity 475
- Acid-base balance, adrenocortical steroids and 708-709

- Acromegaly 83-85
 basal metabolic rate in 84
 diabetes as complication of 84
 growth hormone and 273 573-874
 secretion of corticotropin in 273
 of thyrotropin in 773
 Addison's disease 840-843
 blood adrenocortical steroids and 691
 diagnostic procedure 840-841
 electrolyte metabolism and 841
 etiology 840
 pregnancy and 841-842
 renal clearance and 709
 treatment of 842-843
 effect of cortisone on 697 69 711 713
 of DCA in 105
 Adenohypophysis see Pituitary anterior
 Adenosinetriphosphate
 as energy source in choline acetylation 106
 insulin and 184
 Adipokinin 251 254-256 296
 Adrenal cortex
 activity blood steroids and 691
 secretory control of 290
 corticotropin and 290-291
 anterior pituitary and 219
 atrophy as cause of Addison's disease 840
 following adrenal steroid therapy 101
 cancer 1 ketosteroid excretion and 848
 capillaries and 716
 cardiovascular system and 713-717
 control of 692-693
 β dehydrogenase in 678
 diseases response to cortisone and 847
 effect of ACTH on 675
 of amphenone B on 694
 on blood 744-748
 of corticotropin on 287-290
 in vitro 288-290
 of cortisone on 291
 of DDD on 691-692
 of erucic acid on 692
 of hormones other than corticotropin on following hypophysectomy 278
 growth and 241
 hexokinase and 184
 hormones of see Adrenocortical hormones Adrenocortical steroids and individual compounds
 hydroxylases in 699
 hyperfunction 641-644
 effect on ovaries 842-869
 hypotheses concerning 642ff
 symptoms of 773
 hyperplasia congenital 841
 diffuse effect on production of 11 oxygenated steroids 840
 pathogenesis 850
 17 ketosteroid excretion in 848
 hypofunction see Adrenal cortex in sufficiency
 insufficiency 694-704 840 see also Adrenalectomy
 treatment of 697-698 707
 vasomotor response in 716
 insulin and 184 186
 kidney function and 704 712
 lymphoid tissue and 742-744
 malignant tumors and 750-753
 malnutrition and 136
 morphology of 690-697
 muscle function and 720-721
 nervous system and 717-720
 overactivity see Adrenal cortex hyperfunction
 physiology of 684-819
 response to ACTH liberation 689
 role in carbohydrate metabolism 721-727
 in electrolyte metabolism 688
 in fat metabolism 732-736
 in protein metabolism 127-132
 in salt metabolism 395 873
 in water metabolism 394 395 688 719
 as source of androgens 610 612
 of progesterone 601 602 603
 stimulation systemic effects of 791-792
 effects of cortisone and 791
 of hydrocortisone and 291
 stress and 688-689 757
 Adrenalectomy
 effect of cortisone on electrolyte metabolism following 706
 on animals 695

- on malignancy 756
- on man 696
- response of bone marrow to 756
- survival following pantothenic acid and 738
- Adrenaline 102-103 107-108
- acetylcholine and 131
- action mechanism of 133-134, 195
- reversal of 131-133
- site of 133
- biosynthesis 107
- site of 113 143
- chemical classification 143
- nature 99
- effect on adrenal cortex of hypophysectomized animals 278
- on ACTH release 692
- on corticotropin secretion 283-285
- on eosinophils 745
- on gonadotropic activity of anterior pituitary 330-331
- on histamine toxicity 757
- in invertebrates 81
- in man, 284
- on potassium metabolism 708
- on thyroid function 459
- liver glycogen breakdown and 194
- as neurohormone 99
- noradrenaline and 150 131
- occurrence in invertebrates 80
- in nerve cells 98
- in vertebrates 102
- phosphorylase and 723
- physiological role of 130-131
- receptors for 134
- response to age and 693
- storage in nervous tissue 107
- thyroxine and 497
- transport axonal 113
- Adrenals see also Adrenal cortex
- atrophy following administration of cortisone 277
- following hypophysectomy 277
- cancer of biosynthesis of androgens in 651
- corticosteroid production in 652
- lack of 3 β dehydrogenase in 652
- masculinizing effect 651 652
- urinary dehydroepiandrosterone and 628
- 17 ketosteroids and 612
- cholesterol of ACTH and 735
- cortisone and 735
- Cushing's syndrome and 813
- diseases of 840-852
- effect of ACTH on 275 276-279 691
- of 2,2-bis (p-chlorophenyl) 1,1-dichloroethane on 819
- on reproductive system 784
- function ascorbic acid and 740
- pyridoxine and 743
- human adrenaline and noradrenaline content of 102
- hyperfunction effect of ACTH on 783
- of cortisone on 783-784
- gonads and 783-784
- hyperplasia, diffuse differentiation between adrenocortical tumor and 850-851
- pathogenesis of, 850
- medulla dopa decarboxylase in 107
- phospholipids of ACTH and 735
- relationship between pituitary pancreas and 726-727
- response to ACTH age and 693
- response to stress 701
- size effect of corticotropin on 216-219
- as source of estrogens 593
- ACTH and 595
- of progesterone 602
- of steroid hormones 590 593 595 602
- of steroid hydroxylases 645, 646
- thyroid and 458 495 780-783
- transplantation 693
- vitamins and 736-742
- Adrenocortical extracts
- amino acid metabolism and 723-729
- effect on renal clearance 710
- growth inhibition by 699
- Adrenocortical hormones 672-644 see also Adrenocortical steroids and individual compounds
- administration methods of 696-697
- biosynthesis adrenocortical hyperfunction and 652ff
- ACTH and 259 673 652
- mechanism of 623 677
- sites of 691
- in blood 693-695
- species differences 691
- circulating effect on ACTH action 692

- effect on corticotropin secretion 281-282
- on thyroid function 459
- growth hormone and 241
- metabolism of 559 631-640
- response of eosinophils to 744-745
- secretion corticotropin and 288
- structure 551
- terminology 630
- Adrenocortical steroids see also Adrenocortical hormones and individual compounds
- activity cytotoxic 703
- lymphotoxic 744
- antiinflammatory 760 761-766
- structural requirements for 541
- orientation of side chain and 526
- ascorbic acid and 219 220
- assay of 701-704 742
- atherosclerosis and 716-717
- biologically active 627
- inactive 622
- biosynthesis in adrenogenital syndrome 652-653 654
- in blood control of corticotropin secretion by 281
- diurnal variation in concentration 286
- brain lesions following administration of 719
- effect on ACTH production 281 630
- on anaphylactic reactions 771-773
- species variations in 772
- on antibody synthesis 760
- on antigen-antibody reactions 773-775
- on bone 753-754
- on bone marrow 746-748
- on carbohydrate utilization 750
- on cellular response to infection 660 761
- on cerebral blood flow and metabolism 718
- in chronic leukemia 655
- on electroencephalogram 718
- on experimental cancer 451 752
- on intracellular fluid and electrolytes 710
- on joint inflammation 764-765
- on purines 722
- on red blood cells 748
- on renal clearance 709-710
- species differences in 710
- on resistance to toxins 757-758
- on rheumatoid arthritis 765-766
- on stomach 555-556
- on susceptibility to infection 776-780
- on sweat 712-713
- on tissue cultures 769-770
- on tissue damage 758
- on water excretion 711-712
- on wound healing 768
- enzymes and 723-724 730-732
- eosinopenia following administration of 746
- growth inhibition by 699 700-701
- hydroxylation enzymatic 645-646
- mechanism of 646
- hypersensitivity and 760
- insulin and 726
- maintenance of lactation and 784
- metabolic effects 460
- metabolism differences between in vitro and in vivo 632
- effect of anethole on 692
- microbiological reactions 640-641
- nature of 640ff
- placenta as source of 812
- psychotic reactions following treatment with 717-718
- response to stress and 701
- secretion in Cushing's syndrome 815
- sensitivity reactions and 770-771
- untoward effects of 699-701 717-718
- urinary in toxemia of pregnancy 863
- Adrenocorticotrophic hormone 681 688
- 822 see also Corticotropin(s)
- activity 822
- adrenal ascorbic acid-depleting 423 424 425 742
- galactopoietic 355
- mode of administration and 696
- adrenogenital syndrome and 602 652 653 654
- alkalosis following administration of 709
- amino acid metabolism and 727 728
- antithyroid drugs and 783
- B hormone and 421 423-425
- biosynthesis adrenocortical steroids and 630 652

- in adrenogenital syndrome 652 653 661
- blood adrenogenital syndrome and 602
- in diffuse adrenocortical hyperplasia 860
- clotting and 749
- cellular origin 823
- cortisol and 630
- cortisone and 701
- diabetes and 726
- effect on adrenal cortex 625
 - on adrenals 691 736
 - on bone fractures 767
 - on electrolyte metabolism 706
 - on fibrosis 768
 - on hypopituitarism 826
 - on nephrotic syndrome 704
 - on red blood cells 748
 - on scurvy 741
 - in humans 741-742
 - on skin grafts 775
 - on stomach 755
 - on syphilis 780
 - on thyroid function 459
 - on trichinosis 780
- glutathione and 724 726
- intermedin and 137
- ketogenesis and 735 736
- liberation adrenal response to 688-689
- multiplicity of 822
- pantothenic acid and 737
- pituitary thyrotropin and 783
- placenta as source of 812 861
- polypeptide nature 137
- prevention of thyroid crisis by treatment with 836
- production of androgens and 652 654
 - of adrenocortical steroids and 623 625 627 628 652
- release estrogen and 671
 - mechanism of 692
- response of eosinophils to 45 746
- role in Cushing's syndrome 652 843-844 845
 - in estrogen production by adrenals 595
- secretion in diffuse adrenal hyperplasia 850
- Adrenocorticotropins see Adrenocorticotrophic hormones and Corticotropins
- Adrenogenital syndrome 848-862
 - ACTH and 602 662 653 661
 - biosynthesis of adrenocortical steroids and 663 664
 - of androgens in 661 862
 - differentiation between adrenocortical tumors and adrenocortical hyperplasia in 861-862
 - histologic picture of ovaries in 869
 - excretion of 17 ketosteroids in 848
 - of pregnanetriol in 848-849
 - lack of steroid hydroxylases in 663 654-656
 - manifestations 849
 - salt losing type 851
 - 21 hydroxylation and 663
 - treatment 862
 - symptoms 218 hydroxylase and 663
 - treatment 653 862
- Androstosterone
 - acetate as precursor of 612
 - from cortisol 611
 - from 17 hydroprogesterone 611
 - metabolism 619 621 664
 - metabolites of 619 635
 - microbiological reactions 620
 - sources 610 612
- Alcohols structure molecular rotation and 536 536
- Aldosterone 542-544 622 706-707
 - activity 572 707
 - ACTH and 707
 - biosynthesis 627 629 630-661
 - growth hormone and 631
 - in blood 694
 - effect on electrolyte metabolism 70
 - on salt and water retention 232
 - isolation 572 707
 - metabolic effects 687 688
 - physiology and therapeutic action 846
 - possible origin 107
 - secretion corticotropin and 290
 - structure 572-574 687
- Alkaloids 5 hydroxytryptamine structure in 136
- Alloctanic acid 522
- Allopregnane derivatives uranediol as metabolite of 566
- Allopregnane 3 α 20 α diol
 - from allopregnane 3 20 dione 606
 - metabolism 603

- from prog. sterone 605
sources 605
- Allopr. gnane-3 α 20 β diol
from allopr. regnane-3 α 20 β diol 606
metabolism 603
sources 605
- Allopregnane-3 α 20 β dione
metabolism 603 606
microbiological reactions 643 644
from progesterone 606
sources 601
- Allopregnane-3 β 11 β 17 α 20 β 21 β pentol
6 β
- Allopregnane-3 β 11 β 17 α 20 β 21 β pentol
as metabolite of cortisol 630
- Allopr. gnane-3 β 17 α 20 β 21 β tetrol 62 β
- Allopregnane-3 β 17 α 20 α triol 6 β 2
- Allopregnane-3 β 16 α 20 β triol 562
from tyrogenin 562
- Allopregnane-3 β 16 α 20 α triol 561
configuration 563
- Allopregnane-3 β 16 α 20 β triol 561 563
configuration 562
- Allopregnane-3 β 20 β 21 β triol as metabo-
lite of deoxycorticosterone 638
- Allopregnan 20 β of configuration 535
- Allosan diabetogenic activity 159
- Amenorrhea in myxedema 494
- Amine oxidase liver thyroxine and 497
- Amino acid metabolism effect of adreno-
cortical steroids on 777-779
factors affecting 779
- Amino acids
in B hormone 473
blood parathyroids and 164
in growth hormone of anterior pitui-
tary 210
in insulin 177 178
in oxytocin 403-404
in vasopressin 404
- p-Aminobenzonic acid acetylation in
salin and 180
- Amino-peptides e
ACE and 730
ACTH and 730
- Aminopterin
citrovorum factor and 663
folic acid and 670
progesterin and 669
response of pigeon crop glands to pro-
lactin and 376 377
- p-Amino aliphatic acid 468
antithyroid activity 468
- Amphenone B (1 2 bis (p-aminophenyl)-
2-methylpropanone 1-dihydrochloride 497)
antithyroid activity 469
effect on adrenals 692
- Amphibians
melanin synthesis in B hormone
and 491
metamorphosis effect of thyroid hor-
mone on 491
sex reversal with steroids in 675
water metabolism role of neurohypo-
physis in 396-399
- Anaphylaxis
adrenocortical steroids and 771-773
species variations 772
- Androgens see also individual com-
pounds 610-672
activity metabolic 674
physiological 673-674
extragenital 673-674
renotropic 674
assay 666 667 668
biosynthesis 611-613
ACTH and 657
in adrenogenital syndrome 653 654
8 α 0
cortisol and 630
in blood 610 611
cortisone and 655 674
effect on β glucuronidase 673 676
effect of and related steroids on ad-
renal cortex of hypophysectomized
animals 248
growth effects in male pituitary
dwarfs 740
metabolism 580 613-621
microbiological reactions 620-621 672
role in sex differentiation in verte-
brates 675
secretion fructose content of semen
and 864
testicular tumors and 867
source 665
- Δ^4 Androstadiene-3 17 dione forma-
tion microbiological 607 608 609
671 643 644
- Androstane-3 α 17 β diol
metabolism 614

- as metabolite of testosterone 616
- metabolites of 617 618
- Androstane 3β 17β diol
 - formation microbiological 620
 - metabolism 614
 - metabolites 618
- Androstane 3 17 dione
 - metabolism 614
 - as metabolite of Δ^4 androstene 3 17 dione 618
 - of androsterone 618
 - of epiandrosterone 618
 - metabolites of 616 617
 - microbiological reactions 620 621
 - source 615
- Androstane- 3α 16 17 triol synthesis 561
- Androstan 17β ol configuration 535
- Δ^4 Androstene 3β 17α diol
 - acetate as precursor of 612
 - conversion to Δ^4 androstene 3 17 dione 650
 - to testosterone 650
 - labeled preparation 577
 - metabolism 614
 - as metabolite of dehydroepiandrosterone 617 619
 - metabolites of 616 617
 - microbiological reactions 620
 - reduction 650
 - source 615
- Δ^4 Androstene 3 17 dione 611
 - acetate as precursor of 612
 - from Δ^4 androstene 3β 17β diol 650
 - biosynthesis 624 627 628 630
 - ACTH and 652
 - from dehydroepiandrosterone 613 650
 - from 11 deoxycortisol 611
 - formation microbiological 607 608 620
 - 11β hydroxylation enzymatic 645
 - metabolism 613 614 631
 - as metabolite of dehydroepiandrosterone 617
 - of testosterone 616 618 619
 - metabolites of 616 617 618
 - microbiological reactions 620 621
 - as precursor of 6β (and 11β) hydroxy Δ^4 androstene 3 17 dione 624 626
 - of 6α hydroxy Δ^4 androstene-3 17 dione 626
 - reaction with toluenesulfonic acid 550
 - reduction 650
 - sources 610 611 612 615
- Δ^4 Androstene 3β 16 17 triol urinary synthesis 560
- Androsterone
 - acetate as precursor of 612
 - cholesterol as precursor of, 591 613
 - metabolism 614
 - in vitro 640
 - as metabolite of androgens 613 616 617 618
 - of cortisone 638, 639 640
 - of 11 deoxycortisol 638
 - of 17α hydroxy progesterone 633
 - of testosterone 616
 - metabolites of 618
 - source 615
- Anestrus
 - in sheep gonadotropins and 335
- Anethole effect on metabolism of adrenal steroids 692
- Aniline derivatives
 - antithyroid activity 462 468-470
 - mechanism of 462
- Animal husbandry application of gonadotropins in 338-342
- Animals
 - breeding in anestrus season PMG and 338-340
 - experimental responses elicited by growth hormone preparations in 243
- Anions inorganic antithyroid activity 464
- Annelids
 - central nervous system of adrenaline like substance in 58
 - growth hormone in 59
- Antiauxin(s) see also Auxin inhibitors of
 - activity assay 3
 - mode of 38
 - flowering and 31
 - terminology 2
 - types of 38
- Antibiotics auxins and 25
- Antibodies
 - formation adrenocortical steroids and 753 760 774

- adrenocorticotrophic hormone and 774
 to pituitary hormones 203-206
 precipitating relationship between neutralizing antibodies and 219
 to protein hormones 218-219
- Anticonvulsants
 combination with acetylcholine receptive substance 123
 structure 123
- Antidiuresis see also Antidiuretic hormone
 produced by posterior pituitary extracts 393
 role of anterior pituitary in 399
 thyroxine and 399
- Antidiuretic hormone of posterior pituitary
 assay of 396
 bacterial 396
 effect on kidney tubules 397
 escape of following surgical removal of posterior pituitary 115
 physiological significance 393
 release from neurohypophysis 116
 DFP and 117
 secretion of 395-396
 target organ for 394 399
 urinary excretion 396
- Antiepileptics
 combination with acetylcholine receptor 125
 structure 125
- Antifibrinolysin
 cortisone and 762
- Antigen antibody reactions adrenocortical steroids and 773-775
- Antihistamines effect on amphibian metamorphosis 491
- Antihyaluronidase in blood of cancer patients 751
- Antimony in thyroid 436-437
- Antistine 469
 antithyroid activity 469
- Antithyroid agents see also individual compounds
 action 442 462
 ACTH and 483
 effect on uptake of radioactive iodine 828
- estrogens and 672
 therapeutic effect on ophthalmopathy 834
 on thyrotoxicosis 832
- Antuistrin G 269
- Appetite
 ACTH and 755
 cortisone and 753
- Araneidae* neurosecretory system in 48
- Arginase effect of adrenocortical steroids on 731-732
- Arthritis rheumatoid
 blood glutathione in 724
 fibrinoid and 739
 scurvy and 739
 therapeutic effect of ACTH on 727
 of cortisone on 687 727 763 766
 thyrotropin and 498
 thyroxine and 498
- Arthropods see also Crustaceans
 neurosecretory principles in 78
- Arthus reaction
 ACTH and 773
 cortisone and 773
- Ascorbic acid see also Vitamin C
 adrenal 742
 depletion by ACTH 230 423 424 425
 by B hormone 423 424 425
 thyroid medication and 781
 adrenocortical steroids and 219 220 739
 auxin and 29
 deficiency see also Scurvy
 effect on adrenals 739-740
 stress and 741
- Ascorbic acid oxidase auxin and 41 42
- Astatine (eka iodine element 85) 436
 in mammalian thyroid 436
- Atherosclerosis
 adrenocortical steroids and 716-717
 role of thyroid in 486
- Atropine trophic action 130
- Auxin a 4ff
 chemical nature 4
 molecular weight 5
- Auxin b chemical nature 4
- Auxin(s) see also individual compounds
 action mode of 37-48
 activity 47-48

- configuration of sidechain and 8
 growth promoting 7
 requirements for 7
 metabolic 38-42
 metabolic control of 47
 phosphorylation and 39-40
 plant growth mechanism of 1-2
 requirements for 38
 site of 47
 structure and 8-14 12-16
 effect of side chain length 14
 of ring substitution 12-14
 role of geometrical isomerism 8ff
 10-12
 sulhydril compounds and 39
 antagonists of 14 15-16
 ascorbic acid and 29
 biogenesis 27-28
 bud inhibition by 20ff 32
 antagonists of 22
 nutrition and 22
 as apical dominance 20
 combination with substrate 14 37-38
 cytoplasmic streaming and 46-47
 destruction 27 28-29
 enzymatic 28-29
 inhibitors of 29
 by light 28
 effect on fruit embryos 34-35
 on plant respiration 39 40
 on roots 15 23-24 38
 concentration and 128
 mechanism of 44
 on xylem differentiation 24
 enzymes and 29 41 42
 flowering and 1 18 19 30-32
 inhibitory effect 31 32
 promotive effect 31 32
 response of flower ovaries to 33
 sexuality of flowers and 32
 formation enzymatic pollination and 33
 fruit setting and 33 34
 induction of flowering in pineapple by 30
 possible mechanism 31
 inhibition of root growth by 23-24
 inhibitors of 25-27
 in *Datura* embryo of 27
 clormine and 26
 germination and 25
- natural 4-7
 plant tumors and 35-37
 plant viruses and 37
 role in physiological inhibitions 20-2, shoot 2 3
 synergists of 14 15 24-25
 synthetic 7-16 see also individual compounds
 transport of 16-20
 currents of 19
 in plants 17
 nonpolar 17
 factors influencing 18
 polar 16ff
 flowering and 18 19
 water uptake and 43 45 47
 Auxin protein 29
 Avena coleoptile auxin in 5
- ### B
- BAI see 2 3 Dimercapto 1 propanol
 Bacteria see also Microorganisms
 cortisone and 758 778-779
 extracts effect on adrenal cortex of hypophysectomized animals 278
 formation of antidiuretic substances by 396
 Basal metabolism in acromegaly 874
 Benzoic acids substituted auxin activity of 9-10 12
 Bile acids
 conversion to cortisone 551
 degradation to 20 ketosteroids 566
 labeled preparation 5 7
 11 oxygenated steroids from 577
 side chains orientation 526
- Birds
 effects of prolactin in 3 3 3 4-375
 gonadal function seasonal changes in 317
- Bisdihydrodioxynolic acid estrogenic activity configuration and 541
- Blood
 adrenal cortex and 744-748
 adrenal hormones in 693
 species differences 671
 androgens and 610 612 614
 clotting ACTH and 749
 effect of adrenocortical steroids on 749
 components parathyroids and 164

- effect of estrogens on 641-672
 of thyroxine on 496
 iodine of 449-450
- Blood cells reaction to inflammation
 ~62-~63
- Blood sugar effect of adrenocortical
 steroids on 771-772
- Blood vessels
 action of vasopressin on 137
 permeability hyaluronidase and 63
- Body fluids corticotropin in 283
- Bombyx mori* egg diapause factor 141-
 147
- Bone(s)
 effect of adrenal cortex on 93-104
 fractures effect of ACTH on ~67
 of cortisone on ~64
 parathyroids and 153 155-160
- Bone marrow
 adrenal cortex and ~44
 response to adrenalectomy 45
 to adrenocortical steroids 746-~47
 ~48ff
- Botulinum toxin mode of action 117
- Brain
 effect of adrenocortical steroids on
 19ff
 5 hydroxytryptamine and 104 105
 137 136
 inhibitory hormone in of polychaetes
 99
 as source of hormones in worms 58
- Broodiness in poultry prolactin and 3 4
- Brucellosis effect of cortisone on 7 9
- Brunn effect 396-397
 ecological significance 399
- Buds
 differentiation of nucleic acid syn-
 thesis and 22
 dormant growth inhibitor in 23
 inhibition by auxins 20ff 32
 lateral role of auxin in growth of 70
- Bufoformone possible neurohormonal ac-
 tivity 100
- Burns effect of adrenocortical steroids
 on ~59
- C**
- Calciferol 534
 correlation with glycerol aldehyde 534
- Calcium
 absorption parathyroids and 165
 concentration in mammary gland
 parathyroids and 153
 excretion renal mechanism 155
 parathyroids and 155
 metabolism growth hormone and 200
 role of thyroxine in 487
 serum parathyroid activity and 161-
 162
- Cancer see also individual organs
 experimental effect of adrenocortical
 steroids in ~51-~52
 hormone treatment of 751
- Capillaries adrenal cortex and 716
- Carbohydrate metabolism
 adrenal cortex and 721-~727
 corticotropin and 296
 in crustaceans hormonal control of 166
 Cushing's syndrome and 846
 effect of B hormone on 471
 of growth hormone preparations on
 51
 of pituitary extracts on 25
 in insects corpora allata and 4
 pantothenic acid and ~38
 role of insulin in 184
 thyroid and 484-485
- Carbohydrates transport in plants 17
- Carbon
 isotopic introduction into steroids
 574-576
 into ring A 574-575
 ring D 575-576
 into side chains 576
- Carcinogens
 cortisone and ~53
 effect of thyroxine on 496-497
- Cardiovascular system
 adrenal cortex and 713-714
 thyroxine and 493
- Casein iodinated effect on milk produc-
 tion in cows 495
- Catalase 79
 indoleacetic acid oxidation and 29
 inhibition by sex hormones 676
 occurrence in plants 79
- Cattle response of ovary to gonadotro-
 pic stimulation 338

- Cells,
 neurosecretory, see Neurosecretory cells
 surface
 as site of acetylcholine action 133
 of adrenaline action 133
 carminophil of anterior pituitary as source of prolactin 313 346 347
 α , of islets of Langerhans
 as source of glucagon 192 193
 β of islets of Langerhans
 experimental necrosis 189
 as source of insulin 17a 193
 Leydig see Leydig cells
 mast see Mast cells
 syncytial effect of chorionic gonadotropin on 561
 formation of steroid hormones in 861
 Cephalopods physiologically active substances in 78 79
 Chelidonic acid
 auxins and 24-25
 Chilopoda neurosecretory system in 78
 Chloramphenicol thyroxine and 492
 Chloride metabolism
 effect of adrenocortical steroids on 705
 thyroid and 487
 3 β 6 β Cholestanediol 547
 configuration 546
 Cholestan-3 β ol synthesis 538
 Cholestan 7 α (and 7 β) ol configuration 535
 Cholestenone labeled conversion to progesterone 575
 Cholesterol
 androgen biosynthesis and 613
 biosynthesis 591-593
 possible mechanisms 592-593
 sites of 591
 configuration 532-533
 conversion of diosgenin to 561
 correlation of 20 hydroxy steroids with 532
 degradation 536
 labeled distribution of labeling agent in 578
 preparation 575 516 577
 as precursor of adrenocortical steroids 627
 of androgens 612
 of estrogens 596
 of neutral steroid hormones 591
 of pregnenolone, 649
 of progesterone 602 603
 precursors of, 591-593
 structure 593
 Choline deficiency, cortisone and 758
 Cholinesterases
 esteratic site of, 121
 interaction of acetylcholine and 191
 red cell acetylcholine and 128
 Chordates collection of iodine by thyroid in 435
 Choriocarcinoma of testis 868
 Chorionepithelioma 863
 Chromatophorotropin(s)
 in brain of oligocheles 59
 in crustaceans 61 99-100 138
 action mechanism of 61-62
 differentiation between 61
 molecular weight 80
 neurohormonal activity 99
 release 115
 site of origin, 61
 in insects 76
 invertebrate effect on vertebrate chromatophores 81
 interchangeability of 81
 in *Tachyples* 78
 Chymotrypsin action on corticotropin A 278-229
 Cinnamic acid auxin activity of cis isomer 8
 trans Cinnamic acid as auxin inhibitor 26
 Citric acid serum parathyroids and 168-169
 Citric acid cycle insulin and 187
 Citroglycose, bone parathyroids and 159-160
 Cirrhosis hepatic Cushing like pictures in 844
 Citrovorum factor thyroid and 489
 Coconut milk auxins and 25
 Coenzyme A
 in adrenal cortex 737
 function of 738
 formation thyroid and 481
 Cold stimulating effect on thyroid 459-460

- Collagen fibrinoid and 740
- Colloid thyroidal
 fluorescence of 437
 glycoprotein in 437
 histochemistry of 437
- Compound A see 11 Dehydrocorticosterone
- Compound B see Corticosterone
- Compound F see Cortisone
- Compound F see Cortisol
- Compound S see 11 Deoxycortisol
- Corpora allata 138
 connection with neurosecretory system 67 68 82
 effect on metabolism 74
 endocrine activity 69-70 73 141
 factors affecting 70
 ovaries and 70
 endocrine stimulation of possible mechanism 68
 myotropic principle in 17
 relation to neurosecretory cells of brain 67 69
 as release and storage organ for insect hormones 139
 role in color adaptation 16
 in postembryonic development 71
 as source of juvenile hormone 72
- Corpora cardiaca
 chromatophorotropic activity 76
 connection with neurosecretory system 67 68 82
 endocrine activity 69
 endocrine stimulation of possible mechanisms 68
 formation of hormones in 140
 metabolism and 74
 myotropic factor in 77
 neurosecretory cells in 98
 prothoracic glands and 72
 relation to neurosecretory cells of brain 67 69
 as release and storage organs for insect hormones 72 138 139
 role in postembryonic development 71
 sinus glands and 19
 as storage and release center for prothoracotropic hormone of brain 72
- Corpus luteum
 cysts of 860
- formation role of FSH and luteinizing hormone 602
- of lactation placental prolactin and 310
 prolactin requirement 360
- maintenance factors concerned in 310 371 861
- pregnancy and 360 860 861 862
- progesterone metabolism in control of 311
- prolactin and 359 360 369 370
 as source of progesterone 601 609 603
 of 6 β steroid hydroxylase 616
 types of 370
- Corticoids see Adrenocortical hormones
- Adrenocortical steroids permissive action 351
- Corticosterone 292 699
 assay 703
 biosynthesis 621 626 627 698 630
 ACTH and 652
 progesterone as intermediate in 602
 in blood 691
 cholesterol as precursor 591
 hyaluronidase and 163
 metabolism *in vivo* 635 636
 as metabolite of deoxycorticosterone 645
 secretion corticotropin and 290
 microbiological reaction 642
 therapeutic effect on Addison's disease 843
 on adrenal insufficiency 697
- Corticotropin A 279 280
 activity 222
 amino acid composition 294
 terminal amino acids in 217
 corticotropin B and 292-224 295 296
 empirical formula 295
 isolation 222
 peptic hydrolysis 227-228
- Corticotropin B 219 290 823
 activity 222
 amino acids essential for 295
 amino acid composition 214 225
 corticotropin A and 292-224 295 226
 isolation 221-29
 preparation from ACTH 893
- α Corticotropin 279

Corticotropin(a) 235 275-296,
see also Adrenocorticotrophic hormone
activity 226 235
absorption from injection site and 294
adrenal ascorbic acid depleting 276 277
androgenic 292
diabetogenic, 263 265
factors influencing 292 293
fat mobilizing 254 256
glycotropic 256 257
glycostatic 252
growth 241
of hydrolyzed preparations 294
hypoglycemic 261
means for prolonging, 293
melanophore-expanding 295-296
metabolic 296
method of preparation and 290 293
purification and 293-294
route of administration and 292 293 294-295
assay 220 276 277
in body fluids 295
carbohydrate metabolism and 256
chemical structure 230
chemistry of 219-234
classification 220
composition amino acid 224-226
differences in 279-280
content of anterior pituitary in 245
cortisone and 291
degradation 287
disappearance from blood 280-287
effect on adrenal cortex 287-290
in vitro 288-290
on adrenal 275 276 277-279
on carbohydrate and fat metabolism 268
on pancreas 258
effects extra adrenal 295-296
systemic 291 292
growth hormone and 235-236 262-263 268 275
hydrocortisone and 291
inactivation by blood, 230
by plasma 230
isoelectric point 226
molecular weight 225 226
multiple 276-280

physicochemical properties 224-226
preparation 290-222
activity and 290
refractoriness to 292 293
secretion in acromegaly, 273
adrenal hormones and 281-282
adrenaline and 283-285
control of, 280-285
factors influencing 280
hypothalamus and 282-283
rate of 285-287
sources 221
stability 293
ultraviolet absorption 225 236
Corticotropin actide, see Corticotropin B
Cortisol 523 622 see also Hydrocortisone
activity 630
androgenic metabolites and 590 611 635 665
ACTH and 630 652
biosynthesis 623 624 625 626 627 628 630
in adrenogenital syndrome 654
in Cushing's disease 651
progesterone as intermediate in 609
cholesterol as precursor of 591
configuration 577
conversion of 11 β 17 α dihydroxy-3 20 dione Δ^4 pregnen 21 al to 648 to 17 keto derivative 650 651
from cortisone 550
degradation product synthesis of 538
from epicortisol 559
9 α fluoro activity 560
from 11 α hydroxysteroids 560
metabolism 632 636-637
in vitro 638 639 640
in vivo 634
as metabolite of cortisone 638
microbiological formation 642
reaction 643
preparation 521
review on 521
therapeutic effect on adrenogenital syndrome 653
urinary excretion 631
Cortisone 5 3 627 687
activity androgenic metabolites and 611 635

- antiinflammatory 704 700 61 70
 763
 hyperten. in 714 715
 oral 697
 administration adrenal atrophy fol-
 lowing 27 701
 effect of testosterone and related
 steroids on 278
 alkalosis following 709
 Cushing like pictures following 844
 modes of 696 697
 adrenocorticotrophic hormone and 211
 701
 allergic rhinitis and 73
 amphenone B and 692
 antifibrinolysis and 762
 assay of 703
 asymmetric centers 576
 bacteria and 58 778-779
 from bile acids 551
 biosynthesis 627 626
 in Cushing's disease 661
 carcinomas and 701
 conversion to 1 α 21 dihydroxyprog-
 nane 3 11 20 triene 611
 to 17 keto derivative 650 661
 of sarmentogenin to 661
 diabetes and 725
 effect on adrenals 78 781 291 691
 735
 on amino acid metabolism 721 719
 on antibody formation 713
 on bacterial toxins 58
 on blood proteins 710
 on bone fractures 767
 on brucellosis 779
 on carbohydrate metabolism 721
 722
 on electrocardiogram 713 714
 on electrolyte metabolism 61 705
 706 710-711
 on electroshock threshold 703
 on embryos 639
 on eosinophils 741
 on experimental cancer 751-757
 on eye lesions 76 - 68
 on fat metabolism 732 733
 on fibrosis 76 - 68
 on growth 741 700 701
 on histamine toxicity 5
 in human malignancy 751
 on mast cells 743-744
 on muscle function 720 721
 on myocardial infarction 708-709
 on nucleic acids 71
 on pregnancy 699 700
 on renal clearances 709
 on renal lesions due to choline de-
 ficiency 728
 on respiration of lactating mamma-
 ry tissue 356
 on reticuloendothelial system 743
 on sepsis 740 741
 in humans 742
 on skin 754-755
 on lesions of 766-767
 on spleen 743
 on syphilis 780
 on thymus and lymph nodes 742
 on thyroid diseases 81
 function 459
 on tissue and organ grafts 775-776
 on water excretion 711-717
 effects clinical 688
 systemic 291
 enzymes and 31
 estrogens and 669 784
 in fluoro activity 560
 formation enzymatic 648
 microbiological 642 643
 species differences in 691
 glutathione and 724 725
 growth hormone and 265
 hair growth and 755
 halogen derivatives effect in ad-
 renalectomized animals 695
 hyaluronidase and 763 764
 hyperinsulinism and 725-726
 inactivation hepatic 844
 iodide metabolism and 440 459
 labeled preparation 578
 metabolism 631 632
 in vivo 634
 number of possible stereoisomers 536
 phosphorylase and 723
 pollen sensitivity and 713
 preparation 521
 protein metabolism and 727
 Rb erythroblastosis and 715
 resistance to tumor transplantation
 and 753

- response to adrenocortical diseases and, 847
 review on 521
 structure 537
 susceptibility to infection and 776-780
 synthesis 538 539
 therapeutic effect on Addison's disease 807 842
 on adrenogenital syndrome 653 802
 on diseases of the nervous system 717
 on hypopituitarism 826
 on ovarian dysfunction 860
 on rheumatoid arthritis 687
 on Stein Leventhal syndrome 807
 on thyroid storm 836
 thyroxine and 498
 toxic effects 776
 in man 699
 urinary excretion 634
 viruses and 779
 vitamins B and 737-739
 Cortisone acetate effects systemic 292
 Coumarin
 auxins and 24
 in plants 24
 Creatine metabolism thyroid hormone and 483
 Creatinine metabolism thyroid hormone and 483
 Creatinuria in thyrotoxicosis 835
 Cretinism 836-838
 mental development and 837
 thyrotropin and 837-838
 Crop gland of pigeon prolactin and 358 375-378
 Crown gall bacteria *Pseudomonas* (or *Agrobacterium*) *tumefaciens*
 auxin and 30 37
 virus like agent in 35
 Crown galls bacteria free formation 35
 Crustaceans
 acetylcholine in 61
 attempts at establishing occurrence of sex hormones in 64ff
 castration parasitic 64
 central nervous system neurosecretory cells in 60
 chromatophorotropins of 99-100
 differentiation between 61
 neurohormonal activity 99
 color adaptation 61-62
 role of pigment dispersing enzyme in 62
 decapod action of 5 hydroxytryptamine in 135
 as source of molt inhibiting hormone 63
 of molt promoting hormone 62
 eyestralk sex reversal hormone of 65
 gonads hormonal activity 64-65
 growth factors affecting 142
 neurosecretory products and 140 141 142
 heart action in hormonal principles affecting 67
 hormonal control of physiological processes in 61
 hormones in 58 60-67 90-100
 activity 141
 cyclical release 116-117
 formation site of 138 141
 gonad inhibiting 65
 growth inhibiting 100
 metabolic 66-67
 molt inhibiting 61 67 100
 molt promoting in 61 62
 molting cyclical release 116
 ovarian inhibiting 61
 release of 63 138
 retinal pigment migration and 62 116
 sources of 60
 storage 63 100-111 138 141
 5 hydroxytryptamine in 61 67 105
 metabolism 66-67
 hormonal control of 66
 molting 62-63
 endocrine control of 62-63 72
 factors affecting 61
 molting gland (y organ) 60
 prothoracic gland of insects and 60
 as source of metabolic hormone 65
 myotropic principles in 77
 neurohemal organs in 61
 neurosecretory material of 108 137-141 144 see also Crustaceans hormones
 axoplasmic transport 113
 formation site of 139

- release and storage organs for 138
139
neurosecretory systems of 139
organ system controlling growth and
molting in 63
possible correlation between gonads
and metabolic processes in 67
reproduction 63-66
factors influencing 63-64
retinal pigment migration hormonal
control of 62 116
sinus gland of 60
Cushing's syndrome 843-845
adrenal and 843
ACTH and 292 652 843-844 845
biosynthesis of cortisol and cortisone
in 651 652
course of disease 847
diagnosis 847
electrolyte metabolism in 845-846
hormone secretion in 845
hyalinization of pituitary basophils
in 844
17 hydroxysteroids in blood 845
metabolic disturbances in 845-846
pathogenesis 843-845
secretion of adrenocortical steroids in
652 845
symptoms 651
treatment 848
Cyanoide antithyroid activity 461-462
Cyclohexane stereochemistry 573-524
Cytochrome placental 372
Cysts non neoplastic of ovary 850-857
Cytochrome b₅ in insects 5
Cytochrome c thyroid and 47
Cytochrome oxidase
inhibition by sex hormones 676
plant growth and 43
role in intrathyroidal iodine metabo-
lism 442 443
Cytoplasmic streaming
auxin and 46-47
water uptake and 46-47
- D
- DCA see Deoxycorticosterone acetate
DDD see 2,2-bis (p Dichlorophenyl)
1,1-dichloroethanol
DFP see Diisopropyl fluorophosphate
DOC see Deoxycorticosterone
Dark adaptation B hormone and
420-421
Datura auxin inhibitor in embryos of 27
epi Dehydroandrosterone 523
Dehydroascorbic acid 193
diaprogenic activity 190
11 Dehydrocorticosterone 622
assay 703
in blood 694
effect on thymus and lymph nodes 742
on urinary electrolyte excretion 702
metabolism 637 637
in vitro 635 636
microbiological reactions 642
synthesis 637
11 Dehydrocortisone microbiological
formation 642
 Δ^4 Dehydrocortisone (Diene com-
pound) effect in adrenalectomized
animals 698
 Δ^4 Dehydrocortisone acetate
effect on muscle function 720
on glycogen deposition 722
Dehydroepiandrosterone 523 545 611
acetate as precursor of 612 613
action of 3 β dehydrogenase on 628
biosynthesis 627 628 630
in blood 628
conversion to Δ^4 androstene 3 17
dione 624 627 650
to 11 β hydroxy Δ^4 androstene 3 17
dione 624
16 α hydroxylation enzymatic 646
labeled preparation 576
metabolism 613 614
as metabolite of Δ^4 androstene-3 β 17 β
diol 617
metabolites of 616 617 618 619
microbiological formation 644
reactions 620 621
nature of 545
source 612 615
urinary excretion 613
adrenal cancer and 623 652 848
17 β Δ^4 Dehydroestradiol 539
17 β -dihydroequilenin from 540
17 β -estradiol from 540
 Δ^4 Dehydroestrone 512
conversion to estrone 540

- $\Delta^{11(10)}$ Dehydroestrone 543
 structure 543
 synthesis partial 542 543
- 14β Δ^4 Dehydroestrone 512
 isoequinin 1 and 543
- 3β Dehydrogenase 629
 activity 628
 in adrenal cortex 628
- Dehydrogenases steroid 645 648
 action, 648
 number of 648
 occurrence 648
- Dehydroglucoscorbic acid 189
 diabetogenic activity 190
- Dehydroisoascorbic acid 189
 diabetogenic activity 190
- Δ^1 Dehydrotestolactone formation
 microbiological 609
- Deiodase thyroidal 446 454
 thyrotropic hormone and 446
- Deoxycorticosterone 622
 biosynthesis 627
 progesterone as intermediate in 602
 derivatives therapeutic value in Addison's disease 842-843
 effect on adrenal cortex 278 281
 on enzymes 731
 esters effect in adrenalectomized animals 698
 11β hydroxylation enzymatic 645
 metabolism 632
in vitro 637-638
in vivo 637
 metabolites of 635 636
 microbiological reactions 641 642 643
 as precursor of aldosterone 630
 as precursor of corticosterone 624 626
 of cortisol 626
 of $6\beta, 21$ dihydroxy Δ^4 pregnene 3 20 dione 626
- Deoxycorticosterone acetate
 activity anesthetic 720
 effect of 11 desaturation on 559
 hypertensive 714-715
 adrenocortical atrophy and 701
 effect on brain excitability 719
 on electroshock threshold 702
 on metabolism of amino acids 728
 of electrolytes 702 705 711 846
 in Addison's disease 846
 in Cushing's syndrome 84
 on scurvy 740
 growth inhibition by, 709
 labeled preparation 575
 myocardial necrosis caused by 714
 response to insulin and 720
 role in diseases responding to steroid therapy 707
 secretion effects of increased 689
 structure 569
 therapeutic effect on adrenal insufficiency 697
 on adrenogenital syndrome 802
 toxic effects 714 756 757
 thyrotropic hormone and 707
- Δ^1 Deoxycorticosterone acetate effect in adrenalectomized animals 698
- Deoxycorticosterone glucoside
 effect in adrenalectomized animals 698
 on electrolyte metabolism 700
 mode of administration 697
- 11 Deoxycortisol 523
 acetate as precursor of 612
 androgens from 611
 biosynthesis 620 627
 conversion to 17 keto derivative 600
 11β hydroxylation enzymatic 645 646
 labeled preparation 577
 metabolism 632
in vitro 638 639
in vivo 632-634 638
 microbiological formation 643
 reactions 641 642 643
 as precursor of cortisol 626
 of 11 deoxycorticosterone 624
 source of 611
- 21 Deoxycortisol biosynthesis 627
- 21 Deoxycortisone as precursor of cortisone 625 626
- 11 Deoxy 17 hydrocorticosterone effect in adrenalectomized animals 698
- Deoxylase steroid 645 649
- 17 Deoxysteroids adrenal orientation of side chains 526
- Dermatitis effect of cortisone and hydrocortisone on 770
- Desmolases steroid 645 649
 20 22 action of 649
 number of 649

- Diabetes insipidus
 effect of a adrenocortical steroids on
 water excretion in 712
 neurohypophysis and 313
 Diabetes in lituræ
 a adrenocortical hormones and 721 725
 as complication of acromegaly 571
 experimental production 189-190
 growth hormone and 274
 metabolism of glucose in 182
 of organic phosphorus compounds in
 157
 of proteins and 188
 pituitary and 251 262-265
 Diabetes steroid 816
 Diabetogenic factor pituitary 251 262
 Diapause factor 141
 2,4-Dichloroaniline
 as auxin antagonist 31 35
 2,2-bis (p-Dichlorophenyl) 1,1-dichloro
 ethanol effect on adrenals 691-692
 840
 Dyt
 deficient effect on ovaries 837
 growth and 237-238
 Diethylstilbestrol radioactive in tribo-
 lism 537-538
 Dihydroequilenin urinary configura-
 tion 540
 1 β -Dihydroequilenin
 source 531
 synthesis 540
 3 α -Dihydrotestololactone formation mi-
 crobiological 608
 3 β -21-Dihydroxyallopregnane 11-20 di-
 one 622
 11 β -21-Dihydroxyallopregnane 3-20 di-
 one biosynthesis 625 676
 3 α -6 α -Dihydroxyallopregnan-20-one in
 human pregnancy urine 637
 3 β -1 α -Dihydroxyallopregnan-20-one
 672
 3 α -21-Dihydroxyallopregnan-20-one
 as metabolite of deoxycorticosterone
 638
 of 1-oxycortisone 638
 3 β -21-Dihydroxyallopregnan-20-one
 as metabolite of deoxycorticosterone
 638
 microbiological reaction 641
 3 β -11 β -Dihydroxyandrostan-17-one as
 metabolite of cortisone and cortisol
 640
 11 α -1 β -Dihydroxy Δ^4 -androsten-3-one
 formation microbiological 620
 11 β -17 α -Dihydroxy-3-20-dione Δ^4 -preg-
 nen-21-al conversion to cortisol 648
 3 α -11 β -Dihydroxyetiocholan-17-one
 as metabolite of adrenosterone 619
 621
 of cortisol and cortisone 634 635
 of 11 β -hydroxy Δ^4 -androsten-3-17-
 dione 619 671
 17 α -21-Dihydroxy Δ^4 -pregnadiene
 3-20-dione 11 β -hydroxylation en-
 zymatic 646
 3 α -17 α -Dihydroxypregnane 11-20-dione
 as metabolite of cortisol and corti-
 sone 634 635
 3 α -21-Dihydroxypregnane 11-20-dione
 as metabolite of corticosterone 635
 636
 of 11-dehydrocorticosterone 635
 636
 17 α -21-Dihydroxypregnane-3-20-dione
 metabolism *in vitro* 638
 as metabolite of 11-deoxycortisol 638
 639
 3 α -6 α -Dihydroxypregnan-20-one in hu-
 man pregnancy urine 632
 3 α -17 α -Dihydroxypregnan-20-one as me-
 tabolite of 11-deoxycortisol 633
 3 α -20-Dihydroxypregnan-11-one as me-
 tabolite of 11-dehydrocorticosterone
 635 636
 1 α -21-Dihydroxypregnane 3-11-20 tri-
 one
 conversion of cortisone to 641
 to urocortisone 647
 6 β -21-Dihydroxy Δ^4 -pregnene 3-20 di-
 one
 biosynthesis 676
 microbiological formation 641
 17-21-Dihydroxy Δ^4 -pregnene 3-20-di-
 one see 11-Deoxycortisol
 20 β -21-Dihydroxy Δ^4 -pregnene 3-11-di-
 one 672
 17 α -21-Dihydroxy Δ^4 -pregnene 3-20-
 dione 11 β -hydroxylation enzymatic
 646

- 17,21 Dihydroxy Δ^4 pregnene 3 11 20
trione see Cortisone
- 6 β ,17 α Dihydroxy progesterone micro
biological formation 641
- 11 α 17 α Dihydroxy progesterone micro
biological formation 608 641 642
- 3 5 Duodo-4 hydroxybenzoic acid
derivatives antithyroxine activity
453 475-476
effect on metabolism of thyroxine 453
of triiodothyronine 453
- Duodotyrosine,
conversion to thyroxine 443 444-445
hypophysectomy and, 445
- Dusopropyl fluorophosphate anticho-
linesterase activity 117
- 3 11 Diketopregnane 20 hydroxyster-
oids derived from 529-530
- 2 3 Dimercapto 1 propanol (BAI)
growth inhibiting activity of lac-
tones and 24
- 1 7 Dimethylxanthine as thyroxine an-
tagonist 476
- Diosgenin
conversion to cholesterol 561
to 11 oxygenated steroids 552
structure 551
- Di sec butyleclopentene derivatives
see Auxin a
- Diureis see also Water metabolism
adrenocortical steroids and 711-712
- Dog adrenaline and noradrenaline con-
tent of brain 102
- Doisynolic acid(s) 540-542
configuration 541
estrogenic activity and 540 541
synthesis 541
- Dopa decarboxylase in adrenal medulla
107
- Dopamine see Hydroxytyramine 103
- Drugs
effect on thyroid 461-474 490-49
sympathomimetic structural require-
ment 134
- Dwarfism
pituitary 876-877
symptoms of 827
treatment with growth hormone
preparations 270-273 827
- E
EPS see Exophthalmos producing sub-
stance
- Ectohormone(s)
occurrence 80
of termites 77
- Egg diapause factor of *Bombyx mori* 142
- Egg white toxic effects in rat, adreno-
cortical steroids and 757
- Electrocortin see Aldosterone
- Electroencephalogram effect of adreno-
cortical steroids on 718
- Electrolytes
dietary effect on adrenal cortex of
hypophysectomized animals 218
metabolism adrenal cortex and 688
cortisone and 704-705 707 708
in Cushing's syndrome 845-846
effect of adrenogenital syndrome on
851
thyroid and 486
retention growth hormone and 290
urinary excretion assay of adreno-
cortical steroids based on 702
- Electroshock
mechanism of 719
threshold adrenocortical steroids and
702-703 719
- Elements chemical in thyroid 436
- Endocrine organs effect of environ-
mental changes on 317
- Endocrinology clinical 821-876
- Enteramine see also 5 Hydroxytrypt-
amine 78
in insects 77 82
in intestinal mucosa of tunicates 79
in invertebrates 80
in mollusks 78
as nerve transmitter in crustaceans 67
- Enzyme poisons inhibition of root
growth by 23
- Enzymes
action on growth hormones 204-205
adrenocortical hormones and 723-724
730-732
auxins and 41 42
effects of sex hormones on 673 674
675 ff
activating 677

- inhibitory 676 677
 species differences in 6 6
 eyestalk hormones and 6
 ins et metabolism and 7
 relationship between developmental
 hormones and 7
 mammalian effect on steroid metabo-
 lism 614-619
 pigment dispersing role in color adap-
 tation of crustaceans 62
 in prolactin stimulated pigeon crop
 glands 377
 role in biosynthesis of steroid hor-
 mones 590
 in metabolic estrogen conversions
 598
 thyroid hormone and 477-490
 thyroxine 437 417 406
 action of proteolytic 406
 I α in auxin and 22
 I α inopenia mechanism of 746-747
 Eosinophils effect of adrenocortical
 steroids on 703 744-745 746 47
 Epandrosterone 593
 from Δ androstene 3 17 dione 618
 metabolism 614
 as metabolite of androgens 617 618
 metabolites of 619
 as precursor of androgens 674
 source 615
 synthesis 537-538 539
 Epicorticoesterone microbiological for-
 mation 641 642
 Epicortisol 623
 cortisol from 505
 microbiological formation 642
 Epidermis effect of estrogen on 6 1
 Lphydrocortisone hydrocortisone and
 22
 Epinephrine see Adrenaline
 Epitesterone 573
 metabolism 614
 metabolites of 616
 1 2 Epoxide opening of 525
 Equilin
 cholesterol as precursor of 596
 from Δ dehydroestrone 510
 estrone and 540 596
 source 574
 synthesis total 537
 Equilin
 cholesterol as precursor of 596
 estrone and 540 596
 source 574
 synthesis total 537
 Ergosterol
 calciferol from 574
 conversion to 11 oxygenated steroids
 557
 structure 501
 Ergot
 action on human heart 135
 adrenaline and 131
 Ergothioneine
 antithyroid activity 465
 structure 467
 Erucic acid effect on a renal 672
 Erythroblastosis cortisone and 75
 Erythrocytes effect of adrenocortical
 steroids on 748
 Esterases sex hormones and 612
 Estradiol
 configuration 524
 effect on inflammation 62
 α Estradiol see 1 β Estradiol
 17 α Estradiol source 594
 17 β Estradiol
 in human semen 595
 labeled preparation 576
 metabolic relationships between es-
 trone and 598-600
 metabolism intermediates 596
 in vitro 600
 in vivo 599 601
 radioactive iodo metabolism of 595-
 597
 sources 594 598
 synthesis 540
 Estrane action 598
 Estriol
 in human semen 595
 from 16 keto 17 β -estradiol 598 599
 601
 metabolism in man 601
 occurrence in human tissues and/or
 urine 598
 synthesis 561
 Estrogen(s) 593-601 see also individual
 compounds
 activity 593
 adrenocorticomimetic 605

- lactogenic 359 361 362 368
 mechanism of 368
 progesterone and 363
 structure and, 593
 tumorigenic progestin and 672
 antithyroid substances and 672
 assay 666 667 668
 biosynthesis 595-601
 precursors 596
 routes of 596
 cortisone and 669 784
 effect extragenital 669-672
 on hypophysectomized animals 240
 on luteotropic action of prolactin 370
 of progesterone plus on lactation 361-362
 on pituitary gonadotropic activity 334
 physiological 669-672
 on thyroid function 459
 folic acid and 670
 growth hormone and 241
 hydrocortisone and 669 784
 labeled metabolism 596-598
 preparation of 595-596
 localization in tissues 596-598
 metabolism 596 598-601
 in man 601
 natural activity of epimerization products 541
 configuration 540
 sources 594 784 861 864
 progesterone and 670 730
 secretion by adrenals 853
 effect of ovarian follicle cysts on 855
 in pregnancy 861 864
 testicular tumors and 858
 on secretion of anterior pituitary hormones 368
 sources 593 669
 syntheses partial 539 ff
 therapeutic effect on ovarian agenesis 854
 urinary abortion and 863
 chorionic gonadotropin and 593
 in ovarian agenesis 854
 Istronase 601
 action 600
 Estrone
 cholesterol as precursor of 596
 conversion to 19 nortestosterone 543
 dibromo metabolism of labeled 597
 doisynoic acid and 541
 equilenin and 540 542 596
 equilin and 596
 metabolic relationships between 17 β estradiol and 599-600
 metabolism intermediates 596
 in vitro 600 601
 in vivo 599
 occurrence in human semen 595
 tissues and/or urine 598
 sources 591
 synthesis 537 540 542
 I strone acetate preparation of labeled 57
 Estrone
 estrogenic potency 537
 structure 537
 Estrus
 gonadotropic activity of anterior pituitary and 334-335
 stimulating effect of light on 317
 Δ^4 Δ^5 Ethylene cis hydroxylation 540-532
 Ethyltestosterone activity effect of 11(12) desaturation on 539
 Ethanic acid 532
 Ethioallocholanolic acid see Alloethanic acid
 Ethiocholanone 3 α 17 α diol
 metabolism 614
 as metabolite of ethiocholanolone 618
 of 17 β hydroxyethiocholan-3-one 616
 source 616
 Ethiocholanone 3 α 17 β diol
 metabolism 614
 as metabolite of Δ^4 androstene 3 β 17 β diol 617
 of Δ^4 androstene 3 17 dione 617
 of dehydroepiandrosterone 618
 of ethiocholanolone 618
 of testosterone 616
 source 616
 Ethiocholanone 3 17 dione
 metabolism 614
 as metabolite of ethiocholanolone 618
 source 616
 Ethiocholanone 3 α 16 17 triol synthesis 561
 Ethiocholanolic acid see Ethanic acid

- Liocholanolone**
 acetate as precursor 612
 biosynthesis cholesterol and 611 613
 metabolism 611
 as metabolite of androgens 611 617
 618
 of 11 deoxycortisol 618
 of 1 β hydroxycetolan 3 one 618
 of testosterone 616 617
 metabolites of 618
 source 615
- Lunichoidism**
 with infantile ovaries 884-885
 primary testicular 885
 pituitary hypogonadotropic and 867
 treatment 865
- Exophthalmos**
 experimental 833
 malignant response to cortisone 781
 mucopolysaccharides in edema fluid 833
 pituitary and 833
- Exophthalmos producing substance**
 (EIS) pituitary 210-211 45
 thyrotropin and 211 833 834
- Eye(s)**
 diseases see also Ophthalmopathy
 effect of hydrocortisone on inflammatory 768
 lens effect of cortisone on 767-768
 role of retina in transference of light stimuli to anterior pituitary 319
 320
- Eye stalk of crustaceans** 60
 action on vertebrates 81
 chromatophorotropic factor in 139
 effect on water balance in amphibians 339
 gonads and 66
 hormonal activity 60 61 62 63 65
 enzymes and 67
 neurosecretory cells in 60
 removal effects of 63 65 66 67
 as source of crustacean hormones 14
- F**
- FSH** see Follicle stimulating hormone
 fasting effect on growth hormone secretion 252
- Fat body of insects**
 corpora allata and 74
 corpora cardiaca and 74
 endocrine activity of 68
- Fat metabolism**
 adrenal cortex and 737-738
 corticotropin and 296
 pituitary and 254-256 733 734
 thyroid and 485-486
- Fatty acids biosynthesis insulin and** 188
- Fever effect of cortisone on** 750
- Fibrinogen effect of adrenocortical steroids on** 747
- Fibrinol**
 collagen and 740
 in rheumatoid diseases 753
- Fibrosis**
 effect of ACTH on 768
 of cortisone on 768 63
- Fishes**
 adrenaline and noradrenaline in 103
 effect of thyroid on 491-492
- Flowering**
 auxins and 31
 hormonal stimuli 79-80
 auxin and 30
 photoperiodism and 29 32
 temperature and 31 32-33
- Flowers**
 ovaries of response to auxins 33
 to malate 34
 to pollination 33
 sexuality factors controlling 32
- Fluorescence in thyroid** 497
- Fluorotyrosine antithyroxine activity** 4 6
- Folic acid**
 nucleic acid synthesis and 317
 pigeon crop gland response to prolactin and 376-377
- Follicle stimulating hormone** 211-216
 activity 215 216 303
 standardization 211-213
 amino acid composition 218
 application in animal husbandry 341
 342
 cellular origin 310 314 315 823
 homogeneity 216
 in human pituitary 336 337
 immunological behavior 218-219
 isoelectric point 217

- isolation from sheep 214
 - from swine 214-216
- molecular weight 217
- ovarian function and 832
- physicochemical properties 217-218
- role in corpus luteum formation 602
 - in estrogen production 593
 - in testosterone biosynthesis 611
- secretion body weight and 839
 - during pregnancy 335 336
 - prolactin and 371
- spermatogenesis and 864
- stability, 218
- urinary testicular tumors and 868
 - in infantile ovary syndrome 834
 - in ovarian agenesis 853 854
 - in pituitary hypogonadism 855
- Fruit
 - developing auxin sources for 34
 - embryo effect of auxins on 34-35
 - setting of auxins and 33
- G
- GH see Growth hormone of anterior pituitary
- Galactopoiesis see also Lactation
 - effect of ACTH on 365
 - role of prolactin in 365
- Galactose distribution insulin and 182-183
- Gambogin structure 551
- Gastrointestinal tract thyroid and 495-496
- Giberellin auxin and 25
- α Globulins serum binding of thyroid hormone to 450
- Glucagon 251
 - action 190
 - mode of 194-195
 - site of 193-194
 - amino acid composition 191
 - assay of 194
 - chemical properties 190-191
 - endocrine nature 190
 - formation growth hormone and 193
 - growth effects in hypophysectomized animals 259
 - inulin and 193
 - liberation following growth hormone administration 258 261
 - liver glycogen breakdown and 194
 - molecular weight 190
 - occurrence 190
 - origin 175 197 193
 - purification 190
 - sulfur content 190 191
- Glucocorticoids placenta as source of 861
- Gluconeogenesis, adrenalectomy and 779
- Glucose
 - conversion of glycogen to, 194
 - distribution insulin and 183-184 187
 - metabolism diabetes and, 182 186
 - insulin and 187 180
 - phosphorylation biological 181
 - factors affecting 182
 - utilization adrenal cortex and 77
 - in vitro insulin and 186
- Glucose 6 phosphate
 - formation 181 182
 - factors affecting 182
 - metabolism 181-182
- Glucose 6 phosphate dehydrogenase
 - possible role in lactogenesis 354
- Glucuronic acid conjugation of steroid hormones with 590
- Glutamic aspartic transaminase possible role in lactogenesis 354
- Glutamic dehydrogenase possible role in lactogenesis 354
- Glutathione 724-725
 - ACTH and 724 725
 - auxin inhibitors and 27
 - blood thyroid and 454
 - cortisone and 724 725
- β Glycerinaldehyde configuration 533
- α Glycerophosphatase inhibition by 3 ketosteroids 6,6
- Glycogen
 - depots hypophysectomy and 252
 - formation diabetic serum and 186
 - liver breakdown 194
 - epinephrine and 194
 - glucagon and 194
 - effect of adrenocortical steroids on 93
 - phosphorylation adrenal corticosteroids and 724
- Glycoproteins gonadotropins as 311
- Glyostatic factor pituitary 251 257
 - glytropic factor and 256

- Glycotropic factor pituitary 251 256-257
 glycostatic factor and 256
 Glycyrrhizic acid
 effect on Addison's disease 813
 on adrenal insufficiency 607
 Goat survival following adrenalectomy 695
 Goiter 838-839
 in acromegaly 871
 colloid 838
 etiology 838 839
 treatment 839
 uptake of radioactive iodine by 828
 Gestrogens see also Antithyroid substances and individual compounds
 action mechanism of 463
 Gonadotrophs 314 315
 Gonadotropic hormones 822
 see also individual hormones
 anestrus in sheep and 335
 applications in animal husbandry 338-342
 biosynthesis of testosterone and 611
 cellular distribution 316
 cellular origin 310ff 314 315
 effect on adrenal cortex of hypophysectomized animals 218
 on hypogonadism 866
 glycoprotein nature of 311
 inactivation by gonads 358
 mamalian action on invertebrates 81
 physiology of 309-351
 precursors 314
 release 852-853
 during pregnancy 335-336
 effect of stimuli from reproductive organs on 332
 estrous cycle and 315
 mechanism of 869
 role in cyclic estrogen production 593
 therapeutic effect on eunuchoidism 865
 urinary testis tumors and 868
 Gonadotropin A 217
 Gonadotropin B 217
 Gonadotropin
 chorionic abortion and 862
 assay as basis for pregnancy tests 862
 biosynthesis of testosterone and 612
 corpus luteum of pregnancy and 861
 origin site of 861
 physiological effects 869
 syncytial cells and 861
 chorionic human 217
 effect on urinary excretion of estrogens and 17 ketosteroids 595
 follicle stimulating activity 342
 luteotropic activity 312 370 311 372
 prolactin and 342 371
 physiology of 342-344
 in serum of early pregnancy 313
 urinary excretion in early pregnancy 343
 pregnant mare's serum 216 344-345
 application in animal husbandry 341 342
 effect on heifer ovaries 335
 formation site of 344
 urinary of menopause 345
 Gonadotropins see Gonadotropic hormones and individual gonadotropic hormones
 Gonads
 adrenals and 783-784
 effect of thyroid on 494-495
 on thyroid function 458-459
 hormonal activity in crustaceans 64-65
 in insects 63
 hypofunction pituitary gonadotropin content and 337
 pituitary and 332 338 358
 Grafting
 ACTH and 775
 cortisone and 75-776
 Graves disease
 ophthalmopathy in 834
 therapeutic effect of iodide in 463
 mechanism of 473
 of potassium perchlorate in 462
 Growth
 adrenal cortex and 241
 adrenocortical hormones and 669 700-701
 diet and 737-738
 hypophysectomy and 226-237
 sex hormones and 240-241
 thyroid and 259-240 401-492

Growth hormone of anterior pituitary
 202-206 236-275 822
 action 235 236
 activity adipokinetic 251 255 256
 assay of 202
 chemical structure and 204
 chondrotropic 241
 diabetogenic 257 262-268 366 824
 effect of pH on 264
 method of preparation and, 263-
 264, 265
 potentiating effect of corticotropin
 on 262-263
 species differences in 265 266 268
 galactopoietic 270 366
 glycotropic 256-257
 glycostatic 252
 growth promoting in experimental
 animals 242-246
 hypoglycemic 260 261
 international standard 202
 metabolic 296
 modes of 259
 renotropic 270
 adrenocortical hormones and 241
 alkaline phosphatase of plasma and
 250
 amino acid composition 205
 antigenicity of 206
 biological effects, 242-273
 in central nervous system of annelids
 59
 corticotropin and 235-236 268 275
 cortisone and 270
 degradation 242
 effect on adrenals 277 278
 on biosynthesis of aldosterone 641
 of nucleoproteins and nucleic
 acids 249
 on inflammation 762
 in man 270-273
 diabetogenic 272-273
 metabolic 271-272
 on metabolism of carbohydrates 251
 268
 of electrolytes 200
 of fat 268
 of inorganic substances 249-251
 of proteins, 246-249
 of salt 395
 of water 395

on pancreas 258
 on wound healing 269-270
 estrogens and 241
 extraction 275
 glucagon production and 193
 immunological behavior 205-206
 insulin and 249-249 726
 isolation 202-203 204
 ketogenesis and 735 736
 liver fat and 733 744
 mammary development and 348 351
 molecular weight 204 274
 physicochemical properties 203-205
 274-275
 physiologic role 273-275
 protein nature 271
 purification 203 236
 purity 271-272
 secretion 241-242
 in acromegaly 823-824
 age and, 274
 sedimentation behavior 204
 stability 203-204
 species difference in 204
 thyrotropin and 275
 toxic effects 265 268-269 271
 cortisone and 265

Guinea pig
 effect of growth hormone preparations
 in 245-246

H

HCG *see* Gonadotropin, human chorionic
 Hair
 greying of adrenal cortex and 755
 growth cortisone and 755
 estrogen and 671
 Hamster survival following adrenalectomy 695
 Heart
 effect of adrenocortical steroids on
 713-714
 sympathetic receptor of 134
 vertebrate action of acetylcholine on
 127 128
 Hecogenin structure 551
 Hemopoiesis, parathyroids and 164
 Henle's loop as target organ for anti-
 diuretic hormone 334 339

- Hepatitis acute effect of adrenocortical steroids in 73
- Hesperidine methylchalcone 471
antithyroid activity 41
- Heteroxanthin 4
- Hexokinase
effect of growth hormone preparations on 257
hormones and 257 256-257
inhibitors in anterior pituitary extracts 184-185
in human diabetic serum 185
in plasma 185
insulin and 184-185 187
- Hexosamine in thyroid 437
- Hippulin source 594
- Histamine
toxic effects adrenaline and 75
cortisone and 15
- D Homoandrogens activity 565
- 5 α D Homomrostane 1 17 α dione synthesis 539
- 5 β D Homomrostane 3 17 α dione synthesis 533
- D Homosteroids see also Uraneol
configuration 564ff
synthesis 563-565
as urinary constituents 563
- Hormone
lactogenic see Prolactin
luteinizing see Interstitial cell stimulating hormone
- Hormones see also individual hormones
action mechanism of 67
adrenal see Adrenocortical hormones
Adrenocortical steroids and individual hormones
affecting mammary growth during pregnancy 349 350 351
in crustaceans 60-67
gonadal inhibiting 65-66
metabolic 66-67
molt inhibiting 61 63
molt promoting 61 63
ovarian inhibiting 61
gonadal 665 see also Sex hormones and individual hormones
gonadotropic see Gonadotropic hormones and individual hormones
hexokinase and 256-257
inhibitory in brain of polychaetes 59
in insects 67-77
developmental relationship between enzymes and 75
growth and differentiation hormone 72 73-74 82
juvenile 72 73
prothoracotropic 72
interaction between neurogenic mechanisms controlling pituitary activity and 333
in invertebrates 5-95
action on vertebrates 81
activity concentration and 80
physicochemical properties 80ff
specificity 80
melanophore-expanding 13
stimulating properties 6
in mollusks 8-9
sources of 8-9
ovulating neurogenic stimulus responsible for discharge of 333
pituitary 201-234 389-431 see also individual hormones
placenta as source of 861
plant growth 1-56 see also Auxins and individual compounds
protein antibiotics to 218-219
placenta as source of 861
salt retaining 290
deficiency in hypopituitarism 875
secretion testicular tumors and 867
stimulation of target organs by tropic 360
sugar 841 846 850
in treatment of cancer 151ff
tubular 851-855
in tunicates 79
vertebrates action on invertebrates 81
in worms 58-60
- Hyaluronic acid production by mast cells 763
- Hyaluronidase 61
action effect of steroids on 763-764
possible mechanism of 764
effect on joints 764
on mucopolysaccharides 833
on ophthalmopathy in thyroid diseases 833-834
- Hydatiform mole 863

- Hydrocortisone 291 *see also* Cortisol
 activity antiinflammatory 704 761
 762
 hypertensive 714
 oral 696
 administration, modes of 696 697
 assay 703
 in blood 694
 corticotropin and 230 291
 effect on adrenal cortex 281
 on chick embryos 699
 on eye diseases 768
 effect on metabolism of carbohydrates
 721, 722
 of electrolytes 705
 of proteins 727
 on muscle function 720
 on thymus and lymph nodes 742
 on urinary electrolyte excretion 702
 estrogens and 669 670 784
 growth and 241 701
 growth hormone and 265
 pregnatriol as precursor of 850
 response of eosinophils to 745
 therapeutic effect on adrenal insuffi-
 ciency 697
 on hypopituitarism 820
 on rheumatoid arthritis 765 766
 urinary excretion in Cushing's syn-
 drome 845
- Hydrogen
 isotopic introduction into steroids
 576-578
 by hydrogenolysis 575 577
- Hydrogenases steroid 645 646 649-649
 action 647
 3 β adrenal cancer and 652
- Δ 5 α Hydrogenase 631
 Δ -5 β Hydrogenase 631
 Δ Hydrogenases steroid 631 645 648-
 649
- 11 α Hydroxyallopregnane 3 20 dione
 formation microbiological 608
- 21 Hydroxyallopregnane-3 20-dione
 as metabolite of deoxycorticosterone
 637 638
 metabolites of 638
 as precursor of 11 β 21 dihydroxyallo-
 pregnane-3 20 dione 625 626
- 3 α Hydroxyallopregnane 20 one
 metabolism 603
 sources 604
- 3 β Hydroxyallopregnane 20 one 622
 metabolism 603
 sources 604
- 17 β Hydroxy Δ^4 androstadiene 3 one
 formation microbiological 609
- 3 β Hydroxyandrostan 16 one
 synthesis 561
 in urine of pregnant mares 561
- 3 β Hydroxyandrostan 17 one *see* Epi-
 androsterone
- 6 α Hydroxy Δ^4 androstene 3 17 dione
 biosynthesis 626
 as metabolite of 6 α hydroxy Δ^4 an-
 drostene 3 17 dione 631
- 6 β Hydroxy Δ^4 androstene 3 17 dione
 biosynthesis 624 676
 formation microbiological 608 620
- 11 α Hydroxy Δ^4 androstene 3 17 dione
 formation microbiological 670
- 11 β Hydroxy Δ^4 androstene 3 17 dione
 biosynthesis 624 626 627 630
 ACTH and 652
 formation microbiological 620
 metabolism 619 621 654
 as metabolite of cortisone and corti-
 sol 610
 metabolites of 635
 sources of 610 612
- 16 α Hydroxy Δ^4 androstene 3 17 dione
 formation microbiological 620
- 17 α Hydroxy Δ androsten 3 one *see*
 Epitestosterone
- 3 β Hydroxy Δ^4 androsten 17 one *see*
 Dehydroepiandrosterone
- 11 β Hydroxyandrosterone
 from cortisol 590 634 635
 from cortisone 611 634 635
 as metabolite of adrenosterone 619
 621
 of androsterone 654
 of 11 β hydroxy Δ androstene-3 17
 dione 619 629 654
 urinary in adrenogenital syndrome
 654
- 17 Hydroxycorticosteroids
 in blood 694
 effect of disease on 694
- 17 Hydroxycorticosterone *see* Cortisone

- 6 β Hydroxycortisol as metabolite of cortisol 632 637
- 6 β Hydroxy-3,5 cycloandrostan 1-one 545
isolation from urine 544
synthesis 544
- 11 β Hydroxypandrosterone source 612
- 16 α Hydroxyestrone 601
- 2 α Hydroxyetiocholan-3,17-dione
as metabolite of adrenosterone 619 621
of cortisol and cortisone 634 635
of 11 β hydroxy Δ^4 androstene-3,17-dione 619 621
- 3 α Hydroxyetiocholan 17-one see Etiocholanolone
- 3 β Hydroxyetiocholan 17-one
metabolism 614
as metabolite of testosterone 617
source 616
- 11 β Hydroxyetiocholan 3-one
metabolism 614
metabolites of 618
- 3 α Hydroxy 11 ketopregnane
20 hydroxysteroids derived from 5 α -530
- 17 Hydroxy 20 ketosteroids 527
- 17 α Hydroxy 20 ketosteroids 5,1
rearrangement 561
synthesis 567 569
- Hydroxylases steroid 645-647
action 629-630
adrenogenital syndrome and 653 654-655
number of 629
source of 629
6 α and 6 β 645 646
sources 646
11 β 645-646
action on steroids 645-646
occurrence 645 646
16 646
17 α 646
21 646
21 β adrenogenital syndrome and 653 654
- Hydroxy octopamine
identity with noradrenaline 8
occurrence 78
- 4 (4 Hydroxyphenoxy) 3,5 dihydroxyphenylpropionic acid hormonal activity 475
- 21 Hydroxy Δ^4 pregnadiene-3,20-dione 11 β hydroxylation enzymatic 646
- 16 Hydroxypregnane derivatives urinary 600
- 16 α Hydroxypregnane-3,20-dione for zonation microbiological 608
- 3 α Hydroxypregnan 20-one
labeled preparation 577
metabolism 603
from progesterone 605
sources 604
- 6 β Hydroxy Δ^4 pregnene-3,20-dione for zonation microbiological 607
- 20 β Hydroxy Δ^4 pregnen-3-one
metabolism 603
from progesterone 606
- 3 β Hydroxy Δ^4 pregnen 20-one
isolation from testis 601
metabolism 632
microbiological reactions 643 644
as precursor
of corticosterone 624 625
of cortisol 624 625
of progesterone 602 624 625
- 6 β Hydroxypregesterone biosynthesis 674
- 11 α Hydroxypregesterone formation microbiological 607 608
- 11 β Hydroxypregesterone from 11 keto progesterone 550
- 16 α Hydroxypregesterone microbiological formation 607 608 642
- 1 Hydroxypregesterone
acetate as precursor of 612
androgens from 611
labeled preparation 577
pregnabietol as metabolite of 550
source 611
- 17 α Hydroxypregesterone 672
activity androgenic 654
biosynthesis 675 627
adrenogenital syndrome and 654
conversion to 17 ketoderivative 650
formation microbiological 607
metabolism *in vivo* 632-634
microbiological reactions 641 642 643
as precursor of cortisol 624 625 626

- p Hydroxypropophenone 471
 antithyroid activity 471
 6 Hydroxysteroids 546-547
 configuration 536 546 547
 11 Hydroxysteroids synthesis 550
 11 α Hydroxysteroids synthesis mi
 crobial 550
 11 β Hydroxysteroids
 configuration 558
 preparation 558
 reactivity 548-550
 16 Hydroxysteroids
 configuration 560-563
 synthesis microbial 563
 urinary 560 561
 17 Hydroxysteroids 560
 in blood of patients with Cushing's
 syndrome 845
 configuration chemical reactivity
 and 526-527
 nomenclature revision 526ff
 17 α Hydroxysteroids synthesis 566
 20 Hydroxysteroids
 configuration 528-533 536
 correlation of 529-530
 with cholesterol 532
 natural types of 528-529
 4* 7 β Hydroxysteroids
 configuration 552
 preparation 552
 11 β Hydroxytestosterone biosynthesis
 625
 5 Hydroxytryptamine 78-82 103-105
 134 135-137 143 see also Enteramine
 action on brain 136
 neurohumoral 99 100 135
 peripheral 136-137
 chemical nature 99 143
 in crustaceans 61
 LSD and 136
 occurrence 103 104 105 108 135
 origin of 137
 receptors for 137
 synthesis 104
 Hydroxytyramine
 occurrence 103
 possible neurohormonal activity 99
 100
 Hyperglycemic glycogenolytic (H G)
 factor see Glucagon
 Hyperinsulinism cortisone and 725-726
 Hypersensitivity adrenocortical ster
 oids and 760
 Hypertension adrenocortical steroids
 and 714-716
 Hyperthyroidism
 in acromegaly 273
 effect of ACTH on 781
 on nitrogen metabolism in 481ff
 on tyrosine metabolism 453
 pregnancy and 831
 Hypoglycemic factor see Pancreatropic
 factor
 Hypogonadism 865-867
 classification 865
 pituitary 855
 ovarian agenesis and 853
 Hypometabolism 828
 Hypophysectomy
 effect on conversion of diiodotyrosine
 to thyroxine 445
 on growth and development 236-237
 of insulin following 238-239
 replacement therapy in lactating ani
 mals following 363-365
 Hypopituitarism 825-827
 effect on gonadotropin secretion 825
 hormones affected by 826
 salt hormone deficiency in 825
 therapy of 826
 Hypothalamic neurohypophyseal sys
 tem 323-324
 as source of oxytocin 105 118
 of vasopressin 105 118
 Hypothalamic
 antidiuretic activity adrenocortical
 steroids and 712
 gonadotropin release and 321 869
 lesions precocious puberty due to 869
 pituitary and 282 321 322-333 368
 pituitary ovarian axis and 852

I

- ICSH see Interstitial cell stimulating
 hormone
 Immunization passive cortisone and
 774-775
 Inanition effect on gonadotropic ac
 tivity of anterior pituitary 336 337
 Indole aurins and 25

- Indoleacetaldehyde
 auxin activity " "
 occurrence 4 5 "
- Indole acetic acid
 assay of and derivatives thereof 2-3
 auxin activity 38
 effect on respiration of animal tissues 41
 enzymatic destruction " "
 inhibition of root growth by 23
 occurrence 4 5
 pectin methylesterase and 44
 precursors 27
 structure 6
- Indoleacetic acid oxidase
 action of 28-29
 occurrence in plants 29
- Indoleacetonitrile
 auxins and 6 25
 isolation 6
 occurrence 6
 structure 6
- Indolepyruvic acid 6-"
 auxin activity 7
 molecular weight 7
 occurrence 6-7
 structure 6
- Infection
 bacterial thyroxine and 497-498
 effect of adrenocortical steroids on 760 761
 growth hormone and 2 0
 susceptibility to adrenal steroids and 776-780
- Inflammation
 adrenocortical steroids and 760 761-766
 effect of growth hormone on 762
- Insects
 acetylcholine in 101
 adrenaline in 103
 brain neurosecretory cells in 67 98 138
 corpora allata and 67 69
 corpora cardiaca and 67 69
 metabolism and 74
 endocrine activity 67 68 69
 prothoracic glands and 72
 role in postembryonic development 71
 caste differentiation hormonal control of 77
 color adaptation in 764
 differentiation prothoracic glands and 71
 dormancy hormonal control 71
 embryonic development 70-71
 factors controlling 70
 gonad metabolism and 74
 growth endocrine control of 71 141 142
 hormones in 58 67-77 141 see also Insects neurosecretory materials action on vertebrates 81 formation site of 141 growth and differentiation 72 73 82 action on protozoa 82 physicochemical properties 80 two step reactions 77 83 metabolism 74-76 endocrine control 74ff enzyme systems participating in 75 molting endocrine control 71 prothoracic glands and 72 myotropic principles in 77 neurosecretory materials of 109 137-142 144, see also Insect, hormones in avoplasmic transport 113 formation site of 109 release and storage of 138 139 141 neurosecretory systems of 67 139 postembryonic development 71-74 hormone centers affecting 71 prothoracic (ventral) gland of 60 72 molting gland of crustaceans and 60 reproduction hormonal control of 69-70 sex hormones in 70
- Insulin 145-190
 acetylation of *p* aminobenzoic acid and 189
 action site of 181 185 transforming 184-185 186 translocating 182-184 186 187 adrenal cortex and 185 186 278 adrenocortical steroids and 72 amino acid composition 177 species differences in 178

- anterior pituitary and 185 186 187
 A chain of 179
 amino acid sequences in 180
 structural formula 180
 B chain of 179
 amino acid sequences in 179
 structural formula 180
 chemistry, 176-181
 effect on eosinophils 745
 on glycogen synthesis *in vitro* 186
 on lipogenesis in lactating mammary gland 354
 effects *in vitro* 185-186
in vivo 187
 fatty acid synthesis and 188
 glucagon and 193
 glucose metabolism and 187 189
 growth effects in hypophysectomized animals 238-239
 growth hormone and 248-249 726
 hexokinase and 184-185 187
 homogeneity 177
 inhibition by corticotropin 256
 by pituitary glycotrophic factor 256
 molecular weight 176
 origin 175 193
 oxygen uptake and 187
 physiology of 181-190
 protein nature 177
 reaction with target tissues 186-187
 response to in Cushing's syndrome 847
 deoxycorticosterone acetate and 725
 role in carbohydrate metabolism 184 187 189
 secretion effect of pituitary preparations on 258 261
 structure 176ff
 Intermedin (B) 100 137 355
 acetylcholine and 100
 activity 100 137
 ACTH and 13 225
 effect on rat mammary tissue 355
 in vertebrates 100
 Interstitial cell stimulating hormone 219
 action 369
 amino acid composition 212
 application in animal husbandry 341
 cellular origin 310 315 316 823
 disulfide groups in 316
 growth promoting effect in pigeon 246
 isolation 211
 in human pituitary 336 347
 urine 211
 ovarian function and 852
 production by placenta, 336
 release, mechanism of 671 869
 chemotransmitter effecting 331
 effect of estrogens on 671
 of stimuli from reproductive organs on 332
 role in corpus luteum formation 602
 in estrogen production 593
 by testis 595
 in testosterone biosynthesis 611
 secretion hypothalamus and 321 322
 progesterone and 335
 stimulation of Leydig cell function by 864
 urinary Leydig cell deficiency syndrome and 867
 Intestine vertebrate action of acetylcholine 127 129
 Inulin renal clearance growth hormone and 270
 Invertebrates
 acetylcholine in 101
 hormones in 57-95
 5 hydroxytryptamine in 105
 neurosecretory materials of 105
 neurosecretory systems in 58 82-83
 connection with endocrine organs 82
 demonstration of neurosecretory material in 83
 endocrinological significance 58
 vertebrate hypothalamic hypophyseal system and 82
 occurrence of adrenaline and noradrenaline in 103
 Iodide
 antithyroid activity 463
 mechanism of 473
 metabolism 439ff
 cortisone and 440
 therapeutic effect on Graves disease 473
 on thyrotoxicosis 832

- uptake by thyroid 440-441
 - phases of 441
 - thyroid hormone secretion and 440
- urinary excretion 439-440
- Iodine
 - blood 440-450
 - collection by thyroid in chordates 434
 - in human embryo 438
 - concentration by thyroid tumors 441
 - deficiency as cause of goiter 438
 - dietary thyroid weight and 434
 - metabolism extrathyroidal 439-441
 - intrathyroidal 441-449
 - nonthyroidal 444
 - protein bound cortisone and 743
 - determination of 831
 - distribution of nonthyroid 444
 - thyroid activity and 830
 - radioactive in assay of thyrotropic hormone 20-208
 - uptake by thyroid 440 828-830
 - use in thyrotoxicosis 832-833
 - serum protein bound abortion and 863
 - effect of pregnancy on 450
- Iodocasein toxic effects 489
- Ions effect of water uptake on 46
- Iron plasma adrenocortical steroids and 748
- Iron metabolism thyroid and 487-488
- Irradiation
 - effect on adrenal function 750
 - corticosteroids and 750-760
 - on iodine uptake by thyroid 460
 - on susceptibility to infections 460
- Islets of Langerhans see also α and β cells of pancreas
 - effect of corticotropin on 258 259
 - of growth hormone on 258 259
 - hormones of 175-199 see also Insulin
- Glucagon
- d Isocoumarin 542
- Isoequilin A 543
- Isoprene as precursor of cholesterol 502-503
- Iothyroxine 445
 - hormonal activity 445
- Isotopes
 - in study of estrogen metabolism 546-558
- of pituitary hormone target organs 456
- of thyroid function 434
- J**
- Joints
 - inflammation effect of adrenocortical steroids on 764-765
 - hyaluronidase and 764
 - permeability hyaluronidase and 764
- Juglone as auxin inhibitor 26
- K**
- Kendall's compound E see Cortisone
- Kendall's Compound F see Cortisol
- 11 Ketoandrosterone
 - as metabolite of adrenosterone 619 621 640
 - of cortisol 590 611 634 635
 - of cortisone 634 635 648
 - of 11 β hydroxy Δ^4 androstene 3 17
 - dione 619 621
- 16 Keto 17 β -estradiol conversion to estradiol 598 599 601
- 1 Ketoestrone stability 540
- 16 Ketoestrone
 - metabolism in man 601
 - occurrence in human tissues and/or urine 598
 - source 594
- 3 Ketoetic acid methyl ester synthesis 539
- Δ^4 3 Keto tianic acid methyl ester preparation of labeled 575
- Ketogenic factor 751
- Ketogenesis
 - adrenal cortex and 735-736
 - stress and 736
- Δ^4 3 Keto 17 ketosteroids
 - formation 651
 - as precursors of 17 keto steroids 651
- Ketone catalytic hydrogenation 525
- 18 O Ketopregnanes peroxy acid degradation 527-528
- 11 Ketoprogesterone microbiological reaction 643
- Ketone 183
- Ketosteroids see also individual compound
 - adrenal cytological localization 600
 - catalytic hydrogenation 525

- 3 Ketosteroids
 bromination 548-549
 inhibition of α glycerophosphatase by, 676
- 11 Ketosteroids reduction 548
- 12 Ketosteroids conversion to 11 oxygenated steroids 556-558
- 15 Ketosteroids configuration stability and 640
- 17 Keto teroids
 androgenic secretion in adrenogenital syndrome 848
 a say of, 667
 formation, 650 651
 as metabolites of cortisol and cortisone 638
 of dioxycorticosterone 635
 of progesterone 636
 urinary in adrenal cancer 612
 in adrenocortical hyperfunction 860
 chorionic gonadotropin and 595
 in Cushing's syndrome 816
 in pituitary hypogonadism 866
 of pregnancy 861
- 20 Ketosteroids bromination 566-567
 metabolism differences between *in vitro* and *in vivo* experiments 632
 preparation 566
- Δ^4 3 Ketosteroids 547-551 see also individual compounds
 configuration 548 549
 effect of Δ hydrogenases on 648-649
 formation enzymatic 628-629
 from 3 ketosteroids 548
 labeled preparation 549
 as metabolites of cortisol 637
 preparation 548-549
 protection during 549-551
 reduction of 650-651
 effect of nuclear substitution on 650-651
- Δ^4 3 Ketosteroids synthesis of estrogens from 539
- Kidney
 effect of estrogens on 672
 of pituitary extracts on 39,
 function adrenal cortex and, 704-712
 parathyroids and 153-155
 a source of steroid hydrogenases 647
- Klinefelter syndrome 866
- Krebs cycle adrenal steroids and 730
- LSD see Lysergic acid diethylamide
- LTII see Prolactin
- Lactation see also Galactopoiesis
 Lactogenesis
 inhibition by combined action of estrogen and progesterone 361-362
 maintenance adrenocortical steroids and 784
 hormonal mechanism governing 366
 prolactin and 363-366 366-367
 milk ejection 363
 posterior pituitary and 363 393 399-400
 onset at parturition 359 362-363
 factors affecting 363 369
 theories of 369, 369-363 367
 parathyroids and, 169
 role of pituitary adrenal axis in 351-362 365
- Lactogenesis
 experimental effect of antagonism between ovarian hormone and prolactin on 360
 action of ovary on 369
 prolactin and 361-363
- Lactogenic hormone see Prolactin
- Lactones
 as growth inhibitors 24
 B1L and 24
 unsaturated as germination inhibitors 27
 light and 27
 synergism with auxins, 24
- Leaves mature role in fruit set by auxins or pollen 33-34
- Lecithin
 linkage to protein components of cell surface 170-127
 acetylcholine and 176
- Lecithin adrenaline in 103
- Lipogenic effect of cortisol on 748
- Leucocyte response to adrenal steroids 747-748
- Leukemia effect of adrenocortical steroids on 706 748 751 752-753
- Leukotaxine 761

Leydig cells
 function hypogonadism due to failure
 of 865 866-867
 treatment of 867
 as source of androgens 864 868
 of estrogens 868
 of progesterone 864
 tumors of 867
 Licorice extract beneficial action in
 adrenal insufficiency 697
 Light
 effect on gonadotropic activity of an-
 terior pituitary 317-318
 on reproductive functions of ani-
 mals 317-318
 on thyroid 460
 Limulus brain of hormones in 78
 Lipids
 blood cortisone and 733
 DCA and 733
 Lipogenesis
 in lactating mammary gland 353-354
 insulin and 354
 Liver
 diseases effect on ovarian function
 859
 effect on thyroxine metabolism 451
 452
 estrogens and 598 600 672
 fat adrenals and 733
 growth hormone and 733 744
 as source of steroid hydrogenases 647
 hydroxylases 646
 as target organ of glucagon 193 194
 Lumipandrosterone synthesis 539
 Lunnestrone 531
 Luteotropic hormone see Prolactin
 Luteotropin(s) see also Cyonin Pro-
 lactin
 identity with prolactin 369
 placental 372
 prolactin and 372
 Lymphocytes effect of adrenocortical
 steroids on 703 744
 Lysergic acid diethylamide action on
 Venus heart 135
 central nervous effects 136
 5 hydroxytryptamine and 136
 structure 135
 Lumbricidae clitellogenic factor in 58

M

MCPA see 2 Methyl-4 chlorophenoxy
 acetic acid
 Magnesium blood thyroid and 488
 Malaria effect of cortisone on 780
 Malate growth promoting effect on
 flower ovary 34
 Maleic hydrazide as antiauxin 29 36
 Malnutrition adrenal cortex and 735
 Mammals
 brain photosensitive areas in 319
 effect of steroids on gonads of 675
 metabolic effects of prolactin in 373-
 374
 sexual activity light and 318 320
 Mammary duct system response to pro-
 lactin 348 352 378
 Mammary gland
 action of oxytocin on 137
 calcium concentration by parathy-
 roids and 153
 development prolactin and 347-351
 somatotropin and 348 351
 lactating enzyme systems in 354
 lipogenesis in 353-354 357
 insulin and 354
 oxidative metabolism of 348f
 cortisone and 356 357
 Dickens cycle and 354
 insulin and 355
 prolactin and 357 358
 reponse to prolactin 347-352 353 357
 358
 Mammogen pituitary 347 349
 prolactin and 348
 Mammotropin see Prolactin
 Man
 adrenalectomy in 696 697
 corticosterone therapy and 697-698
 toxic effects of chronic cortisone ther-
 apy in 699
 Manganese in thyroid 456
 Marine worm sessile (*Phoronis*) hor-
 mone in 58
 Mast cells
 action of cortisone on 743-744 763
 production of hyaluronic acid by 763
 Mating behavior
 androgens and 673

- estrogen and 669
- progesterin and 669
- Mecholyl, in insects 77
- Medman structure 136
- Melanin synthesis in amphibians B'
 - hormone and 421
- Melanophore expanding (B) hormone 417, 422
 - activity adrenal ascorbic acid depleting 423 424 425
 - biological assay of 418-419
 - species differences in 422
- ACTH and 421 423-425
- amino acid composition, 423
- origin 390 417 421 425
- pharmacological effects 420
- physicochemical properties 423
- physiological role 419-420
- precursor 422 425
- purification 421-422
- urinary in pregnancy tests 419
- Melanophores
 - effect of posterior pituitary extracts on 330
 - hormone expanding see Melanophore expanding hormone
- Menstrual cycle
 - effect of estrogen on 671
 - of progesterin on 671
- Mental disease 5 hydroxytryptamine and 136
- 2 Mercaptoimidazole
 - antithyroid activity 465
 - structure 465
- Metabolic factor specific 251 253
 - corticotropin and 253
- Metabolism cellular acetylcholine and 129-130
- Metamorphosis of insects endocrine factors participating in 72
- Methadone thyroid and 496
- Methionine labeled incorporation into pituitary proteins 352
- 2 Methyl 4 chlorophenoxyacetic acid weed killing activity 13
- Methylene blue as acetylcholine antagonist 120
- 17 α Methyl estradiol
 - preparation 576
 - radioactive metabolism of 599
- 1 Methylene,
 - activity 510
 - synthesis, 510
- 17- α Methyl D homoandrostan-3 β ol
 - 17 α one synthesis 561
- 4 Methyl 6 hydroxy pyrimidine 2- α thio isobutyric acid
 - antithyroid activity 472
 - structure 472
- 17 Methylprogesterone 566
 - activity 567
 - preparation 566 567
- Methylthiouracil antithyroid activity 476
- Microorganisms
 - reactions with steroid hormones 590-591
 - with adrenocortical steroids 640-644
 - with androgens 670-671
 - with progesterone, 607-609
 - synthesis of 11 oxygenated steroids by 552
- Milk iodinated protein in 459
- Mineralocorticoids 861
- Mitochondria
 - formation of acetylcholine in 106 143
 - as source of steroid hydroxylases 645 646
 - storage of neurohormones and 108
- Mitosis effect of B hormone on 421
- Mollusks
 - hormones in 78-79
 - 5 hydroxytryptamine and 105 135
- Monkey
 - effect of growth hormone preparations on 216
 - survival following adrenalectomy 695
- Morphine sensitivity to thyroid and 436
- Mouse effect of growth hormone preparations on 245
- Mucin action of periodic acid on 311
- Mucopolysaccharide(s)
 - in edema fluid of exophthalmos 833
 - effect of hyaluronidase on 833
- Mucosa intestinal 5 hydroxytryptamine in 136
- Muscles
 - binding of insulin by 187

- effect of progestin on 672
function adrenal cortex and 720-721
- Myasthenia gravis
effect of cortisone on 717
thyrotoxic myopathy and 830
- Myocardial infarction effect of cortisone on 758-759
- Myopathy thyrotoxic 830-836
- Mytolon as acetylcholine antagonist 120
- Myxedema
amenorrhea in 494
effect on cardiovascular system 493
of cortisone on 781
on nervous system 494
of thyroid hormone on extracellular protein in 481
ophthalmopathy in, 833
serum β globulin and 486
- N**
- Naphthalene 1 acrylic acid auxin activity of cis isomer 8
- Naphthoxy compounds
auxin activity optical isomerism and 11 16
as auxin antagonists 11
- Nemertean cerebral organ of 59
- Nephrotic syndrome
action of ACTH on, 704
of adrenocortical steroids on 704
- Nephrocalcinosis parathyroids and 156 157
- Nerves peripheral
effect of adrenocortical steroids on 790
sensory acetylcholine in 101
- Nervous system
action of adrenaline on 131
of noradrenaline on 131
adrenal cortex and 717-720
central control of pituitary gonadotropic function by routes of 320-321
distribution of acetylcholine in 101
nature of links between pituitary and 300ff
progesterone effects and 333
thyroid and 493-494
- Neurohormone(s) 97-151 see also Neurohumors and individual hormones
- action concentration and 129
modes of 118ff 143
site of 143
biosynthesis, site of 106ff 111 114 143
chemical classification 143
nature 99
definition 99
polypeptidic 137
mode of action 143-144
release 143
from active neurons 115
from resting 114-115
chemicals affecting 117-118
cyclical 116-117
from nerve endings 113-118
reflex 115
storage 106-108 143
axon terminations and 109 111
mitochondria and 108
in nervous tissue 108-111
transport 143
axonal 111-113 114
- Neurohumors 99 143
mode of action 143
- Neurohypophysis see Pituitary posterior neural lobe
- Neurons 100-106
adrenergic 98
noradrenaline as principle product of 130
biological significance 142
cholinergic 98
structure 118
- Neurosecretory cells 98
biological significance 98 99 142
of crustaceans 109
hormones arising from 62-63
morphology of 60
products of 108
of hypothalamus 100
as source of oxytocin 100
of vasopressin 105
identification 107
of insects 67
active principles of 68 69 72 109
of worms 58 59
- Neurosecretory centers
of insects 68
endocrine activity 67 68
- Neurospora tryptophan synthesis in 28
- Niacin thyroid and 489

- Nitrogen metabolism thyroid hormone and 480-484
- Nitrogen mustard
antithyroid activity 471
structure 471
- Noradrenaline 102-103 143
action mechanism of 133
neurohormone 99 142
physiological 130-134
pressor cortisone and 714
reversal of, 132, 133
adrenaline and 130 131
biosynthesis 107
site of 113
chemical classification 143
nature 99
as main product of adrenergic neurons 102 130
receptors for 134
response of eosinophils to 745
storage in nervous tissue 107
- 19 Nordeoxycorticosterone activity, 544
- 19 Nor Δ^4 ketosteroids synthesis 543
- 10 Norprogesterone 543
- 19 Norprogesterone
activity, progestational 543 544
synthesis partial 543
- 19 Norsteroids syntheses 539-544
- 19 Nortestosterone
conversion of estrone to 543
testosterone and 544
- Nucleic acids
biosynthesis folic acid and 377
growth hormone and 249
effect of cortisone on 724
metabolism, effect on response of pigeon crop glands to prolactin 376
role of thyroid in 483-484
role in bud differentiation 22
in thyroid 437
- Nucleoproteins biosynthesis growth hormone and 249
- Nutrition thyroid and 460-461
- O
- Octopus enteramine in 78
- Ophthalmopathy in thyroid disease 833-835
hyaluronidase and 833-834
- treatment of 834-835
thyrotropin and 834 835
- Osteoporosis steroid therapy and 292
- Ovarian hormones
assay of 666-668
bovine, action on invertebrates 81
gonadotropic activity of anterior pituitary and 334
physiology of 666-684
prolactin and 360
- Ovary 852-861
agenesis 853, 865
treatment with estrogens 854
of cattle response to gonadotropic stimulation 338
deficiency primary, with high FSH 858
effect of adrenogenital syndrome on 859
of estrogen on, 669
follicle cysts of 855
function cortisone and 784
effect of metabolic disturbances on 859
gonadotropins governing 369
mechanism of, 852
genetic abnormalities 853-854
hypofunction psychogenic 857-858
endocrine patterns of 857-858
treatment 858
infantile with eunuchoidism 854-855
treatment 854-855
urinary T_H in 854
inhibitory effect on lactogenesis 362 369
plant as source of auxin 31
rat as prolactin target tissue 359
- Stein Leventhal syndrome 856-857
differential diagnosis 857
etiology 856
hormone therapy 857
as source of lactation inhibiting hormone 362
of steroid hormones 590ff
of androgens 610 611 612
of estrogens 593 594
- Ovulation
in experimental animals effect of progesterone on 333 334
in farm animals experiments to control 337-342
in hen factors controlling 331

- neurogenic stimulus required for 331
333
inhibitory of 331 333
- Oxidases adrenocortical steroids and 730-731
- Oxytocin 108 143
activity 137 402
amino acid composition 403-404
amino acid sequences in 407 408 412
chemical nature 99 137 143
effect on milk-ejection 400
electrophoretic properties 404-405 409
elementary analysis 404
molecular weight 404
as neurosecretory material 99 143
origin 100 118
physicochemical properties 404-405
polypeptide nature 137, 143 403
purification 401-402
stability, 409
structure 405-409 412 415
synthesis 409-412
vasopressin and 130
- P
- PAS see Periodic acid Schiff reaction
- PBI see Iodine protein bound
- IGH see Growth hormone of anterior pituitary
- PMG see Gonadotropin pregnant mare's serum
- Pain threshold adrenocortical steroids and 718
- Pancreas
effect of estrogens on 642
relationship between pituitary and renals and 726-727
- Pancreatotropic factor pituitary 251 257-261
- Panhypopituitarism effect of cortisone on diuresis in 711
- Pantothenic acid
adrenocortical steroids and 737-738
coenzyme A as functional form of 737
deficiency effect on adrenals 737
stress and 738
- Paper chromatography in study of plant growth hormones 26
- Paralysis familial periodic thyrotoxicosis and 86
- Parathyroids
active principles assay of 166-169
attempts at purification 166
at separation 163
activity 153
biological 164-165
calcium mobilizing 160 166
assay of 167
phosphaturic 153-155 160
assay of 167 168
nature of 165
blood pressure factor of 163
chemistry of 153-154
effect on blood phosphorus 160 161 163 164-165
on bone metabolism 153 155-160
on serum calcium 161-162
on soft tissues 160-161
extract calcium mobilizing effect following nephrectomy 156 157
psychological effects of administration of 164
function test for 155
kidney and 153-155
lactation and 169
physiology of 153
pituitary and 162-163
teeth and 169
thyroid and 163
- Parturition
effect on progesterone level 350
onset of lactation at 350
factor affecting 359-363
- Pectin methylesterase indoleacetic acid and 44
- Pemphigus ACTH and 755
- Penicillin sensitivity to cortisone and 770
- Pentobarbital thyroid and 496
- Pepsin
action on corticotropin 4 227-228
on FSH 218
- Peptone antithyroid 476
- Periodic acid Schiff reaction 312
- Peroxidase role in intrathyroidal iodine metabolism 442 443
- Phenobarbital toxic effects thyroxine and 436
- Phenols aromatic antithyroid activity 470

- Phenothiazine 468
 antithyroid activity 468
- Phenoxy compounds
 auxin activity optical isomerism and 11
 as auxin antagonists 11
- Phenoxyacetic acids substituted auxin activity 10
- Phenylacetic acid, as anti-auxin 38
- Phenylalanine deficiency effect on thyroid 461
- Phenylbutazone 470
 antithyroid activity 470
- γ Phenylbutyric acid as anti-auxin 38
- 2 Phenylcyclopropane 1 carboxylic acid
 auxin activity of *cis* isomer 8
- Phosphatase
 alkaline adrenocorticosteroids and 723
 of plasma growth hormone and 250
- Phosphaturia parathyroids and 153
 154-155 160
- 6 Phosphogluconate dehydrogenase
 possible role in lactogenesis 354
- Phosphorus
 blood parathyroids and 160 162 163
 164-165
 metabolism growth hormone and 249-250
 insulin and 187
 role of thyroid in 487
 radioactive in assay of thyrotropic hormone 207-208
- Phosphorylase
 activation by epinephrine 194 195
 by glucagon 194 195
 adrenaline and 723
 cortisone and 723
- Phosphorylation
 auxin activity and 39 40
 biological insulin and 188
 oxidative role of thyroid hormone in 477
- Photoperiodism 29
 flowering and 29 32
 function of dark period in 31
 sexuality of flowers and 32
- Phycosoma
 internephridial organs 59 60
 endocrine activity of 60
- Pilocarpine trophic action 130
- Pineapple
 flowering induction by auxins 30
 possible mechanism 31
- Pitoin activity diuretic 398 399
- Pitressin
 activity diuretic 398 399
 amino acid composition species differences in 398
 depletion of adrenal ascorbic acid by 712
 salt uresis and 395
- Pituitary anterior 823-827
 activity adrenocorticotrophic hypothalamus and 330
 antidiuretic 399
 gonadotropic 317-342
 chemotransmitter controlling 329
 nature of 330-332
 origin of 329 330
 effect of light on 317-318
 of stimuli from reproductive organs on 332
 estrus cycle and 334-335
 inanition and 336 337
 nutritional factors affecting 337ff
 ovarian hormones and 334
 pituitary stalk and 336
 pregnancy and 335 337
 role of adrenaline in 333
 of central nervous system in 330ff
 of tuber cinereus in 329
 small granule fraction and 316
 vitamins B and 337
- adrenal cortex and 219
- cells of 310-311 823
 acidophilic 311
 as source of luteinizing hormone 310
 sulphhydryl and/or disulfide groups in 316
 basophilic 311 314
 histolization in *Cushing's syndrome* 844
 as source of FSH 310 314
 of ICSH 313 314
 classification 311 312 803
- cytochemistry of 311-316
 diseases of 823-827

- effect of androgens on 613
 of environmental changes on 311
 318
 of estrogen on 611
 of progesterin on 611
 extracts of galactopoietic activity
 360-366
 diabetogenic activity and 360-366
 hexokinase and 184
 galactopoietic complex of 364
 gonadotrophs of 315
 hormones of 872 see also Gonadotropic
 hormones and individual hormones
 cellular origin 310ff
 chemistry of 231-234
 secretion estrogen and 360
 independent 873
 human gonadotropin content 336-337
 hypothalamus and 287-268
 pathways between 322-330
 insulin and 180 186 187
 maintenance of lactation and 363
 pathways of stimulus from retina to
 370
 as possible source of B I melanophore
 expanding hormone 455
 puberty and 869
 receptor for stimuli 317 319 319
 release of ACTH by estrogens and 784
 mechanism of 692
 of corticotropin from rate of 780-787
 role of retina in transference of light
 stimuli to 319
 stalk of 326-330
 lactation and 367
 sympathetic control 320 321
 water metabolism and 34-395
Pituitary extracts
 action on crustacean chromatophores
 81
 activity diabetogenic 262
 fat mobilizing 254 255
 ascorbic acid depleting factor in 277
 effect on carbohydrate metabolism, 257
 growth response to diet and 238
 ketogenic factor in 250-254
 weight factor in 291 277 219
 sources of 381
Pituitary gland
 acidophilic zone of 345
 activity gonadotropic effect of suck-
 ling stimulus on 370
 interaction between hormones and
 neurogenic mechanisms control-
 ling 332
 basophilic zone of 345
 diabetes mellitus and 251
 distribution of nerve fibres in 323
 effect of ACTH therapy on 691
 embryonic origin 359
 fowl brood cells in 347
 exophthalmos producing factor of 417
 function mechanism of 334ff
 gonads and 337 338 358 675
 hormones of see also individual hor-
 mones
 antibodies to 205-206
 interaction between and neurogenic
 mechanism controlling pituitary
 activity 332
 target organs for 456
 hypothalamus and 331-33
 links between central nervous system
 and nature of 330ff
 liver fat and 733 734
 number of corticotropic substances in
 216 296
 of thyrotropic factors in 207 457
 parathyroids and 167-163
 relationship between adrenals pan-
 creas and 776-777
 removal see Hypophysectomy
 similarity between vertebrate and
 invertebrate homologues 87
 stalk of portal system in 324 320 376
 gonadotropic activity of anterior
 pituitary and 376
 transmission of hypothalamic stim-
 uli and 375 327 329 330
 testes and 864
Pituitary posterior 389-90
 extracts action on adrenals of hypo-
 physectomized animals 278
 effects of 390
 hormones of 389-431 see also indi-
 vidual hormones
 secretory elements 390-391
 intermediate lobe of 417
 melanophore expanding (B) hor-
 mone of 390 417-420

- Iphenothiazine 168
 antithyroid activity 468
 Phenox compounds
 auxin activity optical isomerism and 11
 as auxin antagonists 11
 Phenoxycetic acids substituted auxin activity 10
 Phenylacetic acid as antiauxin 38
 Phenylalanine deficiency effect on the thyroid 461
 Phenylbutazone 470
 antithyroid activity 470
 γ Phenylbutyric acid as antiauxin 38
 2 Phenylcyclopropane 1 carboxylic acid auxin activity of cis isomer 8
 Phosphatase
 alkaline adrenocorticosteroids and 723
 of plasma growth hormone and 250
 Phosphatemia parathyroids and 153
 154-155 160
 6 Phosphogluconate dehydrogenase possible role in lactogenesis 354
 Phosphorus
 blood parathyroids and 160 162 163
 164-165
 metabolism growth hormone and 249-250
 insulin and 187
 role of thyroid in 457
 radioactive in assay of thyrotropic hormone 207-208
 Phosphorylase
 activation by epinephrine 194 195
 by glucagon 194 195
 adrenaline and 723
 cortisone and 723
 Phosphorylation
 auxin activity and 39 40
 biological insulin and 168
 oxidative role of thyroid hormone in 477
 Photoperiodism 29
 flowering and 29 32
 function of dark period in 31
 sexuality of flowers and 32
 Phycomsoma
 internephridial organs 59 60
 endocrine activity of 60
 Pilocarpine trophic action 130
 Linapple
 flowering induction by auxins 30
 possible mechanism 31
 Pitocin auxin diuretic 338 399
 Pitressin
 activity diuretic 398 399
 amino acid composition species differences in 398
 depletion of adrenal ascorbic acid by 712
 salt uresis and 395
 Pituitary anterior 822-827
 activity adrenocorticotrophic hypothalamus and 350
 antidiuretic 399
 gonadotropic 317-342
 chemotransmitter controlling 329
 nature of 330-332
 origin of 329 330
 effect of light on 317-318
 of stimuli from reproductive organs on 332
 estrus cycle and 334-335
 inhibition and 336 337
 nutritional factors affecting 33 ff
 ovarian hormone and 334
 pituitary stalk and 326
 pregnancy and 335 337
 role of adrenaline in 333
 of central nervous system in 330ff
 of tuber cinereum in 3-9
 small granule fraction and 316
 vitamin B and 337
 adrenal cortex and 219
 cells of 310-311 823
 acidophilic 311
 as source of luteinizing hormone 310
 sulfhydryl and/or disulfide groups in 316
 basophilic 311 314
 hyalinization in Cushing's syndrome 844
 as source of FSH 310 314
 of ICSH 313 314
 classification 311 312 873
 cytochemistry of 311-316
 diseases of 873-877

- effect of androgens on 673
 of environmental changes on 317
 318
 of estrogen on 671
 of progesterin on 671
 extracts of galactopoietic activity
 365-366
 diabetogenic activity and 365 366
 hexol inase and 184
 galactopoietic complex of 364
 gonadotrophs of 315
 hormones of 899 see also Gonadotropic
 hormones and individual hormones
 cellular origin 310ff
 chemistry of 291-334
 secretion estrogen and 368
 independent 893
 human gonadotropin content 336-337
 hypothalamus and 282 69
 pathways between 329-330
 insulin and 185 186 187
 maintenance of lactation and 563
 pathways of stimulus from retina to
 320
 as possible source of B I melanophore
 expanding hormone 425
 puberty and 869
 receptor for stimuli 317 318 319
 release of ACTH by estrogens and 784
 mechanism of 692
 of corticotropin from rate of 285-287
 role of retina in transference of light
 stimuli to 319
 stalk of 326-330
 lactation and 367
 sympathetic control 320 321
 water metabolism and 394-395
Pituitary extracts
 action on crustacean chromatophores
 81
 activity diabetogenic 962
 fat mobilizing 254 255
 ascorbic acid depleting factor in 217
 effect on carbohydrate metabolism, 257
 growth response to diet and 238
 ketogenic factor in 252-254
 weight factor in 291 277 273
 sources of 389
Pituitary gland
 acidophilic zone of 345
 activity gonadotropic effect of suck-
 ling stimulus on 370
 interaction between hormones and
 neurogenic mechanisms control-
 ling 332
 basophilic zone of 345
 diabetes mellitus and 251
 distribution of nerve fibres in 323
 effect of ACTH therapy on 691
 embryonic origin 389
 fowl broody cells in 347
 exophthalmos producing factor of 457
 function mechanism of 334ff
 gonads and 339 338 505 675
 hormones of see also individual hor-
 mones
 antibodies to 205-206
 interaction between and neurogenic
 mechanism controlling pituitary
 activity 332
 target organs for 456
 hypothalamus and 331-339
 links between central nervous system
 and nature of 330ff
 liver fat and 733 734
 number of corticotropic substances in
 276 286
 of thyrotropic factors in 207 457
 parathyroids and 162-163
 relationship between adrenals pan-
 creas and 796-797
 removal see Hypophysectomy
 similarity between vertebrate and
 invertebrate homologue 82
 stalk of portal system in 324 325 576
 gonadotropic activity of anterior
 pituitary and 326
 transmission of hypothalamic stim-
 uli and 325 327 329 330
 testes and 864
Pituitary posterior 389-390
 extracts action on adrenals of hypo-
 physectomized animals 278
 effects of 590
 hormones of 389-431 see also indi-
 vidual hormones
 secretory elements 390-391
 intermediate lobe of 417
 melanophore expanding (B) hor-
 mone of 390 417-425

- isolation of possibly unitary hormone from 416-417
- neural lobe of, 98 391ff
 - activity, antidiuretic, adrenocortical steroids and 712
 - assay of, 393
 - biological assay of 391-393
 - multihormone hypothesis 416
 - one hormone hypothesis 416-417
 - oxytocic assay of 392-393
 - pressor assay of 392
- hormones of 390-417 *see also* individual hormones
 - chemistry 403-415
 - origin site of 390-391
 - purification 400-403
 - role in milk ejection 363 390 393 399-400
 - structure 98
 - relationship of adrenal cortex and to water metabolism 712
- Pituitary adrenal system 630
 - activation by adrenaline 281
 - role in maintenance of lactation 365
- Pituitary lactogenic complex 351
- Pituitary ovarian axis, hypothalamus and 852
- Placenta 861-862
 - extracts, effect on adrenal cortex of hypophysectomized animals 278
 - function in pregnancy 861
 - gonadotropic activity 335 336
 - luteotropic activity 360 371 372
 - as source of steroid hormones 590 591
 - of adrenocortical steroids 842
 - of ACTH 842
 - of estrogens 593 594 595
 - of progesterone 601 602 603
 - of prolactin 349 360
- Planaria
 - acetylcholine in 101 130
 - hormones in 58
 - pilocarpine in 130
- Plant hormones *see also* Auxins
 - growth 1-56
 - reproduction and 29-35
- Plant(s)
 - disappearance of auxin from 28
 - dormancy, auxin inhibitors and 26
 - sulphydryl compounds and 27
 - growth, cytochrome oxidase and 43
 - mechanism of auxin action in 1-2
 - pathogens, effect on auxin production 35
 - reproduction, hormones and 29-35
 - respiration, auxin and 39 40
 - transport of auxins in 16-20
 - tumors of auxin and 35-37
 - inhibitors of 36
 - water uptake 42-48
 - auxin and 43 45
 - growth and 43 45
 - metabolic control of 43
- Plasma, hexokinase inhibitor in, 185
- Poliomyelitis virus, sensitivity to thyroid and 497
- Pollen sensitivity, cortisone and 773
- Pollination
 - enzymatic auxin formation and 33
 - fruit set and 33
- Polychaetes, inhibitory hormone in brain of 59
- Polypeptides as hormones 137 143-144 405
- Polysaccharides
 - pneumococcal ACTH and 770 775
 - cortisone and 770 773
- Potassium,
 - effect on acetylcholine release from nerve 117
 - metabolism, adrenaline and 708
 - adrenocorticosteroids and 707-709
- Potassium chlorate, effect on Graves disease 462
- Δ^4 ¹¹ Pregnadiene 3 20 dione 11 β h₁
 - hydroxylation, enzymatic 646
- Δ^4 ¹⁸ Pregnadiene 3 20 dione, metabolism 632
- Pregnancy
 - Addison's disease and 841-842
 - chorionic gonadotropin in serum and urine of early 343
 - complications of 862-864
 - abortion 862-863
 - hydramnios, mole and chorionepithelioma 863
 - toxemia 863-864
 - corpus luteum and 862
 - effect of cortisone on 699 700
 - on serum protein bound iodine 459

- estrogens and 862
 - in urine of 593
- function of placenta in 861
- gonadotropic activity of anterior pituitary and 335
- hyperthyroidism and 831
- 17 ketosteroids in urine of 861
- ovarian hyperfunction in early 860
- in precocious puberty 869
- secretion of FSH during 335 336
- Pregnanediol
 - assay of 667
 - biosynthesis of 607
 - urinary abortion and 862
- Pregnane 3α 20α diol
 - in adrenogenital syndrome 602
 - cholesterol as precursor of 601
 - metabolism 603
 - as metabolite of deoxycorticosterone 637
 - of progesterone 631 632
 - from pregnane 3 20 dione 606
 - from pregnenolone 605
 - from progesterone 606
 - sources 605
- Pregnane 3β 20α diol
 - metabolism 603
 - sources 60,
- Pregnanediols urinary configuration 69
- Pregnane 3 20 dione
 - metabolism 603
 - sources 604
- Pregnanetriol
 - excretion in adrenogenital syndrome 848 849
 - as metabolite of 17 hydroxyprogesterone 850
 - as precursor of hydrocortisone 850
- Pregnane 3α 17α 20α triol as metabolite of 17 α hydroxyprogesterone 6 3
- Pregnane 3α 11β 20α triol as metabolite of corticosterone 636
- Δ^4 Pregnene 3β 20α diol
 - from cholesterol 532-533
 - metabolism 603
 - from pregnenolone 603
 - sources 604
- Δ^4 Pregnene 3β 20β diol from cholesterol 532-533
- Δ^4 Pregnene 7β 16α 20α triol 561
 - configuration 563
- Pregnenolone
 - action of 7β dehydrogenase on 678
 - biosynthesis 677 628
 - from cholesterol 649
 - metabolism 603 605 606
 - sources 604
- Pregandrogens 621
 - androgenic metabolites of 611
 - of tissues 611
- Progesterone 601-610
 - activity central nervous system and 333
 - effect of 11(12) desaturation on 559
 - multiplicity of 666
 - progestational 601 602
 - assay of 667 668
 - biosynthesis 602 624 626 628
 - adrenogenital syndrome and 632
 - effect of estrogen plus on lactation 361-62
 - on pituitary gonadotropic activity 334
 - on thyroid function 459
 - on sex cycle of rats 333
 - hypothalamic lesions and 322
 - enzymatic reduction of ring A in 648 649
 - estrogen and 130 640
 - hydroxylation 607
 - enzymatic 645
 - as intermediate in adrenocortical steroid biosynthesis 602
 - isomers structural 629
 - labeled preparation 575 576
 - level effect of parturition on 339
 - in Leydig cells 864
 - localization 609-610
 - metabolism 589 607-607 631 632
 - metabolites of 675
 - microbiological formation 643
 - reactions 607-609 642
 - as precursor of corticosterone 675 626
 - of cortisol 625 676 630
 - of 11 deoxycortisol 675
 - of hydroxyprogesterones 674
 - radioactive metabolism 609
 - reduction products 651
 - release of luteinizing hormone and 335
 - role in pregnancy 862

- isolation of possibly unitary hormone from 416-417
- neural lobe of 98 391ff
- activity antidiuretic adrenocortical steroids and 712
- assay of 393
- biological assay of 391-393
- multihormone hypothesis 416
- one hormone hypothesis 416-417
- oxytocic assay of 392-393
- pressor assay of 392
- hormones of 390-417 see also individual hormones
- chemistry 403-415
- origin site of 390-391
- purification 400-403
- role in milk ejection 363 390 393 399-400
- structure 98
- relationship of adrenal cortex and to water metabolism 712
- Pituitary adrenal system 630
- activation by adrenaline 281
- role in maintenance of lactation 360
- Pituitary lactogenic complex 301
- Pituitary ovarian axis hypothalamus and 852
- Placenta 861-862
- extracts effect on adrenal cortex of hypophysectomized animals 278
- function in pregnancy 861
- gonadotropic activity 370 336
- luteotropic activity 360 371 372
- as source of steroid hormones 500 861
- of adrenocortical steroids 842
- of ACTH 842
- of estrogens 593 594 595
- of progesterone 601 602 603
- of prolactin 349 369
- Planaria
- acetylcholine in 101 130
- hormones in 58
- pilocarpine in 130
- Plant hormones see also auxins
- growth 1-56
- reproduction and 20-35
- Plant(s)
- disappearance of auxin from 28
- dormancy auxin inhibitors and 26
- sulphydryl compounds and 27
- growth cytochrome oxidase and 43
- mechanism of auxin action in 1-2
- pathogens effect on auxin production 30
- reproduction hormones and 29-30
- respiration auxin and 39 40
- transport of auxins in 16-20
- tumors of auxin and 35-37
- inhibitors of 36
- water uptake 42-48
- auxin and 43 40
- growth and 43 45
- metabolic control of 43
- Plasma hexokinase inhibitor in 185
- Poliovirus sensitivity to thyroid and 497
- Pollen sensitivity cortisone and 773
- Pollination
- enzymatic auxin formation and 33
- fruit set and 33
- Polychaetes inhibitory hormone in brain of 59
- Polypeptides as hormones 137 143-144 405
- Polysaccharides
- pneumococcal ACTH and 770 773
- cortisone and 770 773
- Potassium
- effect on acetylcholine release from nerve 117
- metabolism adrenaline and 708
- adrenocorticosteroids and 707-708
- Potassium chlorate effect on Graves' disease 462
- Δ^1 Pregnenolone 3 20 dione 118 by hydroxylation enzymatic 646
- Δ^1 Pregnenolone 3 20 dione metabolism 632
- Pregnancy
- Addison's disease and 841-842
- chorionic gonadotropin in serum and urine of early 343
- complications of 862-864
- abortion 867-868
- hydatiform mole and chorionepithelioma 863
- toxemia 863-864
- corpus luteum and 862
- effect of cortisone on 699 700
- on serum protein bound iodine 459

- estrogens and 862
 - in urine of 873
- function of placenta in 861
- gonadotropic activity of anterior pituitary and 335
- hyperthyroidism and 831
- 17 ketosteroids in urine of 861
- ovarian hyperfunction in early 860
- in precocious puberty 860
- secretion of FSH during 335 336
- Pregnanediol
 - assay of 667
 - biosynthesis of 602
 - urinary abortion and 862
- Pregnane 3α 20α diol
 - in adrenogenital syndrome 602
 - cholesterol as precursor of 591
 - metabolism 603
 - as metabolite of deoxycorticosterone 637
 - of progesterone 631 632
 - from pregnane 3 20 dione 606
 - from pregnenolone 605
 - from progesterone 606
 - sources 605
- Pregnane 3β 20α diol
 - metabolism 603
 - sources 605
- Pregnanediols urinary configuration 529
- Pregnane 3 20 dione
 - metabolism 603
 - sources 604
- Pregnanetriol
 - excretion in adrenogenital syndrome 849-840
 - as metabolite of 17 hydroxyprogesterone 850
 - as precursor of hydrocortisone 850
- Pregnane 3α 17α 20α triol as metabolite of 17α hydroxyprogesterone 633
- Pregnane 3α 11β 20α triol as metabolite of corticosterone 636
- Δ^5 Pregnene 3β 20α diol
 - from cholesterol 532-533
 - metabolism 603
 - from pregnenolone 605
 - sources 604
- Δ^5 Pregnene 3β 20β diol from cholesterol 532-533
- Δ^5 Pregnene 3β 16α 20α triol 561
 - configuration 563
- Pregnenolone
 - action of 3β dehydrogenase on 628
 - biosynthesis 627 643
 - from cholesterol 649
 - metabolism 603 605 606
 - sources 604
- Proandrogens 621
 - androgenic metabolites of 611
 - of tissues 611
- Progesterone 601-610
 - activity central nervous system and 333
 - effect of 11(12) desaturation on 559
 - multiplicity of 666
 - progestational 601 602
 - assay of 667 668
 - biosynthesis 602 624 626 628
 - adrenogenital syndrome and 652
 - effect of estrogen plus on lactation 361-362
 - on pituitary gonadotropic activity 334
 - on thyroid function 459
 - on sex cycle of rats 333
 - hypothalamic lesions and 322
 - enzymatic reduction of ring A in 648 649
 - estrogen and 130 640
 - hydroxylation 607
 - enzymatic 645
 - as intermediate in adrenocortical steroid biosynthesis 602
 - isomers structural 629
 - labeled preparation 575 516
 - level effect of parturition on 359
 - in Leydig cells 864
 - localization 609-610
 - metabolism 589 602-607 631 632
 - metabolites of 635
 - microbiological formation 643
 - reactions 607-609 642
 - as precursor of corticosterone 625 626
 - of cortisol 625 626 630
 - of 11 deoxycortisol 625
 - of hydroxyprogesterones 624
 - radioactive metabolism 609
 - reduction products 651
 - release of luteinizing hormone and 335
 - role in pregnancy 862

- sources 601 601
- therapeutic effect on Stein Leventhal syndrome 857
- Progesterin**
 - aminopterin and 669
 - assay of 666
 - physiological effects 669-672
 - source 665
 - thermogenic effect in men 672
- Progestogens** placenta as source of 461
- Prolactin** 345-378
 - action on invertebrates 81
 - activity 351
 - antigonadal 369 373 374
 - assay of 375-378
 - mammalian 378
 - pigeon crop weight method 375
 - diabetogenic 374
 - galactopoietic 365-366
 - glycotropic 236
 - lactogenic 351-363
 - inhibition by combined estrogen and progesterone administration 360
 - possible mechanism 352ff
 - use of radioiodine in study of 352
 - luteotropic 360 363 368 369 370ff
 - in rat 310 319 321 362 367 370
 - following hypophysectomy 370
 - mammogenic 369
 - in rat 362
 - cellular origin 312 313 316 345-347
 - corpora lutea as target tissue for 352
 - discovery of 351
 - disulfide groups in 316
 - effect on adrenal cortex of hypophysectomized animals 278
 - on luteotropic activity of human chorionic gonadotropin 342
 - on male reproductive system 373
 - identity with luteotropin 369
 - inactivation 358
 - labeled preparation 322
 - metabolic effects in birds 373 374-375
 - in mammals 373-374
 - nomenclature 310
 - ovarian function and 852
 - ovarian hormones and 360
 - physiology of 369
 - pituitary mammogen and 348
 - placental 349 369 372 373
 - corpus luteum of lactation and 370
 - response of mammary duct system to 348 352, 378
 - role in maintenance of lactation 363-365 367
 - in males 864
 - in mammary development 347-351
 - secretion mechanism of 367
 - stimuli affecting 367 368
 - target tissues for 352 358
 - urinary excretion 371
 - Proline oxidase** cortisone and 731
 - Promizole**
 - antithyroid activity 467
 - structure 467
 - Propionylcholine** in ox spleen 100
 - Protein metabolism**
 - adrenal cortex and 727-732
 - adrenalectomy and 729 729
 - adrenocortical steroids and 690 721
 - diabetes and 188
 - effect of growth hormone preparations on 246-249
 - Proteins**
 - biosynthesis insulin and 188 189
 - blood cortisone and 750
 - dietary growth response to pituitary extracts and 238
 - iodinated in milk 439
 - in patients with thyroid cancer 450
 - serum iodine in 450
 - Thoracic glands** of insects 71
 - Corpora cardiaca** and 72
 - dimorphism of in bees 77
 - endocrine activity 68 73
 - hormonal control 141
 - hormonal stimulation by neurosecretory brain cells 72
 - possible mechanism 68
 - role in postembryonic development 71-72
 - as source of growth and differentiation hormone 72 82
 - Thorax** differentiation center in 71
 - Protoanemonin** auxins and 24
 - Protozoa** acetylcholine in 101
 - Pseudocyesis** 860-861
 - hormonal factors and in rats 349

- symptoms 861
 treatment 861
 Pseudopregnancy see Pseudocyesis
 Pseudosapogenin
 structure 561
 synthesis 561
 Puberty 868-869
 anterior pituitary and 869
 precocious 869
 in adrenogenital syndrome 6,2
 pseudoprecocious 869
 Urines effect of adrenocortical steroids
 on 732
 Pyriloxine
 adrenal function and 743
 deficiency cortisone and 738-739
 effect on experimental lympho-
 sarcoma 752
 effects of 738-739
 lymphoid tissue and 743
- ### R
- Radioiodine uptake by thyroid adreno-
 cortical steroids and 782
 Rat
 effect of growth hormone preparations
 in 242-243 244-245
 luteotropic activity of prolactin in
 310-349 351-362 367-369 370
 prolactin like substance in placenta of
 3,2
 Reichstein's Compound S see 11 Deoxy
 cortisol
 Reproductive organs
 male effect of prolactin on 373
 stimuli from effect on gonadotropic
 activity of anterior pituitary 332
 Resorcinol antithyroid activity 4,0
 Respiration corticotropin and 296
 Reticuloendothelial system effect of
 cortisone on 743
 Rhenium in blood and thyroid 4,6
 Rhinitis allergic cortisone and 773
 Riboflavin
 auxin destruction and 28
 thyroid and 499
 Ribonucleic acids auxin and 22-35
 Rodents metabolism of testosterone in
 621-622
 Roots
 in bioassay of auxins 3
 effect of antiauxins on 38
 of auxins on 15-23-24 44-128
 geotropic responses substances affect-
 ing 24
 growth auxins and 23-24 44-128
 growth promoting substances for 3
 Rotifers subcerebral gland of 59
 Rutabaga
 inhibitory effect on thyroid 461
 isolation of goitrogen from 461-838
- ### S
- SHH (Somatotrophic hormone) see
 Growth hormone of anterior pitui-
 tary
 Saline intravenous effect on adrenal
 cortex of hypophysectomized ani-
 mals 278
 Saliva effect of adrenocortical steroids
 on 713
 Salt hormones 844
 adrenogenital syndrome and 8,1
 formation in adrenal cortex of rodents
 851
 Salt metabolism
 role of adrenal cortex in 395
 of neurohypophysis in 394-399
 Sapogenins
 degradation to 20 ketosteroids 466
 11 oxygenated steroids from 5,7
 Sarmetogenin 551
 conversion to cortisone 551
 structure 551
 Schizophrenia
 adrenals and 717
 following LSD intake 136
 Schwartzman phenomena cortisone and
 771
 Scopoletin
 auxin and 29
 inhibition of indoleacetic acid oxidase
 by 29
 in plants 24
 Scurvy
 effect of cortisone on 740-741-742
 rheumatoid arthritis and 739
 Seeds
 response to cold 32
 reversal of 32
 Semen
 estrogens in human 595

- sources 601 604
- therapeutic effect on Stein Leventhal syndrome 857
- Progesterin**
 - aminopterin and 669
 - assay of 666
 - physiological effects 600-672
 - source 665
 - thermogenic effect in men 672
- Progestogens** placenta as source of 561
- Prolactin** 345-378
 - action on invertebrates 81
 - activity 351
 - antigonadal 369 373 374
 - assay of 375-378
 - mammalian 378
 - pigeon crop weight method 375
 - diabetogenic 374
 - galactopoietic 365-366
 - glycotropic 256
 - lactogenic 351-363
 - inhibition by combined estrogen and progesterone administration 360
 - possible mechanism 352ff
 - use of radioiodine in study of 352
 - luteotropic 360 363 368 369 370ff
 - in rat 310 349 351 362 367 370
 - following hypophysectomy 370
 - mammogenic 369
 - in rat 362
 - cellular origin 312 313 316 345-347
 - corpora lutea as target tissue for 352
 - discovery of 351
 - disulfide groups in 316
 - effect on adrenal cortex of hypophysectomized animals 278
 - on luteotropic activity of human chorionic gonadotropin 342
 - on male reproductive system 373
 - identity with luteotropin 369
 - inactivation 358
 - labeled preparation 352
 - metabolic effects in birds 373 374-375
 - in mammals 373-374
 - nomenclature 310
 - ovarian function and 852
 - ovarian hormones and 360
 - physiology of 309
 - pituitary mammogen and 348
 - placental 349 369 372-373
 - corpus luteum of lactation and 340
 - response of mammary duct system to 318 352 378
 - role in maintenance of lactation 363-365 367
 - in males 864
 - in mammary development 347-351
 - secretion mechanism of 367
 - stimuli effecting 367 368
 - target tissues for 342 348
 - urinary excretion 371
- Proline oxidase** cortisone and 731
- Promizole**
 - antithyroid activity 467
 - structure 467
- Propionylcholine** in ov spleen 100
- Protein metabolism**
 - adrenal cortex and 727-732
 - adrenalectomy and 728 729
 - adrenocortical steroids and 690 721
 - diabetes and 188
 - effect of growth hormone preparations on 216-219
- Proteins**
 - biosynthesis insulin and 168 169
 - blood cortisone and 780
 - dietary growth response to pituitary extracts and 238
 - iodinated in milk 439
 - in patients with thyroid cancer 450
 - serum iodine in 450
- Prothoracic glands** of insects 71
 - corpora cardiaca and 72
 - dimorphism of in bees 77
 - endocrine activity 68 73
 - hormonal control 141
 - hormonal stimulation by neurosecretory brain cells 72
 - possible mechanism 68
 - role in postembryonic development 71-72
 - as source of growth and differentiation hormone 72 82
- Prothorax** differentiation center in 71
- Protoanemonin** auxins and 24
- Protozoa** acetylcholine in 101
- Pseudocyesis** 860-861
 - hormonal factors and in rats 349

- Steroids see also Steroid hormones and individual steroids
 activity anesthetic 720
 configuration 520
 absolute 533-536
 of C 17 526-528
 of C 20 528-533
 derived by asymmetric synthesis 534-530
 effect on hyaluronidase action 63-64
 possible mechanism of 764
 esterification 566-571
 hydroxylation 566-571
 enzymatic 645-646
 labeled distribution of labeling agent 578
 with isotopic hydrogen 576ff
 by exchange reactions 576-588
 by hydrogenation of double bonds 576-577
 metabolism effect of mammalian enzymes on 644-649
 hypotheses concerning mechanism of 649-650
 molecular rotation 530-506
 nomenclature 522-523
 numbering of rings in 522
 11 oxygenated from 5 allo Δ^5 dienes 555
 antiinflammatory action 404
 biosynthesis coenzyme A and 738
 proteins metabolism and 727
 effect of diffuse adrenal hyperplasia on formation 800-851
 natural 501
 response to stress and 704
 secretion in Cushing's syndrome 846
 syntheses 501-560
 from Δ^5 dienes 552-506
 from 12 ketones 506-558
 microbial 502 607-609
 starting material 552ff
 16 oxygenated configuration 560-563
 reactions of the side chains 566-571
 ring substituents classification 520-526
 geometry of 526
 sex reversal with in amphibia 670
 stereochemistry of 520-536
 toxicity to embryos 675
 ultraviolet spectra 522
 urinary of man 544
 Δ^4 Steroids labeled preparation 577
 Δ^7 Steroids synthesis 507
 Δ^5 Steroids 544
 Δ^5 Steroids
 preparation 502-503
 synthesis from natural products 507
 of 11 oxygenated steroids from 553-556
 $\Delta^{11,13}$ Steroids biological activity 509
 Sterols side chains orientation of 526
 Stigmasterol
 conversion to 11 oxygenated steroids 502
 structure 501
 Stomach
 effect of adrenocortical steroids on 700-706
 of adrenocorticotrophic hormone on 755
 Spleen effect of cortisone on 743
 Streptococci hemolytic cortisone and 770
 Stress
 adrenal cortex and response to 689-757
 corticotropin secretion and 281-280-280
 effect on adrenals 701-736-737-741
 on thymus 742-743
 on thyroid 460
 on thyroxine metabolism 403
 ketogenesis and 706
 pantothenic acid and 708
 response of eosinophils to 745
 response to adrenocortical steroids and 704
 Substance P 100-137-143
 neurohormonal activity 99
 occurrence 100
 polypeptide structure 137
 Succinic dehydrogenase bone parathyroids and 160
 Sugar hormones 844-846-800
 see also Steroids 11 oxygenated
 Sugars see also Carbohydrates
 structure insulin sensitivity and 183-184
 synergism with auxins 24

- fructose content as measure of androgen secretion 867
- Serini reaction 527-528
- Serotonine see Enteramine & Hydroxytryptamine
- Serum
- diabetic hexokinase inhibitor in human 185
 - inhibitory effect of β lipoprotein fraction on glucose utilization and glycogenesis 186
 - mucoproteins parathyroids and 156
- Sex chromosomal 853
- Sexagens see Sex hormones
- Sex hormones see also individual hormones
- activity dosage and 665
 - assay statistical requirements for valid 668
 - in biological media isolation and purification 667
 - in crustaceans 64
 - effect on enzymes 673, 674 675-677
 - activating 677
 - inhibitory 676-677
 - species differences in 676
 - on pituitary activity 313
 - growth and 240-241
 - in insects 70
 - interaction 666
 - in mollusks 79
 - response to species differences in 665
 - role in sex differentiation in vertebrates 674-675
- Sex organs,
- male effect of androgens on 673
 - effect of corticotropin on of castrated rat 295
- Sheep
- effect of light on breeding season 317-318
 - survival following adrenalectomy 695
- Silkworm (*Bombyx mori*)
- embryonic development 70
 - hormonal control 70-71
 - prothoracic glands isolation of growth and differentiation hormone from 82
- Sinus gland crustacean 138
- chromatophorotropins in 61
 - corpora cardiaca and 139
 - effect of light stimulation of eye on neurosecretory materials in 109
 - origin of 139-140
 - release of hormones by 115, 138-140
 - storage of hormones in 63 138-140
 - structure 98-109
- Skatole antitumor activity 38
- Skin
- amphibian permeability to water 396-398
 - effect of adrenocortical steroids on 754-755
 - sensitivity adrenal steroids and 770-771
- Sodium metabolism effect of adrenocortical steroids on 704-707
- Spermatogenesis FSH and 864
- Squalene as precursor of cholesterol 592-593
- Steroid hormones see also Steroids and individual hormones
- biosynthesis role of enzymes in 590
 - sites of 590
 - of steroid nucleus 591-593
 - chemistry of 521-548
 - conjugation with sulfuric and glucuronic acid 593
 - hydroxylation 624-625 626 629-630
 - enzymatic 645-646
 - microbiological 641-643
 - infrared spectra of 572
 - labeled preparation 574-578
 - introduction of isotopic carbon 574-575
 - of isotopic hydrogen 575-576
 - metabolism 559-664
 - anabolic definition 590
 - catabolic definition 590
 - neutral cholesterol as precursor of 591
 - nuclear substitution products 650
 - reduction products 651
 - nomenclature 572
 - phenolic metabolic interconversions *in vivo* 599
 - origin site of 861
 - stereochemistry 523
 - syntheses partial 530-574
 - total 536-539
 - therapeutic effect on acromegaly 825
 - ultraviolet spectra of 522

- Steroids see also Steroid hormones and individual steroids
 activity anesthetic 720
 configuration 575
 absolute 533-536
 of C 17 526-528
 of C 20 528-533
 derived by asymmetric synthesis 534-535
 effect on hyaluronidase action 763-764
 possible mechanism of 764
 esterification 566-571
 hydroxylation 566-571
 enzymatic 645-646
 labeled distribution of labeling agent 578
 with isotopic hydrogen 576ff
 by exchange reactions 576-588
 by hydrogenation of double bonds 576-577
 metabolism effect of mammalian enzymes on 644-649
 hypotheses concerning mechanism of 649-650
 molecular rotation 535-536
 nomenclature 522-523
 numbering of rings in 572
 11 oxygenated from 5 allo Δ^7 μ ν dienes 555
 antiinflammatory action 704
 biosynthesis coenzyme A and 738
 proteins metabolism and 727
 effect of diffuse adrenal hyperplasia on formation 850-851
 natural 551
 response to stress and 704
 secretion in Cushing's syndrome 846
 syntheses 551-560
 from Δ^7 μ ν dienes 552-556
 from 12 ketones 556-558
 microbial 552 607-609
 starting material 557ff
 16 oxygenated configuration 560-563
 reactions of the side chains 566-571
 ring substituents classification 525 576
 geometry of 576
 sex reversal with in amphibia 675
 stereochemistry of 573-576
 toxicity to embryos 675
 ultraviolet spectra 572
 urinary of man 544
 Δ^8 Steroids labeled preparation 577
 Δ^7 Steroids synthesis 557
 μ Steroids 544
 Δ^7 μ ν Steroids
 preparation 552 553
 synthesis from natural products 557
 of 11 oxygenated steroids from 553-556
 $\Delta^{11(13)}$ Steroids biological activity 559
 Sterols side chains orientation of 576
 Stigmasterol
 conversion to 11 oxygenated steroids 552
 structure 551
 Stomach
 effect of adrenocortical steroids on 755-756
 of adrenocorticotrophic hormone on 755
 Spleen effect of cortisone on 743
 Streptococci hemolytic cortisone and 770
 Stress
 adrenal cortex and response to 689 757
 corticotropin secretion and 281 284-285
 effect on adrenals 701 756 737 741
 on thymus 742-743
 on thyroid 460
 on thyroxine metabolism 453
 ketogenesis and 736
 pantothenic acid and 738
 response of eosinophils to 745
 response to adrenocortical steroids and 704
 Substance P 100 137 143
 neurohormonal activity 99
 occurrence 100
 polypeptide structure 137
 Succinic dehydrogenase bone parathyroids and 160
 Sugar hormones 844 846 850
 see also Steroids 11 oxygenated
 Sugars see also Carbohydrates
 structure insulin sensitivity and 183 184
 synergism with auxins 24

- Sulfaguanidine antithyroid activity 462
 Sulfates, steroidal hydrolysis of 544
 Sulfhydryl compounds
 auxin activity and 39
 dormancy in plants and 27
 Sulfonamides antithyroid activity 469
 Sulfur
 metabolism growth hormone and 250-251
 in thyroid 437
 Sulfuric acid conjugation of steroid hormones with 590
 Superfetation in sheep 1 MG and, 340-341
 Superovulation in cows, gonadotropins and 341-342
 Sweat effect of adrenocortical steroids on 712-713
 Sympathins 102 113
 Synapoidin 806
 Syphilis
 effect of ACTH on 780
 of cortisone on 780
- T**
- T I P (Tumor inducing principle) plant tumors and 35
 TSH (thyroid stimulating hormone) see Thyrotropic hormone of anterior pituitary
 Tapazole(1 methyl 2 mercaptoimida zole)
 therapeutic effect on thyrotoxicosis 832
 toxic effects 832
 thyrotropin and 490
 Teeth parathyroids and 169
 Testis 864-865
 biosynthesis of steroid hormones by 590 593
 effect of androgens on 673
 hormones physiology of 663-664
 hypophysis and 564
 as source of androgens 610 611 612
 of estrogens 593 591 864
 luteinizing hormone and 595
 of progesterone 601 602 603
 tumors of 867-868
 hormone secretion in 867
 Sertoli cell 864
- Testosterone formation microbiological 607 609 613
 Testosterone
 activity effect of 11 desaturation on 559
 gonadotropic 867
 biosynthesis gonadotropic hormones and 611 612
 pathway of 613
 precursors of 612
 configuration of and related compounds 527 528
 effect on adrenal cortex of hypophysectomized animals 278
 on growth of hypophysectomized animals 240
 on thyroid function 459
 hyaluronidase and 763
 labeled preparation 575 577
 19 nortestosterone and 544
 metabolism 613 614
 in rodents 621-622
 as metabolite of Δ^4 androstene-3 17 dione 616 618
 metabolites of 616 617 618 619
 microbiological reactions 620
 as precursor of 11 β hydroxytestosterone 625
 reduction of 650
 enzymatic of ring A in 645
 sources 610 611 612 614 864
 therapeutic effect on hypogonadism 863 866
 Tetrahydro E see Urocortisone
 Tetrahydro F see Urocortisol
 3 α 11 β 17 α 21 Tetrahydroxyallopregnane 11 20 dione 629
 3 β 11 β 17 α 21 Tetrahydroxyallopregnane 20 one 622
 3 α 11 β 17 21 Tetrahydroxypregnane 20 one see Urocortisol
 6 β 11 β 17 α 21 Tetrahydroxy Δ^4 pregnene 3 20 dione biosynthesis 676
 11 β 17 α 20 β 21 Tetrahydroxy Δ^4 pregnene 3 11 dione 622
 3 α (and 3 β) 11 β 17 α 21 Tetrahydroxyallopregnane 20 one as metabolite of cortisol 640
 3 β 17 α 20 β 21 Tetrahydroxyallo

- pregnan 11 one as metabolite of cortisone 639
- 11 β 1 α 20 21 Tetrahydroxyallopregnan-3 one as metabolite of cortisone 640
- 11 β 17 α 20 21 Tetrahydroxy Δ^4 pregnen-3 one as metabolite of cortisone 639
- Tetralidene 1 acetic acid auxin activity of cis isomer 8
- Thiamine thyroid and 488
- Thiazoles
antithyroid activity 460 466
structure 460 466
- Thiocarbamides see also individual compounds
antithyroid activity 462 464-467
mechanism of 467
- Thiocyanate
antithyroid activity 462
mechanism of 462
- Thiopental thyroid and 496
- Thiopiperidone ? 464
antithyroid activity 467
- Thiopyridone ?
antithyroid activity 467
structure 467
- Thiosemicarbazone
antithyroid activity 464
structure 464
- Thiouracil
antithyroid activity 466
mechanism of 462
derivatives therapeutic effect on thyrotoxicosis 832
toxic reactions 837
thyroid tumors and 498
thyrotropin and 499
- Thiourea
antithyroid activity of and derivatives 464 467
auxins and 25
structure 464 467
- Thrombotonin see 5 Hydroxytryptamine
- Thymus
effect of adrenocortical steroids on 742
of stress on 742
- Thyroglobulin 446 447-448
activity 474
iodine content 446 447 448
molecular weight 448
physicochemical properties 448
preparation 446
thyroxine biosynthesis and 440 447
- Thyroid 433-519
activity height of thyroid cells and 430 436
protein bound iodine and 830
proteolytic 447
thyroid stimulating hormone and 447
adrenals and 490 780-783
anatomy chemical 436
gross 434-435
microscopic 430 436
carcinoma of see Thyroid tumors carcinoma
cells electron microscope studies on 430-436
height of and thyroid activity 435 436
chemical elements in 436-437
collection of iodine by of chordates 434
of fetus 438
control of chemical 461-474
conversion of inorganic iodide to organic iodine in 441-442
enzymes catalyzing 442
cytochrome c and 477
diseases of 827-840
diagnostic procedures 827-831
effect on excretion of thyrotropic hormone 457-458
ophthalmopathy in 833-835
drugs and 496-497
effect of environment on 459-460
on gonads 494-495
on oxygen consumption 490-491
of thyroid on 492
embryology of 438-439
enzymes in 437
extracts difference between mammalian and piscine 435
fluorescence in 437
function agents affecting growth and 455-474
hormonal control of 455-459
use of iodine tracers in study of 828-830
gastrointestinal tract and 495-496

- Sulfaguanidine antithyroid activity 462
 Sulfates, steroidal hydrolysis of 544
 Sulfhydryl compounds
 auxin activity and 39
 dormancy in plants and 27
 Sulfonamides antithyroid activity 460
 Sulfur
 metabolism growth hormone and 200-201
 in thyroid 437
 Sulfuric acid conjugation of steroid hormones with, 590
 Superfetation in sheep IMG and 340-341
 Superovulation in cows gonadotropins and 341-342
 Sweat effect of adrenocortical steroids on 712-713
 Sympathins 102, 113
 Synapoidin 866
 Syphilis
 effect of ACTH on 780
 of cortisone on 780
- T**
- T I P (Tumor inducing principle) plant tumors and 35
 TSH (thyroid stimulating hormone) see Thyrotropic hormone of anterior pituitary
 Tapazole(1 methyl 2 mercaptoumidazole)
 therapeutic effect on thyrotoxicosis 832
 toxic effects 832
 thyrotropin and 493
 Teeth parathyroids and 161
 Testis 464 468
 biosynthesis of steroid hormones by 590 593
 effect of androgens on 673
 hormones physiology of 660-684
 hypophysis and 864
 as source of androgens 610 611 612
 of estrogens 593 594 864
 luteinizing hormone and 590
 of progesterone 601 602 603
 tumors of 867-868
 hormone secretion in 867
 Sertoli cell 864
 Testosterone formation microbiological 607, 609 643
 Testosterone,
 activity effect of 11 de auration on 509
 gonadotropic 867
 biosynthesis gonadotropic hormones and 611 612
 pathway of 613
 precursors of 612
 configuration of and related compounds 527 528
 effect on adrenal cortex of hypophysectomized animals 278
 on growth of hypophysectomized animals 240
 on thyroid function 459
 hyaluronidase and 763
 labeled preparation 570 577
 19 nortestosterone and 544
 metabolism 613 614
 in rodents 601-622
 as metabolite of Δ^4 androstene-3 17 dione 616 618
 metabolites of 616 617 618 619
 microbiological reactions 620
 as precursor of 11 β hydroxytestosterone 600
 reduction of 650
 enzymatic of ring A in 648
 sources 610 611 612 614 864
 therapeutic effect on hypogonadism 860 860
 Tetrahydro E see Urocortisone
 Tetrahydro F see Urocortisol
 3 α 11 β 17 α 21 Tetrahydroxypregnan-11 20 dione 622
 3 β 11 β 17 α 21 Tetrahydroxypregnan-20 one 672
 3 α 11 β 17 21 Tetrahydroxypregnan-20 one see Urocortisol
 6 β 11 β 17 α 21 Tetrahydroxy Δ^4 pregnene-3 20 dione biosynthesis 626
 11 β 17 α 20 β 21 Tetrahydroxy Δ^4 pregnene-3 11 dione 622
 3 α (and 3 β) 11 β 17 α 21 Tetrahydroxy allopregnan-20 one as metabolite of cortisol 640
 3 β 17 α 20 β 21 Tetrahydroxy allo-

- synergism with growth hormone 275
 tapazole and 499
 thiouracil and 499
 thyroid function and 455-459
 vitamin C and 499
 Thyrotropin see Thyrotropic hormone
 of anterior pituitary
 Thyroxamine as active thyroid hor-
 mone 454-455
 Thyroxine see also Thyroid hormone
 adrenaline and 496-497
 antagonists 453
 configuration of 4,6
 structural requirements 475-4,6
 antidiuretic activity of anterior pitui-
 tary and 399
 bacterial infection and 497-498
 biosynthesis 441 443
 in chick embryo 438
 in blood 449
 chloramphenicol and 497
 concentration in pituitary 452
 species differences in 452
 cortisone and 498 782
 derivatives as thyroxine antagonists
 47
 effect on adrenal cortex of hypophy-
 sectomized animals 278
 on blood 496
 on carcinogens 496-497
 on cardiovascular system 493
 empirical formula 448
 estrogens and 672
 galactopoietic activity 366
 glucuronide of 451-452
 growth effects in thyroidectomized
 animals 239-240
 iodine content 448
 isolation 449
 metabolism 450-455
 factors affecting 453
 liver and 451 452
 species differences in 452
 vitamins B and 453
 precursors 446
 secretion rate of 449
 role of pituitary in 455
 by thyroid tumors 499
 toxic effects of 492 493
 triiodothyronine and 454
 Tigogenin 567
 conversion to allopregnane 3 β 16 20
 triol 562
 Tissues
 androgens of 610-611
 animal indoleacetic acid and respira-
 tion of 41
 binding of insulin to 186-187
 damage adrenocortical hormones and
 758
 5 hydroxytryptamine content 104
 lymphoid adrenal cortex 742-744
 pyridoxine and 743
 soft effect of parathyroids on 160-161
 sited for biosynthesis of steroid hor-
 mones 590
 Tissue cultures effect of adrenocortical
 steroids on 769-770
 Toxemias of pregnancy 863-864
 hormone excretion in 863
 Toxins
 adrenal cortex and 278 757-758
 bacterial cortisone and 758
 Trauma effect of severe on adrenal cor-
 tex of hypophysectomized animals
 278
 Trichinosis effect of ACTH on 780
 2 4 6 Trichlorophenoxyacetic acid as
 antiauxin 37 38
 3 α 17 α 21 Trihydroxyallopregnane
 11 20 dione as metabolite of corti-
 sone 638 640
 3 β 17 α 21 Trihydroxyallopregnane
 11 20 dione 672
 as metabolite of cortisone 638 640
 11 α 17 α 21 Trihydroxyallopregnane
 3 20 dione microbiological forma-
 tion 643
 3 α 11 β 21 Trihydroxyallopregnan 20
 one as metabolite of corticosterone
 and 11 dehydrocorticosterone 636
 3 β 11 β 20 Trihydroxyallopregnan 20
 one 672
 3 β 17 α 21 Trihydroxyallopregnan 20
 one 672
 as metabolite of 11 deoxycortisol 638
 3 β 7 β 21 Trihydroxyallopregnan 20 one
 microbiological formation 641
 3 α 17 α 21 Trihydroxypregnane 3 20

- growth and, 239-240
 hexosamine content 437
 hormone of *see* Thyroid hormone
 Thyroxine
 inhibitors of 461-474
 therapeutic value 463
 iodide trap of 441
 blockage by monovalent ions 462
 effect of pituitary on 455
 morphology of 434-439
 nucleic acids in 437
 nutrition and 460-461
 parathyroids and 163
 role in atherosclerosis 486
 in lower animals 434-435
 in metabolism of calcium 457
 of carbohydrates 484-485
 of electrolytes 486
 of fat 485-486
 of nucleic acids 483-484
 of vitamins 488-490
 of water 486
 scintigrams of 830
 tumors of 498*f*
 carcinoma effect of thyrotropin on 499
 isolation of iodinated protein from patients with 450
 following treatment with thiouracil or related compounds 498
 physiological aspects of 498-499
 secretion of thyroxine by 499
 uptake of radioiodine by adrenocortical steroids and 782
 weight dietary iodine and 434
 x ray and ultraviolet absorption by 436
- Thyroid crisis 836
 Thyroid hormone *see also* Thyroxine
 activity of 474-498
 mechanism 477
 structure and 474-476
 binding to serum α globulins 450
 circulating 434
 control of amphibian metamorphosis by 491
 definition 474
 enzymes and 477-480
 nitrogen metabolism and 480-484
 release use of iodine tracers in study of 830
 secretion thyroidal uptake of iodine and 440
 synthesis 434
 Thyroiditis 839-840
 effect of TBII on 839
 subacute 839-840
 treatment 839
 uptake of tracer iodine in 839
 Thyronine
 antithyroid activity 476
 iodination biological 443 445
 substituted hormonal activity 444 475
 thyroid hormone activity and 474
 Thyrotoxicosis 831-836
 cardiac symptoms in, 831
 diagnostic procedures 831 835
 effect of psychic factors on 832
 myasthenia gravis and 835
 periodic familial paralysis and 836
 therapy of 832-833
 Thyrotrophs 314 315
 Thyrotropic hormone of anterior pituitary 206-210 872
 activity 456
 radiokinetic 254
 assay of 206-208
 mechanism of 456
 metabolic 206
 standardization 206
 ACTH and 783
 cellular origin 314 823
 cretinism and 837-838
 effect on proteolytic activity of thyroid 447
 on thyroid cancers 499
 on thyroid deiodase 446
 exophthalmos producing substance and 211 933
 fractionation 457
 gonitogens and 463
 isolation 208-210
 metabolism of 457-458
 thyroid disease and 457-458
 multiplicity of 824
 ophthalmopathy and 834 835
 physicochemical properties 210
 role in intrathyroidal iodine metabolism 446
 secretion in acromegaly 273

- Uric acid diabetogenic activity 189
 Urine
 as source of estrogen 203-211
 prolactin in human 211
 steroids in see under individual compounds
 Urocanylecholine in hypophyseal gland of *Murex* 100
 Urocortisol 523
 as metabolite of corticoid 631-63
 Urocortisone 523
 enzymatic conversion of 17 α -21 dihydroxypregnane 3-11-20 triene to 647
 as metabolite of corticoid 631-63
 Uterus
 effect of estrogen on 669-670
 of posterior pituitary extracts on 390
 of progesterin on 669-670
- ### V
- Vagina
 effect of estrogen on 669
 of progesterin on 669
 Vasopressin
 activity 137-403-404
 amino acid composition 404
 species differences in 402-404-414
 amino acid sequences in 413
 chemical nature 99-137-403
 effect on milk ejection 400
 on water diuresis in mice 394
 electrophoretic properties 404-405
 molecular weight 404
 as neurosecretory material 99-143
 origin of 105-118
 oxytocin and 140
 olypeptide structure 127-403
 purification 401-402
 structure 403-406-412-415
 synthesis 414
 Vegetables inhibitory effect on thyroid 461-838
 Venoms 5-hydroxytryptamine in 104
 Venus mercenaria
 heart of 122
 action of acetylcholine on 123
 of acetylcholine analogues on 122ff
 Vernalization
 auxins and 3
 flowering and 32-33
 Vertebrate
 lower melanophore-expanding effect of posterior pituitary extracts in 310
 sex differentiation role of gonadal hormone in 674-675
 L-Valyl 2-thioxazolidone
 goitrogenic activity 461-466-838
 isolation from rutabaga 461-838
 Viruses
 cortisone and 413
 plant anti-auxins and 34
 auxins and 37
 Visual purple B hormone and regeneration of in Japanese toad 490
 Vitamin A thyroid and 488
 Vitamin B₁ thyroid and 489
 Vitamin B₁
 deficiency corticoid and 9
 thyroid and 489
 thyroxine and 489-492
 Vitamin C see also Ascorbic acid
 deficiency atherosclerosis and 717
 thyroid and 437-489-490
 Vitamins see also individual compounds
 adrenal cortex and 736-742
 auxins and 25
 thyroid function and 476
 Vitamins B
 deficiency pituitary gonadotropic activity and 337
 effect on thyroid 488-489
 on thyroxine metabolism 483
- ### W
- Water balance principle (W.B.P.) of posterior pituitary 398
 as phylogenetic precursor of anti-diuretic hormone 398
 Water metabolism
 adrenal cortex and 394-395-688
 anterior pituitary and 394-395
 relationship of adrenal cortex and posterior pituitary to 711-712
 role of neurohypophysis in 330-394-399
 of thyroid in 486

- dione as metabolite of 17 α 21 dihydroxypregnane-3 20 dione 634
 3 α 17 21 Trihydroxypregnane 11 20 dione see Urocortisone
 3 α 11 β 17 α Trihydroxypregnane 20 one as metabolite of cortisol and cortisone, 634
 7 α 11 β 21 Trihydroxypregnane 20 one as metabolite of corticosterone 634 636
 7 α 17 α 21 Trihydroxypregnane 20 one as metabolite of 11 deoxycortisol 633 634 639
 6 β 17 α 21 Trihydroxy Δ^4 pregnene 3 20 dione
 microbiological formation 641
 as precursor of 6 β 11 β 17 α 21 tetrahydroxy Δ^4 pregnene 3 20 dione 626
 8 17 α 21 Trihydroxy Δ^4 pregnene 3 20 dione microbiological formation 641
 11 α 17 21 Trihydroxy Δ^4 pregnene 3 20 dione see Epicortisol
 11 β 17 21 Trihydroxy Δ^4 pregnene 3 20 dione see Cortisol
 14 17 α 21 Trihydroxy Δ^4 pregnene 3 20 dione microbiological formation 642
 17 α 20 β 21 Trihydroxy Δ^4 pregnene 3 11 dione 622
 as metabolite of cortisol 639
 of cortisone 640
 17 α 20 β 21 Trihydroxy Δ^4 pregnene 3 20 dione as metabolite of cortisone 634
 17 α 20 β 21 Trihydroxy Δ^4 pregnene 3 one as metabolite of 11 deoxycortisol 634 639
 2 3 5 Triodobenzoic acid (TIBA)
 as antiauxin 22
 auxins and 15 23
 effect on polar auxin transport 19
 promotion of flowering by 31
 Triodothyronine 490
 activity 473
 metamorphogenic 491
 biosynthesis 443-444
 duodotyrosine a precursor 443
 intermediates 444 445
 in blood 449
 effect on oxygen consumption 490
 metabolism 453 454
 oxidative phosphorylation and 477
 thyroxine and 454
 3 5 3 Triodothyronine
 formation 444
 in thyroid tissue 449
 Triphenylbromomethylene radioactive metabolism 597
 Tryptamine
 as auxin precursor 27
 possible neurohormonal activity 99 100
 Tryptophan as auxin precursor 27-28
Tuber cinereum median eminence of role in gonadotropic activity of anterior pituitary 322
 Tuberculin
 sensitivity cortisone and 498 770
 hydrocortisone and 770
 thyroxine and 498
 Tuberculosis
 as cause of Addison's disease 840
 effects of cortisone on 777-778
 Tubules spermatogenic hormone of 864-865
 hypogonadism due to impaired function of 865-866
 treatment 865 866
 Tumors see also individual tumors
 malignant adrenal cortex and 750-753
 thyroid see Thyroid tumors of
 Tumores neural gland endocrine activity 79
 Tyramine
 possible neurohormonal activity of 99 100
 Tyrosine
 corticotropin B activity and 224
 isolation biological 443
 inhibitors of 462

U

- Umbelliferone auxins and 24
 Urnediol
 precursors 563
 structure 564 of

- Uric acid diabetogenic activity 189
- Urine
 as source of estrogens 571 594
 prolactin in human 711
 steroids in — see under individual compounds
- Urocaryleholone in hypobranchial gland of *Murex* 100
- Urocortisol 593
 as metabolite of cortisol 634 635
- Urocortinone 523
 enzymatic conversion of 1α OH dihydroxypregnane-3 11 20 trione to 647
 as metabolite of cortisone 634 635
- Uterus
 effect of estrogen on 660-670
 of posterior pituitary extracts on 300
 of progesterin on 669-670
- V
- Vagina
 effect of estrogen on 669
 of progesterin on 669
- Vasopressin
 activity 1 403-404
 amino acid composition 404
 species differences in 402 404 414
 amino acid sequences in 413
 chemical nature 99 137 405
 effect on milk ejection 400
 on water diuresis in mice 394
 electrophoretic properties 404-405
 molecular weight 404
 as neurosecretory material 99 143
 origin of 105 115
 oxytocin and 130
 polypeptide structure 1 7 405
 purification 401 407
 structure 405-406 412-415
 synthesis 414
- Vegetables inhibitory effect on thyroid 461 838
- Venoms 5-hydroxytryptamine in 104
- Venus mercenaria*
 heart of 177
 action of acetylcholine on 123
 of acetylcholine analogues on 122 ff
- Vernalization
 auxins and 32
 flowering and 72-73
- Vertebrate
 lower melanophore-expanding effect of posterior pituitary extracts in 300
 sex differentiation role of gonadal hormones in 674-675
- L-5 Vinyl-L-thiooxazolidone
 goitrogenic activity 461 466 838
 isolation from rutabaga 461 838
- Viruses
 cortisone and 749
 plant anti-auxins and 37
 auxins and 37
- Visual purple B hormone and regeneration of in Japanese toad 420
- Vitamin A thyroid and 488
- Vitamin B₆ thyroid and 489
- Vitamin B₁₂
 deficiency cortisone and 739
 thyroid and 489
 thyroxine and 489 492
- Vitamin C see also Ascorbic acid
 deficiency atherosclerosis and 717
 thyroid and 437 489-490
- Vitamins see also individual compounds
 adrenal cortex and 736-742
 auxins and 25
 thyroid function and 476
- Vitamin B
 deficiency pituitary gonadotropic activity and 337
 effect on thyroid 488-489
 on thyroxine metabolism 453
- W
- Water balance principle (W B P) of posterior pituitary 398
 as phylogenetic precursor of anti-diuretic hormone 398
- Water metabolism
 adrenal cortex and 394 395 688
 anterior pituitary and 394 395
 relationship of adrenal cortex and posterior pituitary to 711 712
 role of neurohypophysis in 390 394-399
 of thyroid in 486

- dione as metabolite of 17 α 21 dihydroxypregnane 3 20 dione 638
- 3 α 17 21 Trihydroxypregnane 11 20 dione see Urocortisone
- 7 α 11 β 17 α Trihydroxypregnan 20 one as metabolite of cortisol and cortisone 634
- 3 α 11 β 21 Trihydroxypregnan 20 one as metabolite of corticosterone 635 636
- 3 α 17 α 21 Trihydroxypregnan 20 one as metabolite of 11 deoxycortisol 633 638 639
- 6 β 17 α 21 Trihydroxy Δ^4 pregnene 3 20 dione
microbiological formation 641
as precursor of 6 β 11 β 17 α 21 tetrahydroxy Δ^4 pregnene 3 20 dione 626
- 8 17 α 21 Trihydroxy Δ^4 pregnene 3 20 dione microbiological formation 641
- 11 α 17 21 Trihydroxy Δ^4 pregnene 3 20 dione see Epicortisol
- 11 β 17 21 Trihydroxy Δ^4 pregnene 3 20 dione see Cortisol
- 14 17 α 21 Trihydroxy Δ^4 pregnene 3 20 dione microbiological formation 642
- 17 α 20 β 21 Trihydroxy Δ^4 pregnene 3 11 dione 622
as metabolite of cortisol 639
of cortisone 640
- 17 α 20 β 21 Trihydroxy Δ^4 pregnene 3 20 dione as metabolite of cortisone 638
- 17 α 20 β 21 Trihydroxy Δ^4 pregnen 3 one as metabolite of 11 deoxycortisol 638 639
- 2 3 5 Triiodobenzoic acid (TIBA)
as antiauxin 22
auxins and 15 25
effect on polar auxin transport 19
promotion of flowering by 31
- Triiodothyronine 490
activity 445
metamorphogenic 491
biosynthesis 443-447
diiodotyrosine as precursor 443
intermediates 444 445
in blood 449
effect on oxygen consumption 490
metabolism 453 454
oxidative phosphorylation and 477
thyroxine and 454
- 3 5 3 Triiodothyronine,
formation 445
in thyroid tissue 445
- Triphenylbromoethylene radioactive metabolism 597
- Tryptamine
as auxin precursor 27
possible neurohormonal activity 99 100
- Tryptophan as auxin precursor 27-28
- Tuber cinereum* median eminence of role in gonadotropic activity of anterior pituitary 322
- Tuberculin
sensitivity cortisone and 498 740
hydrocortisone and 770
thyroxine and 408
- Tuberculosis
as cause of Addison's disease 840
effects of cortisone on 777-778
- Tubules spermatogenic hormone of 864-865
hypogonadism due to impaired function of 865-866
treatment 865 866
- Tumors see also individual tumors
malignant adrenal cortex and 750-753
thyroid see Thyroid tumors of
- Tunicates neural gland endocrine activity 79
- Tyramine
possible neurohormonal activity of 99 100
- Tyrosine
corticotropin B activity and 225
iodination biological 443
inhibitors of 462

U

- Umbelliferone auxin and 24
- Urinediol
precursors 565
structure 561 565

Water uptake by plants 42-48

active 45

effect of ions on 46

Weight FSH secretion and body 8,69

Worms

acetylcholine in 59

hormones in 58-60

sources of 58 59

internephridial organs endocrine ac-
tivity 55

Wound healing adrenocortical steroids
and 766-768

X

X ray lethal doses effect of estrogen
pretreatment on survival time 672

